

Reply to "Comment on equivalence between quantum phase transition phenomena in radiation-matter and magnetic systems"

Giuseppe Liberti and Rosa Letizia Zano
 Dipartimento di Fisica, Università della Calabria
 INFN - Gruppo collegato di Cosenza, 87036 Rende (CS) Italy
 (Dated: May 10, 2022)

PACS numbers: 42.50.Fx - Cooperative phenomena in quantum optical systems; 05.70.Jk - Critical point phenomena; 73.43.Ng - Quantum phase transition.

In their Comment [1], Brankov, Tonchev and Zagrebnov, claim that the temperature-dependent effective Hamiltonian derived in Ref. [2] from a radiation-matter Dicke model violate a "rigorous" result that the same authors have obtained 30 years ago [3]. It is clear that Brankov, Tonchev and Zagrebnov have misunderstood the results of Ref. [2] in several ways.

1. First, the temperature-dependent effective Hamiltonian given by

$$H(\beta) = \sum_i a^\dagger_i a_i + \frac{1}{2} S_z - \frac{1}{2N} \coth \frac{\beta}{2} \sum_i S_x^2 \quad (1)$$

is correct for $\beta \gg 1$ and $\beta \ll 1$, i.e. only in the high-temperature limit. It is not permissible to take the limit $\beta \rightarrow 0$ to infer critical properties and some connection with the collective one-dimensional Ising model. For $\beta \rightarrow 0$ this Hamiltonian obviously becomes

$$H = \sum_i a^\dagger_i a_i + \frac{1}{2} S_z - \frac{1}{N} \sum_i S_x^2 \quad (2)$$

2. We have used the Zassenhaus formula [4] in its simplest form, through a perturbative expansion of partition function obtained by decomposing the Hamiltonian into two non-commuting hermitian operators. The results of our analysis are used to show an obvious similarity with the result of Ref. [5] but only in the high-temperature limit, keeping the lowest-order terms of the Polatssek and Becker expansion [6]. It is possible to derive higher-order approximations in a systematic manner but the increasing complexity of the expressions requires numerical calculations in order to derive the thermodynamical properties of the system.

3. We use the term "classical" with respect to the limit reached when $\beta \rightarrow 0$ where the results are exactly those obtained with the Wang and Høie computational method. [7]

-
- [1] J.G. Brankov, N.S. Tonchev and V.A. Zagrebnov, quant-ph/0506207
 [2] G. Liberti and R.L. Zano, Eur. Phys. J. B 44, 535 (2005)
 [3] J.G. Brankov, V.A. Zagrebnov and N.S. Tonchev, Theoret. and Math. Phys. 22, 13 (1975)
 [4] R.M. Wilcox, J. Math. Phys. 8, 962 (1967)
 [5] J. Reslen, L. Quiroga and N.F. Johnson, Europhys. Lett. 69, 8 (2005)
 [6] G. Polatssek and K.W. Becker, Phys. Rev. B 55, 16096 (1997)
 [7] Y.K. Wang and F.T. Høie, Phys. Rev. A 7, 831 (1973)