## Long Range E lectron Transfer A cross -conjugated chain: Role of E lectron C orrelations

S. Lakshmi, Ayan Datta and Swapan K. Pati

Theoretical Sciences Unit and Chemistry and Physics of Materials Unit,

Jawaharlal Nehru Centre for Advanced Scienti c Research, Jakkur Campus, Bangabre 560 064, India.

(D ated: A pril 14, 2024)

We consider a prototype polyene chain: donor- (bridge)-acceptor. The distance between the donor and the acceptor is varied by increasing the number of atom s in the bridge and the rate of electron transfer,  $k_{et}$  is studied for a series of di erent donors,  $D = N H_2$ ,  $O C H_3$ ,  $S C H_3$ , and a xed acceptor,  $A = N O_2$ . We observe a large  $k_{et}$  even at a D-A separation of 45 A, unexpected from the standard noninteracting theories. Such a long range electron transfer is prim arily due to the form ation of bound electron-hole pair (exciton) across the donor-bridge-acceptor atom s. C alculations at various levels, sem i-em pirical (with inclusion of con guration interactions) and m any-body m odels (using density m atrix renorm alization group, DMRG), have been performed to accurately account for such electron correlations.

I. IN TRODUCTION

Electron transfer (ET) between well separated donor and acceptor groups is dom inant in many chem ical and biological processes. These processes include design of electronic switches and wires in molecular level to photosynthesis and respirations in biological system s<sup>1,2</sup>. Typically in m any system s of interest, the spatial separation between the donor and the acceptor ranges from 10 to 30 A<sup>3</sup>. Because of this separation, there is no direct overlap between the donor and the acceptor groups. The electron transfer coupling is mediated via the intervening orbitals, called the bridge states. In most of the studies, the model used is a donor-bridge-acceptor (DBA) com plex, wherein the donor and the acceptor states couple to the bridge in the form of a macrom olecule. The intervening bridge can be water, conjugated organic molecules and polymers, a protein backbone, or even DNA. In many of the system s, these bridge states can lead to substantially altered electron transfer rates K <sub>et</sub> with respect to vacuum tunnelling<sup>4</sup>. The rate is also known to signi cantly depend on the conform ational degrees of freedom of the bridges<sup>5</sup>. However, the exact role of the bridges in controlling electron transfer is very poorly understood particularly for systems with strong electron correlations and weakly bonded species like the hydrogen bonded system  $s^7$ .

For m ost system s, these electron transfer rates fall o exponentially with distance (ln  $k_{\rm et}$  / r). The inverse decay length, , typically varies from 0.2 to 1.5 A  $^1$ , depending on the system s under consideration. Sm all

values have been observed for m any proteins, arti cial peptide chains, DNA segments or a conjugated oligom eric backbone<sup>8,9</sup>. This exponential decay appears in the M arcus theory of non-adiabatic electron transfer<sup>10</sup> that assumes that the rate of ET depends on the electronic coupling element, H<sub>DA</sub> according to

$$k_{DA} (R_{DA}) = \frac{2}{h} H_{DA} f C$$
 (1)

where FC is the Franck-C ondon-weighted average density

of states. The electronic tunnelling matrix element in turn is related to the distance of separation between the D and A as  $H_{DA}$  / exp r=2. The theory assumes that the time scale of the electron transfer process is much faster than the time associated with the reorganization of the solvent nucleus or thermal disordering.

To date, most theoretical work has focused on the properties of the bridging medium and the position of D/A levels with respect to the HOMO and LUMO energies of the bridge<sup>11</sup>. Furtherm ore, these theories assume that the role of the bridge is to provide virtual orbitals that create an e ective electronic coupling between the donor and the acceptor<sup>12,13,14</sup>. This mechanism offen reduces the problem of the whole system consisting of the donor, acceptor and the bridge subspaces to an e ective two-state Ham iltonian consisting only of the donor and the acceptor states. There have been some alternative approach to this charge-transfer problem in recent years. D avison et al. have studied the charge-transfer thorugh a Huckeland extended model bridge using G reen's function form alism  $^{15}$ . On the other hand, a completely new approach for an extended bridge with narrow band donor and acceptor groups predict the charge-transfer m echanism to be long-ranged, involving a K ondo-singlet across the donor (acceptor)-bridge atom s<sup>16</sup>. Most often however, in many of the models, explicit electron-electron interactions in the bridge are completely ignored, while it is well known that electron correlations often lead to qualitatively di erent descriptions.

In this work, we perform a quantitative estimation of the role of the bridge as well as the donor/ acceptor groups for an elective charge transfer across the bridge. The bridge energetics has been found to in uence the strength of the donor-acceptor coupling because of strong mixing between donor/acceptor and bridge orbitals<sup>17,18</sup>. Furthermore, the most frequently used bridges are the -conjugated chains, which are known to be highly correlated system s<sup>19,20</sup>. In the DBA system, the process of

related system  $s^{19/20}$ . In the DBA system, the process of electron transfer takes place with an electron originally at the donor state, transferring to the acceptor state via the bridge states. In the presence of interactions, the

probability of electron transfer to an empty site is larger than it is to a partially occupied site. The probability of the relevant site being empty, partially occupied or doubly occupied depends very strongly on the electron correlation strength. Thus, incorporating electron-electron interactions is crucial to the study of electron-transfer processes.

In the next section, we discuss the electron-transfer process in a -conjugated bridge of varying length with a range of donor strengths. We nd that there is a significant amount of ET for -conjugated system even when the D and A are separated by a very large distance (45 A). In 3rd section, we study the charge-transfer process for a model conjugated chain with and without electron-correlations by employing the density matrix renorm alization group (DMRG) method<sup>21</sup> and show that strong electron correlation is indeed required for long-range ET. We conclude with a summary of all the results in the last section.

## II. SEM I-EM P IR ICAL CALCULATIONS

We consider a series of -conjugated systems shown in Fig. 1. In the top (case 1), the bridge is considered to be a linear polyene chain of length n varying from 2 to 30. The acceptor group is xed as  $A = NO_2$ , and the donor group is varied,  $D = NH_2$ ,  $OCH_3$  and  $SCH_3$ . All the geom etries are optim ized using the AM 1 param eterized Ham iltonian, a part of sem i-em pirical MOPAC package<sup>22</sup>. We have also veried that these geometries correspond to the minimum energy structure by further optim ization at the B 3LY P / 6 - 311G + + (d,p) level<sup>23</sup>. No symmetry constraints were used in the optimizations. In case 2, we consider that the donor and the acceptor are separated by a saturated alkane chain. These two m odel cases serve as a good contrast in the mechanism of electron-transfer (ET). The form er is expected to follow the bridge mediated ET due to the extended conjugation between the donor and the acceptor (through the pz orbitals of the intervening atom s). In the following, we show that the case 1 is indeed very di erent and shows long range ET purely because of electron correlations.

The geom etry optim ized structures were used for conguration interaction (CI) calculations to obtain energies and the dipole moments in the CI basis using the Zemer's INDO method<sup>24</sup>. The levels of CI calculations have been varied but we restrict our discussions to results obtained using singles CI (SCI) only as this gives a reliable estimate of the states involved in charge-transfer. The CI approach adopted here has been extensively used in earlier works, and was found to provide excitation energies and dipole matrix elements in good agreement with experiments<sup>25,26,27</sup>. For the Hartree-Fock determ inant, we use varying number of occupied and unoccupied molecular orbitals to construct the SCI space till a proper convergence is obtained.

The electron transfer coupling element, H  $_{\rm D~A}$  , has been

calculated using the generalized M ulliken-H ush m ethod for two-states, referred as 1 and  $2^{28}\,.$ 

$$H_{DA} = \frac{q}{12} \frac{E_{12}}{12^2 + 4_{12}^2}$$
(2)

where  $_{12}$  (E  $_{12}$ ) is the dipole m om ent (energy) di erence between the two states, and  $_{12}$  is the transition dipole between states 1 and 2. As has been discussed earlier, the m ethod considers only the adiabatic states 1 and 2, and assumes that the contribution from the nonadiabatic states to the transition dipole is small enough since the dipole m om ent operator has a weak distance dependence. Note that, it depends on the calculations whether the nonadiabatic states are well described by the adiabatic states 1 and 2.

The plots of H<sub>DA</sub> with respect to the number of CH unit of the polyacetylene bridge are shown in Fig. 2 for  $D = NH_2$ ,  $OCH_3$ ,  $SCH_3$  and  $A = NO_2$ . All of them have an exponential decay probe for the separation between the D and A.Fitting a single exponential parameter , for these three donors, we get  $= 0.277A^{-1}$ ,  $0.219A^{-1}$  and  $0.237A^{-1}$  for  $D = NH_2$ ,  $OCH_3$ ,  $SCH_3$  respectively. Thus, H<sub>DA</sub> seem s to depend on the donor strength although for all three cases we nd the range of electron transfer quite substantial.

A very important inference that can be drawn from our calculation is that even for a bridge length of 45-50 A, the H  $_{\rm D\,A}\,$  m agnitude is quite signi cant (  $\,$  1.7) and this value is almost independent of the electronegativity (donor ability) of the ligands. This was also evident from the very low decay exponential param eter, 0.25. Such low decay features for ET have been reported for DNA<sup>29</sup> and has triggered a lot of interest in its m olecular electronic properties  $^{30}$  . W e propose here that these D and A separated bridge is also a very good candidate for molecular electronics. W hile polaron assisted ET is believed to be the mechanism in DNA, we suggest the role of electron-electron interactions leading to the form ation of an electron-hole pair (an exciton<sup>31</sup>) responsible for long range charge transfer in conjugated polym eric system s.

The form ation of exciton is evident from our CI calculations. We have analyzed the oscillator strength of the lowest few CI states for various system sizes. At smaller chain lengths, the lowest dipole allowed state (the state that has maximum oscillator strength) lies quite high in energy and very close to the single-electron continuum . However, for large chain length (n > 4), the exciton stabilization is quite large and the dipole allow ed state appears much below the continuum. In fact, as the bridge length increases from n = 2, the lowest dipole allowed state shifts its position in the energy spectrum and above n = 4becomes the lowest energy (fundamental) excitation of the system. For smaller chain length, the exciton form ation is very weak, since such a sm all length scale does not provide room for form ation of a well bound electron-hole pair. In fact, a critical length scale is necessary for a delocalized electron and and a delocalized hole to bind. In

contrast to this feature for the conjugated chains (case 1), for the saturated chains  $H_2N - (CH_2CH_2)_n - NO_2$  (case 2), there are no excitonic bands and the dipole allowed state remains invariant of the chain length enhancements. In fact, the dipole allowed state itself has a sm all oscillator strength (very weak). This is easy to understand as for the saturated chains, the electron and hole over the D and A respectively, remain uncorrelated for any bridge lengths.

Interestingly, the actual decay of ET in  $H_2N$ -(CH=CH)<sub>15</sub>-NO<sub>2</sub> is much smaller. The  $H_{DA}$  at a separation of 1.41 A is 3 (From Fig. 2) and that at 38.5 A is 1.7. Using the equation (1) and assuming that the Franck-C ondon rule to be valid, we get the decay to be only 32% to that for no separation. This is very di erent from the case 2 in Fig. 1, where the D and A are separated by a saturated spacer and the rate decays to zero after n= 3.

Such a feature is very well illustrated in Fig. 3, where we plot the wavefunctions corresponding to highest occupied molecular orbital (HOMO) and lowest unoccupied molecular orbital (LUMO) for two of the model systems: D and A separated by unsaturated and saturated spacers by large distance:  $H_2N - (CH = CH)_{15} - NO_2$  and  $H_2N -$ (CH<sub>2</sub>CH<sub>2</sub>)<sub>15</sub>-NO<sub>2</sub> respectively. The HOMO plot shows the ground state features with electron localized around the donor while the LUMO plot shows the redox process, the lowest charge transfer state. There is a distinct di erence between the two system s. W hile for H  $_2N$  - $(CH_2CH_2)_{15}$ -NO<sub>2</sub>, the electron is localized either on the D or A corresponding to HOMO and LUMO respectively, for H<sub>2</sub>N-(CH<sub>2</sub>CH<sub>2</sub>)<sub>15</sub>-NO<sub>2</sub> shows very large delocalization of charge carriers from the D and A groups into the bridge orbitals.

In Fig. 4, we show the molecular orbitals (MO) coe cients for the atom s in donor and the acceptor groups (N atom in  $NH_2$  group and N and O atom s in  $NO_2$  group).  $_{i}C_{m i}^{y}C_{m i}$  (where i index runs for The MO coe cients, each atom ic site and m indexes the molecular orbitals), are extracted for both HOMO and LUMO states. The HOMO state corresponds to the HF ground state and thus represents the system before charge transfer. On the otherhand, the corresponding LUM 0 coe cients represent the state after charge transfer. As can be seen, the M 0 coe cients before charge transfer are mostly located near the electron donating orbital of N atom in NH2 group. A fter the charge transfer how ever, the electrons are distributed over the  $p_z$  orbitals of three atom s: N and two 0 of the  $NO_2$  group.

## III. QUANTUM MANY BODY CALCULATIONS

For a quantitative understanding of the charge-transfer processes in such systems, we have used quantum manybody models. We consider the bridge to be a onedimensional chain with N-atoms (N even) and the donor(D) and the acceptor(A) atom s are attached to its two ends. The Ham iltonian can be written as

$$H = t \begin{bmatrix} X & X \\ f & (c_{i}^{y} c_{i+1} + hc)g \\ \\ & \\ + U n_{i''} n_{i\#} + V (n_{i} n_{av}) (n_{i+1} n_{av})] \\ \\ & \\ + D n_{D} + A n_{A} + [t_{1} (c_{1}^{y} c_{D} + c_{N}^{y} c_{A} + hc)]$$

where t is the transfer integral, U the Hubbard on-site potential and V is the nearest-neighbor C oulom b interaction parameter in the bridge.  $_D$  and  $_A$  are the on-site energy of donor and acceptor atom s respectively.  $t_1$  is the hopping term between the D (A) atom s with the bridge atom s 1 (N ).  $c_i^y$  creates an electron with spin at site i, and  $n_i = n_{i''} + n_{i\#}$ , is the electron number operator at site i.

The U V m odel or the extended H ubbard m odel is the minimal model to capture the physics of a quasiparticle like exciton, a bound electron-hole pair. In the large U (U = 1 ) lim it, the exciton is stabilized by V , since the Hubbard repulsion U describes the electron-hole continuum. However, this is for an isolated bridge with half-lled band. W hen the donor (D) and the acceptor (A) species are very weakly attached to the bridge, the electron transfer from D (hole transfer from A) can be mediated through the non-interacting electron-hole states or an excitonic state of the bridge. However, depending on the electron correlations in the D and A orbitals, the probability of interm ixing of the D and A states with the bridge states can vary and to account for the electron transfer in such cases, we need to consider the whole system as a macromolecule coupled strongly with its constituents. In our calculations, we have considered two extrem e cases for the donor and acceptors in terms of its electron correlations. Case A with no Coulomb repulsion at the donor and acceptor sites which allows any occupancy of the D /A sites and case B where we have set in nite Coulomb repulsions in D and A <sup>32</sup> , which prevents double occupancy of these sites. Note that for both cases, bridge is described by correlated m odel. The other case with uncorrelated bridge has not been studied, since it has been described elsewhere.<sup>16</sup>

The bridge is assumed to be a half-lled conjugated polymer insulator. From the half-lled extended model literature, it is known that the ground state is a SDW (spin-density wave) state for U > 2V, while it is a CDW (charge-density wave) state for U < 2V. We are interested in charge transferm echanism through a bridge with SDW ground state representing organic conjugated polymer, and thus we set U = 5, V = 1 and t = 1, without loss of generality  $^{\rm 6}$  . The total number of electrons in the D-bridge-A system is N + 1, where N is the number of orbitals in the bridge sites and N + 2 is the total num ber of orbitals of the whole DBA system. This system thus represents a linear correlated chain with one less electron than the half-lled lim it. We present below the charge transfere ciency of this system with bridge chain length varying from 2 to 30.

The charge transfer occurs mostly through the lowest energy excited state of the bridge, which is induced experimentally by interaction with light. The density matrix renormalization group (DMRG) method has been used to study the ground and the excited states of the system <sup>21</sup>. Since the bridge is a one-dimensional chain, the accuracy of the method is extremely good. In DMRG method, we target the ground state and the lowest dipole allowed states, and the density matrix is averaged over these states at every iteration. The dipole allowed state is recognized as the state to which the dipole moment operator acting on the ground state gives nonzero value. In the DMRG basis the transition dipole moment between two states is de ned as

$$_{ij} = h_{i}j^{j}j_{j}i \qquad (4)$$

where  $^{i}$  is the dipole m on ent operator acting between the states  $_{i}$  and  $_{j}$ . The H<sub>DA</sub> is then calculated from the Mulliken-Hush formula given in eqn 2.

W e have plotted the electron-transfer coupling elem ent (H  $_{\rm D A}$  ), de ned earlier, as a function of chain length for the two cases discussed above. Fig 5a shows the H  $_{\rm DA}$ for the case with no C oulom b repulsion at the donor and acceptor sites (case A), for a few representative values of the donor strengths (orbital energies  $_{\rm D}$  ). It can be seen that the H<sub>DA</sub> falls o quite rapidly as expected, but is quite short ranged. An exponential t to the curves give a value of around 0.5 A  $^1$  for , the parameter in the exponential. This type of large exponential fall of electron transfer rate is expected from the well-known M arcus' theory<sup>12</sup>. Fig 5b shows sim ilarplots for the case with large Hubbard repulsion at the donor and acceptor sites (case B). In this case too, the H<sub>DA</sub> falls o exponentially, but the fall is much slower and long ranged. An exponential t to these curves give a value of around  $0.2 \text{ A}^{-1}$ . The value of for case A is larger than that for case B, indicating that the range of transfer is sm aller for the rst case. It is very interesting to note that for the

-conjugated bridge considered in the previous section, we obtained to be of the same order as the case B with correlated bridge. Thus our model calculations are quite justi ed in realizing the ET kinetics for real molecular system s with strong correlations.

To understand the mechanism behind the charge transfer which results in long range and short range transfer for case B and case A respectively, we have plotted the ground state charge densities in Fig. 6 for both cases. Fig 6a shows the ground state charge densities for case A. It can be seen that the presence of a strong donor (a large negative value of  $_{\rm D}$ ) and the absence of any C oulom b repulsion at the donor favors an increase in the charge density at the donor. The acceptor on the other hand, which is always kept at the same strength (with a value of  $_{\rm A} = 0$  as for the bridge), prefers to retain the average charge density that a system with one less electron would have, as a change in the electron num ber at this site, will not result in any change in the total energy. The extra charge density at the donor site is compensated by the

reduction in the charge density at the bridge site next to the donor. The hopping energy between the donor and the st site of the bridge favors the presence of a hole (reduced charge density) at the rst bridge sites, so that the system can be stabilized by this kinetic energy. A further stabilization is e ected by the presence of the nearest neighbor C oulom b repulsion between the donor and this rst bridge site. On the other hand, the last bridge site shows an increased charge density and the last but one bridge site has a reduced charge density. This can be attributed to the fact that hopping between the last bridge site and the acceptor site gets renorm alized because of the absence of any Coulom b repulsion at the acceptor, and hence stabilizes the system . The reduced charge density at the last but one bridge site also allows for hopping between this site and the last bridge site. The presence of the extended Hubbard Coulomb repulsion energy V between these sites further enhances the kinetic stabilization. This excess electron and hole is form ed due to the correlations in the bridge; they are not free but are rather bound. W e refer to this kind of a quasiparticle as \exciton". The exciton is form ed at the ends of the bridge closest to the donor and the acceptor sites and is not spread over the chain. It has rather sm all volum e and is quite localized.

On the otherhand, for the case with large Coulomb repulsion at the donor and acceptor sites (case B), the ground state charge densities show contrasting behavior, as shown in Fig.6b. Here, the charge densities show a dip (reduction) at both the rst and the last bridge sites closest to the donor and the acceptor. In this case, the large Hubbard U ensures that there is only a maximum of one electron which is allowed at the donor and the acceptor sites. No matter how ever large the strength of the donor, the charge at the donor site does not exceed 1. When the donor and the acceptor energies are kept the same as the bridge ( $_{D} = _{B} = _{A} = 0$ ), the charge densities at all sites, except the bridge sites closest to the donor and the acceptor, have an average number of electrons that a system with one less electron from the half-lling would have. This reduction in charge at the rst and last bridge sites is due to the resulting stabilization that would be brought about by the presence of the hopping energy t and the extended Hubbard energy V. A higher donor strength shows a similar feature at the rst and the last sites of the bridge, how ever, a very prominent increase in the charge density can be seen in the rst half of the bridge and a corresponding reduction in the other half of the bridge. This shows that the exciton in this case spreads over the size of alm ost the length of the entire bridge. This large volum e exciton is in fact responsible for the long range electron transfer for the case B.

W e have also looked at the excited state charge densities for both cases. For case A, there is not much difference between the ground and the excited state charge densities indicating that the charge transfer has not been m ediated by the localized exciton, whereas for case B, the excited state charge densities clearly show the movem ent of charge density from the donor to the acceptor with a reverse in the density pattern from that found for the ground state.

W e also have studied the charge-charge correlations functions between the donor/acceptor and the bridge sites:  $hn_{D(A)}n_{B}i$   $hn_{D(A)}ihn_{B}i$ , where  $n_{D(A)}$  and  $n_{B}$ are the charge densities at the donor (acceptor) and the bridge sites respectively. The ground state charge-charge correlations in case A for all strengths of the donor, clearly indicates that the correlations between the donor and the bridge sites are very short ranged and that the exciton is not e ective in mediating the charge transfer between the far separated donor and the acceptor. In contrast to this, the correlations in case B shows features in correspondence with the charge density in Fig.6b, i.e., the donor is highly correlated with the bridge sites and hence transferm ediated by the exciton is the most likely mechanism of charge transfer between the donor and the acceptor.

F inally, to understand the nature of the excitonic state responsible for longer range electron transfer, we have plotted the chain axis component of the transition dipole m om ent between the ground state and the rst and second excited states for the two cases discussed above, for a particular donor strength ( $_{\rm D}$  = 4:0) in Fig.7a and 7b respectively. This transition dipole moment enters into the calculation of H  $_{\rm D\,A}\,$  and hence is a good m easure of the strength of the states carrying the charge for transferring to/from the acceptor end. W hen there is no Coulomb repulsion at the donor and the acceptor sites (case A), the transition dipole moment between the ground state and the rst excited state increases and then falls o for longer chain lengths, whereas that between the ground state and second excited state shows a steady increase, clearly indicating that the state carrying the charge is not a well de ned electron-hole bound pair state. On the other hand, for the case with large Coulomb repulsion at the donor and the acceptor sites (case B), the transition dipole m om ent between the ground and the rst excited state shows an increase with the increase in the chain length, indicating that this is the state which carries the charge through, from one end to the other even for large bridge length. The same between the ground state and the second excited state has zero value for small system sizes and a very negligible value for larger chain lengths, clearly showing that this state does not play any part in the charge transfer between the donor and the acceptor states. This emphatically proves that for case B, the charge is carried by a well de ned

fundam ental quasiparticle for all the chain length resulting in long-ranged electron-transfer.

## IV. CONCLUSIONS

To conclude, we have perform ed sem i-em pirical as well asm any-body DMRG calculations to understand the role of electron-electron interactions in governing the electron transfer in D - A system s. The electron transfer kinetics are truely long-ranged for strongly correlated conjugated polymeric systems and are not a ected much by the strength of the donors. W hat m atters is the extent of interactions in the bridge. It is the mediation of a stable electron-hole pair quasiparticle (exciton) in these system s that is responsible for sm all inverse decay length, . W e have been able to quantify the nature of the exciton from di erent levels of calculations. The CI calculations clearly indicate the formation of the excitonic state for an optimum size of the conjugated bridge length. But for the saturated bridge case, electrons are localized at the respective sites and thus the electron transfer has a much shorter range. From the quantum many-body long-chain DMRG calculations on system swith sm alland large correlations between the donor (acceptor) and the bridge sites, we have been able to distinguish the nature of the quasiparticle responsible for the electron transfer in the two cases. Our results show that when the donor (acceptor) and bridge are weakly correlated, the chargetransfer is purely by uncorrelated electron and hole, leading to short range ET observed in earlier theories. On the other hand, when the donor (acceptor) and bridge sites are strongly correlated, form ation of well-bound electronhole pair spread over the entire chain length is involved in the charge transfer, even for larger chain lengths, giving a very long-range ET. Thus, the exciton form ation is a crucialparam eter in controlling long-range ET . Form ation of such a quasi-particle is known to lead to very interesting optical properties in organic light em itting diodes. The fact that excitons also play an important role in transport mechanism signi es an important contribution towards fabrication of molecular wires where currents can be induced by optical excitations. A lso, for biom olecular systems with known long-range ET like DNA, the identi cation of right kind of quasiparticle responsible for charge and eld carrying processes can lead to novel applications in bio-m olecular optoelectronics. This how ever, requires further study.

- <sup>1</sup> (a) R.A.M arcus, Rev.M od.Phys.65, 599 (1993). (b) R.
   Huber, EM BO J.8, 2125 (1989). (c) J.D eisenhofer and H.
   M ichel, EM BO J.8, 2149 (1989).
- <sup>2</sup> (a) J. Xue and S. R. Forrest, Phys. Rev. B 69, 245322 (2004). (b) J.Muller, J.M. Lupton, A. L. Rogach, J. Feld-

<sup>3</sup> (a) D. Beljonne, G. Pourtois, M. A. Ratner and J. L. Bredas, J. Am. Chem. Soc. 125, 14510 (2003). (b) S. M.

m ann, D .V .Talapin and H .W eller, Phys.Rev.Lett.93, 167402 (2004). (c) J.Fujisawa and T.Ishihara, Phys.Rev. B 70, 113203 (2004).

G asper and G.B.Schuster, J.Am.Chem.Soc.119, 12762 (1997). (c) G.B.Schuster, Acc.Chem.Res.33, 253 (2000). (d) R.N.Barnett, C.L.Cleveland, A.Joy, U.Landm an and G.B.Schuster, Science 294, 567 (2001).

- <sup>4</sup> (a) A.Troisi, M.A.Ratner, M.W.Zimmt, J.Am.Chem. Soc. 126, 2215 (2004).
- <sup>5</sup> (a) M .M .Toutoun ji and M .A .R atner, J.Phys.Chem .A 104, 8566, (2000).
- <sup>6</sup> D. Baeriswyl, D. K. Campbell and S. Mazum dar, in \Conjugated Conducting Polymers", edited by H. Keiss (Springer-Verlag, Berlin, 1992) p.7.
- <sup>7</sup> (a) E. Cukier, S. Daniels, E. Vinson and R. J. Cave, J. Phys. Chem. A 106, 11240, (2002). (b) S. Antonello, F. Form aggio, A. Moretto, C. Toniolo and F. Maran, J. Am. Chem. Soc. 125, 2874 (2003).
- <sup>8</sup> S.S.Isied, M.Y.Ogawa, J.F.W ishart, Chem.Rev.92, 381 (1992); S.S.Isied in \Electron Transfer Reactions", edited by S.S.Isied (Am.Chem.Soc., Washington, 1997), p.331.
- <sup>9</sup> (a) T. Morita and S. Kimura, J. Am. Chem. Soc. 125, 8732 (2003). (b) D. Beljonne, G. Pourtois, C. Silva, E. Hennebioq, L. M. Herz, R. H. Friend, G. D. Scholes, S. Setayesh, K. Mullen and J. L. Bredas, Proc. Natn. Aca. Sci. 99, 10982, (2002).
- <sup>10</sup> A.Am ini and A.Harrim an, J.Photochem.Photobiol.C 4,155, (2003).
- <sup>11</sup> D.N.Beratan, J.N.Betts, J.N.Onuchic, Science, 252, 1285 (1991); S.Priyadarshy, S.M.Risser and D.N.Beraton, J.Phys.Chem. 100, 17678 (1996); D.J.Katz and A. A.Stuchebrukhov, J.Chem.Phys.109, 4960 (1998).
- <sup>12</sup> R.A.M arcus and N.Sutin, Biochim.Biophys.Acta 811, 265 (1985).
- <sup>13</sup> (a) M.Kemp, V.Mujica and M.A.Ratner, J.Chem.Phys. 101, 5172 (1994). (b) (a) M.Kemp, V.Mujica and M.A. Ratner, J.Chem.Phys.101, 6856 (1994).
- <sup>14</sup> M.D.Newton, Chem. Rev. 91, 767 (1991).
- <sup>15</sup> (a) S.G.Davison and J.D.Levine, Solid State Phys., 25,1 (1970). (b) S.K.Pati and S.Ram ashesha, J.Phys.: Condens.M atter, 16, 989 (2004).
- <sup>16</sup> R.G. Endres, D.L.Cox, R.R.P. Singh and S.K. Pati, Phys. Rev. Lett. 88, 166601 (2002).
- <sup>17</sup> N. Lokan, M. N. Paddon-Row, T. A. Smith, M. L. Ross, K. P. Ghiggino and S. Speiser, J. Am. Chem. Soc. 121, 2917 (1999).
- $^{18}$  M .B ixon and J.Jortner, J.Chem .P hys.107, 5154 (1997).
- <sup>19</sup> H. Zhao and S. M azum dar, Phys. Rev. Lett. 93, 157402 (2004).
- <sup>20</sup> L.A.Curtiss and J.R.M iller, J.Phys.Chem .A 102, 160, (1998).
- <sup>21</sup> S.R.W hite, Phys.Rev.Lett. 69, 2863 (1992); Phys.Rev. B.48, 10345 (1993).
- <sup>22</sup> M.J.S.Dewar, E.G. Zoebisch, E.F.Healy and J.J.P. Stewart, J.Am.Chem.Soc.107, 3902 (1985).
- <sup>23</sup> (a) A.D.Becke, J.Chem.Phys.98, 5648, (1993). (b) C.
   Lee, W. Yang, R.G.Parr, Phys. Rev. B 37, 785, (1988).
- <sup>24</sup> J.Ridley, M.C.Zemer, Theor.Chim.Acta 32, 111 (1973);
   A.D.Bacon, M.C.Zemer, Theor.Chim.Acta 53, 21 (1979).
- <sup>25</sup> R.J.Buenker and S.D.Peyerim ho , Theor. Chim . Acta 35, 33 (1974).
- <sup>26</sup> Z. Shuai, D. Beljonne and J. L. Bredas, J. Chem. Phys. 97, 1132 (1992).
- <sup>27</sup> D.Beljonne, Z.Shuai, J.Comil, D.dos Santos and J.L. Bredas, J.Chem. Phys. 111, 2829 (1999).

- <sup>28</sup> Y.K.Shin, M.D.Newton and S.S.Isied, J.Am.Chem. Soc. 125, 3722 (2003).
- <sup>29</sup> (a) M .Pascaly, J.Yoo and J.K .Barton, J.Am .Chem .Soc. 124, 9083 (2002). (b) H.W .Fink and C.Schonenberger, Nature 398, 407 (1999). (c).T.T.W illiam s, D.T.O dom and J.K.Barton, J.Am .Chem .Soc. 122, 9048 (2000).
- <sup>30</sup> (a) H.W ang, J.P. Lew is and O.F. Sankey, Phys. Rev. Lett. 93, 016401 (2004). (b) M. Taniguchi and T.K awai, Phys.Rev.E 70, 011913 (2004). (c) F.Evers, F.W eigend and M.Koentopp, Phys.Rev.B 69, 235411 (2004).
- <sup>31</sup> (a) A. Kadashchuk et al. Phys. Rev. Lett. 93, 066803 (2004). (b) V. Perebeinos, J. Terso and P. Avouris, Phys. Rev. Lett. 92, 257402 (2004). (c) K. Hummer, P. Puschnig and C. A-D raxl, Phys. Rev. Lett. 92, 147402 (2004).
- <sup>32</sup> L. Ley, M. Taniguchi, J. G hijsen, R. L. Johnson, A. Fujin ori, Phys. Rev. B 35, 2839 (1987); A. Fujin ori, N. K in izuka, M. Taniguchi and S. Suga, Phys. Rev. B 36, 6691 (1987); J.O kabayashiet al., Phys. Rev. B 59, R2486 (1999).

1. 
$$D - (CH=CH)_n - A$$
  
 $D = NH_2, OCH_3, SCH_3$   
 $A = NO_2$ 

2. 
$$D - (CH_2 - CH_2)_n - A$$
  
 $D = NH_2$   
 $A = NO_2$ 

FIG.1: Model system s considered for study




FIG.2: ET coupling matrix,  $H_{D,A}$  (in eV) with respect to the increase in the number of ole nic spacers for di erent donors, D and xed  $A = NO_2$ 




FIG.3: Frontier orbital plots with unsaturated and saturated spacers. The shading of the lobes show the parity. A tom symbol: Red: Oxygen, Blue: Nitrogen, Black: Carbon, W hite: Hydrogen. Light green arrow shows the direction of the dipole moment




FIG.4:Coe cients ofm olecular orbital occupancy on the N atom of N H  $_2$  and the N atom and two O atom of N O  $_2$  before (circles) and after (stars) ET .




FIG.5: ET coupling matrix H  $_{\rm D~A}$  with respect to the bridge length for case 1 (above) and case 2 (below) for dimensional error values of the donor strengths ( $_{\rm D}$  = 0:0 to  $_{\rm D}$  = 4:0)




FIG.6: Charge densities at every bridge site for case 1 (above) and case 2 (below) for di erent values of the donor strengths (  $_{\rm D}$  = 0.0 to  $_{\rm D}$  = -4.0)




FIG.7: M odulus of the x component of the transition dipole m om ent,  $j_x jw$  ith respect to chain length for case 1 (above) and case 2 (below) for transition from ground state to rst excited state ( led circles) and ground state to second excited state (diam onds)