

How Multivalency controls Ionic Criticality

Michael E. Fisher, Jean-Noël Aqua,* and Shubho Banerjee†

Institute for Physical Science and Technology, University of Maryland, College Park, Maryland 20742, USA

(Dated: March 23, 2022)

To understand how multivalency influences the reduced critical temperatures, $T_c^*(z)$, and densities, $\rho_c^*(z)$, of $z:1$ ionic fluids, we study equisized hard-sphere models with $z = 1-3$. Following Debye, Hückel and Bjerrum, association into ion clusters is treated with, also, ionic solvation and excluded volume. In good accord with simulations but contradicting integral-equation and field theories, T_c^* falls when z increases while ρ_c^* rises steeply: that 80 – 90% of the ions are bound in clusters near T_c serves to explain these trends. For $z \neq 1$ interphase Galvani potentials arise and are evaluated.

PACS numbers: 05.70.Fh, 61.20.Qg, 64.60.Fr, 64.70.Fx

Multivalent ions play a significant role in condensed-matter, physicochemical, biophysical and, via the plasma transition, astrophysical contexts [1]. The effects of multivalency are, however, often hard to comprehend. One central issue—relevant to electrolyte solutions, molten salts, liquid metals, and dense plasmas [1]—arises in Coulomb-driven phase separation. The most basic model for such ionic fluids consists of $N = \rho V$ hard-core spherical ions of various species σ in a volume V of uniform dielectric constant D , with $N_\sigma = \rho_\sigma V$ ions of diameter a_σ carrying charges $q_\sigma = z_\sigma q_0$, where q_0 is an elementary charge. In the simple equisized $z:1$ charge asymmetric primitive models (C_z APMs), on which we focus here, one has $\sigma = +, -$, $a_+ = a_-$, and $q_+ = zq_0$, $q_- = -q_0$. The basic energy scale and associated reduced temperature and density are then $\varepsilon = zq_0^2/Da$, $T^* = k_B T/\varepsilon$, $\rho^* = \rho a^3$.

Monte Carlo simulations [2] show that (at least for $z \lesssim 5$) the C_z APMs exhibit “gas-liquid” phase separation; furthermore, the critical parameters, $T_c^*(z)$ and $\rho_c^*(z)$, are found to reasonable precision: see Table I and the open circles in Figs. 1 and 2. One observes that $T_c^*(z)$ falls with increasing z , while $\rho_c^*(z)$ rises sharply. But we ask: How can these trends be understood? Or accounted for semiquantitatively? To address this issue we review briefly previous work, including a pioneering field-theoretic attack [3], and then report on a recent study [4] which we believe provides significant insight. This extends an earlier analysis I [5] for the symmetric $z=1$ restricted primitive model (RPM) that was founded on the original Debye-Hückel (DH) approach but incorporated (i) Bjerrum ion pairs and (ii) their solvation in the residual ionic fluid. For $z=2$ and 3 larger ion clusters, trimers and tetramers, must be included [4, 6]; but then explicit results are also obtained for the interphase Galvani potential [7] that appears in any two-phase nonsymmetric ionic system [4, 7].

The field-theoretic analysis of Netz and Orland (NO) [3] was designed to address $z:1$ ionic fluids and colloids ($z \gg 1$) and to include correlations in a systematic manner. The Coulomb interaction, $q_\sigma q_\tau/r$, was transformed to yield a functional integral over an auxiliary potential $\phi(\mathbf{r})$. At the $\langle \phi^2 \rangle$ level the DH effective interaction,




FIG. 1: Reduced critical temperatures for $z:1$ charge asymmetric equisized hard-core primitive model electrolytes (C_z APMs) according to Monte Carlo (MC) simulations [2]; Debye-Hückel (DH) theory; field-theoretic approaches: NO [3] (with a factor $\frac{1}{10}$) and NMF [8]; approximate integral equations: MSA [9], SPB, and MPB [10]; and the present DHBjCI and DHBjCIHC solvated ion-cluster theory [4]; See text.

TABLE I: Monte Carlo (MC) estimates [2] for the reduced critical parameters for $z:1$ equisized hard-sphere electrolytes, values calculated from DHBjCIHC theory (CI) [4], and approximate estimates based on ion cluster statistics: see text.

	critical temp.		$10^2 T_c^*(z)$		critical density		$10^2 \rho_c^*(z)$	
z	MC	CI	E_{DH}	E_{MC}	MC	CI	E_ρ	E_κ
1	4.93 ₃	5.56 ₉	5.45	4.93 ₅	7.50	2.61 ₄	2.72	2.37
2	4.70	4.90 ₇	5.11	4.65	9.3	6.26 ₁	4.27	3.49
3	4.10	4.33 ₄	4.85	4.44	12.5	11.90	6.96	5.40

$v_{DH} \propto e^{-\kappa r}/r$ is captured with

$$\kappa^2(T; \{\rho_\sigma\}) = 4\pi(q_0^2/Dk_B T) \sum_\sigma z_\sigma^2 \rho_\sigma. \quad (1)$$

The reduced free energy density, $\bar{f}(T; \rho) \equiv -F/Vk_B T$, was computed to eighth order in ϕ but a momentum cutoff is essential: NO adopted $|\mathbf{k}_\Lambda| = 2\pi/a$ thereby incorporating the ionic diameter and, for the $z:1$ case, leading to




FIG. 2: Reduced critical densities $\rho_c^*(z)$, for the C_z APM electrolyte as in Fig. 1 (except that the NO plot is not rescaled).

$\kappa^2 a^2 = 4\pi\rho^*/T^*$. Since this treatment of the hard cores is approximate, accurate predictions for $T_c^*(z)$ and $\rho_c^*(z)$ are not expected. Nevertheless, one might anticipate reliable *trends* when z varies in contrast to DH theory which yields *no dependence* on z with (after I)

$$\text{DH: } \kappa_c a = 1, \quad T_c^* = \frac{1}{16}, \quad \rho_c^* = 1/64\pi \simeq 0.005. \quad (2)$$

In fact, as NO report, “the [predicted] deviations from DH theory are pronounced” for $z > 1$: see the bold dashed plots in Figs. 1 and 2.

But evidently the NO results are not merely quantitatively wrong; the *trends* are quite incorrect since T_c^* is asserted to rise rapidly (instead of falling) while ρ_c^* falls sharply for small $z-1$ (instead of rising) and *then* increases but much too slowly. While one may blame the approximate treatment of the hard cores, we believe this is *not* the primary culprit. Indeed, a recent field-theoretic analysis paid closer attention to the ion-ion repulsions [8]; but the subsequent “new mean-field” (NMF) results still exhibit strong increases in T_c^* and an overly weak variation of ρ_c^* : see the NMF plots in Figs. 1 and 2 [8].

Integral equation theories are hardly better: see Figs. 1 and 2. The *mean spherical approximation* (MSA), like DH theory, predicts *no variation* of T_c^* and ρ_c^* with z [9]. A *symmetric Poisson-Boltzmann* (SPB) theory [10] *does* predict the correct falling and rising trends for T_c^* and ρ_c^* , but the degree of variation is woefully inadequate. Moreover, the *modified Poisson-Boltzmann* (MPB) approximation, that the same authors [10] argue should be more reliable, yields the *wrong* trend for T_c^* .

In order to better understand the effects of multivalency we turn to recent calculations [4, 6] based on the *solvated ion-cluster* view [5] of the C_z APM near criticality that is supported ‘pictorially’ by simulations [2]. In brief, the aim is to construct the free energy density, $\bar{f}(T; \{\rho_\sigma\})$, for ionic species σ consisting (i) of + and – *monomers*, i.e., isolated, $n_+ = n_- = 1$ single, unassociated




FIG. 3: Coexistence curves predicted for $z:1$ equisized primitive models by the DHBjCI and DHBjCIHC theories (solid and dashed lines, respectively) together with Monte Carlo estimates (MC) based on [11].

ions of valency $z_+ = z$ and $z_- = -1$; (ii) a set of associated *primary* clusters, $\sigma = 2, 3, \dots$, dimers, trimers, ..., each consisting of one “central” + ion and $m_\sigma = \sigma - 1$ “satellite” counterions for a total of $n_\sigma = m_\sigma + 1$ ions in a cluster of valency $z_\sigma = z - m_\sigma$; up to (iii), the largest primary cluster, the *neutral* or ‘molecular’ $(z+1)$ -mer of one z_+ ion and z negative ions [4, 6].

For each species, \bar{f} contains an ideal-gas term $\bar{f}^{\text{id}}(T, \rho_\sigma)$, and an electrostatic term $\bar{f}_\sigma^{\text{el}}(T, \{\rho_\tau\})$, that, following DH, incorporates *Cluster solvation* in the partially associated *Ionic fluid*: this description is thus dubbed “DHBjCI” [4]. By adding a **H**ard **C**ore free-volume term, $\bar{f}^{\text{HC}}(\{\rho_\sigma\})$, as in I, one may also account for those excluded volume effects not *already* encompassed in the basic solvation and association calculations [4, 5], so generating a “DHBjCIHC” theory [4]. (The effective HC virial coefficient $B_\sigma^{\text{bcc}} = 4a_\sigma^3/3^{3/2}$ has been adopted [4, 5].) Examination of Figs. 1 and 2 reveals that these solvated ion-cluster theories are surprisingly successful! Not only are *both* the downward trend in $T_c^*(z)$ and the rapid rise of $\rho_c^*(z)$ well captured, but the *quantitative* agreement with each of the MC estimates is significantly better than achieved by other approaches.

One must recognize that (all) these theories are of mean-field character: thus 5 to 15% over-estimates of $T_c^*(z)$ are to be expected. Indeed, neglected fluctuations typically depress T_c by such amounts and also flatten the *coexistence curves* as seen in Fig. 3. Second, note that the hard-core terms have a small effect on $\rho_c^*(z)$ while reducing $T_c^*(z)$ values by only 5–10%. Nevertheless, Fig. 3 reveals that the liquid phases, especially for $\rho^* \gtrsim 0.15$, are sensitive to \bar{f}^{HC} : but, recall the discussion in I. In fact, the crucial feature of DHBj-type theories—not represented in field-theoretic or standard integral-equation treatments—is the *chemical equilibrium* maintained be-

tween the cluster species via the Law of Mass Action:

$$\rho_\sigma = K_{m,z}(T) \rho_+ \rho_-^m \exp [\mu_+^{\text{Ex}} + m\mu_-^{\text{Ex}} - \mu_\sigma^{\text{Ex}}], \quad (3)$$

for $\sigma = m + 1 \geq 2$, with the excess chemical potentials $\mu_\sigma^{\text{Ex}} = -(\partial/\partial\rho_\sigma)[\bar{f}^{\text{HC}} + \sum_\sigma \bar{f}_\sigma^{\text{El}}]$, while the *association constants* are taken as [4, 5]

$$K_{m,z}(T; R) = \frac{1}{m!} \prod_{i=1}^m \int_a^R d\mathbf{r}_i \exp \left(-\frac{\mathcal{E}_{m,z}(\{\mathbf{r}_i\})}{k_B T} \right), \quad (4)$$

in which $\mathcal{E}_{m,z}(\{\mathbf{r}_i\})$ is the electrostatic energy of an isolated $(m + 1)$ -mer with satellite coordinates $\{\mathbf{r}_i\}$. The lower limits a and the condition $\mathcal{E}_{m,z} = +\infty$ for $|\mathbf{r}_i - \mathbf{r}_j| < a$, represent hard cores. Following Bjerrum [5], the necessary cut-off radius R is chosen so that $(\partial K_{m,z}/\partial R)$ is minimal. The resulting 3-fold $K_{2,z}$ integral is manageable but the 6-fold integral for $K_{3,3}$ requires a Padé approximant study of the low- T expansion cross-checked to a part in 10^3 by MC evaluations [4]. It transpires, however, that T_c^* and ρ_c^* are *insensitive* to the $K_{m,z}$ values [4].

Lastly, one needs to account for the solvation of *all* the ion species, σ , by the free ions and charged clusters via the electrostatic terms [4, 5, 6]

$$\bar{f}_\sigma^{\text{El}}(T; \{\rho_\tau\}) = \frac{4\pi\rho_\sigma}{Dk_B T} \sum_{l=0}^{\infty} \frac{u_{2l}(\kappa a_\sigma)}{a_\sigma^{2l+1}} \sum_{m=-l}^l \langle |Q_{lm}^\sigma|^2 \rangle, \quad (5)$$

where the $u_{2l}(x)$ are related to the spherical Bessel functions $k_l(x)$ [4]; the second sum requires the cluster electric multipole moments, Q_{lm}^σ , thermally averaged [4] over the ionic configurations that already enter in the $K_{m,z}(T)$.

Finally, a_σ is an *effective cluster diameter*, i.e. the radius of the approximating sphere (centered to minimize $\bar{f}_\sigma^{\text{El}}$) that substitutes for the true, thermally fluctuating, hard-core exclusion domain: see I and [4]. One concludes, as in I, that a most reasonable choice for a_σ is the average over solid angle of the radial distance to the true exclusion surface of the ground-state cluster: this yields $a_2 = (\frac{3}{4} + \frac{3}{8} \ln 3) \simeq 1.162a$, $a_3 = 1.250a$ and $a_4 = 1.375a$. For $z=1$ the values of T_c^* and ρ_c^* vary by less than $\pm 2\%$ over plausible alternatives for a_2 [4]; but the sensitivity to a_3 and a_4 for $z=2$ and 3 is greater. As a result, this hard-to-avoid approximation contributes significantly to the overall quantitative uncertainties.

From the total free energy $\bar{f}(T, \{\rho_\sigma\})$, all thermodynamic properties follow [4, 5]. One may then conclude from Figs. 1 and 2 that the principal defect of the field-theoretic and integral-equation approaches is a failure to account effectively for strong ionic association near criticality. But can the actual *trends* of T_c^* and ρ_c^* with z be demonstrated in a direct, transparent way? To answer, consider the *fractions*, $y_\sigma = n_\sigma N_\sigma / N$, of ions bound in clusters of n_σ ions with $\rho_\sigma = (y_\sigma / n_\sigma) \rho$. The critical point

TABLE II: Inverse screening length κ and fractions, $y_\sigma = n_\sigma N_\sigma / N$, of ions in clusters of n_σ ions at criticality, as percentages, according to DHBjCIHC theory [4].

z	$\kappa_c a$	y_+^c	y_-^c	y_2^c	y_3^c	y_4^c
1	1.04	9.14	9.14	81.72	—	—
2	1.37	1.31	10.33	15.43	72.93	—
3	1.57	0.34	8.04	3.32	11.13	77.17

values that result from DHBjCIHC theory [4, 12] are displayed in Table II. A significant fact is the rapid decrease in y_+^c , the fraction of unassociated z_+ ions, from 9.1 to 1.3 to 0.3%. But more can be learned!

To understand the variation of $T_c^*(z)$ let us regard the electrolyte in the critical region as a mixture of clusters with fixed mole fractions $x_\sigma = (y_\sigma / n_\sigma) / \sum_\tau (y_\tau / n_\tau)$. A pair (σ, τ) will either mutually repel or attract with pairwise binding energies, say, $\varepsilon_{\sigma\tau}$. Thus unlike monomers attract with $\varepsilon_\pm = \varepsilon$. However, a *dimer* attracts only *negative monomers* with $\varepsilon_{2-} = (z - \frac{1}{2})\varepsilon/z$; but repels all $z_+ \geq +2$ ions. Two dimers repel when $z \geq 3$; but one has $\varepsilon_{2,2}/\varepsilon \simeq 0.586$ and 0.345 for $z=1$ and 2. And so on.

To estimate T_c^* for this mixture we adopt a van-der-Waals approach as in [2(d)]. Thus, for the overall cluster density $\hat{\rho} (= \rho \sum_\sigma y_\sigma / n_\sigma)$, we take $p/\hat{\rho}k_B T \simeq Z(B_0 \hat{\rho}) + B_1(T^*) \hat{\rho}$ with $Z(u) = 1 + u + \dots$ in which the second virial coefficient has been decomposed as $B(T^*) = B_0 + B_1(T^*)$ where $B_0 (= b_0 a^3$, say) represents the hard-core repulsions while $B_1(T^*)$ embodies the attractions. Solving $\partial_\rho p = \partial_\rho^2 p = 0$, as usual, yields ρ_c^* and $B_c^* \equiv B_1(T_c^*)/b_0 a^3$. At low T , which is relevant here, one has

$$B_1(T^*) \approx - \sum_{\sigma, \tau} b_{\sigma\tau} a^3 x_\sigma x_\tau \exp(\varepsilon_{\sigma\tau}^* / T^*), \quad (6)$$

where $\varepsilon_{\sigma\tau}^* \equiv \varepsilon_{\sigma\tau} / \varepsilon$, while $b_{\sigma\tau} a^3$ specifies the volume of mutual attractions: this *vanishes* if σ and τ repel.

Now, the $x_+ x_-$ term dominates in $B_1(T^*)$ at low T with corrections of relative order $(x_2^2 / x_+ x_-) e^{-0.414/T^*}$ for $z=1$ and $2(x_2 / x_+) e^{-1/2zT^*}$ for $z \geq 2$. We may then *calibrate* $B_1(T_c^*)/a^3$ by using pure DH theory (2) for which, since association is not considered, $x_+ = x_- = \frac{1}{2}$. Thereby we obtain the E_{DH} estimates

$$T_c^*(z) \simeq 1 / (16 + |\ln 4 x_+^c(z) x_-^c(z)|), \quad (7)$$

in which $x_+^c \propto y_+^c$ and $x_-^c \propto y_-^c$ follow from Table II.

The resulting predictions are listed in Table I under E_{DH} . In light of the heuristic nature of the arguments, they reflect the trend of the MC and CI values surprisingly well. Certainly the contention that association is a prime factor is well confirmed. By replacing 16 by 20.27 (or 17.96) in (7), and the factor 4 by $1/x_+^c(1)x_-^c(1)$, one calibrates $B_1(T_c^*)$ on the MC (or CI) values for the RPM. Column E_{MC} in Table I lists the MC-calibrated values: for $z=2$ and 3 these match the Monte Carlo estimates to within 1% and 8%, respectively [13].




FIG. 4: Reduced Galvani potentials, $\Delta\bar{\phi} = q_0\Delta\phi/k_B T$, vs. T/T_c for $z:1$ electrolytes according to pure DH theory (dotted) and DHBjCI(HC) theories : solid (dashed) plots.

Now, for the critical density, the significance of ion pairing is already clear in *pure* DHBj theory for the RPM [5]. The heavy depletion of the free ions (which, in DHBj theory, drive the transition *alone*) means that to reach criticality the overall density $\rho (= \rho_+ + \rho_- + 2\rho_2)$ must be increased until the DH criterion $\rho_+^* + \rho_-^* = \rho_{DH}^* = 1/64\pi$ is met : see (2). Does the same depletion-by-association mechanism account for the z -dependence of $\rho_c^*(z)$?

To progress, rewrite (1) generally as $\kappa^2 a^2 = 4\pi\rho^\dagger/T^*$, with the *effective, depleted ionic density*

$$\rho^\dagger \equiv \rho^* \sum_{\sigma} z_{\sigma}^2 y_{\sigma}(T; \{\rho_{\sigma}\}) / z n_{\sigma}. \quad (8)$$

If one accepts the DH criterion and uses Table II, the estimates E_{ρ} , in Table I, result. Although these fall short of the Monte Carlo values by 74, 54 and 44% for $z = 1-3$, they reproduce the accelerating increase with z (by factors 1.57, 1.63 vs. 1.24, 1.34).

An alternative approach adopts the DH value $\kappa_c a = 1$: see (2) but note from Table II that DHBjCIHC theory implies that $\kappa_c a$ rises from 1.04 for the RPM to 1.57 for $z=3$. Then using the E_{DH} values for T_c^* , in Table I, leads to the E_{κ} predictions for $\rho_c^*(z)$: these are all rather low but the increases with z , by factors 1.47, 1.55, again reflect the correct behavior.

Finally, we note that the Galvani potential, $\Delta\phi$, that arises between coexisting phases in charge asymmetric fluids is readily calculated [4, 7]. The predictions from pure DH theory are shown dotted in Fig. 4 : one finds $\Delta\phi_{DH} \propto (1 - z^{-1})$. The other plots result from the DHBjCI and DHBjCIHC theories [4]. Surprisingly, the calculations suggest no clear trend with z . It is natural to conjecture that $\Delta\phi$ vanishes as $G_0(T_c - T)^{\beta}$; moreover to the extent that the expected mean-field value $\beta = \frac{1}{2}$ is realized, the present results support this.

In conclusion, we have elucidated the mechanisms underlying how multivalency influences critical behavior.

Specifically, we have summarized briefly analytical calculations for 3:1, 2:1 and 1:1 equisized charged hard-sphere fluids [4] that, for the first time, reasonably reflect the true variation of critical temperatures and densities, $T_c^*(z)$ and $\rho_c^*(z)$ (as revealed by simulations [2]). On that basis, supported by analysis that correlates $T_c^*(z)$ and $\rho_c^*(z)$ with the increasingly depleted populations of free ions and charged clusters as z increases, it is clear that recognizing ionic association is inescapable for a successful theory. Previous treatments [3, 8, 9, 10], lacking allowance for ion clusters fail seriously. The ion-cluster solvation theories also yield quantitative results for the interphase Galvani potentials.

National Science Foundation support via Grants CHE 99-81772 and 03-01101 is gratefully acknowledged.

* Present address: École normale supérieure de Lyon, 69364, Lyon, France

† Present address: Department of Physics, Rhodes College, Memphis TN 38112

- [1] See, (a) articles in *Elementary Processes in Dense Plasmas*, edited by S. Ichimaru and S. Ogata (Addison-Wesley, New-York, 1995); (b) A. A. Likhalev, Phys. Uspekhi **43**, 777 (2000); (c) A. Yu. Grosberg *et al.*, Rev. Mod. Phys. **74**, 329 (2000); (d) H. Weingärtner and W. Schröer, Adv. Chem. Phys. **116**, 1 (2001); (e) Y. Levin, Rep. Prog. Phys. **65**, 1377 (2002).
- [2] (a) P. J. Camp and G. N. Patey, J. Chem. Phys. **111**, 9000 (1999); (b) Q. Yan and J. J. de Pablo, Phys. Rev. Lett. **86**, 2054 (2001); (c) A. Z. Panagiotopoulos and M. E. Fisher, Phys. Rev. Lett. **88**, 045701 (2002); (d) Y. C. Kim and M. E. Fisher, Phys. Rev. Lett. **92**, 185703 (2004).
- [3] R. R. Netz and H. Orland, Europhys. Lett. **45**, 726 (1999).
- [4] J.-N. Aqua, S. Banerjee, and M. E. Fisher, cond-mat/0410692 [submitted for publication].
- [5] (a) M. E. Fisher and Y. Levin, Phys. Rev. Lett. **71**, 3826 (1993); (b) Y. Levin and M. E. Fisher, Physica A **225**, 164 (1996); to be denoted I.
- [6] M. N. Artyomov *et al.*, J. Chem. Phys. **118**, 6394 (2003).
- [7] M. J. Sparnaay, *The Electrical Double Layer* (Pergamon Press, Oxford, 1972), pp. 4, 338; P. B. Warren, J. Chem. Phys. **112**, 4683 (2000); I. Iosilevski and A. Chigvintsev, J. Phys. IV France **10**, Pr5-451 (2000); A. R. Muratov, Zh. Éksp Teor. Fiz. **120**, 104 (2001) [Sov. Phys. JETP **93**, 89 (2001)].
- [8] J.-M. Caillol, Mol. Phys. **103**, 1271 (2005); we quote the (most favorable) “MSA regularization” results.
- [9] E. Waisman and J. L. Lebowitz, J. Chem. Phys. **56**, 3086 (1972); E. Gonzalez-Tovar, Mol. Phys. **97**, 1203 (1999).
- [10] A. K. Sabir *et al.*, Mol. Phys. **93**, 405 (1998); C. W. Outhwaite, Cond. Matt. Phys. **7**, 719 (2004).
- [11] Y. C. Kim, M. E. Fisher, and E. Luijten, Phys. Rev. Lett. **91**, 065701 (2003).
- [12] While the y_{σ}^c are *not* very sensitive to the a_{σ} , they cannot be determined unequivocally via simulations.
- [13] The arguments based on (6) extend naturally to charged

hard dumbbell models and their variants : see J. M. Romero-Enrique *et al.*, Phys. Rev. E **66**, 041204 (2002).