

Statistical mechanics in the context of special relativity II

G. Kaniadakis*

*Dipartimento di Fisica, Politecnico di Torino,
Corso Duca degli Abruzzi 24, 10129 Torino, Italy*

(Dated: May 1, 2019)

The special relativity laws emerge as one-parameter (light speed) generalizations of the corresponding laws of classical physics. These generalizations, imposed by the Lorentz transformations, affect both the definition of the various physical observables (e.g. momentum, energy, etc), as well as the mathematical apparatus of the theory. Here, following the general lines of [Phys. Rev. E **66**, 056125 (2002)], we show that the Lorentz transformations impose also a proper one-parameter generalization of the classical Boltzmann-Gibbs-Shannon entropy. The obtained relativistic entropy permits to construct a coherent and selfconsistent relativistic statistical theory, preserving the main features of the ordinary statistical theory, which recovers in the classical limit. The predicted distribution function is a one-parameter continuous deformation of the classical Maxwell-Boltzmann distribution and has a simple analytic form, showing power law tails in accordance with the experimental evidence. Furthermore the new statistical mechanics can be obtained as stationary case of a generalized kinetic theory governed by an evolution equation obeying the H-theorem and reproducing the Boltzmann equation of the ordinary kinetics in the classical limit.

PACS numbers: 03.30.+p, 05.20.-y

I. INTRODUCTION

In high energy physics, various approaches of statistical mechanics and kinetic theory have been adopted to study cosmological models [1, 2], multiparticle production in hadron-hadron collisions [3], possible Lorentz invariance violation in extension of standard-model [4], neutron star matter [5], black hole models [6, 7, 8, 9] etc. Common to all such approaches is the explicit or implicit assumption of a very specific form of the entropy, i.e. the celebrated Boltzmann-Gibbs-Shannon entropy which, according to Jaynes [10] maximum entropy, principle conducts to the Maxwell-Boltzmann exponential distribution.

On the other hand since long time it is known that there exist open problems in high energy physics which cannot be solved within the ordinary Maxwell-Boltzmann statistics e.g the cosmic rays spectrum [11, 12, 13] and the black hole entropy Bekenstein-Hawking area law [14, 15].

In statistical physics, beside the ubiquitous exponential distribution $\exp(-x)$, also non-exponential distributions have been considered. The most famous ones appear within the quantum statistical mechanics, namely the Fermi-Dirac and the Bose-Einstein distributions, which are forced by the exclusion-inclusion principle. In the last decades a particular attention has been addressed to the statistical distributions presenting power law tails as x^{-a} . The latter distributions have been observed experimentally in many fields of physics, including cosmic rays physics [11, 12, 13], plasma physics [16] and multiparticle production processes [17].

In order to propose theories based on non-exponential statistical distributions, one can follow two different

ways. The first is the entropic one consisting of a proper generalization of the system entropy. After fixing the entropy, the Jaynes maximum entropy principle univocally determines the form of the distribution function. The problem then is essentially reduced to the choice of the right entropic form.

The second way is the kinetic one which is very familiar in non equilibrium statistical mechanics. It is well known that the classical statistical mechanics can be obtained as particular case of the ordinary kinetic theory for $t \rightarrow \infty$, within the Boltzmann or the Fokker-Planck picture. Generalized kinetic theories necessarily conduct to generalized statistical mechanics producing non-exponential distributions.

Both the entropic as well as the kinetic path, conduct to the conclusion that it is possible to develop, beside the classical statistical mechanics, other coherent and self-consistent statistical theories [18, 19, 20, 21, 22, 23, 24].

It is remarkable that the Maxwell-Boltzmann distribution does not emerge within the statistical mechanics. It is more proper to say that the classical statistical mechanics is built starting from the Maxwell-Boltzmann distribution. It is commonly accepted that this distribution emerges within the Newton mechanics. Indeed, numerical simulation of classical molecular dynamics unequivocally conducts to the exponential distribution. At this point, the question if the Maxwell-Boltzmann distribution obtains also in the case when the microscopic dynamics is governed by the special relativity laws, naturally arises.

In the relativistic statistical theory the collision invariants are imposed by the laws of special relativity. Hence the first difference between the classical and the relativistic distribution functions regards the different forms of the collision invariants, in the argument of the distribution function. A second difference could be origi-

*Electronic address: giorgio.kaniadakis@polito.it

nated by the form of the entropy of the relativistic many body system, which could be different with respect to the Boltzmann-Gibbs-Shannon entropy.

In the ordinary relativistic statistical mechanics it is accepted that the entropy has the same form of the classical statistical mechanics. Then the Jaynes principle yields an exponential distribution which reproduces exactly the Maxwell-Boltzmann distribution in the rest frame. The most famous and important relativistic particle system is undoubtedly the cosmic rays [11, 12, 13]. The cosmic rays spectrum has a very large extension (13 decades in energy and 33 decades in particle flux) and presents a power law asymptotic behavior. Unfortunately the Maxwell-Boltzmann distribution fails to explain this experimental spectrum. The same power law asymptotic behavior of the distribution function was observed also in other relativistic systems [16, 17]. Hence for relativistic particle systems, the experimental evidence suggests a non-exponential distribution function with power law tails. This distribution, can be originated exclusively by an entropy manifestly different from the Boltzmann-Gibbs-Shannon one.

We recall that all the classical physical observables (momentum, energy, etc) when considered within the special relativity are properly generalized. Any relativistic formula can be viewed as a one-parameter (light speed) generalization or deformation of the corresponding classical formula. Consequently it is natural to assume that also the entropy of a relativistic system could be a one-parameter generalization of the classical entropy. In this case the logarithm appearing in the expression of Boltzmann-Gibbs-Shannon entropy must be replaced by another one parameter generalized logarithm. Then also the ordinary exponential function appearing in the Maxwell-Boltzmann distribution must be replaced by an one-parameter generalized exponential. These two generalized functions have been proposed heuristically in ref. [18] and are given by

$$\exp_{\{\kappa\}}(t) = \left(\sqrt{1 + \kappa^2 t^2} + \kappa t \right)^{1/\kappa}, \quad (1.1)$$

$$\ln_{\{\kappa\}}(t) = \frac{t^\kappa - t^{-\kappa}}{2\kappa}. \quad (1.2)$$

Note that in the classical limit $\kappa \rightarrow 0$ one easily recovers the ordinary exponential and logarithm namely $\exp_{\{0\}}(t) = \exp(t)$, $\ln_{\{0\}}(t) = \ln(t)$. Successively in ref. [19] it has been shown that the statistical theory, developed starting from these generalized functions, emerges within the framework of the special relativity. The entropy for a relativistic many body system assumes the form

$$S(f) = - \int d^3p f \ln_{\{\kappa\}}(f) = - \langle \ln_{\{\kappa\}}(f) \rangle, \quad (1.3)$$

while the relevant statistical distribution in the rest frame is given by

$$f = \alpha \exp_{\{\kappa\}} \left(- \frac{W - \mu}{\lambda k_B T} \right), \quad (1.4)$$

W being the relativistic kinetic energy, while α and λ are two constants depending on κ . These two latter relationships are one-parameter generalizations of the Boltzmann-Gibbs-Shannon entropy and of the Maxwell-Boltzmann distribution respectively, which are recovered in the classical limit $\kappa \rightarrow 0$.

Main goal of the present contribution is to show that the functions $\exp_{\{\kappa\}}(t)$ and $\ln_{\{\kappa\}}(t)$, leading to the distribution function (1.4) and to the entropy (1.3), emerge naturally and unequivocally within the special relativity theory. These functions replace the ordinary exponential and logarithm, relegating them at the native status of classical functions. It is remarkable that the relativistic statistical theory, based on the entropy (1.3), predicts a power law asymptotic behaviour to the spectrum of relativistic particles according to the experimental evidence.

The paper is organized as follows:

In Sect. II we recall some concepts of relativistic dynamics focusing our attention on the deformations in the mathematical formalism enforced by the finite value of the light speed.

In Sect. III we show that the form of $\exp_{\{\kappa\}}(t)$ can be obtained within the special relativity by using elementary algebraic calculations, starting from the dispersion relation of free relativistic particles.

In Sect. IV we reobtain the exponential $\exp_{\{\kappa\}}(t)$ starting from the concept of Lorentz invariant integration.

In Sect. V starting from the three-dimension κ -sum enforced by the Lorentz transformations we introduce the concept of generalized differential.

In Sect. VI we derive the forms of $\ln_{\{\kappa\}}(t)$ and $\exp_{\{\kappa\}}(t)$ within the special relativity, starting from the additivity law of relativistic momenta.

In Sect. VII we recall some concepts related to maximum entropy principle, in the context of a class of generalized statistical theories.

In Sect. VIII we consider the explicit form of the distribution function within the special relativity.

In Sect. IX we consider the main features of the statistical mechanics and of the kinetic theory based on the entropy (1.3).

In Sect. X we consider some applications of the present theory to specific physical problems and compare the results with the ones obtained within the classical statistical mechanics.

Finally in Sect. XI some concluding remarks are reported, while the main mathematical properties of the functions $\exp_{\{\kappa\}}(t)$ and $\ln_{\{\kappa\}}(t)$ are collected in Appendix.

II. DEFORMED SUMS IN SPECIAL RELATIVITY

In this section we introduce the special relativity in an alternative way, starting from the well known formulas

of hyperbolic trigonometry (pag. 58 of ref. [25])

$$\operatorname{arcsinh} x = \operatorname{arccosh} \sqrt{1+x^2} = \operatorname{arctanh} \frac{x}{\sqrt{1+x^2}} . \quad (2.1)$$

We pose $x = \kappa q$, being q the dimensionless variable corresponding to the momentum and κ a dimensionless deformation parameter. After introducing the functions

$$u(q) = \frac{q}{\sqrt{1+\kappa^2 q^2}} , \quad (2.2)$$

$$\mathcal{W}(q) = \frac{1}{\kappa^2} \sqrt{1+\kappa^2 q^2} - \frac{1}{\kappa^2} , \quad (2.3)$$

corresponding to the dimensionless velocity and kinetic energy respectively we can write Eq. (2.1) in the form

$$\operatorname{arcsinh}(\kappa q) = \operatorname{arccosh}(1 + \kappa^2 \mathcal{W}) = \operatorname{arctanh}(\kappa u) , \quad (2.4)$$

which in the classical limit $\kappa \rightarrow 0$ reproduces the dispersion relation $q = \sqrt{2\mathcal{W}} = u$ of Newton mechanics. Finally, after introducing the dimensionless total energy

$$\mathcal{E}(\mathcal{W}) = \mathcal{W} + \frac{1}{\kappa^2} , \quad (2.5)$$

one can write Eq. (2.1) also in the form

$$\operatorname{arcsinh}(\kappa q) = \operatorname{arccosh}(\kappa^2 \mathcal{E}) = \operatorname{arctanh}(\kappa u) . \quad (2.6)$$

It is immediate to verify the validity of the following formulas

$$q(u) = \frac{u}{\sqrt{1-\kappa^2 u^2}} , \quad (2.7)$$

$$q(\mathcal{W}) = \sqrt{2\mathcal{W} + \kappa^2 \mathcal{W}^2} , \quad (2.8)$$

$$q(\mathcal{E}) = \sqrt{\kappa^2 \mathcal{E}^2 - \frac{1}{\kappa^2}} , \quad (2.9)$$

$$u(\mathcal{W}) = \frac{\sqrt{2\mathcal{W} + \kappa^2 \mathcal{W}^2}}{1 + \kappa^2 \mathcal{W}} , \quad (2.10)$$

$$u(\mathcal{E}) = \frac{\sqrt{\kappa^2 \mathcal{E}^2 - 1/\kappa^2}}{\kappa^2 \mathcal{E}} , \quad (2.11)$$

$$\mathcal{W}(u) = \frac{1}{\kappa^2 \sqrt{1-\kappa^2 u^2}} - \frac{1}{\kappa^2} , \quad (2.12)$$

$$\mathcal{W}(\mathcal{E}) = \mathcal{E} - \frac{1}{\kappa^2} , \quad (2.13)$$

$$\mathcal{E}(q) = \frac{1}{\kappa^2} \sqrt{1+\kappa^2 q^2} , \quad (2.14)$$

$$\mathcal{E}(u) = \frac{1}{\kappa^2 \sqrt{1-\kappa^2 u^2}} . \quad (2.15)$$

At this point one can define the physical variables velocity v , momentum p and total energy E through

$$\frac{v}{u} = \frac{p}{mq} = \sqrt{\frac{E}{m\mathcal{E}}} = \kappa c = v_* , \quad (2.16)$$

and the kinetic energy as $W = E - mc^2$, being c the light speed. We impose that $\lim_{c \rightarrow \infty, \kappa \rightarrow 0} v_* < \infty$, in order

to preserve the validity of the definitions of the physical variable, also in the classical limit. After inserting the physical variables in the above obtained formulas, one recovers all the formulas of the one particle dynamics within the special relativity. For instance, Eq.s (2.7) and (2.14) transform

$$p = \frac{mv}{\sqrt{1-(v/c)^2}} , \quad (2.17)$$

$$E = \sqrt{m^2 c^4 + p^2 c^2} , \quad (2.18)$$

and so on.

Let us consider in the one-dimension frame \mathcal{S} two identical particles of rest mass m . We suppose that the first particle moves towards right with velocity u_1 while the second particle moves towards left with velocity u_2 . The momenta of the two particles are given by $q_1 = q(u_1)$ and $q_2 = q(u_2)$ respectively, where $q(u)$ is defined through Eq. (2.7). The total energies of the two particles are given by $\mathcal{E}_1 = \mathcal{E}(u_1)$ and $\mathcal{E}_2 = \mathcal{E}(u_2)$ respectively, with $\mathcal{E}(u)$ given by Eq. (2.15). Analogously the kinetic energies of the two particles are given by $\mathcal{W}_1 = \mathcal{W}(u_1)$ and $\mathcal{W}_2 = \mathcal{W}(u_2)$ respectively, with $\mathcal{W}(u)$ defined by Eq. (2.12).

We consider now the same particles in a new frame \mathcal{S}' which moves at constant speed u_2 towards left with respect to the frame \mathcal{S} . In this new frame, which is the rest frame of the second particle, the two particles have velocities given by $u'_1 = u_1 \stackrel{\kappa}{\oplus} u_2$ and $u'_2 = 0$ respectively, being

$$u_1 \stackrel{\kappa}{\oplus} u_2 = \frac{u_1 + u_2}{1 + \kappa^2 u_1 u_2} , \quad (2.19)$$

the well known relativistic additivity law for the dimensionless velocities. In the same frame \mathcal{S}' the particle momenta are given by $q'_1 = q(u'_1)$ and $q'_2 = 0$. Analogously in \mathcal{S}' , the particle kinetic energies are given by $\mathcal{W}'_1 = \mathcal{W}(u'_1)$ and $\mathcal{W}'_2 = 0$, while the total particle energies result to be $\mathcal{E}'_1 = \mathcal{E}(u'_1)$ and $\mathcal{E}'_2 = 1/\kappa^2$.

A very interesting result follows from the relation $\mathcal{E}'_2 = 1/\kappa^2$, regarding the physical meaning of the parameter κ . It is evident that the quantity $1/\kappa^2$ represents the dimensionless rest energy of the relativistic particle.

Let us ask now the following questions: is it possible to obtain the value of the momentum q'_1 (or of the total energy \mathcal{E}'_1 , or of the kinetic energy \mathcal{W}'_1) starting directly from the values of the momenta q_1 and q_2 (or of the total energies \mathcal{E}_1 and \mathcal{E}_2 , or of the kinetic energies \mathcal{W}_1 and \mathcal{W}_2) in the frame \mathcal{S} ? The answers to the above questions are affirmative as we will see in the following.

First we consider the case of the relativistic momentum q'_1 . Clearly we can write $q'_1 = q(u'_1) = q(u_1 \stackrel{\kappa}{\oplus} u_2)$. Then following ref. [19] we have

$$q(u_1 \stackrel{\kappa}{\oplus} u_2) = \frac{u_1 \stackrel{\kappa}{\oplus} u_2}{\sqrt{1 - \kappa^2 (u_1 \stackrel{\kappa}{\oplus} u_2)^2}}$$

$$\begin{aligned}
&= \frac{u_1 \oplus_{\kappa} u_2}{\sqrt{1 - \kappa^2 \left(\frac{u_1 + u_2}{1 + \kappa^2 u_1 u_2} \right)^2}} \\
&= (u_1 \oplus_{\kappa} u_2) \frac{1 + \kappa^2 u_1 u_2}{\sqrt{(1 - \kappa^2 u_1^2)(1 - \kappa^2 u_2^2)}} \\
&= \frac{u_1 + u_2}{\sqrt{(1 - \kappa^2 u_1^2)(1 - \kappa^2 u_2^2)}} \\
&= \frac{u_1}{\sqrt{1 - \kappa^2 u_1^2}} \sqrt{1 + \frac{\kappa^2 u_2^2}{1 - \kappa^2 u_2^2}} \\
&+ \frac{u_2}{\sqrt{1 - \kappa^2 u_2^2}} \sqrt{1 + \frac{\kappa^2 u_1^2}{1 - \kappa^2 u_1^2}} \\
&= q_1 \sqrt{1 + \kappa^2 q_2^2} + q_2 \sqrt{1 + \kappa^2 q_1^2} . \quad (2.20)
\end{aligned}$$

Hence we can write

$$q(u_1 \oplus_{\kappa} u_2) = q(u_1) \oplus_{\kappa} q(u_2) , \quad (2.21)$$

being

$$q_1 \oplus_{\kappa} q_2 = q_1 \sqrt{1 + \kappa^2 q_2^2} + q_2 \sqrt{1 + \kappa^2 q_1^2} , \quad (2.22)$$

the κ -sum of relativistic momenta introduced in ref. [18]. In words, the relativistic momentum q'_1 of the first particle, in the rest frame of the second particle \mathcal{S}' , is the κ -deformed sum of the momenta q_1 and q_2 of the two particles in the frame \mathcal{S} . The κ -sum of the relativistic momenta and the relativistic sum of the velocities are intimately related. They reduce to the standard sum as the velocity c approaches to infinity (or equivalently the parameter κ approaches to zero). The deformations in the above sums, in both cases, are relativistic effects and are originated from the fact that c has a finite value.

We proceed now to calculate the total energy \mathcal{E}'_1 of the first particle in the frame \mathcal{S}' starting directly from the values, in the frame \mathcal{S} , of the total energies \mathcal{E}_1 and \mathcal{E}_2 . Preliminarily we recall the following mathematical properties of the inverse hyperbolic trigonometric functions (pag. 58 of ref. [25])

$$\begin{aligned}
&\operatorname{arcsinh} x_1 \pm \operatorname{arcsinh} x_2 \\
&= \operatorname{arcsinh} \left(x_1 \sqrt{1 + x_2^2} \pm x_2 \sqrt{1 + x_1^2} \right) , \quad (2.23)
\end{aligned}$$

$$\begin{aligned}
&\operatorname{arccosh} y_1 \pm \operatorname{arccosh} y_2 \\
&= \operatorname{arccosh} \left(y_1 y_2 \pm \sqrt{(y_1^2 - 1)(y_2^2 - 1)} \right) , \quad (2.24)
\end{aligned}$$

$$\begin{aligned}
&\operatorname{arctanh} z_1 \pm \operatorname{arctanh} z_2 \\
&= \operatorname{arctanh} \frac{z_1 \pm z_2}{1 + z_1 z_2} . \quad (2.25)
\end{aligned}$$

After posing $x = \kappa q$, $y = \kappa^2 \mathcal{E}$, $z = \kappa u$ and taking into account Eq. (2.6), we are able to define the κ -sum of the relativistic total energies as follows

$$\mathcal{E}_1 \oplus_{\kappa} \mathcal{E}_2 = \kappa^2 \mathcal{E}_1 \mathcal{E}_2 + \frac{1}{\kappa^2} \sqrt{(\kappa^4 \mathcal{E}_1^2 - 1)(\kappa^4 \mathcal{E}_2^2 - 1)} . \quad (2.26)$$

After having noticed that

$$\begin{aligned}
\mathcal{E}(u_1) \oplus_{\kappa} \mathcal{E}(u_2) &= \frac{1}{\kappa^2} \frac{1}{\sqrt{1 - \kappa^2 u_1^2}} \oplus_{\kappa} \frac{1}{\kappa^2} \frac{1}{\sqrt{1 - \kappa^2 u_2^2}} \\
&= \frac{1}{\kappa^2} \frac{1}{\sqrt{(1 - \kappa^2 u_1^2)(1 - \kappa^2 u_2^2)}} \\
&+ \frac{1}{\kappa^2} \sqrt{\left(\frac{1}{1 - \kappa^2 u_1^2} - 1 \right) \left(\frac{1}{1 - \kappa^2 u_2^2} - 1 \right)} \\
&= \frac{1}{\kappa^2} \frac{1}{\sqrt{(1 - \kappa^2 u_1^2)(1 - \kappa^2 u_2^2)}} \\
&+ \frac{u_1 u_2}{\sqrt{(1 - \kappa^2 u_1^2)(1 - \kappa^2 u_2^2)}} \\
&= \frac{1}{\kappa^2} \frac{1 + \kappa^2 u_1 u_2}{\sqrt{(1 - \kappa^2 u_1^2)(1 - \kappa^2 u_2^2)}} \\
&= \frac{1}{\kappa^2} \frac{1 + \kappa^2 u_1 u_2}{\sqrt{(1 + \kappa^2 u_1 u_2)^2 - \kappa^2 (u_1 + u_2)^2}} \\
&= \frac{1}{\kappa^2} \frac{1}{\sqrt{1 - \kappa^2 \left(\frac{u_1 + u_2}{1 + \kappa^2 u_1 u_2} \right)^2}} \\
&= \frac{1}{\kappa^2} \frac{1}{\sqrt{1 - \kappa^2 (u_1 \oplus_{\kappa} u_2)^2}} \\
&= \mathcal{E}(u_1 \oplus_{\kappa} u_2) , \quad (2.27)
\end{aligned}$$

we can conclude that the total energy $\mathcal{E}'_1 = \mathcal{E}(u_1 \oplus_{\kappa} u_2)$ of the first particle, in the rest frame of the second particle \mathcal{S}' , is the κ -sum, defined through Eq. (2.26), of the total energies \mathcal{E}_1 and \mathcal{E}_2 of the particles in the frame \mathcal{S} :

$$\mathcal{E}(u_1 \oplus_{\kappa} u_2) = \mathcal{E}(u_1) \oplus_{\kappa} \mathcal{E}(u_2) . \quad (2.28)$$

The κ -sum for the relativistic kinetic energies can be defined starting from

$$\mathcal{W}_1 \oplus_{\kappa} \mathcal{W}_2 = \mathcal{E}_1 \oplus_{\kappa} \mathcal{E}_2 - \frac{1}{\kappa^2} , \quad (2.29)$$

and assumes the form

$$\begin{aligned}
\mathcal{W}_1 \oplus_{\kappa} \mathcal{W}_2 &= \mathcal{W}_1 + \mathcal{W}_2 + \kappa^2 \mathcal{W}_1 \mathcal{W}_2 \\
&+ \sqrt{\mathcal{W}_1 \mathcal{W}_2 (2 + \kappa^2 \mathcal{W}_1)(2 + \kappa^2 \mathcal{W}_2)} . \quad (2.30)
\end{aligned}$$

After taking into account Eqs. (2.12), (2.19) and (2.30) one immediately obtains

$$\mathcal{W}(u_1 \oplus_{\kappa} u_2) = \mathcal{W}(u_1) \oplus_{\kappa} \mathcal{W}(u_2) . \quad (2.31)$$

In words the κ -sum, defined through Eq. (2.30), of the particle kinetic energies \mathcal{W}_1 and \mathcal{W}_2 in the frame \mathcal{S} , gives the kinetic energy of one of the two particles in the rest frame \mathcal{S}' of the other particle.

Besides the relationships (2.21), (2.28) and (2.31), linking the κ -sums defined through Eqs. (2.19), (2.22), (2.26) and (2.30), one can easily obtain:

$$q(u_1) \overset{\kappa}{\oplus} q(u_2) = q \left(u_1 \overset{\kappa}{\oplus} u_2 \right) , \quad (2.32)$$

$$\mathcal{E}(u_1) \overset{\kappa}{\oplus} \mathcal{E}(u_2) = \mathcal{E} \left(u_1 \overset{\kappa}{\oplus} u_2 \right) , \quad (2.33)$$

$$\mathcal{W}(u_1) \overset{\kappa}{\oplus} \mathcal{W}(u_2) = \mathcal{W} \left(u_1 \overset{\kappa}{\oplus} u_2 \right) , \quad (2.34)$$

$$u(q_1) \overset{\kappa}{\oplus} u(q_2) = u \left(q_1 \overset{\kappa}{\oplus} q_2 \right) , \quad (2.35)$$

$$\mathcal{E}(q_1) \overset{\kappa}{\oplus} \mathcal{E}(q_2) = \mathcal{E} \left(q_1 \overset{\kappa}{\oplus} q_2 \right) , \quad (2.36)$$

$$\mathcal{W}(q_1) \overset{\kappa}{\oplus} \mathcal{W}(q_2) = \mathcal{W} \left(q_1 \overset{\kappa}{\oplus} q_2 \right) , \quad (2.37)$$

$$q(\mathcal{E}_1) \overset{\kappa}{\oplus} q(\mathcal{E}_2) = q \left(\mathcal{E}_1 \overset{\kappa}{\oplus} \mathcal{E}_2 \right) , \quad (2.38)$$

$$u(\mathcal{E}_1) \overset{\kappa}{\oplus} u(\mathcal{E}_2) = u \left(\mathcal{E}_1 \overset{\kappa}{\oplus} \mathcal{E}_2 \right) , \quad (2.39)$$

$$\mathcal{W}(\mathcal{E}_1) \overset{\kappa}{\oplus} \mathcal{W}(\mathcal{E}_2) = \mathcal{W} \left(\mathcal{E}_1 \overset{\kappa}{\oplus} \mathcal{E}_2 \right) , \quad (2.40)$$

$$q(\mathcal{W}_1) \overset{\kappa}{\oplus} q(\mathcal{W}_2) = q \left(\mathcal{W}_1 \overset{\kappa}{\oplus} \mathcal{W}_2 \right) , \quad (2.41)$$

$$u(\mathcal{W}_1) \overset{\kappa}{\oplus} u(\mathcal{W}_2) = u \left(\mathcal{W}_1 \overset{\kappa}{\oplus} \mathcal{W}_2 \right) , \quad (2.42)$$

$$\mathcal{E}(\mathcal{W}_1) \overset{\kappa}{\oplus} \mathcal{E}(\mathcal{W}_2) = \mathcal{E} \left(\mathcal{W}_1 \overset{\kappa}{\oplus} \mathcal{W}_2 \right) . \quad (2.43)$$

Let us make some remarks on the meaning of the κ -sums. We note that in the classical limit the κ -sum for the relativistic kinetic energies given by Eq. (2.30) reduces to the following expression of Galileian relativity

$$\mathcal{W}_1 \overset{0}{\oplus} \mathcal{W}_2 = \mathcal{W}_1 + \mathcal{W}_2 + 2\sqrt{\mathcal{W}_1 \mathcal{W}_2} . \quad (2.44)$$

Then a deformed sum appears also in classical physics when we change the frame of observation of the particle system. Clearly this happens only for the kinetic energy. For the classical momenta and velocities hold the ordinary sum.

We emphasize that the four κ -sums given by Eqs. (2.19), (2.22), (2.26) and (2.30), emerge only when we change the frame of observation of the particles from \mathcal{S} to \mathcal{S}' . More precisely these sums give the value of q'_1 (or u'_1 , \mathcal{E}'_1 , \mathcal{W}'_1) of the first particle in the rest frame \mathcal{S}' of the second particle, starting directly from the value of q_1 (or u_1 , \mathcal{E}_1 , \mathcal{W}_1) and q_2 (or u_2 , \mathcal{E}_2 , \mathcal{W}_2) of the two particles in the old frame \mathcal{S} .

III. MINKOWSKI FOUR-VECTORS AND κ -EXPONENTIAL

From the general discussion in the previous section appears clear that any relativistic formula can be viewed as a one-parameter deformation of the corresponding classical formula. For instance the kinetic energy given by

the relativistic dispersion relation

$$W = \sqrt{m^2 c^4 + p^2 c^2} - m c^2 . \quad (3.1)$$

can be viewed as the one-parameter relativistic deformation of the classical parabolic dispersion relation $W_{cl} = p^2/2m$. In the present section we will show that the relativistic dispersion relation imposes a one-parameter deformation to the ordinary exponential function, which presents now as a mathematical tool of classical physics. The new deformed exponential, as we will see in the following sections, replaces the ordinary exponential in the relativistic physics.

We adopt the metric $g^{\mu\nu} = \text{diag}(1, -1, -1, -1)$ in the Minkowski space and recall that the length of any four-vector is Lorentz invariant. In particular for the four-momentum $p^\mu = (E/c, \mathbf{p})$, $p_\mu = (E/c, -\mathbf{p})$ one obtains the relativistic dispersion relation

$$p^\mu p_\mu = m^2 c^2 , \quad (3.2)$$

which in terms of the total energy assumes the form

$$E = \sqrt{m^2 c^4 + p^2 c^2} . \quad (3.3)$$

We can write the latter equation as follows

$$\left(\frac{E}{m c^2} \right)^2 - \left(\frac{p}{m c} \right)^2 = 1 , \quad (3.4)$$

and then

$$\left(\frac{E}{m c^2} - \frac{p}{m c} \right) \left(\frac{E}{m c^2} + \frac{p}{m c} \right) = 1 . \quad (3.5)$$

At this point we can eliminate the variable E by employing once again Eq. (3.3) obtaining

$$\left(\sqrt{1 + \left(\frac{p}{m c} \right)^2} - \frac{p}{m c} \right) \left(\sqrt{1 + \left(\frac{p}{m c} \right)^2} + \frac{p}{m c} \right) = 1 . \quad (3.6)$$

The latter relationship, after introducing the dimensionless momentum q , defined through Eq.(2.16), can be written as

$$\left(\sqrt{1 + \kappa^2 q^2} - \kappa q \right) \left(\sqrt{1 + \kappa^2 q^2} + \kappa q \right) = 1 , \quad (3.7)$$

or equivalently

$$\left(\sqrt{1 + \kappa^2 q^2} - \kappa q \right)^{1/\kappa} \left(\sqrt{1 + \kappa^2 q^2} + \kappa q \right)^{1/\kappa} = 1 . \quad (3.8)$$

We remark that Eq. (3.8) follows directly from the dispersion relation. Interestingly in the classical limit $\kappa \rightarrow 0$ Eq. (3.8) reduces to $\exp(-q) \exp(q) = 1$ while the dispersion relation becomes the classical one $W = p^2/2m$. In this way we obtain a direct link between the dispersion relation of free classical particles and the ordinary exponential function.

In the light of the above result, we reconsider Eq. (3.8) which, after introducing the function

$$\exp_{\{\kappa\}}(q) = \left(\sqrt{1 + \kappa^2 q^2} + \kappa q \right)^{1/\kappa} , \quad (3.9)$$

writes in the form

$$\exp_{\{\kappa\}}(-q) \exp_{\{\kappa\}}(q) = 1 . \quad (3.10)$$

The function $\exp_{\{\kappa\}}(q)$, reproducing the exponential function, $\exp(q) = \exp_{\{0\}}(q)$ in the classical limit $\kappa \rightarrow 0$, represents a one-parameter relativistic generalization of the ordinary exponential.

IV. LORENTZ INVARIANT INTEGRATION AND κ -EXPONENTIAL

In this section we present a second path which unambiguously conducts to the $\exp_{\{\kappa\}}(q)$ function. Let us consider the following integral in the three dimension momentum space, within the classical physics framework

$$I_{cl} = \int d^3q F , \quad (4.1)$$

being F an arbitrary quantity. Whenever F depends only on $q = |\mathbf{q}|$, the above integral can be reduced to the following one dimension integral

$$I_{cl} = \int_0^\infty dq 4\pi q^2 F . \quad (4.2)$$

After introducing the physical variable $\mathbf{p} = mv_*^\infty \mathbf{q}$ the integral (4.1) transforms

$$I_{cl} = \int \frac{d^3p}{(mv_*^\infty)^3} F . \quad (4.3)$$

In the framework of a relativistic theory it is well known that the integral (4.3) must be replaced by a Lorentz invariant integral $I_{cl} \rightarrow I_{rel}$, being

$$I_{rel} = A \int d^4p \theta(p_0) \delta(p^\mu p_\mu - m^2 c^2) F , \quad (4.4)$$

and A a constant. Now we can introduce the variable $\mathbf{q} = \mathbf{p}/mv_*$ and take into account that $p^\mu = (E/c, \mathbf{p})$ and $E = \sqrt{m^2 c^4 + p^2 c^2}$. After properly choosing the value of A , the four-dimension integral (4.4) can be easily reduced to the following three dimension integral

$$I_{rel} = \int \frac{d^3q}{\sqrt{1 + \kappa^2 q^2}} F . \quad (4.5)$$

We remark that in (4.4) the integration element d^4p is a scalar because the Jacobian of the Lorentz transformation is equal to unity. Then I_{rel} transforms as F . For this reason in (4.5) the integration element $d^3q/\sqrt{1 + \kappa^2 q^2}$ is

a scalar. Finally we can reduce (4.5) in the following one dimension integral

$$I_{rel} = \int_0^\infty \frac{dq}{\sqrt{1 + \kappa^2 q^2}} 4\pi q^2 F . \quad (4.6)$$

It is important to note that in classical limit $\kappa \rightarrow 0$, Eqs. (4.5) and (4.6) reproduce the corresponding classical ones given by (4.1) and (4.2) respectively.

We focus now our attention to the classical and relativistic expression of the one dimension integrals given by (4.2) and (4.6) respectively. One immediately observes that the relativistic integral is obtained directly from the classical one, by making the substitution $dq \rightarrow d_{\{\kappa\}}q$, being

$$d_{\{\kappa\}}q = \frac{dq}{\sqrt{1 + \kappa^2 q^2}} , \quad (4.7)$$

the κ -differential. The relativistic one dimension integral can be written in the form

$$I_{rel} = \int_0^\infty d_{\{\kappa\}}q 4\pi q^2 F . \quad (4.8)$$

Clearly the κ -integral $\int d_{\{\kappa\}}q$ is originated from the fact that the light speed has a finite value.

As a working example, involving explicitly the κ -integration, we consider the definition of the kinetic energy. In classical mechanics the kinetic energy, using dimensionless variables, is defined as

$$\mathcal{W}_{cl}(q) = \int_0^q dq q . \quad (4.9)$$

It is straightforward to verify that the extension of the above definition in special relativity is given by

$$\mathcal{W}(q) = \int_0^q d_{\{\kappa\}}q q . \quad (4.10)$$

Then, the replacement of the ordinary integration with by the κ -integration, in the classical definition (4.9), permits us to recover the relativistic expression (2.3) of the kinetic energy.

Let us consider now the κ -derivative related to the above defined κ -integration

$$\frac{d}{d_{\{\kappa\}}q} = \sqrt{1 + \kappa^2 q^2} \frac{d}{dq} . \quad (4.11)$$

Eq. (4.10) linking \mathcal{W} and q can be written now also in the following differential form

$$\frac{d}{d_{\{\kappa\}}q} \mathcal{W}(q) = q , \quad (4.12)$$

which, after integration with the condition $\mathcal{W}(0) = 0$, yields the relativistic expression of the kinetic energy (2.3). In the $\kappa \rightarrow 0$ limit the above differential equation reduces to the classical one $(d/dq)\mathcal{W}(q) = q$.

Taking into account that the ordinary exponential results to be eigenfunction of the ordinary derivative, namely $(d/dq)\exp(q) = \exp(q)$, the question to determine the eigenfunction of the κ -derivative, naturally arises. After some simple calculations one obtains

$$\frac{d}{d_{\{\kappa\}}q} \exp_{\{\kappa\}}(q) = \exp_{\{\kappa\}}(q) \quad , \quad (4.13)$$

so that the κ -exponential arises as eigenfunction of the κ -derivative.

V. RELATIVISTIC SUMS AND κ -DIFFERENTIAL

In this section we explain the physical origin of the mechanism enforcing the deformation in the κ -differential. We consider two identical relativistic particles with rest mass m and velocities \mathbf{v}_1 and \mathbf{v}_2 in the three-dimension frame S . The modulus of the relative velocity V of the particles depends on \mathbf{v}_1 and \mathbf{v}_2 and is given by (page 20 of ref. [26])

$$V(\mathbf{v}_1, \mathbf{v}_2) = \sqrt{(\mathbf{v}_1 \ominus \mathbf{v}_2)^2 - \frac{1}{c^2} \left(\frac{\mathbf{v}_1 \times \mathbf{v}_2}{1 - \mathbf{v}_1 \mathbf{v}_2 / c^2} \right)^2} \quad , \quad (5.1)$$

being

$$\mathbf{v}_1 \ominus \mathbf{v}_2 = \frac{\mathbf{v}_1 - \mathbf{v}_2}{1 - \mathbf{v}_1 \mathbf{v}_2 / c^2} \quad . \quad (5.2)$$

We perform now the calculation of the modulus of relative momentum related to the modulus of the relative velocity of the two particles according to the relativistic formula

$$P(V) = \frac{mV}{\sqrt{1 - V^2/c^2}} \quad . \quad (5.3)$$

Clearly results that $P(V) = P(\mathbf{v}_1, \mathbf{v}_2)$ and after taking into account that

$$\mathbf{v} = \frac{\mathbf{p}/m}{\sqrt{1 + \mathbf{p}^2/m^2c^2}} \quad , \quad (5.4)$$

we can conclude that $P(V) = P(\mathbf{p}_1, \mathbf{p}_1)$. At this point we introduce the dimensionless variables $\mathbf{q} = \mathbf{p}/mv_*$ and $Q = P/mv_*$. After tedious but straightforward calculation, one arrives to the following expression for the modulus of the dimensionless relative momentum

$$Q(\mathbf{q}_1, \mathbf{q}_2) = \sqrt{\frac{\left(\mathbf{q}_1 \ominus_{\kappa} \mathbf{q}_2 \right)^2 - \kappa^2 (\mathbf{q}_1 \times \mathbf{q}_2)^2}{1 - \kappa^4 (\mathbf{q}_1 \times \mathbf{q}_2)^2}} \quad , \quad (5.5)$$

being

$$\mathbf{q}_1 \ominus_{\kappa} \mathbf{q}_2 = \mathbf{q}_1 \sqrt{1 + \kappa^2 \mathbf{q}_2^2} - \mathbf{q}_2 \sqrt{1 + \kappa^2 \mathbf{q}_1^2} \quad , \quad (5.6)$$

the κ -difference introduced in ref. [18]. Clearly $Q(\mathbf{q}_1, \mathbf{q}_2)$ represents the modulus of momentum of the first (second) particle in the rest frame of second (first) particle.

We suppose now that the two particles in the frame S have momenta $\mathbf{q}_1 = \mathbf{q} + d\mathbf{q}$ and $\mathbf{q}_2 = \mathbf{q}$ respectively and calculate the value of $Q(\mathbf{q} + d\mathbf{q}, \mathbf{q})$. One easily obtains

$$Q(\mathbf{q} + d\mathbf{q}, \mathbf{q}) = d_{\{\kappa\}}q \quad , \quad (5.7)$$

being $q = |\mathbf{q}|$ and $d_{\{\kappa\}}q$ given by Eq. (4.7).

The physical meaning of κ -differential $d_{\{\kappa\}}q$ immediately follows. The modulus of the infinitesimal difference of the two particle momenta dq , in the frame S , becomes $d_{\{\kappa\}}q$, if this difference is observed in the rest frame of one of the two particles. The meaning of the κ -derivative follows also readily. If d/dq is the derivative in the frame S , the deformed derivative $d/d_{\{\kappa\}}q$ represents an ordinary derivative in the rest frame of one of the two particles.

VI. RELATIVISTIC SUMS AND κ -DEFORMED LOGARITHM AND EXPONENTIAL

The ordinary logarithm $h(x) = \ln(x)$ is the only existing function (unless a multiplicative constant) which results to be solution of the function equation $h(x_1 x_2) = h(x_1) + h(x_2)$. Let us consider now the generalization of this equation within the special relativity obtained by substituting the ordinary sum by the κ -sum of the dimensionless relativistic momenta

$$h(x_1 x_2) = h(x_1) \oplus_{\kappa} h(x_2) \quad . \quad (6.1)$$

We proceed by solving this equation, which assumes the explicit form

$$h(x_1 x_2) = h(x_1) \sqrt{1 + \kappa^2 h(x_2)^2} + h(x_2) \sqrt{1 + \kappa^2 h(x_1)^2} \quad . \quad (6.2)$$

After performing the substitution $h(x) = \kappa^{-1} \sinh \kappa g(x)$ we obtain that the auxiliary function $g(x)$ obeys the equation $g(x_1 x_2) = g(x_1) + g(x_2)$, and then is given by $g(x) = A \ln x$. The unknown function $h(x)$ becomes

$$h(x) = \frac{\sinh(\kappa \ln x)}{\kappa} \quad , \quad (6.3)$$

where we have set $A = 1$ in order to recover, in the limit $\kappa \rightarrow 0$, the classical solution $h(x) = \ln(x)$. The function given by Eq. (6.3) in the following is denoted by $\ln_{\{\kappa\}}(x)$ and can be written also in the form

$$\ln_{\{\kappa\}}(x) = \frac{x^{\kappa} - x^{-\kappa}}{2\kappa} \quad . \quad (6.4)$$

Holding $\exp_{\{\kappa\}}(\ln_{\{\kappa\}}(x)) = \ln_{\{\kappa\}}(\exp_{\{\kappa\}}(x)) = x$ it results that $\ln_{\{\kappa\}}(x)$ is the inverse function of $\exp_{\{\kappa\}}(x)$ and thus can be viewed as a generalization of the ordinary logarithm, $\ln(x) = \ln_{\{0\}}(x)$, in the framework of special relativity.

It is straightforward to obtain the $\exp_{\{\kappa\}}(x)$, starting directly from κ -sum of the dimensionless relativistic momenta, as solution of the functional equation

$$\exp_{\{\kappa\}}(x_1) \exp_{\{\kappa\}}(x_2) = \exp_{\{\kappa\}}(x_1 \oplus_{\kappa} x_2) , \quad (6.5)$$

which in the classical limit reduces to the functional equation $\exp(x_1) \exp(x_2) = \exp(x_1 + x_2)$ defining the ordinary exponential.

In the Appendix, the main mathematical properties of the functions κ -exponential and κ -logarithm are reported.

VII. MAXIMUM ENTROPY PRINCIPLE

We start by recalling some elements of classical statistical mechanics. We consider the Boltzmann-Gibbs-Shannon entropy

$$S = - \int d^3p f(p) \ln(f(p)) , \quad (7.1)$$

and the constraints functional

$$C = a_0 \left[\int d^3p f(p) - 1 \right] + \mathbf{a} \left[\mathbf{M} - \int d^3p \mathbf{g}(p) f(p) \right] , \quad (7.2)$$

where the constants a_0 and $\mathbf{a} = \{a_1, a_2, \dots, a_l\}$ are the $l + 1$ Lagrange multipliers, while the l moments $\mathbf{M} = \{M_1, M_2, \dots, M_l\}$ are the mean values of the l functions $\mathbf{g}(p) = \{g_1(p), g_2(p), \dots, g_l(p)\}$. The variational equation

$$\frac{\delta}{\delta f} (S + C) = 0 , \quad (7.3)$$

means that the entropy $S(p)$ is maximized under the constraints imposing the conservation of the norm of $f(p)$ and the a priori knowledge of the values of the l moments M_j of $f(p)$, namely

$$\int d^3p f(p) = 1 , \quad M_j = \int d^3p g_j(p) f(p) . \quad (7.4)$$

The maximum entropy principle expressed by the variational equation (7.3) yields the following statistical distribution

$$f(p) = \frac{1}{e} \exp(-\mathbf{a} \cdot \mathbf{g}(p) + a_0) . \quad (7.5)$$

In a classical many body particle system the collisional invariants are essentially the particle number and the kinetic energy $W(p) = p^2/2m$. Then we need to involve two Lagrange multipliers $a_0 = \mu/k_B T$, $a_1 = 1/k_B T$ and one moment $M_1 = \int d^3p g_1(p) f(p) = \int d^3p W(p) f(p) = \langle W \rangle$. In this way the distribution function given by Eq. (7.5) reduces to the Maxwell-Boltzmann one of classical statistical mechanics

$$f(p) = \frac{1}{e} \exp\left(-\frac{W(p) - \mu}{k_B T}\right) . \quad (7.6)$$

Let us consider now the following generalized trace form entropy

$$S = - \int d^3p f(p) \Lambda(f(p)) , \quad (7.7)$$

being $\Lambda(f)$ a function generalizing the ordinary logarithm. The variational equation (7.3) with S and C given by (7.7) and (7.2) respectively, produces the distribution function $f = f(p)$ defined through

$$\frac{\partial}{\partial f} f \Lambda(f) = -\mathbf{a} \cdot \mathbf{g}(p) + a_0 . \quad (7.8)$$

The maximum entropy principle, in the case of the ordinary statistical mechanics, relates the Boltzmann-Gibbs-Shannon entropy and the Maxwell-Boltzmann distribution by means of a twofold link. Firstly, the distribution is obtained by maximizing the entropy under proper constraints. Secondly, both the entropy and the distribution are expressed in terms of the same function in direct and inverse form. Indeed, the Maxwell-Boltzmann distribution is given in terms of the ordinary exponential while the Boltzmann-Gibbs-Shannon entropy is defined starting from the ordinary logarithm. Clearly, this is a very strong interpretation of the maximum entropy principle. In generalized statistical theories, customary, is used a more weak interpretation of the maximum entropy principle by relaxing the second link.

Our next task is to use the strong interpretation of the maximum entropy principle holding for the ordinary statistical mechanics in order to generalize minimally it. The requirement of the twofold link between entropy and distribution function also in generalized statistical theories imposes that the function $\Lambda(f)$ is a solution of the following functional-differential equation

$$\frac{\partial}{\partial f} f \Lambda(f) = \lambda \Lambda(f/\alpha) + \eta , \quad (7.9)$$

being $\{\alpha, \lambda, \eta\}$ three real constants. Indeed, the distribution given by (7.8) assumes now the form

$$f(p) = \alpha \Lambda^{-1}\left(-\frac{\mathbf{a} \cdot \mathbf{g}(p) - a_0 + \eta}{\lambda}\right) , \quad (7.10)$$

being $\Lambda^{-1}(x)$ the inverse function of $\Lambda(x)$. Note that the constants λ and η introduce a scaling in the Lagrange multipliers. Clearly in the classical limit it must result: $\Lambda(x) \rightarrow \ln(x)$, $\Lambda^{-1}(x) \rightarrow \exp(x)$ and $\{\alpha, \lambda, \eta\} \rightarrow \{1/e, 1, 0\}$.

The problem related to the determination of the generalized logarithm $\Lambda(f)$ satisfying Eq. (7.9) and the conditions $\Lambda(1) = 0$, $\Lambda'(1) = 1$ and $0\Lambda(0) = 0$, can be solved easily. Beside the ordinary logarithm, Eq. (7.9) admits other new solutions. The general solution of this equation defines a three-parameter family of generalized logarithms indicated by $\Lambda(f) = \ln_{\{\kappa, r, \varsigma\}}(f)$ being

$$\ln_{\{\kappa, r, \varsigma\}}(f) = \frac{\varsigma^\kappa f^{r+\kappa} - \varsigma^{-\kappa} f^{r-\kappa} - \varsigma^\kappa + \varsigma^{-\kappa}}{(\kappa + r)\varsigma^\kappa + (\kappa - r)\varsigma^{-\kappa}} , \quad (7.11)$$

while $\{\kappa, r, \varsigma\}$ are three free parameters. The constants α , λ and η are given by

$$\alpha = \left(\frac{1+r-\kappa}{1+r+\kappa} \right)^{\frac{1}{2\kappa}}, \quad (7.12)$$

$$\lambda = \frac{(1+r-\kappa)^{\frac{r+\kappa}{2\kappa}}}{(1+r+\kappa)^{\frac{r-\kappa}{2\kappa}}}, \quad (7.13)$$

$$\eta = (\lambda - 1) \frac{\varsigma^\kappa - \varsigma^{-\kappa}}{(\kappa+r)\varsigma^\kappa + (\kappa-r)\varsigma^{-\kappa}}. \quad (7.14)$$

Some specific choice for the parameters $\{\kappa, r, \varsigma\}$, generating particular expressions for the function $\Lambda(f)$ and for its inverse function $\Lambda^{-1}(x) = \exp_{\{\kappa, r, \varsigma\}}(x)$, have been already considered in mathematical statistics, information theory and statistical mechanics [20]. As stressed previously, in ref.s [18, 19, 20, 21, 22, 23], it is shown that it is possible to develop selfconsistent statistical theories starting from any generalized entropy satisfying some standard requirements. Fortunately, all these requirements are satisfied by the entropy defined through (7.7) with $\Lambda(f) = \ln_{\{\kappa, r, \varsigma\}}(f)$.

Let us focus now our attention to the solution (7.11) corresponding to the particular choice $\{\kappa, r = 0, \varsigma = 1\}$ [18]. It is easy to verify that results $\ln_{\{\kappa, 0, 1\}}(x) = \ln_{\{\kappa\}}(x)$ and $\exp_{\{\kappa, 0, 1\}}(x) = \exp_{\{\kappa\}}(x)$ with

$$\ln_{\{\kappa\}}(x) = \frac{x^\kappa - x^{-\kappa}}{2\kappa}, \quad (7.15)$$

$$\exp_{\{\kappa\}}(x) = \left(\sqrt{1 + \kappa^2 x^2} + \kappa x \right)^{1/\kappa}. \quad (7.16)$$

Thus the κ -exponential and κ -logarithm, emerging in the special relativity in the place of the ordinary exponential and logarithm, reappear also in the statistical theory through the maximum entropy principle. It is immediate to verify that the reality and positivity of the parameters α and λ impose that $-1 < \kappa < 1$.

Furthermore, in the case $\{\kappa, 0, \varsigma\}$, Eq. (7.11) produces a two parameter generalized logarithm and exponential which results to be the scaled κ -logarithm and the scaled κ -exponential respectively, defined as

$$\ln_{\{\kappa, \varsigma\}}(x) = \frac{\ln_{\{\kappa\}}(\varsigma x) - \ln_{\{\kappa\}} \varsigma}{\sqrt{1 + \kappa^2 \ln_{\{\kappa\}}^2 \varsigma}}, \quad (7.17)$$

$$\exp_{\{\kappa, \varsigma\}}(x) = \frac{1}{\varsigma} \exp_{\{\kappa\}} \left(x \sqrt{1 + \kappa^2 \ln_{\{\kappa\}}^2 \varsigma} + \ln_{\{\kappa\}} \varsigma \right). \quad (7.18)$$

We remark that the functions $\exp_{\{\kappa\}}(x)$ and $\ln_{\{\kappa\}}(x)$ generalizing the ordinary exponential and logarithm respectively, emerge both in special relativity and independently in many body physics, through the maximum entropy principle, the cornerstone of statistical mechanics. This interesting fact suggests us to consider with a particular attention the statistical theories, enforced by the

entropies related to the generalized logarithms defined through (7.15) and (7.17).

In the next section we consider the statistical theory involving the generalized logarithm (7.15).

VIII. GENERALIZED DISTRIBUTION FUNCTION

Let us consider the entropy $S = - \langle \ln_{\{\kappa\}} f \rangle$ related to the generalized logarithm defined through (7.15). Here the distribution function $f = f(x, p)$ depends on the four-vectors position x and momentum p . The explicit form of this entropy is given by

$$S = - \int d^3 p f \ln_{\{\kappa\}}(f), \quad (8.1)$$

and after maximized under the constrains (7.2) by solving the variational equation

$$\frac{\delta}{\delta f}(S + C) = 0, \quad (8.2)$$

yields the distribution function

$$f = \alpha \exp_{\{\kappa\}} \left(- \frac{\mathbf{a} \cdot \mathbf{g} - a_0}{\lambda} \right), \quad (8.3)$$

with $\mathbf{g} = \mathbf{g}(x, p)$ and

$$\alpha = \left(\frac{1-\kappa}{1+\kappa} \right)^{1/2\kappa}, \quad (8.4)$$

$$\lambda = \sqrt{1 - \kappa^2}, \quad (8.5)$$

being $|\kappa| < 1$. Before to proceed further we explain briefly the meaning of the constants α and λ , related by $1/\lambda = \ln_{\{\kappa\}}(1/\alpha)$. We write the entropy in the form $S = \int_{\mathcal{R}} d^3 p \sigma(f)$ where $\sigma(f) = -f \ln_{\{\kappa\}}(f)$ is the entropy density. It is easy to verify that for $f = \alpha$ the entropy density assumes its maximum value, which is given by $\sigma_{max} = \alpha/\lambda$.

We focus now our attention on the form of $\mathbf{g} = \mathbf{g}(x, p)$. In ref. [27] it is shown that for a relativistic many body system, in presence of external electromagnetic fields, the more general microscopic invariant has the form $(p^\nu + qA^\nu/c) U_\nu + \text{constant}$, being U_ν the hydrodynamic four-vector velocity with $U^\nu U_\nu = c^2$. Then we can pose $g_1 = (p^\nu + qA^\nu/c) U_\nu - mc^2$, $a_1 = 1/k_B T$ and $a_0 = \mu/k_B T$. The distribution (8.3), in the case of relativistic statistical systems, assumes the form

$$f = \alpha \exp_{\{\kappa\}} \left(- \frac{(p^\nu + eA^\nu/c) U_\nu - mc^2 - \mu}{\lambda k_B T} \right), \quad (8.6)$$

and results to be quite different with respect the Juttner distribution of the ordinary relativistic statistical mechanics, where in place of κ -exponential appears the ordinary exponential [27]. The distribution (8.6), in the

global rest frame where $U^\nu = (c, 0, 0, 0)$ and in absence of external forces ($A^\nu = 0$), simplifies as

$$f = \alpha \exp_{\{\kappa\}} \left(-\frac{W - \mu}{\lambda k_B T} \right) , \quad (8.7)$$

being W the relativistic kinetic energy

$$W = \sqrt{m^2 c^4 + \mathbf{p}^2 c^2} - mc^2 , \quad (8.8)$$

and in the $\kappa \rightarrow 0$ limit reduces to the classical distribution (7.6). It is remarkable that the above distribution (8.7) presents a power law asymptotic behavior, namely

$$f \propto W^{-1/\kappa} , \quad (W \rightarrow \infty) , \quad (8.9)$$

in contrast with the ordinary relativistic distribution (Maxwell-Boltzmann distribution where the energy is given by its relativistic expression) which decays exponentially.

IX. GENERALIZED STATISTICS AND KINETICS

In this section we report the main features of statistical theory, based on the entropy (8.1), some of which are already known in the literature.

Equiprobability: The non negative entropy (8.1) achieves its maximum value at equiprobability, $f(p) = 1/\Omega$ for $\forall p$, and this value is $S = \ln_{\{\kappa\}} \Omega$.

Thermodynamic stability: The entropy (8.1) is concave

$$S[tf_1 + (1-t)f_2] \geq tS[f_1] + (1-t)S[f_2] , \quad (9.1)$$

so that the system, in thermodynamic equilibrium, is stable [19].

Lesche stability: The Lesche stability condition is satisfied by any physically meaningful quantity. In particular it is a necessary condition to experimentally detect a physical observable. The entropy given by (8.1), being a physical quantity depending on a probability distribution, should exhibit a small relative error

$$R = \left| \frac{S[f] - S[f']}{\sup(S[f])} \right| , \quad (9.2)$$

with respect to small changes of the probability distribution

$$D = \|f - f'\| . \quad (9.3)$$

Mathematically this implies that: For any $\varepsilon > 0$ there exists $\delta > 0$ such that $R \leq \varepsilon$ holds for all distribution functions satisfying $D \leq \delta$. The Lesche stability condition holds for the Boltzmann-Gibbs-Shannon entropy and is proved also for the entropy (8.1) in ref. [28]. In the thermodynamic limit it is obtained $\delta = \varepsilon^{1/(1-|\kappa|)}$.

Relativistic kinetic equation: The statistical distribution (8.7) can be viewed as stationary case of a distribution function $f = f(x, p)$, describing a relativistic many

body system, governed by the generalized Boltzmann evolution equation [19]

$$p^\nu \partial_\nu f - mF^\nu \frac{\partial f}{\partial p^\nu} = \int \frac{d^3 p'}{p'^0} \frac{d^3 p_1}{p_1^0} \frac{d^3 p'_1}{p_1'^0} G \times \left[a(f' \otimes_{\kappa} f'_1) - a(f \otimes_{\kappa} f_1) \right] . \quad (9.4)$$

In the latter equation the standard notations of the ordinary Boltzmann kinetics are used and $a(f)$ is a positive and increasing arbitrary function, which does not affect the stationary form of f . The factor G is the transition rate, depending only on the nature of the two body particle interaction. The composition law \otimes_{κ} defined through

$$\ln_{\{\kappa\}}(f \otimes_{\kappa} h) = \ln_{\{\kappa\}} f + \ln_{\{\kappa\}} h , \quad (9.5)$$

is a one-parameter generalization of the ordinary product having the following properties

- i) associative law: $(f \otimes_{\kappa} h) \otimes_{\kappa} w = f \otimes_{\kappa} (h \otimes_{\kappa} w)$,
- ii) neutral element: $f \otimes_{\kappa} 1 = 1 \otimes_{\kappa} f = f$,
- iii) inverse element: $f \otimes_{\kappa} (1/f) = (1/f) \otimes_{\kappa} f = 1$,
- iv) commutative law: $f \otimes_{\kappa} h = h \otimes_{\kappa} f$.

Furthermore it results $f \otimes_{\kappa} 0 = 0 \otimes_{\kappa} f = 0$ and the generalized division \oslash_{κ} can be defined through $f \oslash_{\kappa} h = f \otimes_{\kappa} (1/h)$.

H-theorem: The four-vector entropy $S^\nu = (S^0, \mathbf{S})$ it was defined as

$$S^\nu = - \int \frac{d^3 p}{p^0} p^\nu f \ln_{\{\kappa\}} f . \quad (9.6)$$

The temporal component S^0 of this four-vector coincides with the κ -entropy defined previously through Eq. (8.1) while the spatial component \mathbf{S} is the entropy flow. Starting from the evolution equation (9.4) it has been demonstrated the H-theorem, which represents the second law of thermodynamics and states that the entropy production is never negative, and, in equilibrium conditions, there is no entropy production. Namely, the following relation was obtained

$$\partial_\nu S^\nu \geq 0 , \quad (9.7)$$

which is the local formulation of the relativistic H-theorem and represents the second law of the thermodynamics [19].

Entropy transformation: After recalling the identity $d^3 p/p^0 = d^4 p \, 2\theta(p^0) \delta(p^\mu p_\mu - m^2 c^2)$ the definition (9.6) assumes the form

$$S^\nu = - \int d^4 p \, 2\theta(p^0) \delta(p^\mu p_\mu - m^2 c^2) p^\nu f \ln_{\{\kappa\}} f . \quad (9.8)$$

In this expression $d^4 p$ is a scalar because the Jacobian of the Lorentz transformation is equal to unit. Then since p^ν transforms as a four-vector, we can conclude that S^ν transforms as a four-vector like the particle four-flow

$$N^\nu = \int d^4 p \, 2\theta(p^0) \delta(p^\mu p_\mu - m^2 c^2) p^\nu f , \quad (9.9)$$

for which the conservation law of the total particle number imposes

$$\partial_\nu N^\nu = 0 \quad . \quad (9.10)$$

Calculation of the parameter κ : We focus now attention to the general discussion of sec. II. The results obtained in sect. II permit us to consider the problem related to the calculation of the parameter κ from a very different and more physically sound point of view with respect the one of ref. [19]. Clearly, the velocity v_* introduced through $|\kappa| = v_*/c$ does not emerge within the one-particle relativistic theory. In general v_* could depend on c namely $v_* = v_*(c)$ with the condition that $v_*(\infty) < \infty$. The fact that the condition $|\kappa| < 1$ and then $v_* < c$, follows from the maximum entropy principle, supports the supposition that v_* could emerge in the framework of the many body theory. Taking in mind that in statistical mechanics the only emerging velocity is the thermal velocity, in the following we identify v_* with the thermal velocity of the many body system.

The particle thermal energy, being the mean particle kinetic energy $\langle W \rangle$, is related to the thermal velocity $v_* = \kappa c$ according to the relativistic relationship

$$\langle W \rangle = mc^2 \left(\frac{1}{\sqrt{1 - \kappa^2}} - 1 \right) \quad . \quad (9.11)$$

Clearly, in a statistical theory, the mean particle kinetic energy $\langle W \rangle$ is considered a known quantity. Then the expression of κ in terms of $\langle W \rangle$ follows immediately

$$\kappa^2 = 1 - \left(1 + \frac{\langle W \rangle}{mc^2} \right)^{-2} \quad , \quad (9.12)$$

so that the theory does not contain free parameters and in the classical limit $c \rightarrow \infty$, results $\kappa = 0$. From (9.12) follows that $\lim_{c \rightarrow \infty} \kappa c = \sqrt{2 \langle W \rangle / m} < \infty$ as required from the theory developed in sect. II. Note that in the case we know experimentally the distribution function, the value of κ and consequently, by using (9.12), the value of the ratio $\langle W \rangle / mc^2$ can be obtained from the slope of the distribution tails in a log-log plot. The normalization condition for the distribution function f and the definition of the particle mean kinetic energy, starting from f , permit us to determine univocally T and μ . In this way one can link $\langle W \rangle$ and then κ to the temperature of the system.

Bearing in mind the meaning of the parameter κ we rewrite the distribution function (8.7) in the form

$$f = \alpha \exp_{\{\kappa\}} \left(- \frac{W - \mu}{k_B T'} \right) \quad , \quad (9.13)$$

being W the relativistic kinetic energy while the temperature T' is given by

$$T' = \sqrt{1 - \frac{v_*^2}{c^2}} T \quad . \quad (9.14)$$

We remark that the temperature T' is contracted with respect to the temperature T by a factor which, curiously, has the same form of the one entering in the definition of the relativistic temperature suggested by Einstein [29] and Planck [30].

X. APPLICATIONS

In the last four years the present theory has been used successfully to study some systems which manifestly cannot be treated within the ordinary statistical mechanics. Clearly, the most natural applications regard relativistic many body systems.

The first application, considered in ref. [19], concerns the cosmic rays. Since long time it is known that the cosmic rays spectrum, which extends over 13 decades in energy, from a few hundred of MeV (10^8 eV) to a few hundred of EeV (10^{20} eV) and spans 33 decades in particle flux, from 10^4 to 10^{-29} [m^2 s r s GeV] $^{-1}$, is not exponential and, then, it violates the Boltzmann equilibrium statistical distribution $\propto \exp(-E/k_B T)$ [11, 12, 13]. On the other hand, it is known that the particles composing the cosmic rays are essentially the normal nuclei as in the standard cosmic abundance of matter. Then, the cosmic rays can be viewed as an equivalent statistical system of identical relativistic particles with masses near the mass of the proton (938 MeV). These characteristics (relativistic particles with a very large extension both for their flux and energy) yield the cosmic rays spectrum an ideal physical system for a preliminary test of the correctness and predictability of any relativistic theory. In ref. [19], a high quality agreement between the predictions of the present theory and the observed data, in the whole cosmic ray spectrum, has been found. This agreement over so many decades is quite remarkable.

Other physical applications of the theory regard the formation of quark-gluon plasma [31] and kinetic models describing a gas of interacting atoms and photons [32].

On the other hand, the theory has been applied successfully also to the study of natural or artificial systems exhibiting a limiting velocity in the propagation of the information (like the one imposed by light speed in physics) where a mechanism analogous to the relativistic one can emerge, deforming the distribution function. In ref. [33] the problem of the fracture propagation has been considered within the present theory. Others applications have been considered in games theory [34], in economy for study the income distribution [35] and in construct financial models [36], etc.

We come back to the high energy physics being the more proper field of application of the theory. In the following, as further application, we consider the open problem of the black hole physics regarding the intriguing question of the Bekenstein-Hawking area law [14, 15]. This law asserts that the entropy of a black hole scales as the area A of the event horizon namely $S \propto A$. It is well known that the ordinary Boltzmann statistical mechanics

and thermodynamics fails to justify this law and in contrary predicts that the entropy scales as the volume V of spatial region delimited by the event horizon, namely $S \propto V$.

Indeed, if we indicate with N the degrees of freedom or microscopic components in the volume V and with M the number of states accessible by each component, within the Boltzmann statistics the number of microstates Ω of the whole system is given by $\Omega \propto M^N$ with $N \gg 1$. After introducing the notations

$$n(0) = \exp N \quad , \quad m(0) = \ln M \quad , \quad (10.1)$$

the number of microstates can be written also in the form

$$\Omega \propto n(0)^{m(0)} \quad . \quad (10.2)$$

Customary one assumes a uniform distribution of the components in the volume V obtaining $N \propto V$. Thus the entropy

$$S = \ln \Omega \quad , \quad (10.3)$$

of the black hole results to be given by

$$S \propto V \quad . \quad (10.4)$$

This result of the Boltzmann statistics, already known in the literature, is manifestly in contradiction with the Bekenstein-Hawking area law.

Very recently, in ref. [37], has been suggested to treat this problem within non-standard statistics. Even if the authors limit their discussion to the nonextensive [38] aspects of the problem the proposed approach has a general validity and can be adopted also in the present case.

Clearly the black hole is a relativistic system and appears natural to write its entropy in the microcanonical picture as

$$S = \ln_{\{\kappa\}} \Omega \quad . \quad (10.5)$$

Furthermore the present formalism generalizes Eqs. (10.1) and (10.2) as follows

$$n(\kappa) = \exp_{\{\kappa\}} N \quad , \quad m(\kappa) = \ln_{\{\kappa\}} M \quad , \quad (10.6)$$

$$\Omega \propto n(\kappa)^{m(\kappa)} \quad . \quad (10.7)$$

After recalling the property (12.15) the number of microstates becomes

$$\Omega \propto \exp_{\{\kappa/m(\kappa)\}} (m(\kappa)N) \quad , \quad (10.8)$$

while the black hole entropy can be written as

$$S \propto \ln_{\{\kappa\}} \left[\exp_{\{\kappa/m(\kappa)\}} (m(\kappa)N) \right] \quad . \quad (10.9)$$

In the thermodynamic limit $N \gg 1$, then Eqs. (10.8) and (10.9) predict that the number of microstates and the entropy scales as power laws

$$\Omega \propto N^{m(\kappa)/\kappa} \quad , \quad (10.10)$$

$$S \propto N^{m(\kappa)} \quad . \quad (10.11)$$

We remark that the power law behavior of the various physical quantities is a feature of the present theory in contrast with the exponential and logarithmic behaviors in classical statistical mechanics. Finally, we assume, as customary, a uniform distribution of the components in the volume V , namely $N \propto V$ and obtain the following expression of entropy

$$S \propto V^{m(\kappa)} \quad . \quad (10.12)$$

After posing

$$m(\kappa) = 2/3 \quad , \quad (10.13)$$

the Bekenstein-Hawking law, namely $S \propto A$, follows immediately and thus we can conclude that the present statistical theory is consistent with the area law of black hole entropy. This result is in agreement with ref. [37] where the authors propose a relationship analogous to the one given by Eq. (10.13).

XI. EPIMYTHION

We have shown that within the standard framework of the special relativity, the Lorentz transformations lead to a proper one-parameter generalization of the entropy just as for other physically meaningful quantities (e.g. momentum, energy, etc).

The new entropy generates a coherent and self-consistent relativistic statistical theory (both statistical mechanics as well as kinetic theory) preserving the main features (maximum entropy principle, thermodynamic stability, continuity condition, H-theorem, etc) of the ordinary statistical theory which riemerges in the classical limit.

The distribution function predicted by the present theory is a one-parameter, continuous deformation of the classical Maxwell-Boltzmann distribution exhibiting a power law tail in accordance with the experimental evidence in several relativistic many body systems.

XII. APPENDIX

We start by considering the κ -sum for the dimensionless relativistic momenta

$$t_1 \oplus_{\kappa} t_2 = t_1 \sqrt{1 + \kappa^2 t_2^2} + t_2 \sqrt{1 + \kappa^2 t_1^2} \quad , \quad (12.1)$$

and after posing $t_1 = t + dt$ and $t_2 = -t$ we can introduce κ -differential through

$$\begin{aligned} d_{\{\kappa\}} t &= (t + dt) \ominus_{\kappa} t \\ &= (t + dt) \oplus_{\kappa} (-t) \\ &= \frac{dt}{\sqrt{1 + \kappa^2 t^2}} \quad . \end{aligned} \quad (12.2)$$

The κ -derivative $d/d_{\{\kappa\}}t$ admits as eigenfunction the κ -exponential, namely

$$\frac{d}{d_{\{\kappa\}}t} \exp_{\{\kappa\}}(t) = \exp_{\{\kappa\}}(t) , \quad (12.3)$$

being

$$\exp_{\{\kappa\}}(t) = \left(\sqrt{1 + \kappa^2 t^2} + \kappa t \right)^{1/\kappa} , \quad (12.4)$$

$$\exp_{\{\kappa\}}(t) = \exp\left(\frac{1}{\kappa} \operatorname{arcsinh}(\kappa t)\right) . \quad (12.5)$$

The κ -exponential can be viewed as a generalization of the ordinary exponential

$$\exp_{\{0\}}(t) = \exp(t) , \quad (12.6)$$

$$\exp_{\{-\kappa\}}(t) = \exp_{\{\kappa\}}(t) , \quad (12.7)$$

and has the properties

$$\exp_{\{\kappa\}}(t) \in C^\infty(\mathbf{R}) , \quad (12.8)$$

$$\frac{d}{dt} \exp_{\{\kappa\}}(t) > 0 , \quad (12.9)$$

$$\frac{d^2}{dt^2} \exp_{\{\kappa\}}(t) > 0 , \quad (12.10)$$

$$\exp_{\{\kappa\}}(-\infty) = 0^+ , \quad (12.11)$$

$$\exp_{\{\kappa\}}(0) = 1 , \quad (12.12)$$

$$\exp_{\{\kappa\}}(+\infty) = +\infty , \quad (12.13)$$

$$\exp_{\{\kappa\}}(t) \exp_{\{\kappa\}}(-t) = 1 . \quad (12.14)$$

Furthermore it results

$$\left(\exp_{\{\kappa\}}(t) \right)^r = \exp_{\{\kappa/r\}}(rt) , \quad (12.15)$$

$$\exp_{\{\kappa\}}(t_1) \exp_{\{\kappa\}}(t_2) = \exp_{\{\kappa\}}(t_1 \oplus t_2) , \quad (12.16)$$

$$\exp_{\{\kappa\}}(t_1) \otimes_{\kappa} \exp_{\{\kappa\}}(t_2) = \exp_{\{\kappa\}}(t_1 + t_2) , \quad (12.17)$$

where the product \otimes is defined through Eq. (9.5). The following asymptotic relations also hold

$$\exp_{\{\kappa\}}(t) \underset{t \rightarrow 0}{\sim} 1 + t + \frac{t^2}{2} + (1 - \kappa^2) \frac{t^3}{3!} , \quad (12.18)$$

$$\exp_{\{\kappa\}}(t) \underset{t \rightarrow \pm\infty}{\sim} |2\kappa t|^{\pm 1/|\kappa|} . \quad (12.19)$$

Finally, after tedious but straightforward calculations the following formula can be obtained

$$\int_0^\infty t^{r-1} \exp_{\{\kappa\}}(-t) dt = \frac{|2\kappa|^{-r}}{1+r|\kappa|} \frac{\Gamma\left(\frac{1}{|2\kappa|} - \frac{r}{2}\right)}{\Gamma\left(\frac{1}{|2\kappa|} + \frac{r}{2}\right)} \Gamma(r) , \quad (12.20)$$

holding for $r|\kappa| < 1$. Note that this integral is given also in [19] by Eq. (5.31) which unfortunately contains

a typing error (the first minus in the denominator of the right hand side must be replaced by a plus so that after some simple algebra Eq. (5.31) of [19] reproduces the present result).

The κ -logarithm $\ln_{\{\kappa\}}(t)$, defined as the inverse function of the of κ -exponential, namely $\ln_{\{\kappa\}}(\exp_{\{\kappa\}} t) = \exp_{\{\kappa\}}(\ln_{\{\kappa\}} t) = t$, is given by

$$\ln_{\{\kappa\}}(t) = \frac{t^\kappa - t^{-\kappa}}{2\kappa} , \quad (12.21)$$

$$\ln_{\{\kappa\}}(t) = \frac{1}{\kappa} \sinh(\kappa \ln t) , \quad (12.22)$$

and can be viewed as a generalization of the ordinary logarithm

$$\ln_{\{0\}}(t) = \ln(t) , \quad (12.23)$$

$$\ln_{\{-\kappa\}}(t) = \ln_{\{\kappa\}}(t) . \quad (12.24)$$

The κ -logarithm, just as the ordinary logarithm, has the properties

$$\ln_{\{\kappa\}}(t) \in C^\infty(\mathbf{R}^+) , \quad (12.25)$$

$$\frac{d}{dt} \ln_{\{\kappa\}}(t) > 0 , \quad (12.26)$$

$$\frac{d^2}{dt^2} \ln_{\{\kappa\}}(t) < 0 , \quad (12.27)$$

$$\ln_{\{\kappa\}}(0^+) = -\infty , \quad (12.28)$$

$$\ln_{\{\kappa\}}(1) = 0 , \quad (12.29)$$

$$\ln_{\{\kappa\}}(+\infty) = +\infty , \quad (12.30)$$

$$\ln_{\{\kappa\}}(1/t) = -\ln_{\{\kappa\}}(t) . \quad (12.31)$$

Moreover it has also the properties

$$\ln_{\{\kappa\}}(t^r) = r \ln_{\{r\kappa\}}(t) , \quad (12.32)$$

$$\ln_{\{\kappa\}}(t_1 t_2) = \ln_{\{\kappa\}}(t_1) \oplus_{\kappa} \ln_{\{\kappa\}}(t_2) , \quad (12.33)$$

$$\ln_{\{\kappa\}}(t_1 \otimes_{\kappa} t_2) = \ln_{\{\kappa\}}(t_1) + \ln_{\{\kappa\}}(t_2) , \quad (12.34)$$

while the following asymptotic formulas hold

$$\ln_{\{\kappa\}}(1+t) \underset{t \rightarrow 0}{\sim} t - \frac{t^2}{2} + \left(1 + \frac{\kappa^2}{2}\right) \frac{t^3}{3} , \quad (12.35)$$

$$\ln_{\{\kappa\}}(t) \underset{t \rightarrow 0^+}{\sim} -\frac{1}{2|\kappa|} t^{-|\kappa|} , \quad (12.36)$$

$$\ln_{\{\kappa\}}(t) \underset{t \rightarrow +\infty}{\sim} \frac{1}{2|\kappa|} t^{|\kappa|} . \quad (12.37)$$

Furthermore holds the very useful formula

$$\frac{d^2}{dt^2} t \ln_{\{\kappa\}}(t) > 0 , \quad (12.38)$$

and finally one obtains [19]

$$\int_0^1 dt \left(\ln_{\{\kappa\}} \frac{1}{t} \right)^{r-1} = \frac{|2\kappa|^{1-r}}{1+(r-1)|\kappa|} \frac{\Gamma\left(\frac{1}{|2\kappa|} - \frac{r-1}{2}\right)}{\Gamma\left(\frac{1}{|2\kappa|} + \frac{r-1}{2}\right)} \Gamma(r) . \quad (12.39)$$

It results evident that the above used procedure, to introduce the κ -exponential and κ -logarithm, starting from the additivity law of the momenta, can be used to obtain new functions if we start from other generalized additivity laws. For instance, starting from the additivity law of the velocities we have $d_{\{\kappa\}}t = (1 - \kappa^2 t^2)^{-1} dt$, while instead of κ -exponential we obtain the function

$$\varphi(t) = \left(\frac{1 + \kappa t}{1 - \kappa t} \right)^{1/2\kappa}, \quad (12.40)$$

$$= \exp\left(\frac{1}{\kappa} \operatorname{arctanh}(\kappa t)\right), \quad (12.41)$$

whose inverse function assumes the form

$$\varphi^{-1}(t) = \frac{1}{\kappa} \frac{t^\kappa - t^{-\kappa}}{t^\kappa + t^{-\kappa}}, \quad (12.42)$$

$$= \frac{1}{\kappa} \tanh(\kappa \ln t). \quad (12.43)$$

The function $\varphi(t) \in C^\infty(I)$ with $I =] -|\kappa|^{-1}, +|\kappa|^{-1}[$

is connected to $\exp_{\{\kappa\}}(t)$ through

$$\varphi(t) = \exp_{\{\kappa\}}\left(\frac{t}{\sqrt{1 - \kappa^2 t^2}}\right). \quad (12.44)$$

At this point one could be tempted to replace $\exp_{\{\kappa\}}(t)$ with the function $\varphi(t)$ as the right one for generalize the ordinary exponential.

There are several reasons to consider $\exp_{\{\kappa\}}(t)$ as the more proper generalization of the ordinary exponential. Beside the motivations emerging within the one-particle relativistic dynamics, there is the one related to the maximum entropy principle. This principle, the cornerstone of statistical physics, selects without any ambiguity (see sect. VII) just the κ -exponential as the more natural generalization of the ordinary exponential. More precisely the function $\exp_{\{\kappa\}}(t)$ is the only one existing which emerges simultaneously both in the one-particle relativistic dynamics as well as in the many-body physics through the maximum entropy principle.

-
- [1] J. Bernstein, *Kinetic Theory in the expanding Universe* (Cambridge University Press), Cambridge, England, (1988).
- [2] G.M. Kremer, and F.P. Devecchi, Phys. Rev. D **65**, 083515 (2002).
- [3] A. Giovannini, S. Lupia, and R. Ugoccioni, Phys. Rev. D **65**, 094028 (2002).
- [4] D. Colladay, and P. McDonald, Phys. Rev. D **70**, 125007 (2004).
- [5] R. Prix, Phys. Rev. D **69**, 043001 (2004).
- [6] Hyun Kyu Lee, Phys. Rev. D **66**, 063001 (2002).
- [7] V. Husain, Phys. Rev. D **69**, (2004).
- [8] G. Gour, Phys. Rev. D **67**, 127501 (2003).
- [9] R.K. Bhaduri, and M.N. Tran, Phys. Rev. D **69**, (2004).
- [10] E.T. Jaynes, Phys. Rev. **106**, 620 (1957); **108**, 171 (1957).
- [11] V.M. Vasyliunas, J. Geophys. Res. **73**, 2839 (1968).
- [12] P.L. Biermann, and G. Sigl, *Physics and Astrophysics of Ultra-Hight-Energy Cosmic Rays*, Lectures Notes in Physics **576**, Springer-Verlag Berlin (2001), [astro-ph/0202425].
- [13] S.P. Swordy, Cosmic Rays Data, Chicago University, <http://astroparticle.uchicago.edu/announce.htm>.
- [14] J.D. Bekenstein, Phys. Rev. D **7**, 2333 (1972).
- [15] S.W. Hawking, Commun. Math. Phys. **43**, 1999 (1975).
- [16] A. Hasegawa, A.M. Kunioki, and M. Duong-van, Phys. Rev. Lett. **54**, 2608 (1985).
- [17] G. Wilk and, Z. Włodarczyk, Phys. Rev. D **50**, 2318 (1994).
- [18] G. Kaniadakis, Physica A **296**, 405 (2001).
- [19] G. Kaniadakis, Phys. Rev. E **66**, 056125 (2002).
- [20] G. Kaniadakis, M. Lissia and A.M. Scarfone, Phys. Rev. E **71**, 046128 (2005).
- [21] S. Abe, J. Phys. A: Math. Gen. **36**, 8733 (2003).
- [22] J. Naudts, Physica A **316**, 323 (2002).
- [23] P.-H. Chavanis, Physica A **332**, 89-122 (2004).
- [24] A.G. Bashkirov, Phys. Rev. Lett. **93**, 130601 (2004).
- [25] I.S. Gradshteyn and I.M. Ryzhik, *Table of integrals, series, and Product*, Academic Press London (2000).
- [26] C. Cercignani, and G. Medeiros Kremer, *The relativistic Boltzmann equation: theory and applications* (Birkhauser Verlag), Basel-Boston-Berlin (2002).
- [27] S.R. de Groot, W.A. van Leeuwen, and Ch.G. van Weert, *Relativistic Kinetic Theory: Principle and Applications*, North-Holland, Amsterdam (1980).
- [28] G. Kaniadakis and A.M. Scarfone, Physica A **340**, 102 (2004).
- [29] A. Einstein, Jahrb. Radioaktivitat und Electronik **4**, 411 (1907).
- [30] M. Planck, Ann. Phys. **26**, 1 (1908).
- [31] A.M. Teweldeberhan, H.G. Miller, and R. Tegen, Int. J. Mod. Phys. E **12**, 669 (2003).
- [32] A. Rossani and A. M. Scarfone, J. Phys. A: Math. Gen. **37**, 4955 (2004).
- [33] M. Cravero et al., Physica A **340**, 410 (2004).
- [34] F. Topsoe, Physica A **340**, 11 (2004).
- [35] A. Dragulescu, [cond-mat/0307341].
- [36] D. Rajaonarison, D. Bolduc and H. Jayet, Econ. Lett. **86**, 13 (2005).
- [37] M. Botta Cantcheff and J.A.C. Nogales, [gr-qc/0504004].
- [38] M. Gell-Mann and C. Tsallis (Eds.), *Nonextensive Entropy-Interdisciplinary applications*, Oxford University, New York, (2004).