

On electrons and hydrogen-bond connectivity in liquid water

M. V. Fernandez-Serra¹ and Emilio Artacho^{2,3}

¹Laboratoire de Physique de la Matière Condensée et Nanostructures (LPMCN),
Université Claude Bernard Lyon 1, 69622 Villeurbanne, France

²Donostia International Physics Center, Universidad del País Vasco, 20080 San Sebastian, Spain,

³Department of Earth Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EQ, UK
(Dated: January 24, 2022)

The network connectivity in liquid water is revised in terms of electronic signatures of hydrogen bonds (HBs) instead of geometric criteria, in view of recent X-ray absorption studies. The analysis is based on *ab initio* molecular-dynamics simulations at ambient conditions. Even if instantaneous thread-like structures are observed in the electronic network, they continuously reshape in oscillations reminiscent of the r and t modes in ice (~ 170 fs). However, two water molecules initially joint by a HB remain effectively bound over many periods regardless of its electronic signature.

Water is an extremely intriguing liquid that continues to excite the interest of scientists in many disciplines. Many of its anomalous properties [1, 2] originate in the hydrogen bonds (HBs) among water molecules [3, 4]. The concept of a network liquid emerges naturally from this HB connectivity, an intuitive image that has provided interesting insights into the properties of water [1, 5]. Direct structural information to characterize such network structure is hard to obtain experimentally. Diffraction techniques [6, 7] or radial distribution functions (RDFs) vary naturally, but rely on reverse Monte Carlo techniques using force-field models to obtain further structural information [8]. Spectroscopic probes provide a rich source of complementary information. X-ray emission [9, 10] (XES) and X-ray absorption [11, 12, 13] (XAS) spectroscopies explore the electronic states of the liquid right below and above the Fermi level, respectively. In particular, the work by Wernet et al. [13] has recently introduced an extremely interesting new component into the study of liquid water, by relating a pre-edge feature in the XAS spectra with broken HBs. The authors propose to determine connectivity by looking at an electronic-structure signature of the HBs. Their conclusion is daring: the average coordination in liquid water would be 2 instead of the previously accepted value slightly under 4, displaying a lamentous picture, instead of the distorted, partly broken and fluctuating tetrahedral network described in so many papers before [6, 7, 14, 15].

Is it true? This would be the wrong question to ask. The kind of network depends on the definition of the hydrogen bond, furthermore, on deciding whether two given molecules in a given conformation are bonded or not. There is no direct physical HB observable and there is arbitrariness in the choice of what is actually measured. Instead, we address the question of how relevant is the newly proposed network image for the description of the liquid in the sense of the insights it offers. The conventional criterion [5] for HB is based on geometric considerations: an oxygen-oxygen distance within the first peak of the O-O RDF, and an upper critical bend angle (see Fig. 1). This "geometric" definition is based on total

energy considerations in contrast to the newly proposed "electronic" one. In this paper we explore the adequacy of the newly proposed probe, including its time scale, virtually instantaneous as compared with atomic motions.

Electronic structure calculations have been performed based on density-functional theory (DFT), within the generalized-gradient approximation (BLYP) [16, 17]. The Siesta method is used [18, 19] with a basis set of atomic orbitals at the double-polarized level [20]. For liquid water at ambient conditions, *ab initio* molecular-dynamics (AIMD) simulations have been performed in the microcanonical ensemble, based on the DFT forces and the Born-Oppenheimer approximation. Further details are found in Ref. 15. XAS spectra have been calculated for selected conformations (see below). The pseudo-atomic orbitals in the basis set have been PAW-transformed [21] into all-electron atomic orbitals for calculating matrix elements. For our basis set, neglecting the very small inter-molecular matrix elements was found to give an adequate approximation for the purposes of this paper. The strong excitonic effect introduced by the attraction between the core hole and the excited electron is estimated in the $Z + 1$ approximation [22].

Notwithstanding the importance of the XAS experimental data for this and other purposes, the probe provides a rather indirect measure of the electronics of the hydrogen bond, not least because of the mentioned excitonic effect. This and other difficulties [23, 24] (broadening, alignments) make it very difficult to obtain quantitative comparisons for the liquid phase. We find it more useful for our purposes to use a ground-state probe that we validate against XAS data in cleaner systems. This validation is two-sided. On one hand we test our chosen probe, on the other, we test the extent to which the XAS probe reflects the properties of the electronic ground state.

A very natural choice for describing electronic bonding within our method is bond order. In its simplest definition [25], the bond order between two atoms, 1 and 2, is $Q_{1,2} = \frac{1}{2} \sum_P \frac{P_1 P_2}{S_P}$, where \sum_P sums over the basis functions associated to atom 1 (2), and P_1 and S_P are the

density and overlap matrices, respectively. Bond orders depend on the choice of basis, and their arbitrariness has been described at length (see [26] and refs. therein). It is relative changes in that value what we use in this work, and these are shown below to be meaningful enough to support its conclusions. Mulliken's bond orders are compared with Mayer's [27], from which the same conclusions are drawn. We believe that any other electronic-structure signature of the bonding [26] would reflect the same physics.

There has been a controversy [28, 29, 30, 31] on the covalence of the HB and on its bonding or antibonding character, which would seem to affect our choice of electronic probe. The electronic characteristics of the HB were nicely illustrated with maximally localized Wannier functions [30]. They can also be described in terms of an intramolecular polarization (rehybridization within the molecules) and intermolecular polarization or charge transfer (admixture of orbitals of different molecules), if using the language of single-molecule orbitals [32], or of atomic orbitals, in either mono- or multi-determinantal wave-functions [29]. The physics of the interactions behind the HB is, however, quite clear if one avoids the semantic problems that have been partly behind that controversy. In a typical HB there is an important electrostatic attraction that dominates the energetics [30]. In addition, there is a deformation of the electronic cloud around O's accepting lone pair towards the donating H, in response to the field generated by the latter. There are other effects (e.g. quantum fluctuations of the protons) that are energetically less significant in principle, but could still be important for liquid water. They are beyond the scope of this paper. The mentioned polarization of the lone pair happens at the expense of a slight contraction of the electron cloud involved in the O-H bond within the donor molecule, due to Pauli exclusion [30]. This last deformation is the one behind the antibonding character of the HB discussed in the literature [29]. However, the original deformation of the O's lone-pair towards the proton remains clearly bonding. In this study we thus concentrate on Q_{OH} , between the donor H and the acceptor O.

In order to compare XAS and Q_{OH} we have calculated XAS spectra for a four-layer (001) ice slab, with the surface molecules exposing non-donated protons (dangling HBs). Repeated slabs are separated by 15 Å of vacuum. An extra water molecule has been placed on top of one of the surface molecules, and its position has been varied. Figure 1 shows the XAS calculated for the donor surface molecule and Q_{OH} for that HB as a function of HB O-H distance for two different values of the ap angle, (see Fig. 1). 40° corresponds to an ideal tetrahedral arrangement as in ice. A zero ap angle gives a rather unfavorable situation, since the proton faces midway between both lone pairs of O (in the nodal plane of the acceptor's HOMO orbital, of C_{2v} 's B_1 character), which




FIG. 1: XAS peak intensity ratio $I_{\text{edge}}/I_{\text{pre}}$ (a), and bond order Q_{OH} (b) versus intermolecular distance for two values of the ap angle, $\phi = 40^\circ$ and 0° , for the HB between an acceptor molecule and a clean (001) ice surface.

partly inhibits the electronic deformation.

We focus on two distinctive features of the XAS spectrum directly related to the pre-edge observed in the experiments: (i) the relative intensity of the peaks related to edge and pre-edge features, and (ii) the energy difference between both peaks E_s . As discussed by Cavalleri et al. [22], the spectral intensity is taken from the Z approximation (initial-state), while E_s is obtained from the $Z+1$ approximation, given its sensitivity to final-state effects. Fig. 1 shows that $\phi = 0^\circ$ produces a pre-edge twice as large as $\phi = 40^\circ$ for the same d_{OO} . This remarkable effect is closely reproduced by Q_{OH} , as well as the distance dependence. E_s increases with distance in a similar manner (not shown), also well replicated by the bond order (the effect of the ap angle is less noticeable in this case). A detailed study of the dependence of Q_{OH} on intermolecular geometry in a water pair can be found in Ref. 33. It is important to note that both magnitudes (XAS and Q_{OH}) agree in not displaying any obvious feature (discontinuity, zero, minimum) that would define a natural threshold for HB breaking. We will thus refrain from establishing an arbitrary criterion for the moment and explore what can be learnt independent of it.

The Q_{OH} 's are then calculated in an AIMD simulation of liquid water for all water pairs within a first coordination shell, as defined by the first peak in the O-O RDF. The first point that becomes apparent is that every water molecule is mainly donating one strong HB, while the second bond order is 2.2 times weaker in average, partly supporting the one-dimensional network picture proposed in Ref. 13. The asymmetry is, however, not extreme, as can be seen in the distribution of strong to weak bond-order ratio in Fig. 2.




FIG. 2: (a) Distribution of strong to weak bond-order ratio in the liquid during a 25 ps AIMD run. (b) Evolution of the donating Q_{OH} 's for a given molecule [zoom of the inset].

Fig. 2(b) shows a fragment of the time evolution of the Q_{OH} on the two donated HBs for a representative molecule (the complete trajectory is shown in the inset). The figure shows a clear oscillation with a period of 170 fs. It would correspond to the intermolecular r and t vibrations in ice (hindered rotation and translation) [34]. An intermolecular O-H under-damped oscillation with a period of 170 fs has been indeed directly observed in liquid water [35] using ultrafast infrared spectroscopy. The figure shows that the oscillation in Q_{OH} is more pronounced than what expected from Fig. 2 (a), with lower values 5 times smaller than the higher ones in average. It shows that an anti-phase vibration of both donating HBs only accounts for part of the oscillation, the remaining part coming from the in-phase vibration that would weaken (strengthen) both donated HBs simultaneously. The figure also shows that a single HB survives many such extreme cycles before breaking (the average life-time for a HB is a few ps). [5, 33].

HBs with low enough values of instantaneous Q_{OH} are certainly contributing to the measured XAS pre-edge. The XAS probe is thus reflecting a very pronounced electronic effect, a swinging of the electron cloud of the lone pair following the "lapping" and stretching of HBs, which appears to XAS as if many HBs were broken. Considering the anti-phase component, that behavior would give rise to a pulsating 1D lament-like network in age, relevant for electron dynamics. It is, however, irrelevant to the liquid dynamics, since it is apparent in Fig 2 that the molecules in a HB are still effectively bound even when the electron deformation is very small. The inclusion of quantum fluctuations to the nuclear dynamics could even exaggerate the beating effect, since quantum and thermal fluctuations are comparable in scale (the

zero point motion of a 170 fs oscillation is 12 meV).

Figure 3 shows the distribution of Q_{OH} for all the water pairs in the liquid with $r_{OO} < 3.5$ Å, for two different definitions of bond order. In both cases there is a clear minimum at small values of the bond order, with 25 % HBs below that threshold. It offers a natural criterion (albeit still arbitrary) for determining the presence of a HB, namely, $Q_{OH} > Q_{OH}^{min}$ [36]. Interestingly, the width of the distribution for the HBs ($2 Q_{OH} = 0.030$, $Q_{OH} = 0.023$) is comparable to that for a single oscillating HB through time (the one in Fig. 2: $2 Q_{OH} = 0.026$, $Q_{OH} = 0.020$). This means that the very different HB strengths are not so much due to different configurations as the liquid flows (on the time scale of several ps, the HB lifetime) but remarkably related to these 170 fs vibrations.




FIG. 3: Distribution of bond orders in a 25 ps AIMD liquid simulation, using Mulliken's [25] (a) and Mayer's [27] (b).

After monitoring the electronic deformation as signature for connectivity, we finish this study by assessing the effect of such deformation on the liquid structure itself. Having observed how the electronic cloud deforms in response to stretch and ap, it should be expected that the energetics would be affected by the ap, and so would the configurations visited in AIMD trajectories. In Fig. 4 the distribution of ap angles obtained from AIMD and two classical non polarizable models are compared [37]. The angles α_1 and α_2 (as defined in Fig. 1) within the first coordination shell are used to characterize both ap and twist. The distributions show clear differences. The main difference between SPCE and TIP5P is the fact that the latter puts negative charges around the center of the electron lone pairs, which induces a more realistic ap response. It is, however, exaggerated, since the polarization is static. AIMD distributions reflect both the preference for the tetrahedral geometry and the flexibility given by the dynamical response of the electron cloud. It has been argued [13, 22, 38, 39] that, in spite of these differences, AIMD lines up with any force field so far to produce qualitatively wrong configurational sampling of liquid water. It is certainly true that there are clear shortcomings in our scheme (BLYP approximations for exchange and correlation, neglect of protonic quantum effects) as well as in others. What we propose here, however, is a highly plausible explanation of the experimental

results that does not imply the paradigm shift proposed by these authors.




FIG. 4: Distribution of θ_1 and θ_2 angles (as in Fig. 1) in liquid water, for (a) SPCE [40] and (b) TIP5P [4] force fields, and (c) AIMD. Darker regions indicate higher values. The $\theta_1 = \theta_2$ diagonal represents pure water. Moving normal to that diagonal represents twist. The maximum value along the diagonal is for zero twist (abrupt cut for rigid molecules). (d) Isosurface of the AIMD distribution of donating H's around a molecule.

In summary, using our electronic probe for HB connectivity we observe what could be described as 1D filamentous structures, but they are pulsating in a 170 fs period, the geometric connectivity surviving intact for many periods and thus many reshaping of the filaments. Even if this image of pulsating filaments is not relevant for the description of the liquid, it is likely that questions addressing its electronic structure could benefit from it.

We thank J. M. Soler and X. Blase for useful discussions and M. Dawber for help with the figures. EA acknowledges the hospitality at the Donostia International Physics Centre. We acknowledge financial support from the British Engineering and Physical Sciences Research Council, the Natural Environment Research Council through the eMinerals project, the Cambridge European Trust, the Region Rhône-Alpes, and the Comunidad Autónoma de Madrid. The calculations were performed in the Cambridge Crayfield High Performance Computing Facility.

Electronic address: marivi@lpm.cn.univ-lyon1.fr

- [1] J. R. Errington and P. G. Debenedetti, *Nature* 409, 318 (2001).
- [2] R. J. Speedy and C. A. Angell, *J. Chem. Phys.* 65, 851 (1976).
- [3] F. Stillinger, *Science* 209, 451 (1980).
- [4] M. W. Mahoney and W. L. Jorgensen, *J. Chem. Phys.* 112, 8910 (2000).

- [5] A. Luzar, *J. Chem. Phys.* 113, 10663 (2000).
- [6] J. M. Sorenson, G. Hura, R. M. Glaeser, and T. Head-Gordon, *J. Chem. Phys.* 113, 9149 (2000).
- [7] A. K. Soper, *Chem. Phys.* 258, 121 (2000).
- [8] A. K. Soper, *J. Phys. Condens. Matter* 9, 2717 (1996).
- [9] J. H. Guo et al., *Phys. Rev. Lett.* 89, 137402 (2002).
- [10] S. Kashtanov et al., *Phys. Rev. B* 69, 024201 (2004).
- [11] S. Myneni et al., *J. Phys. Condens. Matter* 14, 213 (2002).
- [12] M. Cavalleri et al., *Chem. Phys. Lett.* 364, 363 (2002).
- [13] Ph. Wernet et al., *Science* 304, 995 (2004).
- [14] G. W. Robinson, S.-B. Zhu, S. Singh, and M. W. Evans, "Water in Biology, Chemistry and Physics" (World Scientific, Singapore, 1996).
- [15] M. V. Fernandez-Serra and E. Artacho, *J. Chem. Phys.* 121, 11136 (2004).
- [16] A. D. Becke, *Phys. Rev. A* 38, 3098 (1998).
- [17] C. Lee, W. Yang, and R. G. Parr, *Phys. Rev. B* 37, 785 (1988).
- [18] P. Ordejón, E. Artacho, and J. M. Soler, *Phys. Rev. B* 53, R10441 (1996).
- [19] J. M. Soler et al., *J. Phys. Condens. Matter* 14, 2745 (2002).
- [20] E. Anglada, J. M. Soler, J. Junquera, and E. Artacho, *Phys. Rev. B* 66, 205101 (2002).
- [21] P. E. Blochl, *Phys. Rev. B* 50, 17953 (1994).
- [22] M. Cavalleri et al., *Phys. Chem. Chem. Phys.* (2005), in press.
- [23] J. D. Smith et al., *Science* 306, 851 (2004).
- [24] B. Hetenyi, F. De Angelis, P. Giannozzi, and R. Car, *J. Chem. Phys.* 118, 8632 (2004).
- [25] R. S. Mulliken, *J. Chem. Phys.* 23, 1833 (1955).
- [26] C. Fonseca-Guerra, J.-W. Handgraaf, E. J. Baerends, and F. M. Bickelhaupt, *J. Comput. Chem.* 25, 189 (2003).
- [27] I. Mayer, *Chem. Phys. Lett.* 97, 270 (1983).
- [28] E. D. Isaacs et al., *Phys. Rev. Lett.* 82, 600 (1999).
- [29] T. K. Ghanty, V. N. Staroverov, P. K. Korten, and E. R. Davidson, *J. Am. Chem. Soc.* 122, 1210 (2000).
- [30] A. Romero, P. L. Silvestrelli, and M. Parrinello, *J. Chem. Phys.* 115, 115 (2001).
- [31] A. Nilsson et al., *J. Chem. Phys.* 122, 154505 (2005).
- [32] P. Hunt, M. Sprik, and R. Vuilleumier, *Chem. Phys. Lett.* 376, 68 (2003).
- [33] M. V. Fernandez-Serra, PhD thesis, University of Cambridge, 2004.
- [34] P. L. Silvestrelli and M. Parrinello, *J. Chem. Phys.* 111, 3572 (1999).
- [35] C. J. Fecko et al., *Science* 301, 1698 (2003).
- [36] If one does use the Q_{OH}^{min} criterion, HBs do not "break" while oscillating. In fact, in $Q_{OH} \rightarrow Q_{OH}^{min}$ cases, the XAS curve in Fig. 1 saturates to the broken HB value, supporting this criterion. For $\theta = 40^\circ$ this happens at $d_{OO} = 3.8$ Å (instead of the 3.3 Å of Ref. 13), which would give less broken HBs (average coordination of 3.5 for our AIMD run) even if the configurations were the same.
- [37] For force fields: 64 molecules, 30 ps, NVT ($\alpha = 12.417$ Å, $T = 300$ K).
- [38] M. Cavalleri, M. Odelius, A. Nilsson, and L. G. M. Pettersson, *J. Chem. Phys.* 121, 10065 (2004).
- [39] A. Nilsson et al., *Science* 308, 793a (2005).
- [40] H. J. C. Berendsen, J. R. Grigera, and T. P. Straatsma, *J. Chem. Phys.* 91, 6269 (1987).