

Broad Peak in the $d_{x^2-y^2}$ Superconducting Correlation Length as a Function of Hole Concentration in the Two-Dimensional t - J Model

W. O. Putikka

Physics Department, The Ohio State University, 1680 University Dr., Mansfield, OH 44906

M. U. Luchini

31 Wingate Road, London W6 0UR, United Kingdom

(Dated: October 6, 2018)

We have calculated high temperature series to 12th order in inverse temperature for singlet superconducting correlation functions of the 2D t - J model with s -, $d_{x^2-y^2}$ - and d_{xy} -symmetry pairs. We find the correlation length for $d_{x^2-y^2}$ pairing grows strongly with decreasing temperature and develops a broad peak as a function of doping at $T/J = 0.25$ for $J/t = 0.4$. The correlation lengths for s - and d_{xy} -symmetry remain small and do not display peaks. Antiferromagnetic spin correlations at low doping act to suppress the $d_{x^2-y^2}$ and d_{xy} superconducting correlation lengths. Our results support the hypothesis that the strong electronic correlations found in the CuO_2 planes of high temperature superconductors are the origin of the superconducting order.

Shortly after the discovery of high temperature superconductors by Bednorz and Müller[1], Anderson[2] proposed that the new superconductors should be viewed as single band doped Mott insulators, with a novel, non-phonon origin for their superconductivity. The model used to describe the CuO_2 planes found in these materials is the two-dimensional Hubbard model, which in its strongly correlated limit was restated by Zhang and Rice[3] as the 2D t - J model. Below we show that the 2D t - J model has strongly enhanced $d_{x^2-y^2}$ pairing correlations for physical choices of the model parameters and doping range.

The Hubbard and t - J models have been widely studied by every method available in theorists' arsenals[4]. The results to date for pairing correlations have been at best mixed[5, 6, 7, 8, 9, 10, 11, 12, 22]. Lacking small parameters, strongly correlated models present severe difficulties for analytic calculations and numerical techniques have difficulties with finite size effects and the fermion sign problem. To avoid these problems, we calculate high temperature series to 12th order in inverse temperature β for the pairing correlations in the 2D t - J model. The series are generated by a linked cluster expansion[13], putting four extra operators on each cluster for the two pairs.

The Hamiltonian for the t - J model is

$$H = -tP \sum_{\langle ij \rangle \sigma} (c_{i\sigma}^\dagger c_{j\sigma} + c_{j\sigma}^\dagger c_{i\sigma}) P + J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j, \quad (1)$$

where the sums are over nearest neighbor pairs of sites on a square lattice and the P operators eliminate states with doubly occupied sites from the Hilbert space. The spin operators are given by $\mathbf{S}_i = \sum_{\alpha\beta} c_{i\alpha}^\dagger \boldsymbol{\sigma}_{\alpha\beta} c_{i\beta}$, where $\boldsymbol{\sigma}_{\alpha\beta}$ is a vector of Pauli matrices.

We limit our calculation to spin singlet pairing, with the pair operator for sites i and j given by

$$\Delta_{ij} = \frac{1}{\sqrt{2}} (c_{i\uparrow} c_{j\downarrow} - c_{i\downarrow} c_{j\uparrow}), \quad (2)$$

and the Hermitian adjoint

$$\Delta_{ij}^\dagger = \frac{1}{\sqrt{2}} (c_{i\downarrow}^\dagger c_{j\uparrow}^\dagger - c_{i\uparrow}^\dagger c_{j\downarrow}^\dagger). \quad (3)$$

The Hermitian real space four-point correlation function with operators on sites i, j, k and l is

$$\begin{aligned} P(i, j, k, l) = & \frac{1}{2} \langle \Delta_{ij}^\dagger \Delta_{kl} + \Delta_{lk}^\dagger \Delta_{ji} \rangle \quad (4) \\ & - \frac{1}{4} \left[\langle c_{i\downarrow}^\dagger c_{l\downarrow} \rangle \langle c_{j\uparrow}^\dagger c_{k\uparrow} \rangle + \langle c_{i\downarrow}^\dagger c_{k\downarrow} \rangle \langle c_{j\uparrow}^\dagger c_{l\uparrow} \rangle \right. \\ & + \langle c_{i\uparrow}^\dagger c_{l\uparrow} \rangle \langle c_{j\downarrow}^\dagger c_{k\downarrow} \rangle + \langle c_{i\uparrow}^\dagger c_{k\uparrow} \rangle \langle c_{j\downarrow}^\dagger c_{l\downarrow} \rangle \\ & + \langle c_{i\downarrow}^\dagger c_{i\downarrow} \rangle \langle c_{k\uparrow}^\dagger c_{j\uparrow} \rangle + \langle c_{k\uparrow}^\dagger c_{i\uparrow} \rangle \langle c_{i\downarrow}^\dagger c_{j\downarrow} \rangle \\ & \left. + \langle c_{l\uparrow}^\dagger c_{i\uparrow} \rangle \langle c_{k\downarrow}^\dagger c_{j\downarrow} \rangle + \langle c_{k\downarrow}^\dagger c_{i\downarrow} \rangle \langle c_{l\uparrow}^\dagger c_{j\uparrow} \rangle \right], \end{aligned}$$

where we have removed the single particle contributions to $P(i, j, k, l)$ to focus on pairing fluctuations. For a non-interacting system $P(i, j, k, l) = 0$. Due to the no double occupancy constraint in the t - J model we have two restrictions on the site choices: $i \neq j$ and $k \neq l$. Otherwise the pairs are allowed to take on all arrangements and sizes on the clusters.

From our data we calculate series for the real space correlators $P(\mathbf{r}, \mathbf{r}', \mathbf{R})$, where \mathbf{r} and \mathbf{r}' are vectors giving the orientation and internal size of the two pairs and \mathbf{R} is the distance between the pairs, determined by the distance between the centers of mass. Since the pairs are spin singlets, \mathbf{r} and \mathbf{r}' are limited to the upper half plane and positive x -axis to avoid double counting. All of the distances are measured in terms of the lattice spacing a . Since the original four point correlation function $P(i, j, k, l)$ is identically zero for non-interacting systems, we also have $P(\mathbf{r}, \mathbf{r}', \mathbf{R}) = 0$ for non-interacting systems. The internal symmetry of the pairs is included by multiplying $P(\mathbf{r}, \mathbf{r}', \mathbf{R})$ by phase factors $\phi_i(\mathbf{r})$ and $\phi_i(\mathbf{r}')$ where each phase factor can have the values $+1, -1$ or 0 . We have $\phi_i(\mathbf{r}) = 0$ when \mathbf{r} is aligned with a node. Here the

subscript i labels the different symmetries we consider: s , $d_{x^2-y^2}$ or d_{xy} . The phase factor of a single pair is shown in Fig. 1 for the anisotropic symmetries $d_{x^2-y^2}$ and d_{xy} . An s -symmetry pair has uniform phase, $\phi_s(\mathbf{r}) = +1$ for all \mathbf{r} .

From $P(\mathbf{r}, \mathbf{r}', \mathbf{R})$ and the phase factors we want to construct a correlator $P_i(\mathbf{R})$ which depends only on \mathbf{R} , the separation between the pairs, and the symmetry of the pairs ($i = s, d_{x^2-y^2}$ or d_{xy}). This is then used to measure the strength of the pairing correlations by calculating the $\mathbf{q} = 0$ correlation length via

$$\xi_i^2 = \frac{1}{2d} \frac{\sum_{\mathbf{R}} |\mathbf{R}|^2 P_i(\mathbf{R})}{\sum_{\mathbf{R}} P_i(\mathbf{R})}, \quad (5)$$

with dimension $d = 2$.

One way to calculate $P_i(\mathbf{R})$ is to sum over \mathbf{r} and \mathbf{r}' independently

$$P_i(\mathbf{R}) = \sum_{\mathbf{r}} \phi_i(\mathbf{r}) \sum_{\mathbf{r}'} \phi_i(\mathbf{r}') P(\mathbf{r}, \mathbf{r}', \mathbf{R}). \quad (6)$$

We find that for the anisotropic pair states with $|\mathbf{R}| > 0$ Eq. 6 gives $P_i(\mathbf{R})$'s that are very small and negative. This suggests there is spurious cancellation occurring in our definition of $P_i(\mathbf{R})$.

As an example of the problem that can arise when calculating $P_i(\mathbf{R})$ by Eq. 6, consider the special case[14] of pairs limited in size to nearest neighbors with $d_{x^2-y^2}$ symmetry. This is the type of calculation done in Refs. 5, 6, 7, 8, 9, 10, 11, 12 and Ref. 22. We have two types




FIG. 1: Internal nodes and spatial phases for d -symmetry pairs: a) $d_{x^2-y^2}$, b) d_{xy} . The light lines are the underlying square lattice and the solid black lines are the nodal lines. To determine the phase, consider one operator of a pair at the origin (where the nodes cross) and put the second operator on a different lattice site. The phase associated with that part of the lattice is the internal phase of the pair. The second operator cannot be put on the nodal lines; the pair wave function is zero for those configurations. For s -symmetry there are no nodes determined by symmetry and the phase is always positive.

of correlators to consider (omitting the subtracted single particle terms for clarity):

$$\langle O_+ \rangle = \langle c_{i\downarrow}^\dagger c_{i+\hat{\mathbf{x}}\uparrow}^\dagger c_{i+\mathbf{R}+\hat{\mathbf{x}}\uparrow} c_{i+\mathbf{R}\downarrow} \rangle$$

$$\langle O_- \rangle = \langle c_{i\downarrow}^\dagger c_{i+\hat{\mathbf{x}}\uparrow}^\dagger c_{i+\mathbf{R}+\hat{\mathbf{y}}\uparrow} c_{i+\mathbf{R}\downarrow} \rangle,$$

where $\hat{\mathbf{x}}$ and $\hat{\mathbf{y}}$ are unit vectors in the x and y directions. For $d_{x^2-y^2}$ symmetry we have the phases $\phi_+ = +1$ and $\phi_- = -1$. Now consider what happens to $P_{x^2-y^2}(\mathbf{R})$ above and below T_c as $|\mathbf{R}| \rightarrow \infty$:

$$\langle O_+ \rangle \rightarrow |\Phi_0|^2 + A_< \exp(-|\mathbf{R}|/\xi_{x^2-y^2}) \quad T < T_c$$

$$\rightarrow A_> \exp(-|\mathbf{R}|/\xi_{x^2-y^2}) \quad T > T_c$$

and

$$\langle O_- \rangle \rightarrow -|\Phi_0|^2 + B_< \exp(-|\mathbf{R}|/\xi_{x^2-y^2}) \quad T < T_c$$

$$\rightarrow B_> \exp(-|\mathbf{R}|/\xi_{x^2-y^2}) \quad T > T_c,$$

where $|\Phi_0|^2$ is the magnitude squared of the $d_{x^2-y^2}$ superconducting condensate and the A 's and B 's are non-universal coefficients of the parts of the correlator decaying with correlation length $\xi_{x^2-y^2}$. Below T_c we then have $P_{x^2-y^2}(\mathbf{R}) \sim \langle O_+ \rangle - \langle O_- \rangle \sim |\Phi_0|^2$ as $|\mathbf{R}| \rightarrow \infty$ due to the compensating phases for $\langle O_- \rangle$. This is the definition of long range superconducting order with $d_{x^2-y^2}$ symmetry.

High temperature series, however, are limited to the non-ordered high temperature phase where $P_{x^2-y^2}(\mathbf{R}) \sim (A_> - B_>) \exp(-|\mathbf{R}|/\xi_{x^2-y^2})$. Now if $A_>$ and $B_>$ do not reflect the symmetry of the ordered state we should expect $P_{x^2-y^2}(\mathbf{R})$ to be typically small and of indeterminate sign. Extracting $\xi_{x^2-y^2}$ via Eq. 5 is essentially impossible. In our calculation, $\langle O_+ \rangle$ and $\langle O_- \rangle$ are both positive, giving exactly this problem. This difficulty should be expected if there is anything in a calculation that prevents the system from being in the long range ordered state. Eq. 6 is generally a poor choice for a correlation function outside of the ordered phase since it can be anomalously small for reasons unrelated to the strength of the correlations.

To reduce the possibility of spurious cancellations in the definition of $P_i(\mathbf{R})$ we choose to limit the $P(\mathbf{r}, \mathbf{r}', \mathbf{R})$ we include to having $\mathbf{r} = \mathbf{r}'$. By doing this we hold the internal degrees of freedom of the pairs fixed while we measure the distance dependence \mathbf{R} of the correlation function. We have one pairing operator with a definite internal configuration \mathbf{r} at position zero and a second pairing operator with the same internal configuration at position \mathbf{R} . We then sum over \mathbf{r} since we are only interested in the \mathbf{R} dependence. Calculating a correlator where we allow \mathbf{r} and \mathbf{r}' to vary independently mixes distance correlation with internal degrees of freedom, which isn't the quantity we want to measure. Our definition of $P_i(\mathbf{R})$ is

$$P_i(\mathbf{R}) = \sum_{\mathbf{r}} \phi_i^2(\mathbf{r}) P(\mathbf{r}, \mathbf{r}, \mathbf{R}). \quad (7)$$

With this definition of $P_i(\mathbf{R})$ we are guaranteed to not have cancellation in the \mathbf{r} sum due to trivial symmetry reasons. The effects of different symmetries are still




FIG. 2: Examples of configurations of two pairs on a square lattice. The first pair is represented by \times 's and the second by \circ 's. a), b) and c) are examples of contributions to the $\mathbf{R}=(1, 0)$ correlator. For s -symmetry a), b) and c) all contribute, while for $d_{x^2-y^2}$ -symmetry the nodes eliminate c) and only a) and b) contribute. For d_{xy} -symmetry the nodes eliminate a) and b) so only c) contributes. d) is an example of a possible pair correlation not included in our definition of $P_i(\mathbf{R})$, Eq. 7.

present through the nodes of the pair wave function where $\phi_i(\mathbf{r}) = 0$. The correlator defined in Eq. 7 is more versatile than the correlator defined by Eq. 6 due to Eq. 7 being able to find growing fluctuations in the disordered phase and also show the order parameter in the ordered phase. An important feature of Eq. 7 is that to distinguish s and $d_{x^2-y^2}$ pairing we need to include pairs with $|\mathbf{r}| > 1$. Also, to have ξ_{xy}^2 positive we need to include pairs with $|\mathbf{r}| > 1$.

With $P_i(\mathbf{R})$ defined by Eq. 7, series can be calculated for ξ_i defined by Eq. 5. The resulting series are extrapolated to low temperatures using Padé approximants[15] either in the original expansion variable βJ or in the transformed variable $w = \tanh(\alpha\beta J)$ used by Singh and Glenister[16]. Here α is a numerical factor which can be adjusted to improve the convergence of the Padés. The estimated errors are from the spread of the Padé approximants at low temperatures.

Our results at $T/J = 0.25$ are shown in Fig. 3. Clearly the 2D t - J model does not have s - or d_{xy} -symmetry pairing for the range of parameters shown. However, the correlation length for $d_{x^2-y^2}$ -symmetry pairing $\xi_{x^2-y^2}$ is large and suggests the ground state of the 2D t - J model has long range superconducting order with $d_{x^2-y^2}$ symmetry. Further support for this interpretation is shown in Fig. 4, where $\xi_{x^2-y^2}$ versus doping is plotted for a range of temperatures. The temperature dependence




FIG. 3: Pairing correlation length versus doping. The t - J model parameter ratio is $J/t = 0.4$ and the temperature is $T/J = 0.25$. Results for s -, $d_{x^2-y^2}$ - and d_{xy} -symmetry pairing are shown. The lines are guides for the eye.

shows $\xi_{x^2-y^2}$ grows rapidly with decreasing temperature over most of the doping range shown in Fig. 4, though $\xi_{x^2-y^2}$ decreases with decreasing temperature for $\delta \lesssim 0.04$. This is distinct from s - and d_{xy} -symmetry pairing, which for low temperatures have ξ_s and ξ_{xy} slowly decreasing with decreasing temperature over the whole doping range shown. The temperature scale for growth in $\xi_{x^2-y^2}$ also agrees well with the temperature scales found for two-point correlation functions[17].




FIG. 4: Pairing correlation length for $d_{x^2-y^2}$ -symmetry versus doping for a range of temperatures. The t - J model parameter ratio is $J/t = 0.4$. The data sets are in the order indicated on the plot, with the highest curve corresponding to $T/J = 0.25$. The lines are guides for the eye.

Our results are in accord with experiments on high temperature superconductors[18]. The pairing has $d_{x^2-y^2}$ symmetry, with the doping range where $\xi_{x^2-y^2}$ is large comparable to where T_c is observed. The maximum

in $\xi_{x^2-y^2}$ is shifting to smaller doping as the temperature is reduced, and likely is near $\delta \approx 0.2$ for $T/J \approx 0.1$, though calculations at lower temperatures are needed to show this. Also, $\xi_{x^2-y^2}$ grows very steeply on the underdoped side, while falling off more slowly for overdoping. This is due to $\xi = 0$ at half filling for all symmetries and all temperatures and the strong antiferromagnetic correlations present for underdoping. In experiments T_c frequently jumps from zero to a non-zero value for underdoped samples with a small change in doping.

Band structure estimates[21] for t - J model parameters give a next nearest neighbor hopping term t' roughly the same size as J . This term has not been included in the calculations presented here. We expect that t' would allow holes to move more easily through the antiferromagnetic background present in the t - J model, and thus would move the peak in $\xi_{x^2-y^2}$ to lower doping. Although a t' term is not necessary to have strong superconducting correlations, calculations including a t' term should improve detailed comparisons between experiments on high temperature superconductors and the t - J model.

The 2D superconducting correlation length does not need to grow to an extreme size like ξ_{AF} at half filling[19] to produce true 3D long range order due to the effective interplanar hopping t_\perp driving the 2D to 3D crossover for superconductors being much larger than the interplanar spin coupling J_\perp . The temperature dependence for $\xi_{x^2-y^2}$ shown in Fig. 4 is likely to produce T_c 's in the range observed for high temperature superconductors. Lower temperature results are needed to show this conclusively.

A peak in $\xi_{x^2-y^2}$ has not been observed in previous calculations. The most likely reason is the problem with cancellation in $P_i(\mathbf{R})$ discussed above. Previous calculations[4, 5, 6, 7, 8, 9, 10, 11, 12, 22] the authors have examined have mixed terms for anisotropic pairing correlators. The negative U Hubbard model does not have this problem due to the pairing being predominantly on-site and isotropic[20]. The $\mathbf{q} = 0$ pairing susceptibility was calculated by high temperature series in Ref. 12, with the conclusion that $d_{x^2-y^2}$ pairing is not found in the 2D t - J model. The results of Ref. 12 are influenced by single particle terms we have removed and the cancellation effect. We believe these effects distort the results of Ref. 12 and when properly accounted for the $\mathbf{q} = 0$ pairing susceptibility will show $d_{x^2-y^2}$ correlations are present in the 2D t - J model.

In conclusion, we find strong $d_{x^2-y^2}$ -symmetry pairing correlations in the 2D t - J model for physical values of the model parameters and doping. This suggests the origin of high temperature superconductivity is the strong electronic correlations found in the CuO_2 planes.

This calculation was started while W. O. P. was visiting ETH, supported by the Swiss National Science Foundation and by a faculty travel grant from the Office of

International Studies at The Ohio State University, and continued while taking part in a workshop on high temperature superconductors at KITP, supported by NSF grant PHY99-07949. W. O. P. was also partially supported by NSF grant DMR01-05659. M. U. L. was supported by EPSRC Grant No. GR/L86852. The computations were done on the Itanium II cluster at the Ohio Supercomputer Center. We thank Tin-Lun Ho, Ciriya Jayaprakash, Mohit Randeria, Maurice Rice, Doug Scalapino, Rajiv Singh, David Stroud and John Wilkins for many useful discussions.

Note added in proof: After submission of this paper we received Ref. 22 where calculations related to ours are reported.

-
- [1] J. G. Bednorz and K. A. Müller, Z. Phys. B**64**, 189 (1986).
 - [2] P. W. Anderson, Science **235**, 1196 (1987).
 - [3] F. C. Zhang and T. M. Rice, Phys. Rev. B**37**, 3759 (1988).
 - [4] For reviews see D. J. Scalapino, Phys. Rep. **250**, 329 (1995) and E. Dagotto, Rev. Mod. Phys. **66**, 763 (1994).
 - [5] J. E. Hirsch and H. Q. Lin, Phys. Rev. B**37**, 5070 (1988).
 - [6] S. R. White, *et al.*, Phys. Rev. B**39**, 839 (1989); S. R. White, *et al.*, Phys. Rev. B**40**, 506 (1989).
 - [7] M. Imada, J. Phys. Soc. Jpn. **60**, 2740 (1991).
 - [8] D. J. Scalapino, S. R. White and S. C. Zhang, Phys. Rev. Lett. **68**, 2830 (1992).
 - [9] S. Zhang, J. Carlson and J. E. Gubernatis, Phys. Rev. Lett. **78**, 4486 (1997).
 - [10] C. T. Shih, Y. C. Chen, H. Q. Lin and T. K. Lee, Phys. Rev. Lett. **81**, 1294 (1998).
 - [11] S. Sorella, *et al.*, Phys. Rev. Lett. **88**, 117002 (2002).
 - [12] L. P. Pryadko, S. A. Kivelson and O. Zachar, Phys. Rev. Lett. **92**, 067002 (2004).
 - [13] M. P. Gelfand, R. R. P. Singh and D. Huse, J. Stat. Phys. **59**, 1093 (1990).
 - [14] This example comes from conversations the authors had with T.-L. Ho, C. Jayaprakash and M. Randeria.
 - [15] A. J. Guttmann, in *Phase Transitions and Critical Phenomena*, ed. by C. Domb and J. L. Lebowitz (Academic, San Diego, 1989) Vol. 13, p. 1.
 - [16] R. R. P. Singh and R. L. Glenister, Phys. Rev. B**46**, 11871 (1992).
 - [17] W. O. Putikka, M. U. Luchini and R. R. P. Singh, Phys. Rev. Lett. **81**, 2966 (1998); cond-mat/9912269.
 - [18] *Physical Properties of High Temperature Superconductors*, Vol 1-5, ed. D. M. Ginzberg (World Scientific, Singapore).
 - [19] S. Chakravarty, B. I. Halperin and D. R. Nelson, Phys. Rev. Lett. **60**, 1057 (1988); Phys. Rev. B**39**, 2344 (1989).
 - [20] A. Moreo and D. J. Scalapino, Phys. Rev. Lett. **66**, 946 (1991); A. Moreo, Phys. Rev. B**45**, 5059 (1992).
 - [21] E. Pavarini, I. Dasgupta, T. Saha-Dasgupta, O. Jepsen and O. K. Andersen, Phys. Rev. Lett. **87**, 047003 (2001).
 - [22] T. Koretsune and M. Ogata, J. Phys. Soc. Japan **74**, 1390 (2005).