

# The optical phenomena in multiple-photon interactions (I)

Xiangyao Wu<sup>a 1</sup>

Pi-Feng Gong<sup>a</sup>, Zhi-Yong Yi<sup>b</sup>, Chang-Rui Wu<sup>a</sup>, Xiao-Yan Gu<sup>c</sup> and Zong-Hua Shi<sup>a</sup>

a. *Institute of Physics , Qufu Normal University, Qufu 273165, China*

b. *BeiJing Institute of PetroChemical TECHN ,P.O.Box 8001, BeiJing 102617, China*

c. *Institute of High Energy Physics, P.O.Box 918(4), Beijing 100039, China*

## Abstract

In the recent experiment, the phenomena of superluminal and slow-light propagation in dispersive medium were found, and there are various explanation in theory. We find the phenomenon can be explained by multiple-photon interaction. Otherwise, the multiple-photon interaction can also appear other optical phenomenon: doubling frequency , sum frequency, difference frequency which are found in nonlinear optics.

**PACS number(s): 63.20.Kr, 14.70.Bh, 42.65.-k, 42.50.Gy**

---

<sup>1</sup>E-mail:wuxy2066@163.com

## 1. Introduction

Since laser appeared in 1960's, many new optical phenomena were found in nonlinear optics, e.g. frequency doubling, sum frequency, many-photo absorption, light condense, ultra-short impulse, laser self-interaction and so on [1]. These phenomena are generated when strong light interact with the nonlinear medium. In the past years, the new optical phenomena of superluminal and slow-light were found in the dispersion medium [2][3][4][5][6], and there are various explanation in theory. In this paper, we research these new optical phenomena with multiple-photon interaction, photon-photon interaction belongs to electromagnetic interaction, and it must satisfy some conservation laws, one of them is  $C$  parity conservation. For photon,  $C$  parity  $\eta_c = -1$ . So, photon-photon interaction should satisfy [7]:

- (1) if the results of photon-photon interaction generate photons, then even photons will generate even photons and odd photons will generate odd photons.
- (2) if the results of photon-photon generate material particle ( $m_0 \neq 0$ ), then they will be generate neutral particle  $n_0$  or neutral particle system ( $n + \bar{n}$ ).

In quantum field theory, there isn't direct-acting among photons, and they interact indirectly by Dirac vacuum. More recently, the noncommutative QED theory has been developed [10][11]. In the theory, photon has self-interaction unlike the case of ordinary QED, and Many new predictions [10][11] can be obtained from the noncommutative QED theory: multiple-photon direct-acting, superluminal photon, the photons of bound states and so on. In this paper, we can also obtain many new optical phenomena from multiple-photon interaction, and they are discussed only in kinematics. The research from dynamics will be appeared in the next paper.

## 2. Doubling frequency , sum frequency and difference frequency

In the following, we can obtain the optical phenomena of doubling frequency, sum frequency and difference frequency from three-photon collision. We consider two kinds of collision of three-photon.

case (a):

$$\begin{array}{c} \xrightarrow{h\nu_1} \\ \xrightarrow{h\nu_2} \end{array} \xleftarrow{h\nu_3} = \begin{array}{c} \xrightarrow{h\nu_4} \\ \xrightarrow{h\nu_5} \end{array} \xleftarrow{h\nu_6}$$

where  $\nu_i$  ( $i = 1, 2, \dots$ ) is frequency of  $i$ -th photon, the arrows represent moving direction of photon. In the photon collision, the energy and momentum are conservative.

$$h(\nu_1 + \nu_2 + \nu_3) = h(\nu_4 + \nu_5 + \nu_6) \quad (1)$$

$$h\left(\frac{1}{\lambda_1} + \frac{1}{\lambda_2} - \frac{1}{\lambda_3}\right) = h\left(\frac{1}{\lambda_4} + \frac{1}{\lambda_5} - \frac{1}{\lambda_6}\right) \quad (2)$$

Combining the Eq.(1) and (2)

$$\nu_3 = \nu_6 \quad (3)$$

$$\nu_1 + \nu_2 = \nu_4 + \nu_5 \quad (4)$$

From Eq.(4) if  $\nu_4 = 2\nu_1$  then  $\nu_5 = \nu_2 - \nu_1$  and if  $\nu_4 = 3\nu_1$  then  $\nu_5 = \nu_2 - 2\nu_1$ . So three-photon collision may produce frequency doubling, triplex frequency and difference frequency light phenomena.

case (b):

$$\begin{array}{c} \xrightarrow{h\nu_1} \\ \xrightarrow{h\nu_2} \end{array} \xleftarrow{h\nu_3} = \begin{array}{c} \xleftarrow{h\nu_4} \\ \xleftarrow{h\nu_5} \end{array} \xrightarrow{h\nu_6}$$

From the conservation law of energy and momentum

$$h(\nu_1 + \nu_2 + \nu_3) = h(\nu_4 + \nu_5 + \nu_6) \quad (5)$$

$$h\left(\frac{1}{\lambda_1} + \frac{1}{\lambda_2} - \frac{1}{\lambda_3}\right) = h\left(\frac{1}{\lambda_6} - \frac{1}{\lambda_4} - \frac{1}{\lambda_5}\right) \quad (6)$$

From Eq.(5) and Eq.(6)

$$\nu_6 = \nu_1 + \nu_2 \quad (7)$$

So three-photon collision can also produce sum frequency light. In experiment, frequency doubling, many-times frequency, sum frequency and difference frequency light are usually found in nonlinear medium [1]. We think these optical phenomena can be produced from collision three-photon or more photons.

### 3. Phenomena of superluminal

The superluminal phenomena have been found in many experiments [2]. We think the phenomena can be observed in three-photon collision, as follows:

case (a):

$$\begin{array}{c} \xrightarrow{h\nu_1} \\ \xrightarrow{h\nu_2} \end{array} \leftarrow h\nu_3 = \xrightarrow{h\nu_4}$$

From the conservative laws of energy and momentum

$$E = h(\nu_1 + \nu_2 + \nu_3) \quad (8)$$

$$P = \frac{h}{\lambda_1} + \frac{h}{\lambda_2} - \frac{h}{\lambda_3} \quad (9)$$

Where  $E$  and  $P$  are energy and momentum of photon  $\nu_4$

The speed of photon  $\nu_4$  is  $\tilde{c}$ :

$$\tilde{c} = \frac{E}{P} = \frac{h(\nu_1 + \nu_2 + \nu_3)}{h(\frac{1}{\lambda_1} + \frac{1}{\lambda_2} - \frac{1}{\lambda_3})} = \frac{c(\nu_1 + \nu_2 + \nu_3)}{\nu_1 + \nu_2 - \nu_3} > c$$

case (b):

$$\begin{array}{c} \downarrow h\nu_2 \\ \xrightarrow{h\nu_1} \\ \uparrow h\nu_3 \end{array} = \xrightarrow{h\nu_4}$$

Let  $\nu_2 = \nu_3$ , the energy and momentum of photon  $\nu_4$  is

$$E = h\nu_1 + h\nu_2 + h\nu_3 \quad (10)$$

$$P = \frac{h}{\lambda_1} \quad (11)$$

The speed of photon  $\nu_4$  is

$$\tilde{c} = \frac{E}{P} = \frac{c(\nu_1 + \nu_2 + \nu_3)}{\nu_1} > c$$

Obviously, many-photon ( $n > 3$ ) collision can also produce superluminal photon.

#### 4. Phenomena of slow-light

More recently, the phenomena of slow-light were found in many experiments[6][7][8][9]. We think it can also be observed in many-photon collision.

$$\begin{array}{c} \xrightarrow{h\nu_1} \\ \xrightarrow{h\nu_2} \end{array} \xleftarrow{h\nu_3} = \xrightarrow{h\nu_4} + n_0$$

The photon  $\gamma_4$  energy and momentum are

$$E = h\nu_1 + h\nu_2 + h\nu_3 - mc^2 \quad (12)$$

$$P = \frac{h}{\lambda_1} + \frac{h}{\lambda_2} - \frac{h}{\lambda_3} \pm P_0 \quad (13)$$

where  $mc^2$  and  $P_0$  are energy and momentum of neutral particle  $n_0$ . In Eq.(13), "  $\pm$  " is from the same or opposite momentum direction of photon  $\nu_4$  and particle  $n_0$ .

The speed of photon  $\gamma_4$  is

$$\tilde{c} = \frac{E}{P} = c \frac{\nu_1 + \nu_2 + \nu_3 - \frac{mc^2}{h}}{\nu_1 + \nu_2 - \nu_3 \pm \frac{P_0 c}{h}} \quad (14)$$

if

$$\frac{\nu_1 + \nu_2 + \nu_3 - \frac{mc^2}{h}}{\nu_1 + \nu_2 - \nu_3 \pm \frac{P_0 c}{h}} < 1 \quad (15)$$

then

$$\tilde{c} < c \quad (16)$$

The Eq. (15) is

$$2h\nu_3 < mc^2 \pm P_0 c \quad (17)$$

When the momentum direction of photon  $\nu_4$  and particle  $n_0$  is same, the Eq. (17) is

$$2h\nu_3 < mc^2 - P_0 c \quad (18)$$

On this condition, photon  $\gamma_4$  is slow-light.

When the momentum direction of photon  $\nu_4$  and  $n_0$  is opposite, the Eq. (17) is

$$2h\nu_3 < mc^2 + P_0c \quad (19)$$

On this condition, photon  $\gamma_4$  is slow-light.

We think that the phenomena of slow-light can also been found in multiple-photon interaction, which means there are slow-light photons in the vacuum.

## 5. Particle mass problem

In particle physics, particle mass is from spontaneous symmetry breaking and Higgs mechanism and particles can be produced by photon-photon interaction and they can be annihilation into photons. We can give particle rest mass  $m_0$  in photon-photon interaction. We can consider it as follows:

case (a): two-photon collision

$$\xrightarrow{h\nu_1} \xleftarrow{h\nu_2} = n_0$$

$$\nu_1 + \nu_2 \longrightarrow n_0$$

From the conservative laws of energy and momentum

$$E = h\nu_1 + h\nu_2 \quad (20)$$

$$P = \frac{h}{\lambda_2} - \frac{h}{\lambda_1} \quad (21)$$

Where  $E$  and  $P$  are energy and momentum of particle  $n_0$ . From special relatively, the relation between the total energy  $E$  of particle  $n_0$  and its momentum  $P$  and rest mass  $m_0$  is

$$E^2 = m_0^2c^4 + c^2P^2 \quad (22)$$

From Eq.(20), (21) and (22)

$$E_0^2 = m_0^2c^4 = E^2 - c^2P^2 = h^2(\nu_1 + \nu_2)^2 - c^2\left(\frac{h(\lambda_2 - \lambda_1)}{\lambda_1\lambda_2}\right)^2 = 4h^2\nu_1\nu_2$$

and so

$$m_0 = \frac{2h\sqrt{\nu_1\nu_2}}{c^2} = \frac{2\sqrt{\varepsilon_1\varepsilon_2}}{c^2} \quad (23)$$

The kinetic energy  $T$  of particle  $n_0$  is

$$T = E - E_0 = h\nu_1 + h\nu_2 - 2h\sqrt{\nu_1\nu_2} = h(\sqrt{\nu_1} - \sqrt{\nu_2})^2 \quad (24)$$

case (b): Multiple-photon collision

$$\begin{array}{ccc} \overrightarrow{h\nu_1} & \overleftarrow{h\nu'_1} & \\ \overrightarrow{h\nu_2} & \overleftarrow{h\nu'_2} & \\ \vdots & \vdots & \\ \overrightarrow{h\nu_m} & \overleftarrow{h\nu'_n} & \end{array} = n_0$$

$$(\nu_1 + \nu_2 + \cdots + \nu_m) + (\nu'_1 + \nu'_2 + \cdots + \nu'_n) \longrightarrow n_0$$

The total energy  $E$  and momentum  $P$  of particle  $n_0$  are:

$$E = h\nu_1 + h\nu_2 + \cdots + h\nu_m + h\nu'_1 + h\nu'_2 + \cdots + h\nu'_n \quad (25)$$

$$P = \frac{h}{\lambda_1} + \frac{h}{\lambda_2} + \cdots + \frac{h}{\lambda_m} - \frac{h}{\lambda'_1} - \frac{h}{\lambda'_2} - \cdots - \frac{h}{\lambda'_n} \quad (26)$$

and

$$E^2 = m_0^2 c^4 + c^2 P^2 \quad (27)$$

From Eq. (25), (26), and (27)

$$\begin{aligned} E_0^2 = m_0^2 c^4 &= E^2 - c^2 P^2 \\ &= h^2(\nu_1 + \nu_2 + \cdots + \nu_m + \nu'_1 + \nu'_2 \\ &\quad + \cdots + \nu'_n)^2 - h^2(\nu_1 + \nu_2 + \cdots + \nu_m - \nu'_1 - \nu'_2 - \cdots - \nu'_n)^2 \\ &= 4h^2(\nu_1 + \nu_2 + \cdots + \nu_m)(\nu'_1 + \nu'_2 + \cdots + \nu'_n) \end{aligned}$$

and so

$$\begin{aligned}
m_0 = \frac{E_0}{c^2} &= \frac{2h\sqrt{(\nu_1 + \nu_2 + \cdots + \nu_m)(\nu_{1'} + \nu_{2'} + \cdots + \nu_{n'})}}{c^2} \\
&= \frac{2\sqrt{(\varepsilon_1 + \varepsilon_2 + \cdots + \varepsilon_m)(\varepsilon_{1'} + \varepsilon_{2'} + \cdots + \varepsilon_{n'})}}{c^2}
\end{aligned} \tag{28}$$

where  $\varepsilon_i = h\nu_i$ ,  $\varepsilon_j = h\nu_{j'}$  are energy of photons whose frequency are  $\nu_i, \nu_{j'}$ .  
The kinetic energy of particle  $n_0$  :

$$\begin{aligned}
T = E - E_0 &= h(\nu_1 + \nu_2 + \cdots + \nu_m + \nu_{1'} + \nu_{2'} + \cdots + \nu_{n'}) \\
&\quad - 2h\sqrt{(\nu_1 + \nu_2 + \cdots + \nu_m)(\nu_{1'} + \nu_{2'} + \cdots + \nu_{n'})} \\
&= h(\sqrt{\nu_1 + \nu_2 + \cdots + \nu_m} - \sqrt{\nu_{1'} + \nu_{2'} + \cdots + \nu_{n'}})^2
\end{aligned} \tag{29}$$

So all matter particle ( $m_0 \neq 0$ ) can be produced in photons collisions. Its rest mass  $m_0$  is from the energy of interaction photons, and its kinetic energy is the rest energy of photons after those photons generate matter particle.

## 6. Conclusion

There are a number of new optical phenomena were found in nonlinear medium. We think these phenomena can be explained by multiple-photon interaction. Otherwise, matter particle can be obtained by the energy of interaction photons. These predictions can be checked in photon collider, but the dynamics mechanism of multi-photon interaction will be further research.



## References

- [1] R. W. Boyd, Nonlinear Optics 1-282 Academic press, **INC** , 1992
- [2] L. J. Wang, A. Kuzmich and A.Dogariu, Nature **406**, 277 (2000)
- [3] M.S. Bigelow, N.N. Lepeshkim and W.R. Boyd, Science **301**, 200 (2003)
- [4] A. Kuzmich, A. Dogariu, L.J. Wang, P.W. Milonni and Y.R. Chiao, Phys. Rev. Lett. **86**, 3925 (2001)
- [5] D. R. Solli et al, Phys. Rev. Lett. **91**, 143906 (2003)
- [6] J.Marangos, Nature **406**, 243 (2000)
- [7] L. V. Han, S.E.Harris, Z.Dutton and C.H.Behroozi, Nature **397**, 594 (1999)
- [8] M.M.Kash et al., Phys.Rev. Lett. **82**, 5229 (1999)
- [9] U.Leonhardt and P.Piwnicki, Phys.Rev.Lett. **84**, 822 (2000)
- [10] E.V.Gorbar, Michio Hashimoto and V.A.Miransky Phys. Lett. **B611**,207 (2005)
- [11] Car E.Carlson, Christopher D. Carone and Nahum Zobin Phys. Rev. **D66** 075001 (2002)