

# Asymptotic statistics of the $n$ -sided planar Poisson-Voronoi cell.

## I. Exact results

**H. J. Hilhorst**

Laboratoire de Physique Théorique, Bâtiment 210, Université de Paris-Sud  
91405 Orsay Cedex, France

December 27, 2021

### Abstract

We achieve a detailed understanding of the  $n$ -sided planar Poisson-Voronoi cell in the limit of large  $n$ . Let  $p_n$  be the probability for a cell to have  $n$  sides. We construct the asymptotic expansion of  $\log p_n$  up to terms that vanish as  $n \rightarrow \infty$ . We obtain the statistics of the lengths of the perimeter segments and of the angles between adjoining segments: to leading order as  $n \rightarrow \infty$ , and after appropriate scaling, these become independent random variables whose laws we determine; and to next order in  $1/n$  they have nontrivial long range correlations whose expressions we provide. The  $n$ -sided cell tends towards a circle of radius  $(n/4\pi\lambda)^{1/2}$ , where  $\lambda$  is the cell density; hence Lewis' law for the average area  $A_n$  of the  $n$ -sided cell behaves as  $A_n \simeq cn/\lambda$  with  $c = \frac{1}{4}$ . For  $n \rightarrow \infty$  the cell perimeter, expressed as a function  $R(\phi)$  of the polar angle  $\phi$ , satisfies  $d^2R/d\phi^2 = F(\phi)$ , where  $F$  is known Gaussian noise; we deduce from it the probability law for the perimeter's long wavelength deviations from circularity. Many other quantities related to the asymptotic cell shape become accessible to calculation.

**Keywords:** random graphs, exact results

LPT – ORSAY 05/42

<sup>1</sup>Laboratoire associé au Centre National de la Recherche Scientifique - UMR 8627

# 1 Introduction

Experimental data on cellular structures are often compared to the properties of a *Voronoi diagram* [1], sometimes also called a *Voronoi tessellation*. This well-defined mathematical object is a partitioning of space into convex cells constructed around ‘point particles’ (also called ‘centers,’ ‘seeds,’ or ‘germs’) in such a way that each point of space is in the cell of the particle to which it is closest. When the distribution of the point particles is *uniformly random*, the result is said to be a *Poisson-Voronoi diagram*; it is to naturally occurring tessellations what the ideal gas is to real gases.

In this work we will be interested in planar cellular structures. An example analyzed more than seventy years ago by Lewis [2] is the epidermal epithelium of cucumber, which is composed of many flat, tile-like cells. For other applications of Voronoi diagrams outside of physics we refer to the encyclopedic work by Okabe *et al.* [3].

Within physics, two-dimensional cellular structures appear in many different experimental situations, of which we mention only a few. Cerisier *et al.* [4] studied the cellular patterns in surface-tension driven Bénard convection. Elias *et al.* [5] studied two-dimensional magnetic liquid froth. Sire and Seul [6] analyzed droplet patterns in late-stage coarsening of a two-dimensional binary mixture. Moriarty *et al.* [7] analyzed nanostructured cellular layers by means of the Voronoi construction. Earnshaw and Robinson [8, 9] performed a Voronoi cell analysis of the structure of two-dimensional colloidal aggregates. Much important experimental work on tessellations comes from two-dimensional soap froths, as studied experimentally by Glazier *et al.* [10] and by Stavans and Glazier [11]. In simulations of the two-dimensional solid-liquid transition (recent examples are Zahn *et al.* [12] and Quinn and Goree [13]), Voronoi cells constructed around the particle positions define the nearest-neighbor relations between atoms, which in turn allow the identification of lattice defects. Richard *et al.* [14] used the Voronoi construction for the geometrical characterization of hard sphere systems.

Evidently, there are many different reasons why none of the above systems should be described exactly by a Poisson-Voronoi diagram. In dense fluids the particle positions are correlated and the resulting Voronoi diagrams are not of the Poisson type. Both soap froths and binary mixtures coarsen with time, and the experimentally determined properties of these cellular structures usually refer to a stationary scale invariant distribution (sometimes called the ‘equilibrium distribution’) observed at long times. Clearly such systems are best analyzed, in principle, by a dynamical theory such as developed, *e.g.*, by Marder [15], by Stavans *et al.* [16], and by Flyvbjerg [17]. Nevertheless, comparison of dynamical experimental data to the static Poisson-Voronoi

diagrams may give a clue as to what elements should enter into a correct dynamical theory.

Discrepancies between the experimental data and the predictions of a Poisson-Voronoi diagram are often interpreted as the consequence of some physical, biological (see *e.g.* [18]), or other action that one would like to identify. It is therefore of importance that the properties of Poisson-Voronoi diagrams be understood as well as possible.

In field theory ‘pure’ Voronoi diagrams play a role since attempts were made [19, 20] to construct a theory on a lattice of randomly located sites. The Voronoi construction then again determines the nearest neighbor pairs among the sites. Such a random lattice has the advantage of being statistically invariant under arbitrary translations and rotations while still preserving a short-distance cutoff. Field theory provides still another reason for interest in Voronoi diagrams: Godrèche *et al.* [21] showed that the ensemble of cells with trivalent vertices in a plane is related to the problem of counting planar Feynman diagrams (‘Feynman foam’) of a field theory with cubic interaction.

## 1.1 Statistics of planar Poisson-Voronoi cells

Planar Poisson-Voronoi diagrams are characterized by their statistical properties. The first and foremost of these, whether in experimental studies, in theoretical work, or in simulations, is the probability  $p_n$  that a cell have  $n$  sides. Others include the average number  $\langle n \rangle$  of cell sides; the average area  $A_n$  of an  $n$ -sided cell; the statistics of the cell perimeter and of its angles; and correlations between neighboring or nearby cells. The topology of the plane imposes that  $\langle n \rangle = 6$ . One may readily obtain several of the other statistical properties analytically by expressing them as integrals on the positions of the point particles [22, 3]. The calculation of the fraction  $p_n$ , however, has still defied solution [20, 3]. The distribution  $p_n$  peaks at  $n = 6$  and falls off rapidly for large  $n$ . For  $n = 3$  Hayen and Quine [23] have expressed  $p_3$  as a fivefold integral which they evaluated numerically to seven decimal places. For  $n = 4, 5, \dots$  the values for  $p_n$  obtained from Monte Carlo simulations have been tabulated. All Monte Carlo studies cited in reference [3] (see *e.g.* [24, 25, 26]) were restricted to  $n \lesssim 14$ , which is the largest number of sides observed in simulations by conventional algorithms: one has  $p_{14} \approx 10^{-6}$ .

As early as 1984 Drouffe and Itzykson [27], by a then novel method, estimated the probability  $p_n$  for  $n$  up to 50 by Monte Carlo simulations. This work is still today the main reference for the large  $n$  behavior. The presently known values of  $p_n$  have their uncertainty typically in the third digit for  $4 \leq n \lesssim 10$  and in the second digit for  $10 \lesssim n \lesssim 16$ . Very large cells with  $n$  values higher than 60 were observed by Lauritsen *et al.* [28] in simulations of a ‘Voronoi cell Hamiltonian’ which, although of uncertain physical relevance, is interesting because it favors the appearance many-sided

cells; this procedure does not, however, provide estimates for  $p_n$ .

In view of its importance as a system of reference, we study in this work the planar Poisson-Voronoi diagram. We obtain a great many new analytic results for  $n$ -sided cells in an asymptotic expansion for large  $n$ .

Expansions around idealized models are common in physics and as such our approach needs no specific justification. The fact that  $n = \infty$  is not physically accessible in any realization known of today, is of no concern, since the expansion goes into the physical domain  $n < \infty$ . After the work is done, we will point out the relevance of our results for finite  $n$ . In particular, the vast body of experimental and numerical results has led to two empirical laws that are satisfied to a good extent by many systems: Lewis' law [2] and Aboav's law [29]. The first one will be discussed at the end of this paper; the second one will be investigated separately [30] as an extension of our present work.

An announcement of part of our results, including a concise indication of their derivation, has appeared earlier [31]. Here we present the full calculation together with many more results. This is Part I of a series of two papers. It presents our analytic work and the results that derive immediately from it. In Part II [32] we will develop arguments which, with the benefit of hindsight, will help to understand our results heuristically and to employ them in other applications.

## 1.2 Method

We consider a square  $L \times L$  subdomain of the plane  $\mathbb{R}^2$ . In this domain  $N$  point particles are placed at random positions, chosen independently and with uniform probability. We consider the limit  $N, L \rightarrow \infty$  with the particle density  $N/L^2 = \lambda$  fixed. In mathematical terminology this is the *Poisson point process of intensity  $\lambda$*  in the plane. The density  $\lambda$  may be trivially scaled away. The Voronoi cells of this point particle system are necessarily convex polygons; their number of sides  $n$  may be any integer  $\geq 3$ .

We ask about the statistical properties of the Voronoi cell of an arbitrarily selected particle, whose position is taken as the origin of the coordinate system. Let the other particles, indexed by a label  $a$ , have position vectors  $2\mathbf{R}_a$ . The perpendicular bisectors of these vectors pass through the *midpoints*  $\mathbf{R}_a$ , and the line segments that constitute the cell sides lie on these bisectors. Out of the full set of bisectors only a finite number  $\nu$  that pass sufficiently close to the origin will actually contribute a segment to the perimeter of the cell containing the origin. We denote these 'contributing' midpoints as  $\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_\nu$ . They determine the positions of the cell vertices, which we indicate by  $\mathbf{S}_1, \mathbf{S}_2, \dots, \mathbf{S}_\nu$ . The angles between two successive midpoint vectors  $\mathbf{R}_{m-1}$  and  $\mathbf{R}_m$  [vertex vectors  $\mathbf{S}_{m-1}$  and  $\mathbf{S}_m$ ] will be denoted as  $\xi_m$

[as  $\eta_m$ ]. The probability  $p_n$  that we have  $\nu = n$  for a given  $n$ , *i.e.*, that the selected Voronoi cell have exactly  $n$  sides, can be expressed as a  $2n$ -dimensional integral on  $\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_n$ .

We evaluate this integral for asymptotically large  $n$  by a multidimensional steepest descent method. Although the principle of this approach is simple, its execution is beset by numerous interdependent complications. The necessity to deal with these is the reason for the length of this paper. We were initially guided by the idea of describing the perimeter in terms of a Markov process as a function of the polar angle; the Markov aspect of our description will, however, stay in the background and we will not seek to elaborate on it.

We write  $p_n = \text{Tr } e^{-\mathbb{H}}$ , where  $\mathbb{H}$  is a ‘Hamiltonian’ of purely geometrical origin and  $\text{Tr}$  stands for the integrations. For a suitably chosen ‘free’ or ‘noninteracting’ Hamiltonian  $\mathbb{H}_0$  we set  $\mathbb{H} = \mathbb{H}_0 + \mathbb{V}$  and write

$$p_n = (\text{Tr } e^{-\mathbb{H}_0}) \langle e^{-\mathbb{V}} \rangle_0, \quad (1.1)$$

where  $\langle \dots \rangle_0$  denotes the average with respect to  $e^{-\mathbb{H}_0}$ . We will refer to the two factors,  $\text{Tr } e^{-\mathbb{H}_0}$  and  $\langle e^{-\mathbb{V}} \rangle_0$ , as the ‘noninteracting’ (or ‘free’) and the ‘interacting’ problem, respectively. For clarity of presentation, rather than for technical reasons, we study them separately.

In the course of our analysis the appropriate  $\mathbb{H}_0$  appears to be a sum of noninteracting terms in the angles  $\xi_m$  and  $\eta_m$ : for  $n \rightarrow \infty$  these variables decouple. The evaluation of  $\text{Tr } e^{-\mathbb{H}_0}$  amounts to a  $2n$ -dimensional steepest descent problem in the positive orthant (*i.e.* the subspace with all  $\xi_m, \eta_m > 0$ ), with contributions concentrated in a boundary layer of width of order  $n^{-1}$ . The answer can be obtained exactly up to corrections that vanish exponentially as  $n \rightarrow \infty$ .

The result obtained for  $\text{Tr } e^{-\mathbb{H}_0}$  is meaningful only if we are able to show that the calculation of  $\langle e^{-\mathbb{V}} \rangle_0$  actually leads to a controlled perturbation series in negative powers of  $n$ . The expression for the ‘interaction’  $\mathbb{V}$  is so opaque that *a priori* it is not clear whether it lends itself to any kind of large  $n$  expansion, let alone how it will contribute to the  $n$  dependence of the final result.

We overcome this problem by considering the perimeter segments as a set of vectors  $\mathbf{S}_m - \mathbf{S}_{m-1}$  which, by analogy to the theory of elasticity, we expand about their values in a regular  $n$ -sided polygon. An analysis of considerable difficulty shows that we may write  $\mathbb{V} = \mathbb{V}_1 + n^{-\frac{1}{2}}\mathbb{V}_2$ , where  $\mathbb{V}_1$  is a quadratic form in terms of the ‘elastic deviations’, and  $n^{-\frac{1}{2}}\mathbb{V}_2$  collects together a great number of higher order terms. The maximum of  $e^{-\mathbb{V}_1}$  is located in the positive orthant, *inside* the boundary layer identified in the free problem. The ‘elastic’ interaction  $\mathbb{V}_1$  has a coupling constant proportional to  $n^{-1}$ , but is long-ranged along the perimeter in the same way as a one-dimensional

Coulomb interaction. There is a full separation of spatial scales:  $\mathbb{H}_0$  is a Hamiltonian of independent ‘microscopic’ variables; and  $\mathbb{V}_1$  has its weight concentrated on an effectively finite number of wavelengths at the scale of the cell.

The integrations in the interacting problem involve the confluence of the boundary layer maximum and the maximum of  $e^{-\mathbb{V}_1}$ . They can be dealt with via a Hubbard-Stratonovich transformation. Due to the long-rangedness of the interaction and the weakness of the coupling, an expansion in negative powers of  $n$  becomes possible. In the end we find that the leading order term in the interacting problem is equal to a finite constant:  $\langle e^{-\mathbb{V}} \rangle_0 \simeq C$  as  $n \rightarrow \infty$ .

Various technical obstacles surge in the course of our analysis. To conclude this subsection, we mention three of them.

First, in previous approaches [27, 33, 34] the domain of integration in phase space was always defined in terms of a set of  $n$  inequalities. Here we choose variables of integration such that it becomes fully explicit. After suitable transformation the integration bears on only one radial variable and  $2n - 1$  angles. The radial integration can be carried out exactly and the remaining angular integrations constitute the true problem. There is no single set of angular variables in terms of which the integrand takes a simple form.

Secondly, the angular variables satisfy sum rules stemming from various different origins. It is indispensable to strictly keep track of each of them. To that end we denote them as ‘geometrical sum rules,’ a ‘gauge condition,’ and a ‘no-spiral’ constraint; their meanings will be explained as they are encountered.

Thirdly, the asymptotic expansion for  $n \rightarrow \infty$  requires, as usually, that we postulate at the outset the appropriate scaling with  $n$  for the variables of integration. But since  $n$  is also the typical number of terms in many summations that occur, our scaling analysis has to deal with these sums, too. Some remarkable cases of canceling leading orders need particular attention.

Throughout this work we use only those methods of mathematical physics that ordinarily lead to exact results.

### 1.3 Results

We obtain a detailed understanding of the statistical properties of the  $n$ -sided Voronoi cell when  $n$  is large. The main qualitative results are summarized as follows.

- (i) The probability  $p_n$  for a Voronoi cell to have  $n$  sides behaves asymp-

totically as

$$p_n = \frac{C}{4\pi^2} \frac{(8\pi^2)^n}{(2n)!} [1 + \mathcal{O}(n^{-\frac{1}{2}})], \quad n \rightarrow \infty, \quad (1.2)$$

where

$$\begin{aligned} C &= \prod_{q=1}^{\infty} \left(1 - \frac{1}{q^2} + \frac{4}{q^4}\right)^{-1} \\ &= 0.344\,347\dots \end{aligned} \quad (1.3)$$

and where  $\mathcal{O}(n^d)$  denotes a term which for  $n \rightarrow \infty$  is to leading order proportional to  $n^d$ . Alternatively this may be written as

$$\log p_n = -2n \log n + n \log(2\pi^2 e^2) - \frac{1}{2} \log(2^6 \pi^5 C^{-2} n) + \mathcal{O}(n^{-\frac{1}{2}}). \quad (1.4)$$

In (1.2) the factors  $(4\pi^2)^{-1}(8\pi^2)^n/(2n)!$  and  $C$  originate from the free and the interacting problem, respectively, *i.e.*, from the factors  $\text{Tr } e^{-\mathbb{H}_0}$  and  $\langle e^{-\mathbb{V}} \rangle_0$  in (1.1). The constant  $C$  carries a great deal of meaning. Its factor with index  $q$  stems from the ‘elastic’ modes with wavelength  $2\pi/q$ . Hence the contributions to  $C$  are seen to come essentially from the large scale, long wavelength modes. With the decrease of scale they damp out as  $\sim 1/q^2$ .

The remaining results, (ii)-(vii) below, arise as byproducts of the calculation of  $p_n$ . They have, nevertheless, an independent interest equal to that of  $p_n$ , if not greater.

(ii) As shown in figure 1, we write  $\xi_m$  for the angle by which the perimeter turns at the  $m$ th vertex (it is also the angle between two successive midpoint vectors); and  $\eta_m$  for the angle by which it advances as it passes from the  $m$ th to the  $(m+1)$ th vertex. The  $\xi_m$  are positive and add up to  $2\pi$ , and the  $\eta_m$  similarly. Because of these sum rules they have a weak ‘background’ correlation which in the limit  $n \rightarrow \infty$  vanishes as  $\sim n^{-1}$  (here and henceforth, the sign  $\sim$  denotes asymptotic proportionality). In that limit the  $\xi_m$  and  $\eta_m$  become independent random variables with the  $\xi_m$  identically distributed according to the law

$$u(\xi) = \frac{n^2 \xi}{\pi^2} \exp\left(-\frac{n\xi}{\pi}\right) \quad (1.5)$$

and the  $\eta_m$  according to

$$v(\eta) = \frac{n}{2\pi} \exp\left(-\frac{n\eta}{2\pi}\right). \quad (1.6)$$

These distributions have the averages  $\bar{\xi} = \bar{\eta} = 2\pi n^{-1}$ , as had to be the case. The background correlations have contributions not only from the sum rules, but also from the interaction  $\mathbb{V}$ , which renders them nontrivial. We find the




Figure 1: Heavy line: the perimeter of the Voronoi cell of the particle in the origin. Dashed and dotted lines connect the origin to the midpoints and vertices, respectively. Right angles have been marked. The figure defines the angles  $\xi_m$  and  $\eta_m$  discussed in this section as well as  $\beta_m$  and  $\gamma_m$  to be introduced in section 2.

expressions for these correlations in sections 7 and 8 but will not state the results here.

(iii) Let the mean value of the midpoint distances  $R_1, R_2, \dots, R_n$  be denoted by  $R_{\text{av}} = n^{-1} \sum_{m=1}^n R_m$  and let

$$R_{\text{av}} = R_c + \delta R_{\text{av}}, \quad R_c = \left( \frac{n}{4\pi\lambda} \right)^{\frac{1}{2}}. \quad (1.7)$$

In the limit  $n \rightarrow \infty$  we have  $\langle R_{\text{av}} \rangle = R_c$  and  $\delta R_{\text{av}}$  has the Gaussian distribution

$$\text{p}(\delta R_{\text{av}}) = (8\lambda)^{\frac{1}{2}} \exp(-8\pi\lambda \delta R_{\text{av}}^2). \quad (1.8)$$

*i.e.*, its width is independent of  $n$ .

(iv) The fluctuations of the  $R_m$  around their average  $R_{\text{av}}$  have the  $n$  independent variance

$$\langle (R_m - R_{\text{av}})^2 \rangle \simeq \frac{3C_{\text{av}}}{4\pi\lambda}, \quad n \rightarrow \infty, \quad (1.9)$$

with  $C_{\text{av}} = \sum_{q=1}^{\infty} q^{-4} (1 - q^{-2} + 4q^{-4})^{-1} = 0.333418\dots$ . Here and henceforth  $\langle \dots \rangle$  indicates an average over all Voronoi diagrams and  $\simeq$  stands for asymptotic equality. Equations (1.7) and (1.9) together tell us that the  $R_m$  all scale as  $n^{\frac{1}{2}}$  and that the line segments linking the midpoints  $\mathbf{R}_m$  run mostly inside an imaginary tube of width  $\sim n^0$  along a circle of radius  $R_c = (4\pi\lambda)^{-\frac{1}{2}} n^{\frac{1}{2}}$ . The cell *perimeter* is the polygon having not the  $\mathbf{R}_m$ , but the  $\mathbf{S}_m$  as vertices;



however, we show that the distance between the two polygons vanishes as  $\sim n^{-\frac{1}{2}}$ . Therefore the cell perimeter approaches a circle as  $n \rightarrow \infty$ .

This approach to circularity when the *number of sides* tends to infinity is distinct from other approaches to circularity, such as occur for cells whose *area* tends to infinity (see Hug, Reitzner, and Schneider [35]), or whose *in-radius* (*i.e.*, the radius of the largest inscribed circle centered at the origin) tends to infinity (see Calka [36]; Calka and Schreiber [37]). In Part II [32] we compare the two types of limits in greater detail.

(v) Let  $2\pi mn^{-1} = \phi$  and  $R_m = R(\phi)$ . In the limit  $n \rightarrow \infty$  the variable  $\phi$  becomes continuous and may be identified, up to corrections that vanish as  $n^{-\frac{1}{2}}$ , with the polar angle of the perimeter. We set

$$R(\phi) = R_c + \delta R(\phi). \quad (1.10)$$

and refer to  $\delta R(\phi)$  as the ‘excess midpoint distance.’ On scales large compared to the angular scale  $2\pi n^{-1}$  of the individual perimeter segments the following holds. The effect of the Poisson point process on the perimeter  $R(\phi)$  may be represented by a stochastic second order differential equation for  $\delta R(\phi)$ ,

$$\frac{d^2 \delta R}{d\phi^2} = F(\phi), \quad 0 < \phi < 2\pi, \quad (1.11)$$

in which  $F(\phi)$  is zero average Gaussian noise of autocorrelation

$$\langle F(\phi)F(\phi') \rangle = 12\lambda \left[ \delta(\phi - \phi') - \frac{1 - \Gamma(\phi - \phi')}{2\pi} \right], \quad (1.12)$$

where  $\Gamma(\phi)$  is a  $2\pi$ -periodic function whose integral on  $[0, 2\pi]$  vanishes. The solution of (1.11) is subject to a periodicity condition and an integral condition, *viz.*

$$\delta R(0) = \delta R(2\pi), \quad \frac{1}{2\pi} \int_0^{2\pi} d\phi \delta R(\phi) = \delta R_{\text{av}}, \quad (1.13)$$

respectively, with  $\delta R_{\text{av}}$  distributed according to (1.8). Hence in the  $n \rightarrow \infty$  limit  $\delta R(\phi)$  is governed by Gaussian statistics. The correlation function (1.12) determines the probability law of the noise  $F(\phi)$  and hence, by equations (1.11), (1.13), and (1.8), of  $\delta R(\phi)$ . The explicit result is given in section 8.3.

(vi) According to Lewis’ empirical law [2, 3] the average area  $A_n$  of an  $n$ -sided cell grows linearly with  $n$ . For Poisson-Voronoi cells, in spite of attempts, this linearity with  $n$  has not so far been proved [38, 39]. Now the results (iii) and (iv) above imply directly that

$$A_n \simeq \pi R_c^2 = \frac{n}{4\lambda}, \quad n \rightarrow \infty. \quad (1.14)$$

Hence we have derived here from first principles that Lewis’ law holds in the asymptotic regime and has coefficient  $\frac{1}{4}$ .

## 2 Writing $p_n$ as a $2n$ -fold angular integral

In this section we start from the basic  $2n$ -fold integral for  $p_n$  and subject it to a sequence of transformations that will cast it in a form amenable to an asymptotic large  $n$  expansion.

### 2.1 Basic integral for $p_n$

An  $n$ -sided cell is completely defined by  $n$  midpoint positions  $\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_n$  (see figure 1). These are uniformly distributed with density  $4\lambda$ , and  $p_n$  may be written directly as a  $2n$ -fold integral on their coordinates [27, 20, 33, 34],

$$p_n = \frac{(4\lambda)^n}{n!} \int d\mathbf{R}_1 \dots d\mathbf{R}_n \chi(\mathbf{R}_1, \dots, \mathbf{R}_n) e^{-4\lambda \mathcal{A}(\mathbf{R}_1, \dots, \mathbf{R}_n)}, \quad (2.1)$$

where the functions  $\chi$  and  $\mathcal{A}$  to be defined shortly enforce two conditions that must be satisfied if the  $n$  points are to define a valid Voronoi cell. In order to state these conditions we use the polar coordinate representation  $\mathbf{R}_m = (R_m, \Phi_m)$ . In the integrand of (2.1) we may set one of the angles, say  $\Phi_n$ , to zero if we compensate by an extra factor  $2\pi$ , and we may order the remaining angles according to

$$\Phi_n = 0 < \Phi_1 < \Phi_2 < \dots < \Phi_{n-1} < 2\pi \quad (2.2)$$

if a compensating factor  $(n-1)!$  is introduced. We will denote the  $n$  angular differences  $\xi_m$  between successive midpoint vectors as

$$\begin{aligned} \xi_m &= \Phi_m - \Phi_{m-1}, & m &= 1, \dots, n-1, \\ \xi_n &= 2\pi - \Phi_{n-1}. \end{aligned} \quad (2.3)$$

The geometry constrains the  $\xi_m$  to the interval  $(0, \pi)$  and they must satisfy the geometrical sum rule

$$\sum_{m=1}^n \xi_m = 2\pi. \quad (2.4)$$

We find it convenient henceforth to scale the distances such that

$$4\lambda = 1 \quad (2.5)$$

during the calculations, and restore this factor in the final results. Symmetrizing in the angular difference variables by introducing a Dirac delta function we obtain from (2.1)

$$p_n = \frac{2\pi}{n} \int_0^\infty R_1 dR_1 \dots R_n dR_n \int_0^\pi d\xi_1 \dots d\xi_n \delta(\xi_1 + \dots + \xi_n - 2\pi) \chi e^{-\mathcal{A}}. \quad (2.6)$$

We will also need the positions  $\mathbf{S}_m = (S_m, \Psi_m)$  of the cell vertices (see figure 1). These are auxiliary variables in the sense that they can be expressed in terms of the  $\xi_\ell$  and  $R_\ell$ . We are now ready to state the two conditions.

*Condition 1.* The first condition is that for  $a \neq 1, \dots, n$  none of the bisectors passing through the midpoints  $\mathbf{R}_a$  have an intersection with the Voronoi cell. This condition is fulfilled if these midpoints stay outside of a region of the plane equal to the union of  $n$  disks centered at  $\frac{1}{2}\mathbf{S}_m$  and of radius  $\frac{1}{2}S_m$  [40]. If this region has area  $\mathcal{A}$ , then the probability for it to be empty of midpoints (other than the  $n$  that determine the cell perimeter) is equal to  $\exp(-\mathcal{A})$  times its area. An explicit expression for  $\mathcal{A}$  in terms of the  $R_m$  and  $\xi_m$  may be found by straightforward geometry (see Calka [33, 34]).

*Condition 2.* The second condition defines the domain of integration. It says that each of the  $n$  midpoints  $\mathbf{R}_m$  should have its corresponding perpendicular bisector contribute a nonzero segment to the perimeter. This condition was shown by Calka [33, 34] to read explicitly

$$R_{m-1} \sin \xi_{m+1} + R_{m+1} \sin \xi_m > R_m \sin(\xi_m + \xi_{m+1}), \quad m = 1, \dots, n, \quad (2.7)$$

with the conventions  $R_0 = R_n$ ,  $R_1 = R_{n+1}$ , and  $\xi_1 = \xi_{n+1}$ . The function  $\chi$  in (2.1) and (2.6) is the indicator of the domain in phase space where (2.7) is satisfied.

This fully defines the integral (2.6) for  $p_n$ . In being a sum on  $n$  equivalent interacting objects, *viz.* the midpoint vectors  $\mathbf{R}_m$ , this integral is analogous to a partition function in statistical mechanics. But although interesting, this observation provides no direct key as to how to go about its evaluation.

## 2.2 Angular variables

There is no single set of variables in which the calculations of this work take a simple form. We will, throughout, employ simultaneously different sets of angular variables, to be introduced in this subsection.

The polar angles  $\Psi_m$  of the vertex vectors  $\mathbf{S}_m$  satisfy an ordering analogous to (2.2),

$$0 < \Psi_1 < \Psi_2 < \dots < \Psi_n < 2\pi. \quad (2.8)$$

There is, however, no particular ordering relation between the two sets  $\{\Psi_m \mid m = 1, \dots, n\}$  and  $\{\Phi_m \mid m = 1, \dots, n\}$ . We will not use the  $\{\Psi_m \mid m = 1, \dots, n\}$  any more, but instead, by analogy to equation (2.3), introduce the angular differences

$$\begin{aligned} \eta_m &= \Psi_m - \Psi_{m-1}, & m = 2, \dots, n, \\ \eta_1 &= \Psi_1 - \Psi_n + 2\pi, \end{aligned} \quad (2.9)$$

which are constrained to  $(0, \pi)$  and satisfy the geometrical sum rule

$$\sum_{m=1}^n \eta_m = 2\pi. \quad (2.10)$$




Figure 2: A configuration of midpoints  $\mathbf{R}_m$  and  $\mathbf{R}_{m+1}$  leading to a negative angle  $\gamma_m$ . The relation  $\beta_m + \gamma_m = \xi_m$  remains valid. The negativity of  $\gamma_m$  is accompanied by a counterclockwise outward spiraling of the perimeter. The legend is the same as for figure 1.

Next, as shown in figure 1, the set  $\{\beta_m, \gamma_m \mid m = 1, \dots, n\}$  will represent the angles between a vertex vector and a midpoint vector. They are defined in terms of the  $\Phi_m$  and  $\Psi_m$  by

$$\begin{aligned} \beta_m &= \Psi_m - \Phi_{m-1}, & m &= 2, \dots, n, \\ \beta_1 &= \Psi_1, \\ \gamma_m &= \Phi_m - \Psi_m, & m &= 1, \dots, n. \end{aligned} \quad (2.11)$$

The  $\beta_m$  and  $\gamma_m$  are geometrically constrained to the interval  $(-\frac{\pi}{2}, \frac{\pi}{2})$ , and hence, contrary to the  $\xi_m$  and  $\eta_m$ , may be negative. An example of a negative  $\gamma_m$  is shown in figure 2. Whereas this potential negativity may at this point not seem worthy of much attention, it will be seen in section 5 to have important consequences for the scaling of these angles with  $n$ . Because of the geometrical sum rule

$$\sum_{m=1}^n (\beta_m + \gamma_m) = 2\pi, \quad (2.12)$$

the set  $\{\beta_m, \gamma_m \mid m = 1, \dots, n\}$  contains one redundant angle. Hence, leaving out  $\gamma_n$ , the set of  $2n - 1$  variables

$$\{\beta_1, \gamma_1, \dots, \beta_{n-1}, \gamma_{n-1}, \beta_n\} \quad (2.13)$$

is complete in the sense that, given  $\Phi_0 = 0$ , all the other angles  $\Phi_m$  and  $\Psi_m$  are easily expressed in terms of the angles of the set (2.13).

The set  $\{\xi_m, \eta_m \mid m = 1, \dots, n\}$ , due to the sum rules (2.4) and (2.10), contains two redundant variables and is not complete: it specifies only the angles of the  $\mathbf{R}_m$  between themselves and of the  $\mathbf{S}_m$  between themselves, but not the relative angle between the two vector systems. To fix their relative orientation, one other angle is needed, for which we will take  $\beta_1$ . A second complete set of  $2n - 1$  variables is therefore

$$\{\xi_1, \eta_1, \dots, \xi_{n-1}, \eta_{n-1}, \beta_1\}. \quad (2.14)$$

The sets (2.13) and (2.14) have  $\beta_1$  in common. The transformation relation between their  $2n - 2$  remaining variables is

$$\xi_m = \beta_m + \gamma_m, \quad \eta_m = \gamma_m + \beta_{m+1}, \quad m = 1, \dots, n-1, \quad (2.15)$$

and its nonlocal inverse

$$\begin{aligned} \beta_m &= \beta_1 - \sum_{\ell=1}^{m-1} (\xi_\ell - \eta_\ell), \quad m = 2, \dots, n, \\ \gamma_m &= -\beta_1 + \sum_{\ell=1}^{m-1} (\xi_\ell - \eta_\ell) + \xi_m, \quad m = 1, \dots, n-1. \end{aligned} \quad (2.16)$$

where by definition  $\sum_{\ell=1}^0 = 0$  for any summand. It is easy to verify that this transformation has unit Jacobian. Relations (2.15) are valid also for  $m = n$  with the convention that  $\beta_{n+1} = \beta_1$ ; and relations (2.16) are valid for  $m = 1$  and  $m = n$ , respectively.

## 2.3 Making the domain of integration explicit

### 2.3.1 Rewriting the indicator function

From the elementary observation that

$$R_{m-1} = S_m \cos \beta_m, \quad R_m = S_m \cos \gamma_m \quad (2.17)$$

we deduce the relation

$$\frac{R_m}{R_{m-1}} = \frac{\cos \gamma_m}{\cos \beta_m}, \quad m = 1, \dots, n. \quad (2.18)$$

Hence, when  $R_1, \dots, R_n$  are given,  $\beta_m$  determines  $\gamma_m$  and *vice versa*. We now turn to the inequality (2.7) and divide it by  $R_m \sin \xi_m \sin \xi_{m+1}$ . The ratios of the midpoint distances that then appear may be eliminated: we express them in terms of the angular variables by means of relation (2.18). After

some trigonometry  $\xi_m$  and  $\xi_{m+1}$  cancel out of the inequality, which takes the form  $\tan \gamma_m + \tan \beta_{m+1} > 0$  for  $m = 1, 2, \dots, n$ . Since  $-\frac{\pi}{2} < \gamma_m, \beta_{m+1} < \frac{\pi}{2}$ , it follows that the much simpler and geometrically necessary positivity condition

$$\gamma_m + \beta_{m+1} > 0, \quad m = 1, 2, \dots, n, \quad (2.19)$$

is *equivalent to condition* (2.7). Therefore (2.19) defines the domain of integration in phase space.

### 2.3.2 Transforming from $\xi_m$ to $\beta_m$ and $\gamma_m$

Next, considering the outer integration variables  $R_1, \dots, R_n$  in (2.6) as simple parameters, we will transform the integrals on the angles  $\xi_m$  in two successive steps. First, we consider  $\gamma_m$  as a function of  $\beta_m$  defined by (2.18), so that in view of (2.15a) we have  $\xi_m = \beta_m + \arccos[(R_m/R_{m-1}) \cos \beta_m]$ . We transform from  $\xi_m$  to the new integration variable  $\beta_m$ , taking due account of the Jacobian  $j_m = d\xi_m/d\beta_m$ , which is equal to

$$j_m = \frac{\sin(\beta_m + \gamma_m)}{\cos \beta_m \sin \gamma_m}. \quad (2.20)$$

We order the  $\beta_m$  integrations such that, going from the outermost one inward, we encounter successively those on  $\beta_1, \dots, \beta_n$ . We integrate  $\beta_1$  on its full interval  $(-\frac{\pi}{2}, \frac{\pi}{2})$ . For given  $\beta_1$  equation (2.18) fixes  $\gamma_1$ . Inside the  $\beta_1$  integral, in order to satisfy condition (2.19) for  $m = 1$ , we integrate  $\beta_2$  from  $-\gamma_1$  to  $\frac{\pi}{2}$ . For given  $\beta_2$  equation (2.18) fixes  $\gamma_2$ , and so on. In this way the conditions (2.19) are all satisfied, except the one at the end which requires  $\gamma_n + \beta_1 > 0$  and which we will therefore incorporate below by means of a Heaviside theta function. After all this equation (2.6) becomes

$$\begin{aligned} p_n = & \frac{2\pi}{n} \int_0^\infty R_1 dR_1 \dots R_n dR_n \int_{-\pi/2}^{\pi/2} d\beta_1 \int_{-\gamma_1}^{\pi/2} d\beta_2 \dots \int_{-\gamma_{n-1}}^{\pi/2} d\beta_n \\ & \times \theta(\gamma_n + \beta_1) j_1 j_2 \dots j_n \delta\left(\sum_{m=1}^n (\beta_m + \gamma_m) - 2\pi\right) e^{-\mathcal{A}}, \end{aligned} \quad (2.21)$$

in which  $\gamma_m$ , wherever it occurs, is a function of  $\beta_m$  given by (2.18), and where the indicator  $\chi$  is no longer needed. The passage from (2.6) to (2.21) has broken the symmetry between the  $\beta_m$  and the  $\gamma_m$  and introduced an orientation in the cell perimeter. We have moreover broken the cyclic invariance in the index  $m$  by imposing a fixed order of integration of the angular variables. These symmetry breakings first appeared in an initial approach [41] where we constructed the perimeter segments by a Markov process with transition probabilities between successive vertices. We will not elaborate on this here.

We wish to introduce in (2.21) the  $\gamma_1, \gamma_2, \dots, \gamma_n$  as additional independent variables of integration. One easily verifies that

$$\int_{-\beta_m}^{\pi/2} d\gamma_m \frac{\sin \gamma_m}{\cos \beta_m} R_{m-1} \delta\left(R_m - \frac{\cos \gamma_m}{\cos \beta_m} R_{m-1}\right) = 1. \quad (2.22)$$

We now take the product on  $m$  of the LHS of this expression and insert it in the integrand of (2.21). Still defining [42]

$$T_m = \frac{\sin(\beta_m + \gamma_m)}{\cos^2 \beta_m} \quad (2.23)$$

we find that (2.21) becomes

$$\begin{aligned} p_n &= \frac{2\pi}{n} \int_0^\infty R_1 dR_1 \dots R_n dR_n \int d\beta d\gamma \\ &\times \left[ \prod_{m=1}^n T_m R_{m-1} \delta\left(R_m - \frac{\cos \gamma_m}{\cos \beta_m} R_{m-1}\right) \right] e^{-\mathcal{A}} \end{aligned} \quad (2.24)$$

with our earlier convention  $R_0 \equiv R_n$  and where we employ the shorthand notation

$$\begin{aligned} \int d\beta d\gamma &= \int_{-\pi/2}^{\pi/2} d\beta_1 \int_{-\beta_1}^{\pi/2} d\gamma_1 \int_{-\gamma_1}^{\pi/2} d\beta_2 \int_{-\beta_2}^{\pi/2} d\gamma_2 \dots \int_{-\gamma_{n-1}}^{\pi/2} d\beta_n \int_{-\beta_n}^{\pi/2} d\gamma_n \\ &\times \theta(\gamma_n + \beta_1) \delta\left(\sum_{m=1}^n (\beta_m + \gamma_m) - 2\pi\right). \end{aligned} \quad (2.25)$$

The operator  $\int d\beta d\gamma$  represents the integral on all possible shapes of the Voronoi cell without any reference to its radial scale. The fact that the domain of integration appears here fully explicitly constitutes the main achievement of this subsection.

An expression for  $\mathcal{A}$  in terms of the variables of integration  $R_m, \beta_m$ , and  $\gamma_m$  may be obtained directly from its geometrical definition as the area of the union of  $n$  disks. After the necessary rewriting one finds

$$\mathcal{A} = \frac{1}{4} \sum_{m=1}^n R_m^2 [a(\gamma_m) + a(\beta_{m+1})], \quad (2.26)$$

where  $a$  is defined by

$$a(\omega) = \omega + \tan \omega + \omega \tan^2 \omega, \quad (2.27)$$

which is equivalent, and close in form, to the expression used by Drouffe and Itzykson [27] in their simulations.

## 2.4 Radial integrations

### 2.4.1 Introducing the mean midpoint distance $R_{\text{av}}$

The average value of the  $n$  midpoint distances  $R_1, R_2, \dots, R_n$  will be denoted by

$$R_{\text{av}} = n^{-1} \sum_{m=1}^n R_m. \quad (2.28)$$

The circle of radius  $R_{\text{av}}$  will act as a circle of reference and we proceed on the hypothesis that the area difference  $\mathcal{A} - \pi R_{\text{av}}^2$  will be small in a sense still to be determined [43]. Let us now normalize the  $R_m$  according to

$$R_m = R_{\text{av}} \rho_m, \quad (2.29)$$

so that because of (2.28) the ‘reduced midpoint distances’  $\rho_m$  satisfy

$$n^{-1} \sum_{m=1}^n \rho_m = 1. \quad (2.30)$$

Instead of (2.29) other choices of normalization would have been possible. Therefore we will henceforth refer to (2.30) and equivalent relations as the ‘gauge’ condition; this in order to distinguish it from other sum rules that play a role.

We insert a factor  $\delta(R_{\text{av}} - n^{-1} \sum_{m=1}^n R_m)$  in the integrand of the RHS of (2.24) and apply an extra integration on  $R_{\text{av}}$  to it. The radial integrations then take the form

$$\begin{aligned} \int_0^\infty dR_1 \dots dR_n &= \int_0^\infty dR_{\text{av}} \int_0^\infty dR_1 \dots dR_n \delta(R_{\text{av}} - n^{-1} \sum_{m=1}^n R_m) \\ &= \int_0^\infty dR_{\text{av}} R_{\text{av}}^{n-1} \int_0^\infty d\rho_1 \dots d\rho_n \delta(1 - n^{-1} \sum_{m=1}^n \rho_m) \end{aligned} \quad (2.31)$$

and as a result (2.24) becomes

$$\begin{aligned} p_n &= \frac{2\pi}{n} \int_0^\infty dR_{\text{av}} R_{\text{av}}^{2n-1} \int_0^\infty d\rho_1 \dots d\rho_n \delta(1 - n^{-1} \sum_{m=1}^n \rho_m) \\ &\quad \times \int d\beta d\gamma \left[ \prod_{m=1}^n \delta(\rho_m - \frac{\cos \gamma_m}{\cos \beta_m} \rho_{m-1}) \right] \left[ \prod_{m=1}^n \rho_m^2 T_m \right] e^{-\mathcal{A}}, \end{aligned} \quad (2.32)$$

where by convention  $\rho_0 = \rho_n$  and in which the angular integrations are defined by (2.25).



### 2.4.2 Integrating on the $R_m$ at fixed $R_{av}$

The  $n - 1$  integrations on  $\rho_1, \dots, \rho_{n-1}$  may be carried out successively and, due to the delta functions in the integrand, cause  $\rho_m$ , wherever it occurs, to be replaced by

$$\rho_m = \frac{\cos \gamma_m \cos \gamma_{m-1} \dots \cos \gamma_1}{\cos \beta_m \cos \beta_{m-1} \dots \cos \beta_1} \rho_n, \quad m = 1, \dots, n-1. \quad (2.33)$$

There remains one delta function inside the product on  $m$  in (2.32). We rewrite it as

$$\delta\left(\rho_n - \frac{\cos \gamma_1 \dots \cos \gamma_n}{\cos \beta_1 \dots \cos \beta_n} \rho_n\right) = \frac{\delta(G)}{2\pi\rho_n}, \quad (2.34)$$

where

$$G = \frac{1}{2\pi} \sum_{m=1}^n (\log \cos \gamma_m - \log \cos \beta_m). \quad (2.35)$$

The integration on  $\rho_n$  in (2.32), finally, is easily performed due to the delta function constraint in the first line of that equation and leads to the cancellation of the factor  $\rho_n^{-1}$  in (2.34). The result is that (2.32) becomes

$$p_n = \frac{1}{n} \int d\beta d\gamma \delta(G) \left[ \prod_{m=1}^n \rho_m^2 T_m \right] \int_0^\infty dR_{av} R_{av}^{2n-1} e^{-A}, \quad (2.36)$$

in which now the reduced midpoint distances  $\rho_1, \dots, \rho_n$  should be viewed as functions of the integration variables determined by the  $n - 1$  equations (2.33) together with condition (2.30).

It appears from (2.36) that the  $2n$  angles of integration satisfy a nontrivial supplementary constraint  $G = 0$ . This constraint is easily interpreted: for an arbitrary set of angles  $(\beta, \gamma) \equiv \{\beta_m, \gamma_m \mid m = 1, \dots, n\}$  the perimeter will spiral instead of close onto itself after a full turn of  $2\pi$ ; the factor  $\delta(G)$  enforces its closure. We will refer to it as the ‘no-spiral’ constraint. We see that equation (2.34) imposes that (2.33) hold for  $m = n$ . Hence requiring the validity of (2.33) for all  $m = 1, \dots, n$  is another way of incorporating the no-spiral constraint.

In what follows we will extensively use an auxiliary variable  $\tau_m$  representing the excess length of the reduced midpoint distances  $\rho_m$  over their mean value,

$$\rho_m = 1 + \tau_m, \quad m = 1, \dots, n. \quad (2.37)$$

From (2.30) one deduces that they satisfy

$$\sum_{m=1}^n \tau_m = 0, \quad (2.38)$$

which is an alternative expression for the gauge condition.

### 2.4.3 Integrating on the mean midpoint distance $R_{\text{av}}$

Equations (2.29), (2.30), and (2.33) allow us to express  $R_m$  as  $R_{\text{av}}$  times an  $m$  dependent function of the angles which, if desired, may be made explicit. It is then clear from (2.26) that  $\mathcal{A}$  is  $R_{\text{av}}^2$  times a function only of the angles. Bearing in mind that we expect  $\mathcal{A}$  to be close to the area  $\pi R_{\text{av}}^2$  enclosed by the circle of reference, and anticipating later developments, we write

$$\mathcal{A} = \pi R_{\text{av}}^2 (1 + n^{-1}V), \quad (2.39)$$

which defines  $V$ . Integrating on  $R_{\text{av}}$  then converts (2.36) into

$$p_n = \frac{(n-1)!}{2n} \int d\beta d\gamma \delta(G) \left[ \prod_{m=1}^n \rho_m^2 T_m \right] [\pi(1 + n^{-1}V)]^{-n}. \quad (2.40)$$

The integrations that remain in the RHS of (2.40) bear exclusively on the angular variables, *i.e.* on the shape of a cell with average midpoint distance scaled to unity.

For completeness and for use in section 6 we now determine  $V$  explicitly. Substituting (2.37) in (2.29) and subsequently (2.29) and (2.27) in (2.26), comparing to (2.39), and still using the geometrical sum rule (2.12), we obtain

$$\begin{aligned} V(\beta, \gamma) = & \frac{n}{4\pi} \sum_{m=1}^n (1 + \tau_m)^2 \left[ \tan \gamma_m - \gamma_m + \tan \beta_{m+1} - \beta_{m+1} \right. \\ & \left. + \gamma_m \tan^2 \gamma_m + \beta_{m+1} \tan^2 \beta_{m+1} \right] \\ & + \frac{n}{2\pi} \sum_{m=1}^n (2\tau_m + \tau_m^2)(\gamma_m + \beta_{m+1}). \end{aligned} \quad (2.41)$$

We note that (2.41) is antisymmetric under a change of sign of all the angles.

### 2.4.4 The probability distribution of $R_{\text{av}}$

In this subsection we sidestep and make an observation that will be exploited only in section 9. Although the outcome (2.40) of the integral on  $R_{\text{av}}$  is known, something more may be learned by applying a steepest descent analysis to it. Assuming, as is suggested by the notation  $n^{-1}V$  in (2.39) and as will be confirmed in section 6.3.2, that  $V(\beta, \gamma)$  remains of order  $n^0$  when  $n \rightarrow \infty$ , we then find that the integral on  $R_{\text{av}}$  draws its main contribution from a Gaussian centered at the saddle point  $R_{\text{av},c}$  and having a variance  $\sigma_{\text{av},c}^2$  given by

$$R_{\text{av},c}(\beta, \gamma) = \left[ \frac{2n-1}{2\pi(1+n^{-1}V)} \right]^{\frac{1}{2}} = \left( \frac{n}{\pi} \right)^{\frac{1}{2}} (1 + \mathcal{O}(n^{-1})), \quad (2.42)$$

$$\sigma_{\text{av},c}^2(\beta, \gamma) = [4\pi(1+n^{-1}V)]^{-1} = \frac{1}{4\pi} (1 + \mathcal{O}(n^{-1})). \quad (2.43)$$

The results (2.42) and (2.43) are to leading order in  $n$  independent of  $(\beta, \gamma)$ . The  $\mathcal{O}(n^{-1})$  corrections in these equations do depend on  $(\beta, \gamma)$  but will play no role in this work. Hence, provided we confirm the scaling of  $V$ , we have found here that the  $n$ -sided Voronoi cell has a characteristic radial scale of order  $n^{\frac{1}{2}}$  [44]. Furthermore, if we define

$$R_c = \pi^{-\frac{1}{2}} n^{\frac{1}{2}}, \quad R_{\text{av}} = R_c(1 + n^{-\frac{1}{2}} r_{\text{av}}), \quad (2.44)$$

then it follows from the above that in the limit  $n \rightarrow \infty$  the variable  $r_{\text{av}}$  has the probability distribution

$$p_{\text{av}}(r_{\text{av}}) = 2^{\frac{1}{2}} \pi^{-\frac{1}{2}} \exp(-2r_{\text{av}}^2) + \mathcal{O}(n^{-\frac{1}{2}}). \quad (2.45)$$

With  $\delta R_{\text{av}} = n^{-\frac{1}{2}} R_c r_{\text{av}}$  and with the midpoint density  $4\lambda$  restored this becomes equation (1.8) of the introduction.

## 2.5 Passing to the angles of integration $\xi_m$ and $\eta_m$

So far the  $\beta_m$  and  $\gamma_m$  [the set (2.13)] have been the variables most convenient to work with. However, in order to prepare for the large  $n$  expansion, it is now necessary that we pass to the set of variables (2.14) consisting of the  $\xi_m$  and  $\eta_m$  as well as of  $\beta_1$ . Employing the notation  $(\xi, \eta) = \{\xi_m, \eta_m \mid m = 1, \dots, n\}$ ,  $\xi = \{\xi_m \mid m = 1, \dots, n\}$ , and  $\eta = \{\eta_m \mid m = 1, \dots, n\}$ , we now proceed as follows.

By writing  $G(\xi, \eta; \beta_1)$  we indicate explicitly that we want to consider  $G$  as a function of  $\xi$ ,  $\eta$ , and  $\beta_1$ . The prime on  $G'$  stands for differentiation with respect to  $\beta_1$ . Let furthermore  $\beta_*(\xi, \eta)$  denote the solution of the no-spiral constraint

$$G(\xi, \eta; \beta_*) = 0. \quad (2.46)$$

In (2.40) one easily converts the integration on  $(\beta, \gamma)$ , defined in (2.25), to one on the new variables  $(\xi, \eta)$  by shifting the integration intervals appropriately (except the outermost one, for  $\beta_1$ , which remains unchanged) and employing the fact that  $\gamma_n + \beta_1 = 2\pi - \sum_{m=1}^{n-1} \eta_m$ . With the aid of the identity

$$\theta(2\pi - \sum_{m=1}^{n-1} \eta_m) = \int_0^{2\pi} d\eta_n \delta(\sum_{m=1}^n \eta_m - 2\pi) \quad (2.47)$$

and using the occasion to rearrange some factors in the integrand we recast (2.40) in the form

$$\begin{aligned} p_n &= \frac{(n-1)!}{2n} \int_{-\pi/2}^{\pi/2} d\beta_1 \int d\xi d\eta \delta(\sum_{m=1}^n \xi_m - 2\pi) \delta(\sum_{m=1}^n \eta_m - 2\pi) \\ &\quad \times \frac{\delta(\beta_1 - \beta_*)}{|G'(\xi, \eta; \beta_*)|} \left[ \prod_{m=1}^n \rho_m^2 T_m \right] [\pi(1 + n^{-1}V)]^{-n} \end{aligned} \quad (2.48)$$

where

$$\begin{aligned} \int d\xi d\eta &= \int_0^{\pi/2+\beta_1} d\xi_1 \int_0^{\pi/2+\gamma_1} d\eta_1 \int_0^{\pi/2+\beta_2} d\xi_2 \dots \\ &\dots \int_0^{\pi/2+\gamma_{n-1}} d\eta_{n-1} \int_0^{\pi/2+\beta_n} d\xi_n \int_0^{2\pi} d\eta_n. \end{aligned} \quad (2.49)$$

The notation is hybrid; the variables  $\gamma_1, \beta_2, \gamma_3, \dots, \beta_n$  in the upper integration limits in (2.49) should be viewed as functions of  $\xi, \eta$ , and  $\beta_1$  given by (2.16). Each limit of integration depends only on the variables of more outward integrations, as of course it should be.

We note that it is not possible at this stage to carry out the integral on  $\beta_1$ , since  $\beta_*$  depends on  $\xi$  and  $\eta$  and the integration limits of the integrals  $\int d\xi d\eta$  in turn depend on  $\beta_1$ .

## 2.6 Free and interacting problem

### 2.6.1 Defining a free problem and an interaction

In order to relax the constraints (2.4) and (2.10) enforced by the first two delta functions in (2.48), we will exploit the integral representation  $\delta(X) = \int_{c-i\infty}^{c+i\infty} ds \exp(-sX)$ , valid for any constant  $c$ . Upon introducing a parameter  $\kappa$  which at this point is arbitrary and scaling conveniently with  $n$  we can write

$$\begin{aligned} \delta(\Sigma_m \xi_m - 2\pi) \delta(\Sigma_m \eta_m - 2\pi) &= \\ &= \delta(\kappa \Sigma_m \xi_m + (1 - \kappa) \Sigma_m \eta_m - 2\pi) \delta(\Sigma_m \xi_m - \Sigma_m \eta_m) \\ &= \left(\frac{n}{2\pi i}\right)^2 \int_{c-i\infty}^{c+i\infty} ds \int_{-i\infty}^{i\infty} dt e^{2\pi n s - n s [\kappa \Sigma_m \xi_m - (1-\kappa) \Sigma_m \eta_m] - n t (\Sigma_m \xi_m - \Sigma_m \eta_m)}, \end{aligned} \quad (2.50)$$

where we choose  $c > 0$ . Substituting (2.50) in (2.48) we find [45]

$$\begin{aligned} p_n &= \left(\frac{n}{2\pi i}\right)^2 \int_{c-i\infty}^{c+i\infty} ds \int_{-i\infty}^{i\infty} dt \int_{-\pi/2}^{\pi/2} d\beta_1 e^{2\pi n s} \int d\xi d\eta \delta(\beta_1 - \beta_*) e^{-\mathbb{H}(s,t)} \\ &\equiv \text{Tr} e^{-\mathbb{H}(s,t)}, \end{aligned} \quad (2.51)$$

in which the last line defines the  $\text{Tr}$  operator, and the ‘Hamiltonian’  $\mathbb{H}$  is given as the sum of a ‘free’ Hamiltonian  $\mathbb{H}_0$  and an ‘interaction’  $\mathbb{V}$ ,

$$\mathbb{H}(s, t) = \mathbb{H}_0(s, t) + \mathbb{V}, \quad (2.52)$$

where

$$e^{-\mathbb{H}_0(s,t)} = \pi^{-n} \left[ \prod_{m=1}^n \xi_m e^{-ns[\kappa\xi_m + (1-\kappa)\eta_m] - nt(\xi_m - \eta_m)} \right], \quad (2.53)$$

$$e^{-\mathbb{V}} = |G'(\xi, \eta; \beta_*)|^{-1} \left[ \prod_{m=1}^n \rho_m^2 T_m \xi_m^{-1} \right] (1 + n^{-1}V)^{-n}. \quad (2.54)$$

The factor  $\xi_m$  inserted in (2.53) and compensated in (2.54) serves to ensure that  $T_m \xi_m^{-1}$  has a finite limit as  $\xi_m \rightarrow 0$ . The Hamiltonian  $\mathbb{H}_0$  defined by (2.53) is obviously is sum of  $2n$  independent single-variable terms. However, the reason why the splitting (2.52) should be exactly as defined by (2.53) and (2.54), and not something else, will have to be borne out by the calculation.

With equations (2.51)-(2.54) we have achieved the purpose of this section. They have nothing of the simple elegance of other model Hamiltonians in statistical mechanics and are at first sight hardly suggestive of any particular analytical approach. However, the work of this section has served to prepare the ground for what follows.

So far our calculation is valid for all finite  $n = 3, 4, \dots$ . The initial problem (2.1) is exactly equivalent to problem (2.51). However, we cannot continue beyond the present point: it is impossible to carry out the integral on  $\beta_1$  in (2.51) since  $\beta_*$  depends on the  $\xi_m$  and  $\eta_m$  whose upper integration limits in turn depend on  $\beta_1$ . This entanglement will be lifted in section 3 as soon as we start considering the limit  $n \rightarrow \infty$ .

### 2.6.2 Splitting the calculation

Ensuant on equation (2.52) we have the formal factorization

$$\text{Tr } e^{-\mathbb{H}} = (\text{Tr } e^{-\mathbb{H}_0}) \langle e^{-\mathbb{V}} \rangle_0, \quad (2.55)$$

whence follows the splitting

$$\log p_n = \log p_n^{(0)} + \log \langle e^{-\mathbb{V}} \rangle_0, \quad (2.56)$$

where we employ the alternative notation  $p_n^{(0)} = \text{Tr } e^{-\mathbb{H}_0}$ . We will refer to the calculation of  $p_n^{(0)}$  as the ‘free’ problem and to that of  $\langle e^{-\mathbb{V}} \rangle_0$  as the ‘interacting problem.’ Although technically not strictly necessary, it will greatly enhance the clarity of presentation to treat these problems in two successive steps.

In the course of our work we will also be interested in correlation functions of the angles. We write  $\langle F \rangle_0$  for the ‘noninteracting’ average of any function  $F$  of the angular coordinates, *i.e.*,

$$\langle F \rangle_0 = \frac{\text{Tr } F e^{-\mathbb{H}_0}}{\text{Tr } e^{-\mathbb{H}_0}}. \quad (2.57)$$

A full average  $\langle F \rangle$  may be reexpressed in terms of noninteracting averages according to

$$\langle F \rangle = \frac{\text{Tr } F e^{-\mathbb{H}}}{\text{Tr } e^{-\mathbb{H}}} = \frac{\langle F e^{-\mathbb{V}} \rangle_0}{\langle e^{-\mathbb{V}} \rangle_0}. \quad (2.58)$$

In section 4 we will solve the noninteracting problem in the limit  $n \rightarrow \infty$ . The study of the interaction  $\mathbb{V}$  will be taken up in section 6. It will require considerable work to show that the second term in (2.56),  $\log \langle e^{-\mathbb{V}} \rangle_0$ , has a well-controlled expansion in powers of  $n^{-\frac{1}{2}}$ . Correlation functions will be considered in sections 7 and 8.

### 3 Scaling

#### 3.1 Scaling of $\xi_m$ and $\eta_m$ for large $n$

In the preceding section, where  $n$  was an arbitrary integer, we have established the starting point for an expansion for asymptotically large  $n$ . We will now make the first steps in this expansion. A major problem is to know in advance how the various quantities will scale with  $n$ . For  $n \rightarrow \infty$  we expect all angles to become small and we begin by introducing

$$\xi_m = n^{-1} x_m, \quad \eta_m = n^{-1} y_m, \quad m = 1, \dots, n. \quad (3.1)$$

This scaling of  $\xi_m$  and  $\eta_m$  is certainly expected, since they constitute two sets of  $n$  positive variables adding up to  $2\pi$ .

#### 3.2 Scaling the integral for $p_n$

The integral  $\int d\xi d\eta$  included in the trace in (2.51) is defined in (2.49). It there appears that the upper integration limits of the  $\xi_m$  and  $\eta_m$ , given in terms of the  $\beta_\ell$  and  $\gamma_\ell$ , are all positive. Hence after passing to the scaled variables  $x_m$  and  $y_m$  we may replace these upper integration limits by  $\infty$ . Since the integrals converge exponentially, the error should be expected to vanish exponentially with  $n$ . It then becomes possible to commute the integration on  $\beta_1$  through those on the  $x_m$  and  $y_m$  and carry it out with the aid of the factor  $\delta(\beta_1 - \beta_*)$ . Henceforth the angle  $\beta_1$ , wherever it still occurs (such as in expressions for  $\mathbb{V}$  to be given later), should be viewed as the function  $\beta_*(\xi, \eta)$  given by the solution of (2.46).

As a result the integral (2.51) takes the simplified form

$$\begin{aligned}
p_n &\simeq \left(\frac{n}{2\pi i}\right)^2 \frac{(n-1)!}{2n} \pi^{-n} n^{-3n} \int_{c-i\infty}^{c+i\infty} ds \int_{-i\infty}^{i\infty} dt e^{2\pi n s} \\
&\times \int_0^\infty \left[ \prod_{m=1}^n dx_m dy_m \right] \left[ \prod_{m=1}^n x_m e^{-s\kappa x_m - s(1-\kappa)y_m - t(x_m - y_m)} \right] \\
&\times \exp(-\mathbb{V}(x, y)),
\end{aligned} \tag{3.2}$$

where we indicated explicitly that  $\mathbb{V}$ , given by (2.54), depends on the  $x_m$  and  $y_m$ . We use the  $\simeq$  sign to indicate that this is an asymptotic equality.

Provided that  $\mathbb{V}$  be sufficiently well-behaved, the integrals on the  $x_m$  and  $y_m$  will converge at the condition that

$$\operatorname{Re}(\kappa s + t) > 0, \quad \operatorname{Re}((1 - \kappa)s - t) > 0. \tag{3.3}$$

Since  $\operatorname{Re} t = 0$  and  $\operatorname{Re} s = c > 0$ , this condition is satisfied if we choose  $\kappa$  in the interval  $0 < \kappa < 1$ . It is now of great advantage to transform to variables of integration  $u$  and  $v$  defined by

$$u = \kappa s + t, \quad v = (1 - \kappa)s - t. \tag{3.4}$$

In terms of these equation (3.2) becomes

$$\begin{aligned}
p_n &\simeq \left(\frac{n}{2\pi i}\right)^2 \frac{(n-1)!}{2n} \pi^{-n} n^{-3n} \int du e^{2\pi n u} \int dv e^{2\pi n v} \\
&\times \int_0^\infty \left[ \prod_{m=1}^n dx_m x_m e^{-u x_m} \right] \int_0^\infty \left[ \prod_{m=1}^n dy_m e^{-v y_m} \right] \\
&\times \exp(-\mathbb{V}(x, y)),
\end{aligned} \tag{3.5}$$

where  $u$  and  $v$  run independently from  $-i\infty$  to  $i\infty$  while passing to the right of the origin. The difficulty of evaluating (3.5) is now hidden in the expression for  $\mathbb{V}$ .

The final remarks of this section concern practical procedure. Since (3.5) differs from (2.51) only by terms that are exponentially small in  $n$  and since we are preparing for a perturbation series in inverse powers of  $n$ , the scaled integral (3.5) together with expression (2.54) for  $\mathbb{V}$  is as good a starting point as the original integral (2.51) together with (2.52)-(2.54). In particular, (i) setting  $\mathbb{V} = 0$  in (3.5) yields  $p_n^{(0)}$ ; and (ii) the average  $\langle F \rangle_0$  defined in (2.57) is, up to exponentially small differences, equal to expression (3.5) with  $F$  inserted and  $\mathbb{V}$  set to zero, and divided by  $p_n^{(0)}$ .

## 4 Free problem

### 4.1 Calculation of $p_n^{(0)}$

We will solve the noninteracting problem by calculating  $p_n^{(0)}$  starting from expression (3.5) in which we set  $\mathbb{V} = 0$ . The  $2n$  integrals on the  $x_m$  and  $y_m$ , as well as those on  $u$  and  $v$ , then factorize. After carrying out the former we find

$$\begin{aligned}
 p_n^{(0)} &\simeq \frac{(n-1)!}{2n} \pi^{-n} n^{-3n} \left( \frac{n}{2\pi i} \int du \frac{e^{2\pi n u}}{u^{2n}} \right) \left( \frac{n}{2\pi i} \int dv \frac{e^{2\pi n v}}{v^n} \right) \\
 &= \frac{(n-1)!}{2n} \pi^{-n} n^{-3n} \times \frac{n^{2n} (2\pi)^{2n-1}}{(2n-1)!} \times \frac{n^n (2\pi)^{n-1}}{(n-1)!} \\
 &= \frac{(8\pi^2)^n}{4\pi^2 (2n)!},
 \end{aligned} \tag{4.1}$$

where in the first step we did the integrations by closing their contours around the poles in the origin. Additive corrections to (4.1) should be expected to vanish exponentially with  $n$ .

It is instructive and useful for later to analyze the integrals also by the steepest descent method. In the case of the  $u$  integral the integrand may be written as  $\exp(-2n \log u + 2\pi n u)$ , which has a saddle point for  $u = u_c \equiv \pi^{-1}$ . Upon taking its second derivative, it appears that contributions come from a neighborhood of extension  $\sim n^{-\frac{1}{2}}$  around the saddle point. A similar analysis holds for the  $v$  integral, for which the saddle point is at  $v = v_c \equiv (2\pi)^{-1}$ . The steepest descent analysis allows us to conclude that any function  $h(u, v)$  independent of  $n$  and inserted in the integrand contributes an extra factor  $h(u_c, v_c)$  to the value of the integral. This rule will be applied in section 7.1.

### 4.2 Averages and correlations

The averages of the scaled variables  $x_m$  and  $y_m$  are necessarily in any order equal to

$$\langle x_m \rangle \equiv \bar{x} = 2\pi, \quad \langle y_m \rangle \equiv \bar{y} = 2\pi. \tag{4.2}$$

One obtains the noninteracting results for arbitrary multivariate moments of the  $x_m$  and  $y_m$  by starting from (3.5) with the appropriate insertions, setting again  $\mathbb{V} = 0$ , and going through calculational steps similar to those above. We state the results only for the second moments,

$$\langle x_m^2 \rangle_0 = \frac{12\pi^2 n}{2n+1}, \quad \langle y_m^2 \rangle_0 = \frac{8\pi^2 n}{n+1}. \tag{4.3}$$

Due to the symmetry of the noninteracting expression [equation (3.5) with  $\mathbb{V} = 0$ ], the two-point correlations  $\langle x_\ell x_m \rangle$  and  $\langle y_\ell y_m \rangle$  must be index inde-



pendent when  $\ell \neq m$ . This observation combined with the geometrical sum rules (2.4) and (2.10) yields

$$\langle x_\ell x_m \rangle_0 = \frac{8\pi^2 n}{2n+1}, \quad \langle y_\ell y_m \rangle_0 = \frac{2\pi^2 n}{n+1}, \quad \ell \neq m. \quad (4.4)$$

Furthermore  $\langle x_\ell y_m \rangle_0 = \langle x_\ell \rangle_0 \langle y_m \rangle_0$  for all  $\ell$  and  $m$ . Let us define

$$\delta x_m = x_m - \bar{x}, \quad \delta y_m = y_m - \bar{y}. \quad (4.5)$$

From what precedes we then have, still expanding (4.3) in powers of  $n^{-1}$ ,

$$\begin{aligned} \langle \delta x_\ell \delta x_m \rangle_0 &= 2\pi^2 (\delta_{\ell m} - n^{-1}) (1 + \mathcal{O}(n^{-1})), \\ \langle \delta y_\ell \delta y_m \rangle_0 &= 4\pi^2 (\delta_{\ell m} - n^{-1}) (1 + \mathcal{O}(n^{-1})), \\ \langle \delta x_\ell \delta y_m \rangle_0 &= 0. \end{aligned} \quad (4.6)$$

These variances obey the geometrical sum rules (2.4) and (2.10). But although they include terms of order  $n^{-1}$ , we will see in sections 7 and 8 that the interaction  $\mathbb{V}$  generates additional order  $n^{-1}$  contributions.

### 4.3 Probability distribution of $\xi_m$ and $\eta_m$

The probability for the angle  $\xi_m$  to be between  $\xi$  and  $\xi + d\xi$  is given by  $\langle \delta(\xi_m - \xi) \rangle d\xi$ . We set  $\xi = n^{-1}x$  and Fourier represent according to

$$\delta(x_m - x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\mu e^{i\mu(x_\ell - x)}. \quad (4.7)$$

We insert this in (3.5), commute the  $\mu$  integral to the outside, and pass through the same steps that led from (3.5) to the first line of (4.1). The result is similar to (4.1) but with an insertion  $u^2(u - i\mu)^{-2}$ ,

$$\begin{aligned} p_n^{(0)} \langle \delta(x_m - x) \rangle_0 &\simeq \frac{1}{2\pi} \int_{-\infty}^{\infty} d\mu e^{-i\mu x} \\ &\times \frac{(n-1)!}{2n} \pi^{-n} n^{-3n} \left( \frac{n}{2\pi i} \int du \frac{e^{2\pi n u}}{u^{2n}} \left( \frac{u}{u - i\mu} \right)^2 \right) \left( \frac{n}{2\pi i} \int dv \frac{e^{2\pi n v}}{v^n} \right). \end{aligned} \quad (4.8)$$

Because of our discussion on insertions at the end of section 4.1, the second line in (4.8) is equal to  $p_n^{(0)} (u_c / (u_c - i\mu))^2$ . Upon dividing by  $p_n^{(0)}$  and setting  $u_c = 1/\pi$  we obtain the probability distribution of  $x$  in the limit  $n \rightarrow \infty$ ,

$$\begin{aligned} \langle \delta(x_m - x) \rangle_0 &= \frac{1}{2\pi} \int_{-\infty}^{\infty} d\mu \frac{e^{-i\mu x}}{(1 - i\pi\mu)^2} \\ &= \frac{x}{\pi^2} \exp\left(-\frac{x}{\pi}\right). \end{aligned} \quad (4.9)$$

In terms of the original variable  $\xi = n^{-1}x$  this becomes the distribution  $u(\xi)$  announced in equation (1.5) of the introduction. In a fully analogous way an insertion  $\delta(\eta_m - \eta)$  produces

$$\langle \delta(y_m - y) \rangle_0 = \frac{1}{2\pi} \exp\left(-\frac{y}{2\pi}\right), \quad (4.10)$$

from which  $v(\eta)$  of equation (1.6) follows.

## 5 Further scaling

### 5.1 Scaling of $\beta_m$ and $\gamma_m$ for large $n$

The noninteracting calculations of section 4 required no scaling other than the one postulated in (3.1), which concerns the  $\xi_m$  and  $\eta_m$ . We will show now how the scaling of all further variables follows from this initial postulate *plus* what we have learned about the independence of the  $\xi_m$  and  $\eta_m$ .

The first one of equations (2.16) expresses  $\beta_m - \beta_1$  as a sum of  $2m - 2$  terms  $\xi_\ell$  and  $\eta_\ell$  which, in the noninteracting case, are all independent. Since  $m$  is typically of order  $n$ , it follows that  $\beta_m - \beta_1$  scales as  $n^{-\frac{1}{2}}$ . But since  $\beta_1$  is in no way statistically different from the other  $\beta_m$ , they must all scale as  $n^{-\frac{1}{2}}$ . In a similar way one deduces from the second one of equations (2.16) that the  $\gamma_m$  scale with  $n^{-\frac{1}{2}}$ . We therefore set

$$\beta_m = n^{-\frac{1}{2}}b_m, \quad \gamma_m = n^{-\frac{1}{2}}c_m, \quad m = 1, \dots, n, \quad (5.1)$$

as well as

$$\beta_*(\xi, \eta) = n^{-\frac{1}{2}}b_*(x, y). \quad (5.2)$$

Hence the scaling factor for the angles  $\xi_m$  and  $\eta_m$  is different from the one for the angles  $\beta_m$  and  $\gamma_m$ . By symmetry and because of the geometrical sum rule (2.12) we must have  $\langle \beta_m \rangle = \langle \gamma_m \rangle = \pi/n$ . The difference between the scaling of these averages and of the typical values of  $\beta_m$  and  $\gamma_m$ , which are  $\sim n^{-\frac{1}{2}}$ , can arise only because the  $\beta_m$  and  $\gamma_m$  may take negative values, as illustrated in figure 2.

It will be useful to have at hand the scaled equivalent of (2.16). Rendering explicit the fact that here  $\beta_1 = \beta_*(\xi, \eta)$  we find

$$b_m = b_*(x, y) - n^{-\frac{1}{2}} \sum_{\ell=1}^{m-1} (x_\ell - y_\ell), \quad (5.3)$$

$$c_m = -b_*(x, y) + n^{-\frac{1}{2}} \sum_{\ell=1}^{m-1} (x_\ell - y_\ell) + n^{-\frac{1}{2}}x_m, \quad (5.4)$$

valid for  $m = 1, \dots, n$ .

The two sums  $\sum_{\ell=1}^m x_\ell$  and  $\sum_{\ell=1}^m y_\ell$  may be viewed as the displacements of two independent random walks, each built up out of independent increments. Figure 1 shows that these displacements describe the progression of the polar angles of the midpoint vector  $\mathbf{R}_m$  and the vertex vector  $\mathbf{S}_m$ , respectively. Hence the cell perimeter may be viewed [31] as the result of a construction based on two entangled random walks.

## 5.2 Cancellations of orders in $n^{-\frac{1}{2}}$

The nontriviality of this scaling picture comes from the fact that various frequently occurring combinations of the angular variables exhibit cancellations. For later use we collect some of them here, employing a hybrid notation in which on the one hand the  $b_m$  and  $c_m$ , and on the other hand the  $x_m$  and  $y_m$  appear. First of all there are the sums of odd powers of the type

$$\begin{aligned}\gamma_m + \beta_{m+1} &= y_m n^{-1}, \\ \gamma_m^3 + \beta_{m+1}^3 &= (\gamma_m + \beta_{m+1}) \left[ \frac{3}{2}(\gamma_m^2 + \beta_{m+1}^2) - \frac{1}{2}(\gamma_m + \beta_{m+1})^2 \right] \\ &= \frac{3}{2}y_m(c_m^2 + b_{m+1}^2)n^{-2} - \frac{1}{2}y_m^3n^{-3},\end{aligned}\tag{5.5}$$

and analogously

$$\begin{aligned}\beta_m + \gamma_m &= x_m n^{-1}, \\ \beta_m^3 + \gamma_m^3 &= \frac{3}{2}x_m(b_m^2 + c_m^2)n^{-2} - \frac{1}{2}x_m^3n^{-3}.\end{aligned}\tag{5.6}$$

Secondly, we will encounter differences of even powers,

$$\begin{aligned}\beta_\ell^2 - \gamma_\ell^2 &= (\beta_\ell + \gamma_\ell)(\beta_\ell - \gamma_\ell) \\ &= x_\ell(b_\ell - c_\ell)n^{-\frac{3}{2}}, \\ \beta_\ell^4 - \gamma_\ell^4 &= [(\beta_\ell + \gamma_\ell)^2 - 2\beta_\ell\gamma_\ell](\beta_\ell^2 - \gamma_\ell^2) \\ &= -2x_\ell b_\ell c_\ell(b_\ell - c_\ell)n^{-\frac{5}{2}} + x_\ell^3(b_\ell - c_\ell)n^{-\frac{7}{2}}.\end{aligned}\tag{5.7}$$

These relations are easily extended, if needed, to sums and differences of higher powers of the angles.

## 5.3 Solution of $G(\xi, \eta; \beta_*) = 0$

The solution  $\beta_*(\xi, \eta)$  of the no-spiral constraint  $G(\xi, \eta; \beta_*) = 0$  has not been needed explicitly. Since  $\beta_*$  plays an important role, we will show in this subsection how it may be obtained in the scaling limit. The derivation also establishes its uniqueness within the large  $n$  perturbation expansion. Finally,

because of equivalence by symmetry, the study of  $\beta_* = \beta_1$  gives us access to the statistics of all angles  $\beta_m$  and  $\gamma_m$ .

From (2.35) we have straightforwardly

$$\begin{aligned} G(\xi, \eta; \beta_*) &= \frac{1}{4\pi} \sum_{m=1}^n \left( \beta_m^2 - \gamma_m^2 + \mathcal{O}(\beta_m^4 - \gamma_m^4) \right) \\ &= \frac{1}{4\pi n} \sum_{m=1}^n (b_m^2 - c_m^2) + \mathcal{O}(n^{-\frac{3}{2}}), \end{aligned} \quad (5.8)$$

The leading order term is  $\sim n^{-\frac{1}{2}}$ . Its scaling, as well as that of the correction term, may be determined with the aid of equations (5.7) and taking into account the sum on  $m$ . For convenience of notation we rewrite (5.4) just for this occasion as  $b_m = b_* + \tilde{b}_m$  and  $c_m = -b_* + \tilde{c}_m$ , substitute it in the RHS of (5.8), set  $G = 0$ , and solve for  $b_*(x, y)$  using the geometrical sum rule (2.12). The result is that

$$\begin{aligned} b_*(x, y) &= -\frac{1}{4\pi n^{\frac{1}{2}}} \sum_{m=1}^n (\tilde{b}_m^2 - \tilde{c}_m^2) + \mathcal{O}(n^{-1}) \\ &= \frac{1}{4\pi n^{\frac{3}{2}}} \left[ 2 \sum_{m=1}^n \sum_{\ell=1}^{m-1} x_m (x_\ell - y_\ell) + \sum_{m=1}^n x_m^2 \right] + \mathcal{O}(n^{-1}), \end{aligned} \quad (5.9)$$

where in the second line we returned to standard notation. Inside the square brackets of (5.9) the double sum is of order  $n^{\frac{3}{2}}$  and the single sum of order  $n$ . By taking the average  $\langle \dots \rangle_0$  on both sides of (5.9) and using (4.2)-(4.4) one finds, after cancellation of the leading order terms, that  $\langle b_* \rangle_0 = \pi n^{-\frac{1}{2}}$ . This relation should in fact hold to all orders in  $n$  and leads to  $\langle \beta_m \rangle_0 = \langle \gamma_m \rangle_0 = \pi n^{-1}$ , as had to be the case. The distribution of  $b_*$  around its average is on a scale which is larger by a factor  $n^{\frac{1}{2}}$ . Restricting ourselves now to this larger scale, we substitute  $x_m = \bar{x} + \delta x_m$  in (5.9) and, reasoning as before, keep only the leading order. The result,

$$b_*(x, y) = -n^{-\frac{3}{2}} \sum_{m=1}^n m(\delta x_m - \delta y_m) + \mathcal{O}(n^{-\frac{1}{2}}), \quad (5.10)$$

shows that  $b_*(x, y)$  is a weighted sum of  $n$  weakly dependent random variables and therefore, in the limit of large  $n$ , has a Gaussian distribution. Its noninteracting variance may be calculated from (5.10) and (4.6) and is given by  $\langle b_*^2 \rangle_0 \simeq \frac{1}{2} \pi^2$ . However, the interaction  $\mathbb{V}$  yields an additional  $\mathcal{O}(n^0)$  contribution to this variance. Its explicit value to order  $n^0$  may be found analytically with the aid of the interacting theory of the next sections, but we will not pursue this calculation.

## 6 Interaction

The total Hamiltonian  $\mathbb{H}$  was split in equation (2.52) into a noninteracting term  $\mathbb{H}_0$  and an interaction  $\mathbb{V}$ , with an ensuing splitting for  $\log p_n$  in (2.56). In section 4 we dealt with the noninteracting problem and obtained, in equation (4.1), the first term of (2.56). This result is useful only if we can also control the second term in (2.56), which requires the evaluation of  $\langle e^{-\mathbb{V}} \rangle_0$ . Furthermore, if we wish to study correlations,  $\langle F e^{-\mathbb{V}} \rangle_0$  is needed. We will now consider these quantities.

### 6.1 General

The interaction  $\mathbb{V}$  is defined by expression (2.54), which is particularly opaque. We now assert that  $\mathbb{V}$  is in fact of order  $n^0$ . We will show this by constructing an analytically tractable expression  $\mathbb{V}_1$  such that [46]

$$\mathbb{V} = \mathbb{V}_1 + n^{-\frac{1}{2}}\mathbb{V}_2 \quad (6.1)$$

with  $\mathbb{V}_1$  of order  $n^0$  and  $n^{-\frac{1}{2}}\mathbb{V}_2$  representing a collection of higher order terms. Expanding in a power series in  $n^{-\frac{1}{2}}\mathbb{V}_2$  we obtain from (6.1)

$$\langle e^{-\mathbb{V}} \rangle_0 = \langle e^{-\mathbb{V}_1} \rangle_0 - n^{-\frac{1}{2}} \langle \mathbb{V}_2 e^{-\mathbb{V}_1} \rangle_0 + \dots \quad (6.2)$$

Substitution of (6.2) in (2.56) yields for  $\log p_n$  the expansion

$$\log p_n = \log p_n^{(0)} + \log \langle e^{-\mathbb{V}_1} \rangle_0 - n^{-\frac{1}{2}} \langle \mathbb{V}_2 \rangle_1 + \dots, \quad (6.3)$$

where for any function  $F$  of the angular variables we define the *first order average*

$$\langle F \rangle_1 = \frac{\langle F e^{-\mathbb{V}_1} \rangle_0}{\langle e^{-\mathbb{V}_1} \rangle_0}. \quad (6.4)$$

The full average of a function  $F$  is given by

$$\langle F \rangle = \langle F \rangle_1 - n^{-\frac{1}{2}} [ \langle F \mathbb{V}_2 \rangle_1 - \langle F \rangle_1 \langle \mathbb{V}_2 \rangle_1 ] + \dots \quad (6.5)$$

If we can substantiate our claim that  $\mathbb{V}_1$  is of order  $n^0$  and  $n^{-\frac{1}{2}}\mathbb{V}_2$  of higher order, then (6.3) and (6.5) are perturbation series in  $n^{-\frac{1}{2}}$ .

In this section we study  $\mathbb{V}$  with the sole purpose of isolating an appropriate expression  $\mathbb{V}_1$ . This will require an important amount of work. In section 7 we will show the anticipated result, *viz.* that for  $n \rightarrow \infty$  the average  $\langle e^{-\mathbb{V}_1} \rangle_0$  tends to a finite constant  $C$ . In view of equation (6.3) this then establishes the asymptotic expansion of  $\log p_n$  up to terms that vanish as  $n \rightarrow \infty$ . In sections 7 and 8 we will determine first order expressions for various correlation functions.

## 6.2 An analog: the theory of elasticity

Our determination of the interaction  $\mathbb{V}_1$  will be based on the following consideration. The regular  $n$ -sided polygon is an obvious point of symmetry in phase space. In fact, Drouffe and Itzykson [27] constructed an exact lower bound for  $p_n$  by expanding the vertex vectors  $\mathbf{S}_m$  about their regular polygon positions. Such an expansion preserves, however, the long range order of the regular polygon and is therefore likely to cover only a small fraction of the effectively contributing phase space. Therefore we will proceed here as in the theory of elasticity, where one considers the deviations not of the *positions*, but of the *interatomic distances*, from their regular lattice values. The analogs of the interatomic distances are the angles  $\xi_m$  between the mid-point vectors and  $\eta_m$  between the vertex vectors. Hence we will expand  $\mathbb{V}$  in powers of  $\delta x_m$  and  $\delta y_m$  defined in (4.5). This procedure allows, in principle, for large deviations from the regular  $n$ -sided polygon. In the nontrivial and rather lengthy analysis of this section we will step by step identify the order in  $n$  of the terms that we encounter, relegate to  $\mathbb{V}_2$  those of order  $n^{-\frac{1}{2}}$  or smaller, and retain the others. The outcome of this will be expression (6.39) for  $\mathbb{V}_1$ . The terms that contribute to  $\mathbb{V}_1$  are all quadratic in the  $\delta x_m$  and  $\delta y_m$ .

## 6.3 The interaction $\mathbb{V}$ as a series in the angles

Definition (2.54) of  $\exp(-\mathbb{V})$  is a product of factors  $G'$ ,  $\rho_m$ ,  $T_m \xi_m^{-1}$ , and  $1 + n^{-1}V$ . We will successively carry out the analysis of the orders in  $n$  appearing in each of these factors. Conceptually two different steps must be distinguished, even though in practice we have not strictly separated them. First, since the angles  $\beta_m, \gamma_m, \xi_m$ , and  $\eta_m$  are all small, even though not all of the same order, it will be convenient to expand all relevant expressions as power series in these angles. This is a matter of practical convenience rather than of principle. Second, we will convert the resulting series to an expansion in the  $\delta x_m$  and  $\delta y_m$ , retaining only the leading order.

### 6.3.1 The factors $G'$ , $\rho_m$ , and $T_m \xi_m^{-1}$

We first consider the function  $G$ . It is given by (2.35) in terms of the  $\beta_\ell$  and  $\gamma_\ell$  which, in turn, are expressed by (2.16) in terms of  $\beta_1$  and of the  $\xi_m$  and  $\eta_m$ . When differentiating  $G$  with respect to  $\beta_1$  at fixed  $\xi$  and  $\eta$  we obtain

$$\begin{aligned} G'(\xi, \eta; \beta_1) &= \frac{1}{2\pi} \sum_{m=1}^n (\tan \beta_m + \tan \gamma_m) \\ &= 1 + \frac{1}{6\pi} \sum_{m=1}^n (\beta_m^3 + \gamma_m^3) + \dots, \end{aligned} \quad (6.6)$$

where in the second step we used the geometrical sum rule (2.12) to cancel the linear terms in the angles, and where the dots stand for terms of fifth and higher order in  $\beta_m$  and  $\gamma_m$ . According to (5.6) the sum  $\beta_m^3 + \gamma_m^3$  scales as  $\sim n^{-2}$ . Since the sum on  $m$  contributes an extra factor  $n$ , we deduce from (6.6) that

$$G'(\xi, \eta; \beta_*) = 1 + \mathcal{O}(n^{-1}), \quad (6.7)$$

*i.e.*, this derivative is to leading order independent of its arguments. The  $\mathcal{O}(n^{-1})$  corrections in (6.7) may be treated perturbatively (*i.e.*, they belong to  $\mathbb{V}_2$ ), but will not be dealt with in the present work.

Secondly, we obtain from (2.37) and (2.38) the expansion

$$\prod_{m=1}^n \rho_m^2 = \exp \left( - \sum_{m=1}^n \tau_m^2 + \frac{2}{3} \sum_{m=1}^n \tau_m^3 + \dots \right). \quad (6.8)$$

Expanding (2.33) and using (2.37) yields for  $\tau_m$  the series

$$\tau_m = \tau_n + \frac{1}{2} \sum_{\ell=1}^m (\beta_\ell^2 - \gamma_\ell^2) - \frac{1}{24} \sum_{\ell=1}^m (\beta_\ell^4 - \gamma_\ell^4) + \dots, \quad (6.9)$$

which is valid for  $m = 1, 2, \dots, n$ ; we recall that for  $m = n$  it is equivalent to the no-spiral constraint.

Equation (5.7) tells us that  $\beta_\ell^2 - \gamma_\ell^2$  and  $\beta_\ell^4 - \gamma_\ell^4$  scale as  $\sim n^{-\frac{3}{2}}$  and  $\sim n^{-\frac{5}{2}}$ , respectively. When  $m$  is of the order of  $n$ , as is typically the case, the sum on  $m$  contributes a factor  $n$  to the scaling, so that  $\tau_m - \tau_n$  is to leading order proportional to  $n^{-\frac{1}{2}}$ . This leads us to define the auxiliary scaled variables  $r_m$  by

$$\tau_m = n^{-\frac{1}{2}} r_m, \quad m = 1, \dots, n, \quad (6.10)$$

which allow us to write the gauge condition as

$$\sum_{m=1}^n r_m = 0. \quad (6.11)$$

From (2.29), (2.37), (6.10), and (2.44) we find that  $r_m$  is related to the original midpoint distance  $R_m$  by

$$\begin{aligned} R_m &= R_{\text{av}}(1 + n^{-\frac{1}{2}} r_m) \\ &= R_c + \pi^{-\frac{1}{2}}(r_{\text{av}} + r_m) + \mathcal{O}(n^{-\frac{1}{2}}). \end{aligned} \quad (6.12)$$

In terms of scaled variables (6.9) becomes, if we use both equations (5.7),

$$r_m = r_n + \frac{1}{2n} \sum_{\ell=1}^m x_\ell (b_\ell - c_\ell) + \frac{1}{12n^2} \sum_{\ell=1}^m x_\ell b_\ell c_\ell (b_\ell - c_\ell) + \dots, \quad (6.13)$$

valid for  $m = 1, 2, \dots, n$ , and where the dots stand for higher order terms. Let us now consider (6.8). Since  $\tau_m^2 \sim n^{-1}$  and since the sum on  $n$  contributes an extra factor  $n$ , the first term in the exponent in (6.8) is of order  $n^0$ . We will keep it. However,  $\tau_m^3$  is of order  $n^{-\frac{3}{2}}$ , so that its sum on  $m$  will vanish for  $n \rightarrow \infty$  and can be treated perturbatively. Hence the contribution to  $\exp(-\mathbb{V})$  is a factor

$$\prod_{m=1}^n \rho_m^2 = \left[ \exp \left( - \sum_{m=1}^n \tau_m^2 \right) \right] (1 + \mathcal{O}(n^{-1})). \quad (6.14)$$

Thirdly we consider the product of the  $T_m \xi_m^{-1}$ . Symmetrizing in  $\beta_m$  and  $\gamma_m$  with the aid of the no-spiral constraint  $G = 0$  we may write this product as

$$\begin{aligned} \prod_{m=1}^n T_m \xi_m^{-1} &= \prod_{m=1}^n \frac{\sin(\beta_m + \gamma_m)}{(\beta_m + \gamma_m) \cos \beta_m \cos \gamma_m} \\ &= \exp \sum_{m=1}^n \left( \frac{1}{2}(\beta_m^2 + \gamma_m^2) - \frac{1}{3}(\beta_m + \gamma_m)^2 + \dots \right). \end{aligned} \quad (6.15)$$

The sum  $\sum_m (\beta_m^2 + \gamma_m^2)$  contains  $n$  terms of order  $(n^{-\frac{1}{2}})^2$ , is therefore itself of order  $n^0$  and must be kept. Since  $\beta_m + \gamma_m = \xi_m$  scales as  $n^{-1}$ , the sum  $\sum_m (\beta_m + \gamma_m)^2$  also scales as  $n^{-1}$  and may be treated perturbatively. The higher order terms in the sum in the exponential in (6.15) are at least quartic in the angles and may again be treated by perturbation theory. Hence, to leading order,

$$\prod_{m=1}^n T_m \xi_m^{-1} = \left[ \exp \frac{1}{2} \sum_{m=1}^n (\beta_m^2 + \gamma_m^2) \right] (1 + \mathcal{O}(n^{-1})) \quad (6.16)$$

is the contribution of this product to  $\exp(-\mathbb{V})$ .

### 6.3.2 The factor $(1 + n^{-1}V)^{-n}$

When writing (2.39) as we did, we anticipated that  $V$  thus defined would scale as  $\sim n^0$  for  $n \rightarrow \infty$ . We will show here that this is indeed the case. We expand

$$(1 + n^{-1}V)^{-n} = \exp \left( -V + \frac{1}{2}n^{-1}V^2 + \dots \right) \quad (6.17)$$

and next wish to find  $V$  as a series in the angles. The decomposition (2.39) of  $\mathcal{A}$  was motivated by the fact that in this way, and since the  $\tau_m$  are quadratic in the angles, the series for  $V$  contains terms only from third order up. We



expand  $V$  as given by (2.41) and find

$$\begin{aligned}
V &= \frac{n}{\pi} \sum_{m=1}^n (1 + \tau_m)^2 \left[ \frac{1}{3}(\gamma_m^3 + \beta_{m+1}^3) + \dots \right] \\
&\quad + \frac{n}{\pi} \sum_{m=1}^n \tau_m (\gamma_m + \beta_{m+1}) + \frac{n}{2\pi} \sum_{m=1}^n \tau_m^2 (\gamma_m + \beta_{m+1}). \quad (6.18)
\end{aligned}$$

In the first line of (6.18) the dominant term is the one where  $\frac{1}{3}(\gamma_m^3 + \beta_{m+1}^3)$  is summed without any factor  $\tau_m$ . To estimate its order in  $n$  we may invoke (5.5); we will however perform a slightly refined analysis by using that  $\gamma_m + \beta_{m+1} = \eta_m$ , setting as before  $\eta_m = \bar{\eta} + \delta\eta_m$  with  $\bar{\eta} = 2\pi n^{-1}$ , and writing

$$\begin{aligned}
\gamma_m^3 + \beta_{m+1}^3 &= \frac{3}{2}\eta_m(\gamma_m^2 + \beta_{m+1}^2) - \frac{1}{2}\eta_m^3 \\
&= \frac{3}{2}\bar{\eta}(\gamma_m^2 + \beta_{m+1}^2) + \frac{3}{2}\delta\eta_m(\gamma_m^2 + \beta_{m+1}^2) - \frac{1}{2}\eta_m^3. \quad (6.19)
\end{aligned}$$

Because of the scaling (5.1) we have  $\bar{\eta}(\gamma_m^2 + \beta_{m+1}^2) \sim n^{-2}$ . Hence, taking into account the effect of  $n$  times the sum on  $m$  in (6.18), we conclude that the first term on the RHS of (6.19) yields a contribution of order  $n^0$  to  $V$  and should therefore be kept. The term  $-\frac{1}{2}\eta_m^3$  in (6.19) contributes to order  $n^{-1}$  to  $V$  and can be treated perturbatively. The term  $\frac{3}{2}\delta\eta_m(\gamma_m^2 + \beta_{m+1}^2)$  on the RHS of (6.19) requires more attention. The factor  $\delta\eta_m$  is of the same order as  $\bar{\eta}$  in the first term, *viz.*  $\sim n^{-1}$ . However,  $\delta\eta_m$  is a zero average random variable and therefore [47] the sum on  $m$  will raise the order by only  $n^{\frac{1}{2}}$  instead of  $n$ , whence it follows that the resulting contribution to  $V$  can be treated perturbatively. Putting these arguments together we get for the first line of (6.18)

$$\frac{n}{\pi} \sum_{m=1}^n (1 + \tau_m)^2 \left[ \frac{1}{3}(\gamma_m^3 + \beta_{m+1}^3) + \dots \right] = \sum_{m=1}^n (\gamma_m^2 + \beta_{m+1}^2) + \mathcal{O}(n^{-\frac{1}{2}}). \quad (6.20)$$

In the second line of equation (6.18) we perform a similar analysis on the first sum. Using the sum rule (2.38) we can write

$$\frac{n}{\pi} \sum_{m=1}^n \tau_m (\gamma_m + \beta_{m+1}) = \frac{n}{\pi} \sum_{m=1}^n \tau_m \delta\eta_m. \quad (6.21)$$

Since  $\delta\eta_m \sim \eta_m \sim n^{-1}$ , the product  $\delta\eta_m \tau_m$  is of order  $n^{-\frac{1}{2}}$ , and when summed on  $m$  with the prefactor  $n/\pi$  in (6.18), it would appear that it contributes a term of order  $n^{\frac{1}{2}}$  to  $V$ . This again overestimates the order. Since the  $\delta\eta_m$  are zero average random variables, we expect [47] in fact that the sum will raise the order by a factor of only  $n^{\frac{1}{2}}$ , and hence that this term produces an order  $n^0$  contribution to  $V$ . Therefore we keep this term.

In the second sum in the second line of (6.18) we write  $\tau_m^2(\gamma_m + \beta_{m+1}) = 2\pi n^{-1}\tau_m^2 + \tau_m^2\delta\eta_m$ . Both these terms are of order  $(n^{-\frac{1}{2}})^2 \times n^{-1} = n^{-2}$  but the summation on  $m$  increases the order of the first one by  $n$  and of the second one, which is random [47], by only  $n^{\frac{1}{2}}$ . Hence only the first term leads to a contribution of order  $n^0$  to  $V$  and we get

$$\frac{n}{2\pi} \sum_{m=1}^n \tau_m^2(\gamma_m + \beta_{m+1}) = \sum_{m=1}^n \tau_m^2 + \mathcal{O}(n^{-\frac{1}{2}}). \quad (6.22)$$

Finally, we must consider higher orders in  $V$  in the expansion (6.17). Since we concluded that  $V$  is of order  $n^0$ , it follows that all higher order terms  $n^{-1}V^2$ ,  $n^{-2}V^3$ ,  $\dots$  in the expansion of the logarithm can be treated perturbatively. Hence the contributions to  $(1 + n^{-1}V)^{-n}$  that must be kept are given by (6.20), (6.21), and (6.22). Upon adding them together we are led to

$$\begin{aligned} (1 + n^{-1}V)^{-n} &= \exp\left(-\sum_{m=1}^n (\gamma_m^2 + \beta_{m+1}^2) - \sum_{m=1}^n \tau_m^2 - \frac{n}{\pi} \sum_{m=1}^n \tau_m \delta\eta_m\right) \\ &\quad \times (1 + \mathcal{O}(n^{-\frac{1}{2}})), \end{aligned} \quad (6.23)$$

which completes our discussion of the factors composing  $\exp(-\mathbb{V})$ .

### 6.3.3 Intermediate result for $\mathbb{V}$

Upon collecting the contributions identified above in equations (6.7), (6.14), (6.16), and (6.23) we find

$$\begin{aligned} \mathbb{V} &= \frac{1}{2} \sum_{m=1}^n (\beta_m^2 + \gamma_m^2) + 2 \sum_{m=1}^n \tau_m^2 + \frac{n}{\pi} \sum_{m=1}^n \tau_m \delta\eta_m \\ &= \frac{1}{n} \sum_{m=1}^n \left(\frac{1}{2}(b_m^2 + c_m^2) + 2r_m^2\right) + \frac{1}{\pi n^{\frac{1}{2}}} \sum_{m=1}^n r_m \delta y_m + \mathcal{O}(n^{-\frac{1}{2}}), \end{aligned} \quad (6.24)$$

where in the second line we have passed to scaled variables. Although this expression has the appearance of a quadratic form, we recall that  $r_m$  is itself a series in the angles starting out with a quadratic term.

The important facts established at this point are (i) that  $\mathbb{V}$  given by (6.24) is of order  $n^0$  as  $n \rightarrow \infty$ ; and (ii) that the terms neglected in (6.24) may be classified by their order in  $n^{-\frac{1}{2}}$  and can hence be arranged, if desired, in a perturbation series.

A word of caution is in place concerning the averages  $\langle F \rangle$ . To arrive at  $\mathbb{V}$  of equation (6.24), we have used twice in section 6.3.2 that the sum of  $n$  sufficiently weakly correlated random variables has a random part which

is of relative order order  $n^{-\frac{1}{2}}$  with respect to its systematic part. For a two-point correlation at a distance  $k$  (measured in units of the midpoint index  $m$ ) we expect that in an analogous calculation sums of  $k$  terms will intervene, for which the random part is of relative order  $k^{-\frac{1}{2}}$  with respect to the systematic part. Hence the resulting expression (6.24) can be used for correlation functions at distances  $k \sim n$ , but in the absence of further analysis *not* for  $k \sim n^0$ . We speculate that in fact it is valid for correlations at all distances  $k \sim n^\epsilon$  for any  $\epsilon > 0$ , but our further work will not depend on that.

## 6.4 Expressing $\mathbb{V}$ in terms of $\delta x_m$ and $\delta y_m$

We return to the study of expression (6.24) for  $\mathbb{V}$ . In order to prepare for integrating  $\exp(-\mathbb{V})$  on the  $x_m$  and  $y_m$ , as required by the average  $\langle \exp(-\mathbb{V}) \rangle_0$  in (6.3), we first wish to express the  $b_m$ ,  $c_m$ , and  $r_m$  that appear in (6.24) in terms of the  $\delta x_m$  and  $\delta y_m$ .

### 6.4.1 Recursion relation

In principle, this should be done as follows. We may deduce from (6.13) the recursion relation

$$r_m - r_{m-1} = n^{-1} x_m u_m + \mathcal{O}(n^{-2}), \quad m = 1, \dots, n, \quad (6.25)$$

with  $r_0 = r_n$  as an additional condition expressing the no-spiral constraint and with the  $u_m$  defined by

$$u_m = \frac{1}{2}(b_m - c_m), \quad m = 1, \dots, n. \quad (6.26)$$

From (6.26) and (5.4) it follows that the  $u_m$  satisfy

$$u_m - u_{m-1} = n^{-\frac{1}{2}} f_{m-1}, \quad m = 2, \dots, n, \quad (6.27)$$

in which the  $f_m$  are defined as

$$f_{m-1} = -\frac{1}{2}(\delta x_m + \delta x_{m-1} - 2\delta y_{m-1}), \quad m = 1, \dots, n \quad (6.28)$$

with the conventions  $\delta x_0 = \delta x_n$ ,  $\delta y_0 = \delta y_n$ , and  $f_0 = f_n$ . The extension of (6.27) to  $m = 1$  defines  $u_0$ . Because of the geometrical sum rules (2.4) and (2.10) we have  $u_0 = u_n$  as well as

$$\sum_{m=1}^n f_m = 0. \quad (6.29)$$

If initial values  $r_0$  and  $u_0$  are given, the system of equations (6.25) and (6.27) may be solved recursively in the midpoint index  $m$ , yielding the  $r_m$  and  $u_m$

in terms of the  $f_m$ , and hence in terms of the  $x_m$  and  $y_m$ . In the end the initial values are determined by two conditions, *viz.* the no-spiral constraint  $r_n = r_0$  and the gauge condition  $\sum_{m=1}^n r_m = 0$ . Proceeding this way gives a host of terms many of which contain multiple sums on the midpoint index. Their order in  $n$  may be determined by the methods of section 6.3. This route, however, is cumbersome as too much detailed information is carried along. We will therefore adopt a different but equivalent approach which selects the leading order terms more easily. It also reestablishes the cyclic invariance in the index  $m$  and the symmetry under reversal of orientation.

In (6.25) we substitute  $x_m = \bar{x} + \delta x_m$ . Of course  $\delta x_m$  is of the same order as  $\bar{x}$ . But since the  $\delta x_m$  are independent zero average random variables, by a reasoning analogous to the one advanced in subsection (6.3.2), the random part, when summed on  $m$  [as for example in (6.13)], is of order  $n^{-\frac{1}{2}}$  relative to the systematic part. It may therefore be neglected to lowest order. Upon combining (6.25) and (6.27) and setting  $\bar{x} = 2\pi$  we then find for  $r_m$  the second order recursion

$$r_{m+1} - 2r_m + r_{m-1} = 2\pi n^{-\frac{3}{2}} f_m, \quad (6.30)$$

which holds the key to much of what follows.

#### 6.4.2 Solution in terms of Fourier transforms

In order to solve the recursion (6.30) we define the Fourier transforms

$$\hat{r}_q = n^{-1} \sum_{m=1}^n e^{2\pi i q m/n} r_m, \quad \hat{u}_q = n^{-1} \sum_{m=1}^n e^{2\pi i q (m-\frac{1}{2})/n} u_m, \quad (6.31)$$

and

$$\hat{f}_q = \frac{1}{2\pi n^{\frac{1}{2}}} \sum_{m=1}^n e^{2\pi i q m/n} f_m, \quad (6.32)$$

where  $q = 0, \pm 1, \pm 2, \dots, \pm(\frac{1}{2}n - \frac{1}{2})$  for  $n$  odd and  $q = 0, \pm 1, \pm 2, \dots, \pm(\frac{1}{2}n - 1), \frac{1}{2}n$  for  $n$  even. Here  $q$  is the ‘wavenumber’ and  $2\pi/q$  the ‘wavelength’ along the cell perimeter, the scaled index  $2\pi m/n$  being equal to the polar angle in the plane up to corrections of order  $n^{-\frac{1}{2}}$ . In Fourier language the unknowns  $u_m$  and  $r_m$  of (6.27) and (6.30) are readily solved in terms of the forces  $f_m$  and one finds

$$\hat{u}_q = -\frac{\pi}{in \sin \frac{\pi q}{n}} \hat{f}_q \simeq -\frac{1}{iq} \hat{f}_q, \quad q \neq 0, \quad (6.33)$$

$$\hat{r}_q = -\frac{\pi^2}{n^2 \sin^2 \frac{\pi q}{n}} \hat{f}_q \simeq -\frac{1}{q^2} \hat{f}_q, \quad q \neq 0. \quad (6.34)$$

The no-spiral constraint and the gauge condition become  $\hat{u}_0 = 0$  and  $\hat{r}_0 = 0$ , respectively. The  $\simeq$  signs in (6.33) and (6.34) indicate the limit  $n \rightarrow \infty$  at fixed  $q$  and amount to neglecting higher orders in  $q/n$ ; these are small if we restrict ourselves to  $q$  values on a scale  $\sim n^{1-\epsilon}$ , *i.e.* to spatial distances which, in units of the midpoint index  $m$ , scale as  $\sim n^\epsilon$  with  $\epsilon > 0$ .

Next, Fourier transforms of  $\delta x_q$  and  $\delta y_q$  will be defined as

$$\hat{X}_q = \frac{1}{2\pi n^{\frac{1}{2}}} \sum_{m=1}^n e^{2\pi i q m/n} \delta x_m, \quad \hat{Y}_q = \frac{1}{2\pi n^{\frac{1}{2}}} \sum_{m=1}^n e^{2\pi i q m/n} \delta y_m. \quad (6.35)$$

From (6.28) and (6.35) it then follows that

$$\hat{f}_q = -\frac{1}{2}(1 + e^{-2\pi i q/n})\hat{X}_q + \hat{Y}_q \simeq -(\hat{X}_q - \hat{Y}_q). \quad (6.36)$$

We remark that the factor  $n^{-\frac{1}{2}}$  in (6.35) is the usual one for a Fourier transform of  $n$  independent variables; it guarantees that  $\hat{X}_q$  and  $\hat{Y}_q$  are of order  $n^0$ . The factor  $n^{-1}$  in the Fourier transforms (6.31) and (6.32) then follows from the requirement that in each of the equations (6.33), (6.34), and (6.32) all members be of the same order in  $n$ .

### 6.4.3 Defining $\mathbb{V}_1$

Employing (6.31) and (6.35) we are able to express  $\mathbb{V}$  of equation (6.24) as

$$\mathbb{V} = \sum_{q \neq 0} [\hat{u}_q \hat{u}_{-q} + 2\hat{r}_q \hat{r}_{-q} + 2\hat{r}_q \hat{Y}_{-q}] + \mathcal{O}(n^{-\frac{1}{2}}), \quad (6.37)$$

where  $q$  runs through all nonzero integers and where we Fourier transformed using that  $b_m$  and  $-c_m$  are both equal to  $u_m + \mathcal{O}(n^{-\frac{1}{2}})$ . We now insert the solutions (6.33) and (6.34) in (6.37). We relegate to the term  $n^{-\frac{1}{2}}\mathbb{V}_2$  in (6.1) also the higher orders in  $q/n$  discussed above and arrive at

$$\mathbb{V} = \mathbb{V}_1 + \mathcal{O}(n^{-\frac{1}{2}}) \quad (6.38)$$

where we finally decide to define  $\mathbb{V}_1$  as

$$\mathbb{V}_1 = \sum_{q \neq 0} \left[ \left( \frac{1}{q^2} + \frac{2}{q^4} \right) (\hat{X}_q - \hat{Y}_q)(\hat{X}_{-q} - \hat{Y}_{-q}) + \frac{2}{q^2} (\hat{X}_q - \hat{Y}_q) \hat{Y}_{-q} \right]. \quad (6.39)$$

In (6.39) the sum on  $q$  converges due to the presence of the factors  $1/q^2$  and  $1/q^4$ . This confirms again that  $\mathbb{V}_1$  is of order  $n^0$ . The contributions to  $\mathbb{V}_1$  come essentially from wavelengths at the scale of the cell itself and are fully separated in scale from the independent local degrees of freedom that entered the noninteracting calculation.

In this section we have extracted from  $\mathbb{V}$ , defined by (2.54), its leading order term  $\mathbb{V}_1$ , given by (6.39). With this expression at our disposal we can now go ahead and calculate the consequences of the interaction. Expression (6.39) could be considered as an elastic Hamiltonian in the harmonic approximation, if it were not for the fact that the  $\hat{X}_q$  and  $\hat{Y}_q$  are non-Gaussian variables. It will appear below that the deviations from Gaussian statistics affect the results of interest only to higher order in  $n^{-\frac{1}{2}}$ .

## 7 First order

At the end of section 6 we concluded that  $\mathbb{V}$  is of order  $n^0$ . This means, more precisely, that it is  $\sim n^0$  in the region of phase space anticipated to contribute dominantly to the average  $\langle \exp(-\mathbb{V}_1) \rangle_0$ . We therefore expect this average also to be of order  $n^0$ . By showing that below we will accomplish an essential step. The development of this and the following section are standard mathematical physics.

### 7.1 Calculation of $p_n$

We consider the quadratic form  $\mathbb{V}_1$  given by (6.39) above. It is useful to set

$$\hat{\mathbf{Z}}_q = (\hat{X}_q, \hat{Y}_q) \quad (7.1)$$

and to write (6.39) more compactly as

$$\mathbb{V}_1 = \sum_{q \neq 0} \hat{\mathbf{Z}}_q \cdot \mathbf{V}_q \cdot \hat{\mathbf{Z}}_{-q}^T, \quad (7.2)$$

where the superscript T indicates transposition and where

$$\mathbf{V}_q = \begin{pmatrix} q^{-2} + 2q^{-4} & -2q^{-4} \\ -2q^{-4} & -q^{-2} + 2q^{-4} \end{pmatrix}. \quad (7.3)$$

The exponential  $\exp(-\mathbb{V}_1)$  with  $\mathbb{V}_1$  given by (7.2) has to be substituted for  $\exp[-\mathbb{V}(x, y)]$  in (3.5). The variables of integration  $x_m$  and  $y_m$  are now coupled by  $\mathbb{V}_1$ . In order to decouple them we apply a Hubbard-Stratonovich transformation. We write

$$\hat{\boldsymbol{\psi}}_q = (\hat{\psi}_q^x, \hat{\psi}_q^y) \quad (7.4)$$

and represent  $\exp(-\mathbb{V}_1)$  as the integral

$$e^{-\mathbb{V}_1} = \int \left[ \prod_{q \neq 0} d\hat{\boldsymbol{\psi}}_q \frac{(\det \mathbf{V}_q^{-1})^{\frac{1}{2}}}{4\pi} \right] \exp \left[ -\frac{1}{4} \sum_{q \neq 0} \hat{\boldsymbol{\psi}}_q \cdot \mathbf{V}_q^{-1} \cdot \hat{\boldsymbol{\psi}}_{-q}^T + i \sum_{q \neq 0} \hat{\boldsymbol{\psi}}_q \cdot \hat{\mathbf{Z}}_{-q}^T \right]. \quad (7.5)$$

We have that  $\det \mathbf{V}_q = -q^{-4}$  is negative and the paths of integration of the  $\hat{\psi}_q^x$  and  $\hat{\psi}_q^y$  in the complex plane should be chosen such that the integrals in (7.5) exists. Let  $\psi_m^x$  and  $\psi_m^y$  be related to  $\hat{\psi}_q^x$  and  $\hat{\psi}_q^y$  in the same way that  $\delta x_m$  and  $\delta y_m$  are related to  $\hat{X}_q$  and  $\hat{Y}_q$ , *i.e.* by (6.35). We set  $\hat{\psi}_0^x = \hat{\psi}_0^y = 0$ . Then

$$\begin{aligned} \sum_{q \neq 0} \hat{\psi}_q \cdot \hat{\mathbf{Z}}_{-q}^T &= \sum_{q \neq 0} (\hat{\psi}_q^x \hat{X}_{-q} + \hat{\psi}_q^y \hat{Y}_{-q}) \\ &= \frac{1}{(2\pi)^2} \sum_{m=1}^n (\psi_m^x \delta x_m + \psi_m^y \delta y_m). \end{aligned} \quad (7.6)$$

We insert (7.6) in (7.5) and the integral representation (7.5) in (3.5). We commute the  $\hat{\psi}_q$  integrations to the outside and are then able to perform the integrals on the  $x_m$  and  $y_m$ . The result is

$$\begin{aligned} p_n &\simeq \int \left[ \prod_{q \neq 0} d\hat{\psi}_q \frac{(\det \mathbf{V}_q^{-1})^{\frac{1}{2}}}{4\pi} \right] \exp \left[ -\frac{1}{4} \sum_{q \neq 0} \hat{\psi}_q \cdot \mathbf{V}_q^{-1} \cdot \hat{\psi}_{-q}^T \right] \\ &\times \frac{(n-1)!}{2n} \pi^{-n} n^{-3n} \left[ \frac{n}{2\pi i} \int du \frac{e^{2\pi n u}}{u^{2n}} \prod_{m=1}^n \left( 1 - \frac{i\psi_m^x}{4\pi^2 u} \right)^{-2} \right] \\ &\times \left[ \frac{n}{2\pi i} \int dv \frac{e^{2\pi n v}}{v^n} \prod_{m=1}^n \left( 1 - \frac{i\psi_m^y}{4\pi^2 v} \right)^{-1} \right]. \end{aligned} \quad (7.7)$$

Hence one sees that the  $u$  and  $v$  integrals factorize again. The products on  $m$  represent an insertion of the type  $h(u, v)$  discussed in section 4.1, which may be taken at the saddle points  $u = u_c = \pi^{-1}$  and  $v = v_c = (2\pi)^{-1}$ . Doing so and exponentiating these products we get

$$\begin{aligned} p_n &\simeq p_n^{(0)} \int \left[ \prod_{q \neq 0} d\hat{\psi}_q \frac{(\det \mathbf{V}_q^{-1})^{\frac{1}{2}}}{4\pi} \right] \exp \left[ -\frac{1}{4} \sum_{q \neq 0} \hat{\psi}_q \cdot \mathbf{V}_q^{-1} \cdot \hat{\psi}_{-q}^T \right] \\ &\times \exp \left[ -\frac{1}{4} \sum_{q \neq 0} \hat{\psi}_q^x \hat{\psi}_{-q}^x - \sum_{m=1}^n 2\mathcal{R}(\psi_m^x) \right] \\ &\times \exp \left[ -\frac{1}{2} \sum_{q \neq 0} \hat{\psi}_q^y \hat{\psi}_{-q}^y - \sum_{m=1}^n \mathcal{R}(2\psi_m^y) \right], \end{aligned} \quad (7.8)$$

where  $p_n^{(0)}$  was given in (4.1) and the ‘remainder’  $\mathcal{R}(x)$  stands for the infinite series

$$\mathcal{R}(x) = \sum_{k=3}^{\infty} \frac{1}{k} \left( \frac{ix}{4\pi} \right)^k. \quad (7.9)$$

We will delay to section 7.2.2 the proof that in the limit  $n \rightarrow \infty$  the remainders in (7.8) contribute only to relative order  $n^{-1}$ . Admitting this here and

setting

$$\mathbf{E} = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix} \quad (7.10)$$

we therefore have

$$\begin{aligned} p_n &\simeq p_n^{(0)} \int \left[ \prod_{q \neq 0} d\hat{\psi}_q \frac{(\det \mathbf{V}_q^{-1})^{\frac{1}{2}}}{4\pi} \right] \exp \left[ -\frac{1}{4} \sum_{q \neq 0} \hat{\psi}_q \cdot (\mathbf{E} + \mathbf{V}_q^{-1}) \cdot \hat{\psi}_{-q}^T \right] \\ &= p_n^{(0)} \prod_{q \neq 0} \frac{(\det \mathbf{V}_q^{-1})^{\frac{1}{2}}}{[\det(\mathbf{E} + \mathbf{V}_q^{-1})]^{\frac{1}{2}}} \\ &= p_n^{(0)} \prod_{q=1}^{\infty} \Lambda_q^{-1}, \end{aligned} \quad (7.11)$$

where we abbreviated

$$\begin{aligned} \Lambda_q &= \det(\mathbf{1} + \mathbf{V}_q \mathbf{E}) \\ &= 1 - q^{-2} + 4q^{-4} \end{aligned} \quad (7.12)$$

with  $\mathbf{1}$  standing for the  $2 \times 2$  unit matrix. Upon combining (7.11) and (7.12) with expression (4.1) for  $p_n^{(0)}$  one finds for  $p_n$  the result announced in (1.2)-(1.3) of the introduction,

$$p_n = \frac{C}{4\pi^2} \frac{(8\pi^2)^n}{(2n)!} \left[ 1 + \mathcal{O}(n^{-\frac{1}{2}}) \right], \quad (7.13)$$

where

$$C = \prod_{q=1}^{\infty} (1 - q^{-2} + 4q^{-4})^{-1}. \quad (7.14)$$

In the course of the calculations of the previous sections we have encountered correction terms of relative order  $n^{-\frac{1}{2}}$ . In equation (7.13) we have included these  $\mathcal{O}(n^{-\frac{1}{2}})$  terms again, although strictly speaking we have not shown that their net effect does not happen to vanish.

This is the first analytic determination of the large  $n$  behavior of  $p_n$ . Conjectures in earlier work (see [3]) have almost invariably had the form of decaying exponentials, whether or not multiplied by a power of  $n$ , or sums of such expressions. Only Drouffe and Itzykson [27] come close to our analytic expression: they assume that to leading order  $\log p_n \simeq \alpha n$  and estimate  $\alpha \approx -2$ , a value confirmed by (7.13). We refer to [31] for a plot of the asymptotic expression (7.13) and a comparison with Drouffe and Itzykson's simulation data.



## 7.2 Correlations to first order

The calculational technique that led to the expression for  $p_n$  in the preceding subsection, may also be used to find correlation functions at long distances.

### 7.2.1 Long distance correlations along the perimeter

We will study the correlations between the angles  $\xi_\ell$  and  $\eta_m$  in terms of their scaled counterparts  $x_\ell$  and  $y_m$ . Passing to Fourier transforms with the aid of (6.35) we have

$$\langle \delta x_\ell \delta x_m \rangle = \frac{(2\pi)^2}{n} \sum_{q \neq 0} e^{2\pi i(k-\ell)q/n} \langle \hat{X}_q \hat{X}_{-q} \rangle \quad (7.15)$$

and analogous relations for  $\langle \delta y_\ell \delta y_m \rangle$  and the cross correlation  $\langle \delta x_\ell \delta y_m \rangle$ . From definition (6.4) and equations (7.5) and (7.6) it is clear that we obtain the first order expression  $\langle \hat{X}_q \hat{X}_{-q} \rangle_1$  by inserting in the integrand of (7.5) a second derivative  $-\partial^2 / \partial \hat{\psi}_q^x \partial \hat{\psi}_{-q}^x$  that acts on the factor  $\exp(-i \sum_{q \neq 0} \hat{\psi}_q \cdot \hat{\mathbf{Z}}_{-q}^T)$ . Carrying the calculation through we find that we have to differentiate the exponential factors in the second line of (7.8), whence we have, for  $q \neq 0$ ,

$$\langle \hat{X}_q \hat{X}_{-q} \rangle_1 = \frac{1}{2} - \frac{1}{4} \langle \hat{\psi}_q^x \hat{\psi}_{-q}^x \rangle_G + \dots \quad (7.16)$$

Here  $\langle \dots \rangle_G$  is a Gaussian average with respect to the weight  $\text{cst} \times \exp \left[ -\frac{1}{4} \sum_{q \neq 0} \hat{\psi}_q \cdot (\mathbf{E} + \mathbf{V}_q^{-1}) \cdot \hat{\psi}_{-q}^T \right]$  and the dots represent a well-defined series of quartic and higher Gaussian averages coming from the expansion of the remainder  $\mathcal{R}$  in (7.8). When evaluated, these higher order terms contain factors  $n^{-1}$ , as will be shown in section 7.2.2. After completing the calculation for the other basic correlation functions we obtain

$$\begin{aligned} \langle \hat{X}_q \hat{X}_{-q} \rangle_1 &\simeq \frac{1}{2} - \frac{1}{4} \langle \hat{\psi}_q^x \hat{\psi}_{-q}^x \rangle_G, \\ \langle \hat{Y}_q \hat{Y}_{-q} \rangle_1 &\simeq 1 - \langle \hat{\psi}_q^y \hat{\psi}_{-q}^y \rangle_G, \\ \langle \hat{X}_q \hat{Y}_{-q} \rangle_1 &\simeq -\frac{1}{2} \langle \hat{\psi}_q^x \hat{\psi}_{-q}^y \rangle_G, \quad q \neq 0, \end{aligned} \quad (7.17)$$

where the  $\simeq$  sign indicates, as before, asymptotic equality for  $n \rightarrow \infty$ . In view of (6.5) we may drop the index 1 on the averages in (7.17). The three Gaussian averages in (7.17) are due to the interaction  $\mathbb{V}_1$ . They come in addition to the constants ( $\frac{1}{2}$ , 1, and 0) that represent the Fourier transformed noninteracting averages (4.6). Straightforward calculation yields

$$\begin{aligned} \langle \hat{\psi}_q^x \hat{\psi}_{-q}^x \rangle_G &= 2\Lambda_q^{-1} q^{-2}, \\ \langle \hat{\psi}_q^y \hat{\psi}_{-q}^y \rangle_G &= -2\Lambda_q^{-1} (q^{-2} - q^{-4}), \\ \langle \hat{\psi}_q^x \hat{\psi}_{-q}^y \rangle_G &= -4\Lambda_q^{-1} q^{-4}, \quad q \neq 0, \end{aligned} \quad (7.18)$$

From (6.5), (7.18), and (7.17) we obtain for the basic correlation functions the following set of first order expressions,

$$\begin{aligned}\langle \hat{X}_q \hat{X}_{-q} \rangle &\simeq \frac{1}{2} - \frac{1}{2} \Lambda_q^{-1} q^{-2}, \\ \langle \hat{Y}_q \hat{Y}_{-q} \rangle &\simeq 1 + 2 \Lambda_q^{-1} (q^{-2} - q^{-4}), \\ \langle \hat{X}_q \hat{Y}_{-q} \rangle &\simeq 2 \Lambda_q^{-1} q^{-4}, \quad q \neq 0,\end{aligned}\tag{7.19}$$

which represent the main result of this subsection. The terms with the factor  $\Lambda_q^{-1}$  in (7.19) have their origin in the interaction  $\mathbb{V}_1$ . The important point about the  $\hat{\psi}\hat{\psi}$  correlations is that they all decay at least as  $\sim 1/q^2$  when  $q \rightarrow \infty$ . This will turn out to be an essential element in our analysis below. Equations (7.19) may be inverse Fourier transformed according to (7.15) and its analogs so as to yield expressions for the spatial correlations  $\langle \delta x_\ell \delta x_m \rangle$ ,  $\langle \delta y_\ell \delta y_m \rangle$ , and  $\langle \delta x_\ell \delta y_m \rangle$ . We will work these out in detail only for the more restricted correlation function  $\langle (\hat{X}_q - \hat{Y}_q)(\hat{X}_{-q} - \hat{Y}_{-q}) \rangle = \langle \hat{f}_q \hat{f}_{-q} \rangle$  to be considered in section 8. Because of our discussions on spatial scales at the end of section 6.3.3 and following (6.33)-(6.34), the real space correlations so obtained are valid only at distances large compared to the individual perimeter segments.

### 7.2.2 The remainders $\mathcal{R}$

Having gained control of the correlation functions, we can now prove our claim that the remainders  $\mathcal{R}$  in the exponentials in (7.8) contribute only to higher order in  $n^{-1}$ . We may expand these exponentials as perturbation series in the  $\psi_m^x$  and  $\psi_m^y$ . The odd terms vanish because of symmetry. The lowest order contributing terms are therefore proportional to  $\sum_m (\psi_m^x)^4$  and  $\sum_m (\psi_m^y)^4$ . Let us consider the latter. It leads to a correction term which, relative to the term calculated, is of order  $\sum_m \langle (\psi_m^y)^4 \rangle_G$ . We therefore evaluate

$$\begin{aligned}\sum_m \langle (\psi_m^y)^4 \rangle_G &= \frac{(2\pi)^4}{n} \sum_{q_1, q_2, q_3, q_4 \neq 0} \delta_{q_1+q_2+q_3+q_4, 0} \langle \hat{\psi}_{q_1}^y \hat{\psi}_{q_2}^y \hat{\psi}_{q_3}^y \hat{\psi}_{q_4}^y \rangle_G \\ &= \frac{3(2\pi)^4}{n} \left[ \sum_{q \neq 0} \langle \hat{\psi}_q^y \hat{\psi}_{-q}^y \rangle_G \right]^2,\end{aligned}\tag{7.20}$$

where we used Wick's theorem. The fact that  $\langle \hat{\psi}_q^y \hat{\psi}_{-q}^y \rangle_G$  decays as  $q^{-2}$  is crucial here. It means that the sum on  $q$  inside the brackets in (7.20) converges, and hence  $\sum_m \langle (\psi_m^y)^4 \rangle_G$  is of order  $n^{-1}$ . We therefore had the right to neglect the  $\mathcal{R}$  terms in (7.8). The other terms in the perturbation series in the  $\psi_m^x$  and  $\psi_m^y$  can be argued in a similar way to be at least of order  $n^{-1}$ .

### 7.2.3 Debye-Hückel

We remark that in order to arrive at the above results, as well as those that will follow in section 8, one may bypass the Hubbard-Stratonovich transformation by considering the variables of integration  $\hat{X}_q$  and  $\hat{Y}_q$  as zero mean Gaussian variables of variances determined by those of  $\langle \delta x_m^2 \rangle_0$  and  $\langle \delta y_m^2 \rangle_0$  given in (4.2), *i.e.* with weight

$$\exp\left(-\sum_{q \neq 0} \hat{\mathbf{Z}}_q \cdot \mathbf{E}^{-1} \cdot \hat{\mathbf{Z}}_{-q}\right) = \exp\left(-\sum_{q \neq 0} \hat{X}_q \hat{X}_{-q} - \frac{1}{2} \sum_{q \neq 0} \hat{Y}_q \hat{Y}_{-q}\right). \quad (7.21)$$

The procedure of replacing non-Gaussian variables by Gaussian ones of the same mean and variance is known in plasma physics as the Debye-Hückel approximation; it is exact in the limit of high temperature. This limit happens to apply to our case, where the role of the temperature is played by  $n$ : the interaction ‘energy’  $\mathbb{V}_1$  would be extensive (*i.e.* proportional to  $n$ ) in an ordinary thermodynamical system, but is here of order  $n^0$ , which is equivalent to having an inverse-temperature prefactor equal to  $n^{-1}$ .

In the preceding subsection we have chosen to go through the full Hubbard-Stratonovich transformation. This has allowed us to show that the procedure is exact and to prepare the way for the calculation of higher order terms, if desired, in the power series expansion in  $n$ .

## 8 Weight functional for the perimeter

### 8.1 Weight function for the $\hat{f}_q$

Given that for large  $n$  the perimeter approaches a circle, it is natural to ask what the probability distribution is for its deviations from circularity. This question has prompted work, in particular, in the mathematical literature [35, 36, 37], but within the framework of large-cell limits different from ours. Deviations from circularity may be characterized in various ways, *e.g.* as the difference between the radii of the smallest circumscribed and the largest inscribed circle [35, 37]. Here we will exhibit in full generality the weight of a specific trajectory of the perimeter, characterized by the set of scaled ‘excess midpoint distances’  $\{r_1, r_2, \dots, r_n\}$ . The Fourier transforms  $\hat{r}_q$  of these distances are related to the *differences*  $\hat{X}_q - \hat{Y}_q$  by  $q^2 \hat{r}_q = -\hat{f}_q = \hat{X}_q - \hat{Y}_q$ , which follows from (6.34) and (6.36). Hence one of the variables  $\hat{X}_q$  and  $\hat{Y}_q$  is redundant for the description at which we are aiming. The probability distribution of the  $\hat{X}_q$  and  $\hat{Y}_q$  is, up to normalization, equal to the weight  $\exp(-\mathbb{V}_1)$  with  $\mathbb{V}_1$  given by (6.39). In order to derive from it the marginal distribution  $P_1(\{\hat{f}_q\})$  for the  $\hat{f}_q$ , we short-circuit the Hubbard-Stratonovich procedure in the way indicated in section 7.2.3. Upon integrating  $\exp(-\mathbb{V}_1)$

directly with the appropriate noninteracting Gaussian weight (7.21) we find the first order probability law

$$P_1(\{\hat{f}_q\}) \simeq \text{cst} \times \exp\left(-\frac{1}{3} \sum_{q \neq 0} \Lambda_q \hat{f}_q \hat{f}_{-q}\right) \quad (8.1)$$

with  $\Lambda_q$  given by (7.12) and where, because of (6.29), it is understood that  $\hat{f}_0 = 0$ . The law (8.1) leads to the correlation

$$\langle \hat{f}_q \hat{f}_{q'} \rangle \simeq \frac{3}{2} \Lambda_q^{-1} \delta_{q+q', 0}, \quad q, q' \neq 0, \quad (8.2)$$

which may alternatively be derived from (7.19). We remark parenthetically that the weight (8.1) is suitable for computer simulation of the large scale structure of Voronoi cells. The rest of this section elaborates on equations (8.1) and (8.2).

## 8.2 Differential equation for $R(\phi)$

For  $n \rightarrow \infty$  it is possible on those scales to consider the angle

$$\phi = \frac{2\pi m}{n} \quad (8.3)$$

as a continuous variable [49] and to write

$$r_m = r(\phi), \quad f_m = \frac{2\pi}{n^{\frac{1}{2}}} f(\phi). \quad (8.4)$$

Recursion (6.30) relating the  $r_m$  to the  $f_m$  then becomes the second order differential equation

$$r''(\phi) = f(\phi). \quad (8.5)$$

Here  $f(\phi)$  is Gaussian noise whose correlation function  $\langle f(\phi) f(\phi') \rangle$  we will now determine. To this end we set  $\phi' = 2\pi m' n^{-1}$  and consider  $\langle f_m f_{m'} \rangle$ , which with the aid of (6.32) may be written as

$$\begin{aligned} \langle f_m f_{m'} \rangle &= \frac{(2\pi)^2}{n} \sum_q \sum_{q'} e^{-2\pi i(qm + q'm')/n} \langle \hat{f}_q \hat{f}_{q'} \rangle \\ &\simeq \frac{3(2\pi)^2}{2n} \sum_{q \neq 0} e^{-iq(\phi - \phi')} \Lambda_q^{-1}, \end{aligned} \quad (8.6)$$

where to arrive at the second line we used (8.2). Passing from  $f_m$  to  $f(\phi)$  and using (7.12) we obtain from (8.6)

$$\begin{aligned} \langle f(\phi) f(\phi') \rangle &\simeq \frac{3}{2} \left[ \sum_{q \neq 0} e^{-iq(\phi - \phi')} + \Gamma(\phi - \phi') \right] \\ &= \frac{3}{2} [2\pi \delta(\phi - \phi') - 1 + \Gamma(\phi - \phi')], \end{aligned} \quad (8.7)$$

where the  $2\pi$ -periodic function  $\Gamma(\phi)$  is given by

$$\Gamma(\phi) = 2 \sum_{q=1}^{\infty} \frac{q^{-2} - 4q^{-4}}{1 - q^{-2} + 4q^{-4}} \cos q\phi. \quad (8.8)$$

This correlation function result merits a comment. It is instructive to calculate, for comparison to (8.6) and (8.7), also the simpler noninteracting average  $\langle f_m f_{m'} \rangle_0$ . To this end one expresses  $f_m$  and  $f_{m'}$  in terms of the  $\delta x_\ell$  and  $\delta y_\ell$  using (6.28) and employs equations (4.6). In the continuum limit it then appears that in (8.7) the terms  $2\pi\delta(\phi - \phi') - 1$  represent the noninteracting result and  $\Gamma(\phi - \phi')$  the contribution generated by the interaction. The expression for this contribution was derived under the restrictive condition that  $|\phi - \phi'| \gg 2\pi n^{-1}$ . We speculate that (8.7) is valid, in fact, for  $|\phi - \phi'|$  scaling as  $\sim n^{-1+\epsilon}$  with arbitrary  $\epsilon > 0$ . Finally we remark that expressions similar to (8.7) for the correlations  $\langle \delta x(\phi) \delta x(\phi') \rangle$ ,  $\langle \delta x(\phi) \delta y(\phi') \rangle$ , and  $\langle \delta y(\phi) \delta y(\phi') \rangle$  may be found in a fully analogous way.

In the continuum limit the Fourier transform of  $\Lambda_q$  becomes

$$\sum_{q \neq 0} e^{iq(\phi - \phi')} \Lambda_q = 2\pi\delta(\phi - \phi') - 1 + U(\phi - \phi') \quad (8.9)$$

with the  $2\pi$ -periodic function  $U(\phi)$  given by

$$U(\phi) = -2 \sum_{q=1}^{\infty} (q^{-2} - 4q^{-4}) \cos q\phi. \quad (8.10)$$

Similarly, in (8.9) the term  $U(\phi - \phi')$  represents the effect of the interaction. Both  $\Gamma(\phi)$  and  $U(\phi)$  are of zero average on the interval  $[0, 2\pi]$ .

The sums on  $q$  in equations (8.8) and (8.10) may be carried out analytically [see reference [50], equations (1.443.3), (1.443.6), and (1.445.2)]. One finds that (8.10) leads to the quartic polynomial

$$U(\phi) = -\frac{1}{6}\pi^2\left(\frac{7}{15}\pi^2 - 1\right) + \frac{1}{2}\left(\frac{2}{3}\pi^2 - 1\right)(\phi - \pi)^2 - \frac{1}{6}(\phi - \pi)^4, \quad 0 \leq \phi \leq 2\pi, \quad (8.11)$$

and its  $2\pi$ -periodic continuation. We have written it such as to bring out its symmetry under the replacement  $\phi \mapsto 2\pi - \phi$ . The potential  $U(\phi)$  in (8.10) consists of two terms which merit separate discussion. The first term is the one-dimensional electrostatic interaction but with opposite sign, so that it is attractive for like charges; and the second one is repulsive. Near the origin we have the expansion

$$U(\phi) = U(0) + \pi|\phi| - \left(\frac{1}{2} + \frac{2}{3}\pi^2\right)\phi^2 + \dots, \quad (8.12)$$

in which the linear term comes from the electrostatic interaction. As a result  $U(\phi)$  has maxima at  $\phi = \pm\phi_U$ , where  $\phi_U = \pi[1 - \{1 - 3/(2\pi^2)\}^{\frac{1}{2}}] = 0.079\dots\pi$ . We will not exhibit the analytic expression of  $\Gamma(\phi)$  but have represented its graph in figure 3. It has minima at  $\phi = \pm\phi_R$ , where  $\phi_R \approx 0.139\dots\pi$ .




Figure 3: The functions  $U(\phi)$  and  $\Gamma(\phi)$  of equations (8.11) and (8.8), respectively.

### 8.3 Weight functional for $R(\phi)$

Using (8.9) we subject the quadratic form in the exponential of (8.1) to an inverse Fourier transformation. In the continuum limit the probability distribution  $P_1(\{\hat{f}_q\})$  becomes a functional of  $f(\phi)$ , or, because of (8.5), of  $r''(\phi)$ . We find

$$P_1[r(\phi)] \simeq \text{cst} \times \exp(-H[r''(\phi)]), \quad (8.13)$$

$$\begin{aligned} H[r''(\phi)] = & \frac{1}{6\pi} \int_0^{2\pi} d\phi [r''(\phi)]^2 \\ & + \frac{1}{12\pi^2} \int_0^{2\pi} d\phi_1 \int_0^{2\pi} d\phi_2 r''(\phi_1) U(\phi_1 - \phi_2) r''(\phi_2). \end{aligned} \quad (8.14)$$

Here the last line, which is the contribution from the interaction  $\mathbb{V}$ , is analogous to the Coulomb energy of a line charge density  $r''(\phi)$  in a periodic one-dimensional system. Equations (8.13) and (8.14) are valid for functions  $r(\phi)$  coarse grained to a scale much larger than the distance between the individual vertices, *i.e.*, to angle differences  $\Delta\phi \gg 2\pi n^{-1}$ . Moreover, the sum rules encountered during its derivation now lead to three restrictions on the functions  $r(\phi)$  that enter (8.14):

(i) The geometrical sum rule (6.29), taken in the continuum limit and combined with the differential equation (8.5), implies the ‘charge neutrality’ condition  $\int_0^{2\pi} d\phi r''(\phi) = r'(2\pi) - r'(0) = 0$ . It follows that we may add an arbitrary constant to the potential  $U$  in (8.14).

(ii) The no-spiral constraint becomes  $\int_0^{2\pi} d\phi r'(\phi) = r(2\pi) - r(0) = 0$ .

(iii) The gauge condition takes the form  $\int_0^{2\pi} d\phi r(\phi) = 0$ .

This completes our discussion of the first order probability functional  $P_1$ .

We now wish to reconvert the above findings to results for the original midpoint distances  $R_m = R(\phi) = R_c + \delta R(\phi)$ . From (6.12) taken in the continuum limit and combined with (1.10) we find

$$\delta R(\phi) = \pi^{-\frac{1}{2}} [r_{\text{av}} + r(\phi)] + \mathcal{O}(n^{-\frac{1}{2}}). \quad (8.15)$$

It follows that the probability distribution of  $R(\phi)$  is determined by the product of  $p_{\text{av}}(r_{\text{av}})$  given in (2.45) and  $P_1[r(\phi)]$  given in (8.13)-(8.14). Still setting

$$F(\phi) = \pi^{-\frac{1}{2}} f(\phi) \quad (8.16)$$

we are led to the results (1.11)-(1.13) announced in the introduction, in which the factors  $4\lambda$  have been restored.

## 8.4 An application: fluctuations of $R_m$ around $R_{\text{av}}$

As an application we determine how, within a single realization of an  $n$ -sided cell, the midpoint distances  $R_m$  deviate from their average  $R_{\text{av}}$ . This extends the considerations of section 2.4.4, where we showed that  $R_{\text{av}}$  is itself within order  $n^0$  from its ensemble average  $R_c = \pi^{-\frac{1}{2}} n^{\frac{1}{2}}$ .

We have straightforwardly

$$\begin{aligned} n^{-1} \sum_{m=1}^n \langle (R_m - R_{\text{av}})^2 \rangle &= n^{-2} R_{\text{av}}^2 \sum_{m=1}^n \langle r_m^2 \rangle \\ &= n^{-1} R_{\text{av}}^2 \sum_{q \neq 0} \langle \hat{r}_q \hat{r}_{-q} \rangle \\ &= n^{-1} R_{\text{av}}^2 \sum_{q \neq 0} q^{-4} \langle \hat{f}_q \hat{f}_{-q} \rangle \\ &= \frac{3}{\pi} \sum_{q=1}^{\infty} q^{-4} (1 - q^{-2} + 4q^{-4})^{-1}, \quad (8.17) \end{aligned}$$

where the first step is based on the first line of (6.12), in the third step we used (6.34), and in the last step we inserted the explicit expressions for  $R_{\text{av}}$  and  $\langle \hat{f}_q \hat{f}_{-q} \rangle$ . Equation (8.17) yields the result announced in (1.9).

## 8.5 Convergence of $R(\phi)$ and $S(\phi)$

The set of midpoints  $\{\mathbf{R}_m | m = 1, \dots, n\}$  and the set of vertices  $\{\mathbf{S}_m | m = 1, \dots, n\}$  each define an  $n$ -sided polygon. In the limit of large  $n$  these polygons become the curves  $R(\phi)$  and  $S(\phi)$ , respectively. Of these,  $S(\phi)$  is the

true perimeter. We show here that for  $n \rightarrow \infty$  the curve  $R(\phi)$  converges rapidly to  $S(\phi)$  so that in the discussion above we were justified in referring to  $R(\phi)$  as the perimeter.

Two points  $\mathbf{R}_m = (R_m, \Phi_m)$  and  $\mathbf{S}_m = (S_m, \Psi_m)$  have angular coordinates such that  $|\Phi_m - \Psi_m| = |\gamma_m| = \mathcal{O}(n^{-\frac{1}{2}})$ . Since  $\Phi_\ell$  increases by steps of order  $\sim n^{-1}$  when  $\ell$  increases by 1, there exists a third point  $\mathbf{R}_{m'} = (R_{m'}, \Phi_{m'})$ , with  $m'$  depending on  $m$ , such that

$$|\Phi_{m'} - \Psi_m| = \mathcal{O}(n^{-1}). \quad (8.18)$$

Typically we will have  $|m' - m| = \mathcal{O}(n^{\frac{1}{2}})$ . In order to estimate how the distance between  $\mathbf{R}_{m'}$  and  $\mathbf{S}_m$  scales with  $n$  we write

$$|\mathbf{R}_{m'} - \mathbf{S}_m|^2 \simeq (R_{m'} - S_m)^2 + R_c^2(\Phi_{m'} - \Psi_m)^2, \quad (8.19)$$

where we used that to leading order  $R_{m'} \simeq S_m \simeq R_c$ . In order to deal with the first term on the RHS of (8.19) we consider the triangle inequality

$$|R_{m'} - S_m| \leq |R_{m'} - R_m| + |R_m - S_m|. \quad (8.20)$$

Since  $R_m = S_m \cos \gamma_m$ , we have

$$|R_m - S_m| \simeq \frac{1}{2} R_c \gamma_m^2 = \mathcal{O}(n^{-\frac{1}{2}}). \quad (8.21)$$

Furthermore, from (6.25) it follows that

$$\begin{aligned} r_{m'} - r_m &\simeq 2\pi n^{-1} \sum_{\ell=m+1}^{m'} u_\ell \\ &\simeq 2\pi n^{-1} |m' - m| u_m, \end{aligned} \quad (8.22)$$

where in the last step we used that for  $\ell$  in an interval of length  $|m' - m| \sim n^{\frac{1}{2}}$  the value of  $u_\ell$  varies negligibly with respect to its typical value. Hence (8.22) leads to  $|r_{m'} - r_m| = \mathcal{O}(n^{-\frac{1}{2}})$ , which together with (6.12) gives

$$|R_{m'} - R_m| = R_c^2 n^{-\frac{1}{2}} |r_{m'} - r_m| = \mathcal{O}(n^{-\frac{1}{2}}). \quad (8.23)$$

Upon using (8.21) and (8.23) in (8.20) we find that  $|R_{m'} - S_m| \leq \mathcal{O}(n^{-\frac{1}{2}})$ . When substituting this in (8.19) and using (8.18) one obtains that  $|\mathbf{R}_{m'} - \mathbf{S}_m| = \mathcal{O}(n^{-\frac{1}{2}})$ . This says that any vertex on the polygon through the  $S_m$  is, typically, within a distance  $\sim n^{-\frac{1}{2}}$  of a vertex of the polygon through the  $R_m$ . In the limit  $n \rightarrow \infty$  we may therefore indifferently refer to either of them as the perimeter.



## 9 Lewis' law

The average area  $A$  of a Voronoi cell is equal to  $A = \lambda^{-1}$ , where, here and henceforth, we restore the particle density  $\lambda$ . Its average number of sides  $\langle n \rangle$  is topologically constrained to  $\langle n \rangle = 6$ . One may now investigate the average area  $A_n$  of an  $n$ -sided Voronoi cell. Lewis' empirical law [2] states that  $A_n$  is linear in  $n$ . The law is often expressed with a single free parameter  $c$ ,

$$A_n = [c(n - 6) + 1]\lambda^{-1}, \quad (9.1)$$

so as to imply that  $A_6 = \lambda^{-1}$ , *i.e.*, the average area of a cell having  $\langle n \rangle$  sides is equal to the average area of all cells; this appealing identity is not, however, a mathematical necessity. The law (9.1) accounts very well for data on cell areas coming from a wide range of different experimental situations [3], despite deviations often found for the small- $n$  values  $n = 3$  and  $n = 4$ . Lewis' law also seems to apply [25, 17, 51, 52, 53] to models different from Poisson-Voronoi diagrams, whether dynamical or not.

The numerical estimates of the coefficient  $c$  presented in reference [3] are in the range  $0.199 - 0.257$ . In particular, Lauritsen *et al.* [28] estimate  $c = 0.23$ ; Drouffe and Itzykson [27], on the basis of their Monte Carlo simulations for large  $n$ , observe that  $c$  is 'very close' to  $\frac{1}{4}$ ; and Miles and Maillardet [54], on the basis of a comparison of the cell area with the area of its fundamental domain [40], conjecture that  $c$  'does not differ greatly' from  $\frac{1}{4}$ . It has been noticed [38] that when the 'natural' boundary condition  $A_2 = 0$  is imposed on (9.1), one finds by coincidence  $c = \frac{1}{4}$  exactly, whence

$$A_n = \frac{1}{4}(n - 2)\lambda^{-1}. \quad (9.2)$$

Various 'derivations' of Lewis' law (9.1) have been proposed [38, 39, 55], mostly involving the maximization of an entropy defined as a functional of the probability distribution  $\{p_j | j = 3, 4, \dots\}$ . None of these derivations, however, is based on first principles, and Chiu [56, 57] has shown that entropy arguments of this kind do not allow firm conclusions. Moreover, numerical data on  $A_3$  and  $A_4$  for Poisson-Voronoi diagrams clearly show deviations from linearity and hence rule out laws such as (9.1) and (9.2) for the full (nonasymptotic) regime.

It suffices to extend the arguments of the preceding sections only very slightly to arrive at a new analytic result. Let us look at the midpoint distances  $R_m$ . By combining equations (2.29), (2.37), and (6.10) we find that  $R_m = R_{av}[1 + \mathcal{O}(n^{-\frac{1}{2}})]$ . Furthermore the discussion of subsection 2.4.3 has shown that  $R_{av} = R_c[1 + \mathcal{O}(n^{-\frac{1}{2}})]$ . It follows that  $R_m = R_c[1 + \mathcal{O}(n^{-\frac{1}{2}})]$  for all  $m$ , where we recall that  $R_c = (n/4\pi\lambda)^{\frac{1}{2}}$ . Hence the cell perimeter stays within a distance of order unity from a circle of radius  $\sim n^{\frac{1}{2}}$ . Therefore the area of the  $n$ -sided Voronoi cell is sharply peaked around an average  $A_n$

for which we find the asymptotic relation

$$A_n \simeq \pi R_c^2 = \frac{n}{4\lambda}. \quad (9.3)$$

Hence we have demonstrated from first principles that Lewis' law holds to leading order in an asymptotic expansion for large  $n$ , and that its coefficient is  $c = \frac{1}{4}$ .

## 10 Comments and conclusion

By setting out to find the probability  $p_n$  for a Voronoi cell to have  $n$  sides, we ended up unraveling the detailed statistical structure of the many-sided Voronoi cell. We have obtained the asymptotic expansion of  $\log p_n$  up to terms that vanish as  $n \rightarrow \infty$ , and have shown that a perturbation series may be set up for the higher order terms. We found that the problem is characterized by two spatial scales which in the limit  $n \rightarrow \infty$ , and to the order that we calculated, become completely disjoint. The first one is the microscopic scale and the second one the scale of the system size.

The exact methods and results of this work suggest, in retrospect, several heuristic considerations. These may be expected to lend to our arguments a flexibility that will allow them to be applied in more general situations. We reserve the heuristics, together with further extensions of the present results, to a companion paper [32]. Our work also opens the way to efficient Monte Carlo simulations of large Voronoi cells; we also leave this and other numerical applications for future work.

Voronoi cells in spatial dimensions higher than two have likewise received a great deal of attention in the literature [3]. We do not see how to calculate an analytic result analogous to our equations (1.2)-(1.3) in dimensions above two. The reason is that the idea of constructing a Markov process along the cell perimeter, which was at the root of the present work, has no obvious extension to higher dimensions. Nevertheless, we believe that it is possible without detailed calculation to transpose many of the scaling relations found in this work to higher dimensions.

The problem of random points in a plane considered here (the Poisson point process) is twin to that of random lines in a plane: the *Poisson line process*. Instead of Voronoi cells, random lines define *Crofton cells*, about whose statistics which one may ask similar questions. In recent mathematical literature Voronoi and Crofton cells have often been discussed together [34, 35, 37]. In physics random lines in a plane were introduced by Goudsmit [58], who was interested in the statistics of tracks in a cloud-chamber. He asked, specifically, about the probability that three or more independent tracks intersect at almost the same point and converted this question into

one about the area of the Crofton cell. We are confident that the methods of this paper can be made to work for the description of the large  $n$ -sided Crofton cell.

In metallography large cells were observed by Aboav [29] in the arrangement of grains in a polycrystal. Aboav's empirical law (also called the Aboav-Weaire law) states that the neighbor of an  $n$ -sided cell has itself on average  $m_n = a_0 + a_1 n^{-1}$  neighbors, where  $a_0$  and  $a_1$  are constants. This law has been widely applied to most of the experimental and theoretical systems discussed in the introduction. The law is nevertheless violated, although only slightly, by numerical results on Voronoi diagrams [25, 3]. On the basis of the analysis of the present work we will in a future study [30] find the true expression for  $m_n$  when  $n$  is asymptotically large.

## Acknowledgments

The author thanks Pierre Calka for a discussion and for pointing out some important literature on Voronoi tessellations. He thanks Bernard Jancovici for discussions on Coulomb systems. This work has benefitted from a six month sabbatical period (CRCT) awarded to the author by the French Ministry of Education. The Laboratoire de Physique Théorique of the Université de Paris-Sud is associated with the Centre National de la Recherche Scientifique as research unit UMR 8627.

## References

- [1] G. Voronoi, *J. Reine Angew. Math.* **134**, 198 (1908).
- [2] F.T. Lewis, *Anatomical Records* **38**, 341 (1928); **47**, 59 (1930); **50**, 235 (1931).
- [3] A. Okabe, B. Boots, K. Sugihara, and S. N. Chiu, *Spatial tessellations: concepts and applications of Voronoi diagrams*, second edition (John Wiley & Sons Ltd., Cichester, 2000).
- [4] P. Cerisier, S. Rahal, and N. Rivier, *Phys. Rev. E* **54**, 5086 (1996).
- [5] F. Elias, C. Flament, J.-C. Bacri, O. Cardoso, and F. Graner, *Phys. Rev. E* **56**, 3310 (1997).
- [6] C. Sire et M. Seul, *J. Phys. I France* **5**, 97 (1995).
- [7] P. Moriarty, M.D.R. Taylor, and M. Brust *Phys. Rev. Lett.* **89**, 248303 (2002).

- [8] J.C. Earnshaw and D.J. Robinson, *Phys. Rev. Lett.* **72**, 3682 (1994).
- [9] J.C. Earnshaw and D.J. Robinson, *Physica A* **214**, 23 (1995).
- [10] J.A. Glazier, S.P. Gross, and J. Stavans *Phys. Rev. A* **36**, 306 (1986).
- [11] J. Stavans and J.A. Glazier, *Phys. Rev. Lett.* **62**, 1318 (1989).
- [12] K. Zahn, R. Lenke, and G. Maret, *Phys. Rev. Lett.* **82**, 2721 (1999).
- [13] R.A. Quinn and J. Goree, *Phys. Rev. E* **64**, 051404 (2001).
- [14] P. Richard, L. Oger, J.-P. Troadec, and A. Gervois, *Phys. Rev. E* **60**, 4551 (1999).
- [15] M. Marder, *Phys. Rev. A* **36**, 438 (1987).
- [16] J. Stavans, E. Domany, and D. Mukamel, *Europhys. Lett.* **15**, 479 (1991).
- [17] H. Flyvbjerg, *Phys. Rev. E* **47**, 4037 (1993).
- [18] B. Jeune and D. Barabé, *Annals of Botany* **82**, 577 (1998).
- [19] N.H. Christ, R. Friedberg, and T.D. Lee, *Nuclear Physics* **B202**, 89 (1982); **B210**, 310; 337 (1982).
- [20] C. Itzykson and J.M. Drouffe, *Statistical field theory* (Cambridge University Press, Cambridge, 1989), Vol. 2, chapter 11.
- [21] C. Godrèche, I. Kostov, and I. Yekutieli, *Phys. Rev. Lett.* **69**, 2674 (1992).
- [22] J. L. Meijering, *Philips Research Reports* **8**, 270 (1953).
- [23] A. Hayen and M.P. Quine, *Adv. in Appl. Probab.* **32**, 67 (2000).
- [24] M. P. Quine and D. F. Watson, *J. Appl. Probab.* **21**, 548 (1984).
- [25] G. Le Caër and J. S. Ho, *J. Phys. A* **23**, 3297 (1990).
- [26] S. Kumar and S. K. Kurtz, *Materials Characterization* **31**, 55 (1993).
- [27] J.M. Drouffe and C. Itzykson, *Nuclear Physics* **B235** [FS11], 45 (1984).
- [28] K. B. Lauritsen, C. Moukarzel, and H. J. Herrmann, *J. Phys. I France* **3**, 1941 (1993).
- [29] D.A. Aboav, *Metallography* **3**, 383 (1970).
- [30] H.J. Hilhorst, in preparation.

- [31] H.J. Hilhorst, *J. Stat. Mech.* L02003 (2005).
- [32] H.J. Hilhorst, in preparation.
- [33] P. Calka, *Adv. in Appl. Probab.* **35**, 551 (2003).
- [34] P. Calka, *Adv. in Appl. Probab.* **35**, 863 (2003).
- [35] D. Hug, M. Reitzner, and R. Schneider, *Adv. in Appl. Probab.* **36**, 667 (2004).
- [36] P. Calka, *Adv. in Appl. Probab.* **34**, 702 (2002).
- [37] P. Calka and T. Schreiber, *to appear in Ann. Probab.*
- [38] N. Rivier and A. Lissowski, *J. Phys. A* **15**, L143 (1982).
- [39] N. Rivier, *Phil. Mag. B* **52**, 795 (1985).
- [40] This region, when enlarged by a factor of two in both directions, is sometimes referred to as the ‘fundamental domain’ of the Voronoi cell. It is then the region that should be free of point particles.
- [41] H.J. Hilhorst, unpublished.
- [42] The definition of  $T_m$  in reference [31] contains an extra factor  $\cos \gamma_m / \cos \beta_m$ . Only  $\Pi_m T_m$  intervenes in the calculation and the condition  $G = 0$  guarantees that the product of the extra factors is unity.
- [43] In the special case of a regular  $n$ -sided polygon,  $R_{av}$  is the radius of its inscribed circle and  $\mathcal{A} - \pi R_{av}^2$  tends to zero as  $\sim n^{-2}$  when  $n \rightarrow \infty$ . However, in the general case we cannot expect a similar decay with  $n$ .
- [44] The strong implications of this fact will be pursued further in reference [32].
- [45] In reference [31] our method was presented more compactly with an integral only on the single variable  $s$ .
- [46] The term  $\mathbb{V}_1$  was called  $\mathbb{V}_0$  in reference [31].
- [47] Provided that  $\delta\eta_m$  is sufficiently weakly correlated with the expressions that multiply it:  $\gamma_m^2 + \beta_{m+1}^2$ ,  $\tau_m$ , and  $\tau_m^2$ , respectively, in the discussion of section 6.3.2. The weakness of these correlations may be shown within the present theory; however, we will not elaborate on this point.
- [48] In reference [31] at the end of section 3 it should say ‘inverse temperature’ instead of ‘temperature.’

- [49] It is easy to verify that, up to corrections that vanish as  $\sim n^{-\frac{1}{2}}$ , the angle  $\phi$  is equal to the polar angles  $\Phi(\phi) \equiv \Phi_m$  and  $\Psi(\phi) \equiv \Psi_m$ .
- [50] I.S. Gradshteyn and I.M. Ryzhik, *Tables of Integrals, Series, and Products*, 4th ed., Academic Press, New York (1965)
- [51] D. Segel, D. Mukamel, O. Krichevsky, and J. Stavans, *Phys. Rev. E* **47**, 812 (1993).
- [52] M.A. Fortes, *J. Phys. A* **28**, 1055 (1995).
- [53] M.A. Fortes and P.I.C. Teixeira, *Eur. Phys. J. E* **6**, 133 (2001).
- [54] R.E. Miles and R.J. Maillardet, *J. Appl. Probab.* **19A**, 97 (1982).
- [55] S.F. Edwards and K.D. Pithia, *Physica A* **205**, 577 (1994).
- [56] S. N. Chiu, *J. Phys. A* **28**, 607 (1995).
- [57] S. N. Chiu, *Materials Characterization* **34**, 149 (1995).
- [58] S. Goudsmit, *Rev. Mod. Phys.* **17** (1945) 321.