

SPIN GLASSES

Francesco Guerra*

*Dipartimento di Fisica, Università di Roma “La Sapienza”
INFN, Sezione di Roma1, Piazzale A. Moro 2, 00185 Roma, Italy*

May 1, 2019

1 Introduction

From a physical point of view, spin glasses, as dilute magnetic alloys, are very interesting systems. They are characterized by such features as exhibiting a new magnetic phase, where magnetic moments are frozen into disordered equilibrium orientations, without any long-range order. See for example [1] for general reviews, and also [2] for a very readable account about the physical properties of spin glasses. The experimental laboratory study of spin glasses is a very difficult subject, because of their peculiar properties. In particular the existence of very slowly relaxing modes, with consequent memory effects, makes difficult to realize the very basic physical concept of a system at thermodynamical equilibrium, at a given temperature.

From a theoretical point of view some models have been proposed, which try to capture the essential physical features of spin glasses, in the frame of very simple assumptions.

The basic model has been proposed by Edwards and Anderson [3] many years ago. It is a simple extension of the well known nearest neighbour Ising model. On a large region Λ of the unit lattice in d dimensions, we associate an Ising spin $\sigma(n)$ to each lattice site n , and then we introduce a lattice Hamiltonian

$$H_{\Lambda}(\sigma, J) = - \sum_{(n,n')} J(n, n') \sigma(n) \sigma(n'). \quad (1)$$

Here, the sum runs over all couples of nearest neighbour sites in Λ , and J are quenched random couplings, assumed for simplicity to be independent

* e-mail: francesco.guerra@roma1.infn.it

identically distributed random variables, with centered unit Gaussian distribution. The quenched character of the J means that they do not participate to thermodynamic equilibrium, but act as a kind of random external noise on the coupling of the σ variables. In the expression of the Hamiltonian, we have indicated with σ the set of all $\sigma(n)$, and with J the set of all $J(n, n')$. The region Λ must be taken very large, by letting it invade all lattice in the limit. The physical motivation for this choice is that for real spin glasses the interaction between the spins dissolved in the matrix of the alloy oscillates in sign according to distance. This effect is taken into account in the model through the random character of the couplings between spins.

Even though very drastic simplifications have been introduced in the formulation of this model, as compared to the extremely more complicated nature of physical spin glasses, nevertheless a rigorous study of all properties emerging from the static and dynamic behavior of a thermodynamic system of this kind is far from being complete. In particular, with reference to static equilibrium properties, it is not possible yet to reach a completely substantiated description of the phases emerging in the low temperature region. Even physical intuition gives completely different guesses for different people.

In the same way as a mean field version can be associated to the ordinary Ising model, so it is possible for the disordered model described by (1). Now we consider a number of sites $i = 1, 2, \dots, N$, and let each spin $\sigma(i)$ at site i interact with all other spins, with the intervention of a quenched noise J_{ij} . The precise form of the Hamiltonian will be given in the following.

This is the mean field model for spin glasses, introduced by David Sherrington and Scott Kirkpatrick more than thirty years ago [4], [5]. It is a celebrated model. Hundreds and hundreds of articles have been devoted to its study during the years, appearing in the theoretical physics literature.

The relevance of the model stems surely from the fact that it is intended to represent some important features of the physical spin glass systems, of great interest for their peculiar properties, at least at the level of the mean field approximation.

But another important source of interest is connected with the fact that disordered systems, of the Sherrington-Kirkpatrick type, and their generalizations, seems to play a very important role for theoretical and practical assessments about hard optimization problems, as it is shown for example by Mark Mézard, Giorgio Parisi and Riccardo Zecchina in [6].

It is interesting to remark that the original paper was entitled “Solvable Model of a Spin-Glass”, while a previous draft, as told by David Sherrington, contained even the stronger denomination “Exactly Solvable”. However, it turned out that the very natural solution devised by the authors is valid only at high temperatures, or for large external magnetic fields. While, at low

temperatures, the proposed solution exhibits a nonphysical drawback given by a negative entropy, as properly recognized by the authors in their very first paper.

It took some years to find an acceptable solution. This was done by Giorgio Parisi in a series of papers, by marking a radical departure from the previous methods. In fact, a very deep method of “spontaneous replica symmetry breaking” was developed. As a consequence the physical content of the theory was encoded in a functional order parameter of new type, and a remarkable structure emerged for the pure states of the theory, a kind of hierarchical, ultrametric organization. These very interesting developments, due to Giorgio Parisi, and his coworkers, are explained in a brilliant way in the classical book [7]. Part of this structure will be recalled in the following.

It is important to remark that Parisi solution is presented in the form of an ingenious and clever *Ansatz*. Until few years ago it was not known whether this *Ansatz* would give the true solution for the model, in the so called thermodynamic limit, when the size of the system becomes infinite, or it would be only a very good approximation for the true solution.

The general structures offered by the Parisi solution, and their possible generalizations for similar models, exhibit an extremely rich and interesting mathematical content. Very appropriately, Michel Talagrand has inserted a strongly suggestive sentence in the title to his recent book [8]: “Spin glasses: a challenge for mathematicians”.

As a matter of fact, how to face this challenge is a very difficult problem. Here we would like to recall the main features of a very powerful method, yet extremely simple in its very essence, based on a comparison and interpolation argument on sets of Gaussian random variables.

The method found its first simple application in [9], where it was shown that the Sherrington-Kirkpatrick replica symmetric approximate solution was a rigorous lower bound for the quenched free energy of the system, uniformly in the size. Then, it was possible to reach a long waited result [10]: the convergence of the free energy density in the thermodynamic limit, by an intermediate step where the quenched free energy was shown to be subadditive in the size of the system.

Moreover, still by interpolation on families of Gaussian random variables, the first mentioned result was extended to give a rigorous proof that the expression given by the Parisi *Ansatz* is also a lower bound for the quenched free energy of the system, uniformly in the size [11]. The method gives not only the bound, but also the explicit form of the correction in a quite involved form. As a recent and very important result, along the task of facing the challenge, Michel Talagrand has been able to dominate these correction terms, showing that they vanish in the thermodynamic limit. This milestone

achievement was firstly announced in a short note [12], containing only a synthetic sketch of the proof, and then presented with all details in a long paper to be published on *Annals of Mathematics* [13].

The interpolation method is also at the basis of the far reaching generalized variational principle proven by Michel Aizenman, Robert Sims and Shannon Starr in [14].

In our presentation, we will try to be as self-contained as possible. We will give all definitions, explain the basic structure of the interpolation method, and show how some of the results are obtained. We will concentrate mostly on questions connected with the free energy, its properties of subadditivity, the existence of the infinite volume limit, and the replica bounds.

For the sake of comparison, and in order to provide a kind of warm up, we will recall also some features of the standard elementary mean field model of ferromagnetism, the so called Curie-Weiss model. We will concentrate also here on the free energy, and systematically exploit elementary comparison and interpolation arguments. This will show the strict analogy between the treatment of the ferromagnetic model and the developments in the mean field spin glass case. Basic roles will be played in the two cases, but with different expressions, by positivity and convexity properties.

Then, we will consider the problem of connecting results for the mean field case to the short range case. An intermediate position is occupied by the so called diluted models. They can be studied through a generalization of the methods exploited in the mean field case, as shown for example in [15].

The organization of the paper is as follows. In Section 2, we introduce the ferromagnetic model and discuss behavior and properties of the free energy in the thermodynamic limit, by emphasizing, in this very elementary case, the comparison and interpolation methods that will be also exploited, in a different context, in the spin glass case.

Section 3 is devoted to the basic features of the mean field spin glass models, by introducing all necessary definitions.

In Section 4, we introduce, for generic Gaussian interactions, some important formulae, concerning the derivation with respect to the strength of the interaction, and the Gaussian comparison and interpolation method.

In next Section 5 we give simple applications to the mean field spin glass model, in particular to the existence of the infinite volume limit of the quenched free energy [10], and to the proof of general variational bounds, by following the useful strategy developed in [14].

Section 6 will briefly recall the main features of the Parisi representation, and will state the main theorem concerning the free energy.

In Section 7 we will make some mention about results for diluted models.

Finally, in Section 8, we attack the problem of connecting the results for

the mean field case to the more realistic short range models.

Section 9 will be devoted to conclusions and outlook for future foreseen developments.

Our treatment will be as simple as possible, by relying on the basic structural properties, and by describing methods of presumably very long lasting power. The emphasis given to the mean field case reflects the status of research. May be that after some years from now this review would be written according to completely different patterns.

2 A warm up. The mean field ferromagnetic model. Structure and results.

The mean field ferromagnetic model is among the simplest models of statistical mechanics. However, it contains very interesting features, in particular a phase transition, characterized by spontaneous magnetization, at low temperatures. We refer to standard textbooks, for example [16], for a full treatment, and a complete appreciation of the model in the frame of the theory of ferromagnetism. Here we consider firstly some properties of the free energy, easily obtained through comparison methods.

The generic configuration of the mean field ferromagnetic model is defined through Ising spin variables $\sigma_i = \pm 1$, attached to each site $i = 1, 2, \dots, N$.

The Hamiltonian of the model, in some external field of strength h , is given by the mean field expression

$$H_N(\sigma, h) = -\frac{1}{N} \sum_{(i,j)} \sigma_i \sigma_j - h \sum_i \sigma_i. \quad (2)$$

Here, the first sum extends to all $N(N-1)/2$ site couples, and the second to all sites.

For a given inverse temperature β , let us now introduce the partition function $Z_N(\beta, h)$ and the free energy per site $f_N(\beta, h)$, according to the well known definitions

$$Z_N(\beta, h) = \sum_{\sigma_1 \dots \sigma_N} \exp(-\beta H_N(\sigma, h)), \quad (3)$$

$$-\beta f_N(\beta, h) = N^{-1} E \log Z_N(\beta, h). \quad (4)$$

It is also convenient to define the average spin magnetization

$$m = \frac{1}{N} \sum_i \sigma_i. \quad (5)$$

Then, it is immediately seen that the Hamiltonian in (2) can be equivalently written as

$$H_N(\sigma, h) = -\frac{1}{2}Nm^2 - h \sum_i \sigma_i, \quad (6)$$

where an unessential constant term has been neglected. In fact we have

$$\sum_{(i,j)} \sigma_i \sigma_j = \frac{1}{2} \sum_{i,j;i \neq j} \sigma_i \sigma_j = \frac{1}{2}N^2m^2 - \frac{1}{2}N, \quad (7)$$

where the sum over all couples has been equivalently written as one half the sum over all i, j with $i \neq j$, and the diagonal terms with $i = j$ have been added and subtracted out. Notice that they give a constant because $\sigma_i^2 = 1$.

Therefore, the partition function in (3) can be equivalently substituted by the expression

$$Z_N(\beta, h) = \sum_{\sigma_1 \dots \sigma_N} \exp\left(\frac{1}{2}\beta Nm^2\right) \exp\left(\beta h \sum_i \sigma_i\right), \quad (8)$$

which will be our starting point.

Our interest will be in the $\lim_{N \rightarrow \infty} N^{-1} \log Z_N(\beta, h)$. To this purpose, let us establish the important subadditivity property, holding for the splitting of the big N site system in two smaller N_1 site and N_2 site systems, respectively, with $N = N_1 + N_2$,

$$\log Z_N(\beta, h) \leq \log Z_{N_1}(\beta, h) + \log Z_{N_2}(\beta, h). \quad (9)$$

The proof is very simple. Let us denote, in the most natural way, by $\sigma_1, \dots, \sigma_{N_1}$ the spin variables for the first subsystem, and by $\sigma_{N_1+1}, \dots, \sigma_N$ the N_2 spin variables of the second subsystem. Introduce also the subsystem magnetizations m_1 and m_2 , by adapting the definition (5) to the smaller systems, in such a way that

$$Nm = N_1m_1 + N_2m_2. \quad (10)$$

Therefore, we see that the large system magnetization m is the linear convex combination of the smaller system ones, according to the obvious

$$m = \frac{N_1}{N}m_1 + \frac{N_2}{N}m_2. \quad (11)$$

Since the mapping $m \rightarrow m^2$ is convex, we have also the general bound, holding for all values of the σ variables

$$m^2 \leq \frac{N_1}{N}m_1^2 + \frac{N_2}{N}m_2^2. \quad (12)$$

Then, it is enough to substitute the inequality in the definition (8) of $Z_N(\beta, h)$, and recognize that we achieve factorization with respect to the two subsystems, and therefore the inequality $Z_N \leq Z_{N_1} Z_{N_2}$. So we have established (9). From subadditivity, the existence of the limit follows by a simple argument, as explained for example in [17]. In fact, we have

$$\lim_{N \rightarrow \infty} N^{-1} \log Z_N(\beta, h) = \inf_N N^{-1} \log Z_N(\beta, h). \quad (13)$$

Now we will calculate explicitly this limit, by introducing an order parameter M , a trial function, and an appropriate variational scheme. In order to get a lower bound, we start from the elementary inequality $m^2 \geq 2mM - M^2$, holding for any value of m and M . By inserting the inequality in the definition (8) we arrive at a factorization of the sum over σ 's. The sum can be explicitly calculated, and we arrive immediately to the lower bound, uniform in the size of the system,

$$N^{-1} \log Z_N(\beta, h) \geq \log 2 + \log \cosh \beta(h + M) - \frac{1}{2} \beta M^2, \quad (14)$$

holding for any value of the trial order parameter M . Clearly it is convenient to take the supremum over M . Then we establish the optimal uniform lower bound

$$N^{-1} \log Z_N(\beta, h) \geq \sup_M (\log 2 + \log \cosh \beta(h + M) - \frac{1}{2} \beta M^2). \quad (15)$$

It is simple to realize that the supremum coincides with the limit as $N \rightarrow \infty$. To this purpose we follow the following simple procedure. Let us consider all possible values of the variable m . There are $N + 1$ of them, corresponding to any number K of possible spin flips, starting from a given σ configuration, $K = 0, 1, \dots, N$. Let us consider the trivial decomposition of the identity, holding for any m ,

$$1 = \sum_M \delta_{mM}, \quad (16)$$

where M in the sum runs over the $N + 1$ possible values of m , and δ is Kronecker delta, being equal to 1 if $M = m$, and zero otherwise. Let us now insert (16) in the definition (8) of the partition function inside the sum over σ 's, and invert the two sums. Because of the forcing $m = M$ given by the δ , we can write $m^2 = 2mM - M^2$ inside the sum. Then if we neglect the δ , by using the trivial $\delta \leq 1$, we have an upper bound, where the sum over σ 's can be explicitly performed as before. Then it is enough to take the upper

bound with respect to M , and consider that there are $N + 1$ terms in the now trivial sum over M , in order to arrive at the upper bound

$$N^{-1} \log Z_N(\beta, h) \leq \sup_M (\log 2 + \log \cosh \beta(h + M) - \frac{1}{2}\beta M^2) + N^{-1} \log(N + 1). \quad (17)$$

Therefore, by going to the limit as $N \rightarrow \infty$, we can collect all our results in the form of the following theorem giving the full characterization of the thermodynamic limit of the free energy.

Theorem 1. *For the mean field ferromagnetic model we have*

$$\lim_{N \rightarrow \infty} N^{-1} \log Z_N(\beta, h) = \inf_N N^{-1} \log Z_N(\beta, h) \quad (18)$$

$$= \sup_M (\log 2 + \log \cosh \beta(h + M) - \frac{1}{2}\beta M^2). \quad (19)$$

This ends our discussion about the free energy in the ferromagnetic model.

Other properties of the model can be easily established. Introduce the Boltzmann-Gibbs state

$$\omega_N(A) = Z_N^{-1} \sum_{\sigma_1 \dots \sigma_N} A \exp(\frac{1}{2}\beta N m^2) \exp(\beta h \sum_i \sigma_i), \quad (20)$$

where A is any function of $\sigma_1 \dots \sigma_N$.

The observable $m(\sigma)$ becomes self-averaging under ω_N , in the infinite volume limit, in the sense that

$$\lim_{N \rightarrow \infty} \omega_N((m - M(\beta, h))^2) = 0. \quad (21)$$

This property of m is the deep reason for the success of the strategy exploited before for the convergence of the free energy. Easy consequences are the following. In the infinite volume limit, the Boltzmann-Gibbs state becomes a factor state

$$\lim_{N \rightarrow \infty} \omega_N(\sigma_1 \dots \sigma_s) = M(\beta, h)^s. \quad (22)$$

A phase transition appears in the form of spontaneous magnetization. In fact, while for $h = 0$ and $\beta \leq 1$ we have $M(\beta, h) = 0$, on the other hand, for $\beta > 1$, we have the discontinuity

$$\lim_{h \rightarrow 0^+} M(\beta, h) = - \lim_{h \rightarrow 0^-} M(\beta, h) \equiv M(\beta) > 0. \quad (23)$$

Fluctuations can also be easily controlled. In fact, one proves that the rescaled random variable $\sqrt{N}(m - M(\beta, h))$ tends in distribution, under ω_N , to a centered Gaussian with variance given by the susceptibility

$$\chi(\beta, h) \equiv \frac{\partial}{\partial h} M(\beta, h) \equiv \frac{\beta(1 - M^2)}{1 - \beta(1 - M^2)}. \quad (24)$$

Notice that the variance becomes infinite only at the critical point $h = 0$, $\beta = 1$, where $M = 0$.

Now we are ready to attack the much more difficult spin glass model. But it will be surprising to see that, by following a simple extension of the methods here described, we will arrive to similar results.

3 The basic definitions for the mean field spin glass model

As in the ferromagnetic case, the generic configuration of the mean field spin glass model is defined through Ising spin variables $\sigma_i = \pm 1$, attached to each site $i = 1, 2, \dots, N$.

But now there is an external quenched disorder given by the $N(N-1)/2$ independent and identical distributed random variables J_{ij} , defined for each couple of sites. For the sake of simplicity, we assume each J_{ij} to be a centered unit Gaussian with averages $E(J_{ij}) = 0$, $E(J_{ij}^2) = 1$. By quenched disorder we mean that the J have a kind of stochastic external influence on the system, without participating to the thermal equilibrium.

Now the Hamiltonian of the model, in some external field of strength h , is given by the mean field expression

$$H_N(\sigma, h, J) = -\frac{1}{\sqrt{N}} \sum_{(i,j)} J_{ij} \sigma_i \sigma_j - h \sum_i \sigma_i. \quad (25)$$

Here, the first sum extends to all site couples, and the second to all sites. Notice the \sqrt{N} , necessary to ensure a good thermodynamic behavior to the free energy.

For a given inverse temperature β , let us now introduce the disorder dependent partition function $Z_N(\beta, h, J)$ and the quenched average of the free energy per site $f_N(\beta, h)$, according to the definitions

$$Z_N(\beta, h, J) = \sum_{\sigma_1 \dots \sigma_N} \exp(-\beta H_N(\sigma, h, J)), \quad (26)$$

$$-\beta f_N(\beta, h) = N^{-1} E \log Z_N(\beta, h, J). \quad (27)$$

Notice that in (27) the average E with respect to the external noise is made *after* the log is taken. This procedure is called quenched averaging. It represents the physical idea that the external noise does not participate to the thermal equilibrium. Only the σ 's are thermalized.

For the sake of simplicity, it is also convenient to write the partition function in the following equivalent form. First of all let us introduce a family

of centered Gaussian random variables $\mathcal{K}(\sigma)$, indexed by the configurations σ , and characterized by the covariances

$$E(\mathcal{K}(\sigma)\mathcal{K}(\sigma')) = q^2(\sigma, \sigma'), \quad (28)$$

where $q(\sigma, \sigma')$ are the overlaps between two generic configurations, defined by

$$q(\sigma, \sigma') = N^{-1} \sum_i \sigma_i \sigma'_i, \quad (29)$$

with the obvious bounds $-1 \leq q(\sigma, \sigma') \leq 1$, and the normalization $q(\sigma, \sigma) = 1$. Then, starting from the definition (25), it is immediately seen that the partition function in (26) can be also written, by neglecting unessential constant terms, in the form

$$Z_N(\beta, h, J) = \sum_{\sigma_1 \dots \sigma_N} \exp(\beta \sqrt{\frac{N}{2}} \mathcal{K}(\sigma)) \exp(\beta h \sum_i \sigma_i), \quad (30)$$

which will be the starting point of our treatment.

4 Basic formulae of derivation and interpolation

We work in the following general setting. Let U_i be a family of centered Gaussian random variables, $i = 1, \dots, K$, with covariance matrix given by $E(U_i U_j) \equiv S_{ij}$. We treat the index i now as configuration space for some statistical mechanics system, with partition function Z and quenched free energy given by

$$E \log \sum_i w_i \exp(\sqrt{t} U_i) \equiv E \log Z, \quad (31)$$

where $w_i \geq 0$ are generic weights, and t is a parameter ruling the strength of the interaction.

It would be hard to underestimate the relevance of the following derivation formula

$$\begin{aligned} \frac{d}{dt} E \log \sum_i w_i \exp(\sqrt{t} U_i) &= \\ \frac{1}{2} E(Z^{-1} \sum_i w_i \exp(\sqrt{t} U_i) S_{ii}) & \\ - \frac{1}{2} E(Z^{-2} \sum_i \sum_j w_i w_j \exp(\sqrt{t} U_i) \exp(\sqrt{t} U_j) S_{ij}). & \quad (32) \end{aligned}$$

The proof is straightforward. Firstly we perform directly the t derivative. Then, we notice that the random variables appear in expressions of the form $E(U_i F)$, where F are functions of the U 's. These can be easily handled through the following integration by parts formula for generic Gaussian random variables, strongly reminiscent of the Wick theorem in quantum field theory,

$$E(U_i F) = \sum_j S_{ij} E\left(\frac{\partial}{\partial U_j} F\right). \quad (33)$$

Therefore, we see that always two derivatives are involved. The two terms in (32) come from the action of the U_j derivatives, the first acting on the Boltzmann factor, and giving rise to a Kronecker δ_{ij} , the second acting on Z^{-1} , and giving rise to the minus sign and the duplication of variables.

The derivation formula can be expressed in a more compact form by introducing replicas and suitable averages. In fact, let us introduce the state ω acting on functions F of i as follows

$$\omega(F(i)) = Z^{-1} \sum_i w_i \exp(\sqrt{t} U_i) F(i), \quad (34)$$

together with the associated product state Ω acting on replicated configuration spaces i_1, i_2, \dots, i_s . By performing also a global E average, finally we define the averages

$$\langle F \rangle_t \equiv E \Omega(F), \quad (35)$$

where the subscript is introduced in order to recall the t dependence of these averages.

Then, the equation (32) can be written in a more compact form

$$\frac{d}{dt} E \log \sum_i w_i \exp(\sqrt{t} U_i) = \frac{1}{2} \langle S_{i_1 i_1} \rangle - \frac{1}{2} \langle S_{i_1 i_2} \rangle. \quad (36)$$

Our basic comparison argument will be based on the following very simple theorem.

Theorem 2. *Let U_i and \hat{U}_i , for $i = 1, \dots, K$, be independent families of centered Gaussian random variables, whose covariances satisfy the inequalities for generic configurations*

$$E(U_i U_j) \equiv S_{ij} \geq E(\hat{U}_i \hat{U}_j) \equiv \hat{S}_{ij}, \quad (37)$$

and the equalities along the diagonal

$$E(U_i U_i) \equiv S_{ii} = E(\hat{U}_i \hat{U}_i) \equiv \hat{S}_{ii}, \quad (38)$$

then for the quenched averages we have the inequality in the opposite sense

$$E \log \sum_i w_i \exp(U_i) \leq E \log \sum_i w_i \exp(\hat{U}_i), \quad (39)$$

where the $w_i \geq 0$ are the same in the two expressions.

Considerations of this kind are present in the mathematical literature of some years ago. Two typical references are [18] and [19].

The proof is extremely simple and amounts to a straightforward calculation. In fact, let us consider the interpolating expression

$$E \log \sum_i w_i \exp(\sqrt{t}U_i + \sqrt{1-t}\hat{U}_i), \quad (40)$$

where $0 \leq t \leq 1$. Clearly the two expressions under comparison correspond to the values $t = 0$ and $t = 1$ respectively. By taking the derivative with respect to t , with the help of the previous derivation formula, we arrive to the evaluation of the t derivative in the form

$$\begin{aligned} \frac{d}{dt} E \log \sum_i w_i \exp(\sqrt{t}U_i + \sqrt{1-t}\hat{U}_i) = \\ \frac{1}{2} E(Z^{-1} \sum_i w_i \exp(\sqrt{t}U_i)(S_{ii} - \hat{S}_{ii}) \\ - \frac{1}{2} E(Z^{-2} \sum_i \sum_j w_i w_j \exp(\sqrt{t}U_i) \exp(\sqrt{t}U_j)(S_{ij} - \hat{S}_{ij}). \end{aligned} \quad (41)$$

From the conditions assumed for the covariances, we immediately see that the interpolating function is nonincreasing in t , and the theorem follows.

The derivation formula and the comparison Theorem are not restricted to the Gaussian case. Generalizations in many directions are possible. For the diluted spin glass models and optimization problems we refer for example to [20], and to [15], and references quoted there.

5 The thermodynamic limit and the variational bounds

We give here some striking applications of the basic comparison Theorem. In [10] we have given a very simple proof of a long waited result, about the convergence of the free energy per site in the thermodynamic limit. Let us show the argument. Let us consider a system of size N and two smaller

systems of sizes N_1 and N_2 respectively, with $N = N_1 + N_2$, as before in the ferromagnetic case. Let us now compare

$$E \log Z_N(\beta, h, J) = E \log \sum_{\sigma_1 \dots \sigma_N} \exp(\beta \sqrt{\frac{N}{2}} \mathcal{K}(\sigma)) \exp(\beta h \sum_i \sigma_i), \quad (42)$$

with

$$\begin{aligned} & E \log \sum_{\sigma_1 \dots \sigma_N} \exp(\beta \sqrt{\frac{N_1}{2}} \mathcal{K}^{(1)}(\sigma^{(1)})) \exp(\beta \sqrt{\frac{N_2}{2}} \mathcal{K}^{(2)}(\sigma^{(2)})) \exp(\beta h \sum_i \sigma_i) \\ & \equiv E \log Z_{N_1}(\beta, h, J) + E \log Z_{N_2}(\beta, h, J), \end{aligned} \quad (43)$$

where $\sigma^{(1)}$ are the $(\sigma_i, i = 1, \dots, N_1)$, and $\sigma^{(2)}$ are the $(\sigma_i, i = N_1 + 1, \dots, N)$. Covariances for $\mathcal{K}^{(1)}$ and $\mathcal{K}^{(2)}$ are expressed as in (28), but now the overlaps are substituted with the partial overlaps of the first and second block, q_1 and q_2 respectively. It is very simple to apply the comparison theorem. All one has to do is to observe that the obvious

$$Nq = N_1q_1 + N_2q_2, \quad (44)$$

analogous to (10), implies, as in (12),

$$q^2 \leq \frac{N_1}{N} q_1^2 + \frac{N_2}{N} q_2^2. \quad (45)$$

Therefore, the comparison gives the superadditivity property, to be compared with (9),

$$E \log Z_N(\beta, h, J) \geq E \log Z_{N_1}(\beta, h, J) + E \log Z_{N_2}(\beta, h, J). \quad (46)$$

From the superadditivity property the existence of the limit follows in the form

$$\lim_{N \rightarrow \infty} N^{-1} E \log Z_N(\beta, h, J) = \sup_N N^{-1} E \log Z_N(\beta, h, J), \quad (47)$$

to be compared with (13).

The second application is in the form of the Aizenman-Sims-Starr generalized variational principle. Here, we will need to introduce some auxiliary system. The denumerable configuration space is given by the values of $\alpha = 1, 2, \dots$. We introduce also weights $w_\alpha \geq 0$ for the α system, and suitably defined overlaps between two generic configurations $p(\alpha, \alpha')$, with $p(\alpha, \alpha) = 1$.

A family of centered Gaussian random variables $\hat{\mathcal{K}}(\alpha)$, now indexed by the configurations α , will be defined by the covariances

$$E(\hat{\mathcal{K}}(\alpha)\hat{\mathcal{K}}(\alpha')) = p^2(\alpha, \alpha'). \quad (48)$$

We will need also a family of centered Gaussian random variables $\eta_i(\alpha)$, indexed by the sites i of our original system and the configurations α of the auxiliary system, so that

$$E(\eta_i(\alpha)\eta_{i'}(\alpha')) = \delta_{ii'}p(\alpha, \alpha'). \quad (49)$$

Both the probability measure w_α , and the overlaps $p(\alpha, \alpha')$ could depend on some additional external quenched noise, that does not appear explicitly in our notation.

In the following, we will denote by E averages with respect to all random variables involved.

In order to start the comparison argument, we will consider firstly the case where the two σ and α systems are not coupled, so to appear factorized in the form

$$\begin{aligned} & E \log \sum_{\sigma_1 \dots \sigma_N} \sum_{\alpha} w_\alpha \exp(\beta \sqrt{\frac{N}{2}} \mathcal{K}(\sigma)) \exp(\beta \sqrt{\frac{N}{2}} \hat{\mathcal{K}}(\alpha)) \exp(\beta h \sum_i \sigma_i) \\ & \equiv E \log Z_N(\beta, h, J) + E \log \sum_{\alpha} w_\alpha \exp(\beta \sqrt{\frac{N}{2}} \hat{\mathcal{K}}(\alpha)). \end{aligned} \quad (50)$$

In the second case the \mathcal{K} fields are suppressed and the coupling between the two systems will be taken in a very simple form, by allowing the η field to act as an external field on the σ system. In this way the σ 's appear as factorized, and the sums can be explicitly performed. The chosen form for the second term in the comparison is

$$\begin{aligned} & E \log \sum_{\sigma_1 \dots \sigma_N} \sum_{\alpha} w_\alpha \exp(\beta \sum_i \eta_i(\alpha) \sigma_i) \exp(\beta h \sum_i \sigma_i) \\ & \equiv N \log 2 + E \log \sum_{\alpha} w_\alpha (c_1 c_2 \dots c_N), \end{aligned} \quad (51)$$

where we have defined

$$c_i = \cosh \beta(h + \eta_i(\alpha)), \quad (52)$$

as arising from the sums over σ 's.

Now we apply the comparison Theorem. In the first case, the covariances involve the sums of squares of overlaps

$$\frac{1}{2}(q^2(\sigma, \sigma') + p^2(\alpha, \alpha')). \quad (53)$$

In the second case, a very simple calculation shows that the covariances involve the overlap products

$$q(\sigma, \sigma')p(\alpha, \alpha'). \quad (54)$$

Therefore, the comparison is very easy and, by collecting all expressions, we end up with the useful estimate, as in [14], holding for any auxiliary system as defined before,

$$N^{-1}E \log Z_N(\beta, h, J) \leq \quad (55)$$

$$\log 2 + N^{-1}(E \log \sum_{\alpha} w_{\alpha}(c_1 c_2 \dots c_N) - E \log \sum_{\alpha} w_{\alpha} \exp(\beta \sqrt{\frac{N}{2}} \hat{\mathcal{K}}(\alpha))).$$

6 The Parisi representation for the free energy

We refer to the original paper [21], and to the extensive review given in [7], for the general motivations, and the derivation of the broken replica *Ansatz*, in the frame of the ingenious replica trick. Here we limit ourselves to a synthetic description of its general structure, independently from the replica trick

First of all, let us introduce the convex space \mathcal{X} of the functional order parameters x , as nondecreasing functions of the auxiliary variable q , both x and q taking values on the interval $[0, 1]$, *i.e.*

$$\mathcal{X} \ni x : [0, 1] \ni q \rightarrow x(q) \in [0, 1]. \quad (56)$$

Notice that we call x the function, and $x(q)$ its values. We introduce a metric on \mathcal{X} through the $L^1([0, 1], dq)$ norm, where dq is the Lebesgue measure.

For our purposes, we will consider the case of piecewise constant functional order parameters, characterized by an integer K , and two sequences $q_0, q_1, \dots, q_K, m_1, m_2, \dots, m_K$ of numbers satisfying

$$0 = q_0 \leq q_1 \leq \dots \leq q_{K-1} \leq q_K = 1, \quad 0 \leq m_1 \leq m_2 \leq \dots \leq m_K \leq 1, \quad (57)$$

such that

$$x(q) = m_1 \text{ for } 0 = q_0 \leq q < q_1, \quad x(q) = m_2 \text{ for } q_1 \leq q < q_2, \\ \dots, x(q) = m_K \text{ for } q_{K-1} \leq q \leq q_K. \quad (58)$$

In the following, we will find convenient to define also $m_0 \equiv 0$, and $m_{K+1} \equiv 1$. The replica symmetric case of Sherrington and Kirkpatrick corresponds to

$$K = 2, \quad q_1 = \bar{q}, \quad m_1 = 0, \quad m_2 = 1. \quad (59)$$

Let us now introduce the function f , with values $f(q, y; x, \beta)$, of the variables $q \in [0, 1]$, $y \in R$, depending also on the functional order parameter x , and on the inverse temperature β , defined as the solution of the nonlinear antiparabolic equation

$$(\partial_q f)(q, y) + \frac{1}{2}(\partial_y^2 f)(q, y) + \frac{1}{2}x(q)(\partial_y f)^2(q, y) = 0, \quad (60)$$

with final condition

$$f(1, y) = \log \cosh(\beta y). \quad (61)$$

Here, we have stressed only the dependence of f on q and y .

It is very simple to integrate Eq. (60) when x is piecewise constant. In fact, consider $x(q) = m_a$, for $q_{a-1} \leq q \leq q_a$, firstly with $m_a > 0$. Then, it is immediately seen that the correct solution of Eq. (60) in this interval, with the right final boundary condition at $q = q_a$, is given by

$$f(q, y) = \frac{1}{m_a} \log \int \exp(m_a f(q_a, y + z\sqrt{q_a - q})) d\mu(z), \quad (62)$$

where $d\mu(z)$ is the centered unit Gaussian measure on the real line. On the other hand, if $m_a = 0$, then (60) loses the nonlinear part and the solution is given by

$$f(q, y) = \int f(q_a, y + z\sqrt{q_a - q}) d\mu(z), \quad (63)$$

which can be seen also as deriving from (62) in the limit $m_a \rightarrow 0$. Starting from the last interval K , and using (62) iteratively on each interval, we easily get the solution of (60), (61), in the case of piecewise order parameter x , as in (58), through a chain of interconnected Gaussian integrations.

Now we introduce the following important definitions. The trial auxiliary function, associated to a given mean field spin glass system, as described in Section 3, depending on the functional order parameter x , is defined as

$$\log 2 + f(0, h; x, \beta) - \frac{\beta^2}{2} \int_0^1 q x(q) dq. \quad (64)$$

Notice that in this expression the function f appears evaluated at $q = 0$, and $y = h$, where h is the value of the external magnetic field. This trial expression should be considered as the analog of that appearing in (14) for the ferromagnetic case.

The Parisi spontaneously broken replica symmetry expression for the free energy is given by the definition

$$-\beta f_P(\beta, h) \equiv \inf_x (\log 2 + f(0, h; x, \beta) - \frac{\beta^2}{2} \int_0^1 q x(q) dq), \quad (65)$$

where the infimum is taken with respect to all functional order parameters x . Notice that the infimum appears here, as compared to the supremum in the ferromagnetic case.

In [11], by exploiting a kind of generalized comparison argument, involving a suitably defined interpolation function, we have established the following important result.

Theorem 3. *For all values of the inverse temperature β , and the external magnetic field h , and for any functional order parameter x , the following bound holds*

$$N^{-1}E \log Z_N(\beta, h, J) \leq \log 2 + f(0, h; x, \beta) - \frac{\beta^2}{2} \int_0^1 q x(q) dq,$$

uniformly in N . Consequently, we have also

$$N^{-1}E \log Z_N(\beta, h, J) \leq \inf_x (\log 2 + f(0, h; x, \beta) - \frac{\beta^2}{2} \int_0^1 q x(q) dq),$$

uniformly in N .

However, this result can be understood also in the frame of the generalized variational principle established by Aizenman-Sims-Starr and described before.

In fact, one can easily show that there exist an α systems such that

$$N^{-1}E \log \sum_{\alpha} w_{\alpha} c_1 c_2 \dots c_N \equiv f(0, h; x, \beta), \quad (66)$$

$$N^{-1}E \log \sum_{\alpha} w_{\alpha} \exp(\beta \sqrt{\frac{N}{2}} \hat{\mathcal{K}}(\alpha)) \equiv \frac{\beta^2}{2} \int_0^1 q x(q) dq, \quad (67)$$

uniformly in N . This result stems from previous work of Derrida, Ruelle, Neveu, Bolthausen, Sznitman, Aizenman, Contucci, Talagrand, Bovier, and others, and in a sense is implicit in the treatment given in [7]. It can be reached in a very simple way. Let us sketch the argument.

First of all, let us consider the Poisson point process $y_1 \geq y_2 \geq y_3 \dots$, uniquely characterized by the following conditions. For any interval A , introduce the occupation numbers $N(A)$, defined by

$$N(A) = \sum_{\alpha} \chi(y_{\alpha} \in A), \quad (68)$$

where $\chi() = 1$, if the random variable y_{α} belongs to the interval A , and $\chi() = 0$, otherwise. We assume that $N(A)$ and $N(B)$ are independent if the

intervals A and B are disjoint, and moreover that for each A , the random variable $N(A)$ has a Poisson distribution with parameter

$$\mu(A) = \int_a^b \exp(-y) dy, \quad (69)$$

if A is the interval (a, b) , *i.e.*

$$P(N(A) = k) = \exp(-\mu(A))\mu(A)^k/k!. \quad (70)$$

We will exploit $-y_\alpha$ as energy levels for a statistical mechanics systems with configurations indexed by α . For a parameter $0 < m < 1$, playing the role of inverse temperature, we can introduce the partition function

$$v = \sum_{\alpha} \exp\left(\frac{y_{\alpha}}{m}\right). \quad (71)$$

For m in the given interval it turns out that v is a very well defined random variable, with the sum over α extending to infinity. In fact, there is a strong inbuilt smooth cutoff in the very definition of the stochastic energy levels.

From the general properties of Poisson point processes it is very well known that the following basic invariance property holds. Introduce a random variable b , independent of y , subject to the condition $E(\exp b) = 1$, and let b_α be independent copies. Then, the randomly biased point process $y'_\alpha = y_\alpha + b_\alpha$, $\alpha = 1, 2, \dots$ is equivalent to the original one in distribution. An immediate consequence is the following. Let f be a random variable, independent of y , such that $E(\exp f) < \infty$, and let f_α be independent copies. Then the two random variables

$$\sum_{\alpha} \exp\left(\frac{y_{\alpha}}{m}\right) \exp(f_{\alpha}), \quad (72)$$

$$\sum_{\alpha} \exp\left(\frac{y_{\alpha}}{m}\right) E(\exp(mf))^{\frac{1}{m}} \quad (73)$$

have the same distribution. In particular they can be freely substituted under averages.

The auxiliary system which gives rise to the Parisi representation according to (66) (67), for a piecewise constant order parameter, is expressed in the following way. Now α will be a multi-index $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_K)$, where each α_a runs on $1, 2, 3, \dots$. Define the Poisson point process y_{α_1} , then, independently, for each value of α_1 processes $y_{\alpha_1\alpha_2}$, and so on up to $y_{\alpha_1\alpha_2\dots\alpha_K}$. Notice that in the cascade of independent processes $y_{\alpha_1}, y_{\alpha_1\alpha_2}, \dots, y_{\alpha_1\alpha_2\dots\alpha_K}$, the last

index refers to the numbering of the various points of the process, while the first indexes denotes independent copies labelled by the corresponding α 's.

The weights w_α have to be chosen according to the definition

$$w_\alpha = \exp \frac{y_{\alpha_1}}{m_1} \exp \frac{y_{\alpha_1 \alpha_2}}{m_2} \dots \exp \frac{y_{\alpha_1 \alpha_2 \dots \alpha_K}}{m_K}. \quad (74)$$

The cavity fields η and \mathcal{K} have the following expression in terms of independent unit Gaussian random variables $J_{\alpha_1}^i, J_{\alpha_1 \alpha_2}^i, \dots, J_{\alpha_1 \alpha_2 \dots \alpha_K}^i, J'_{\alpha_1}, J'_{\alpha_1 \alpha_2}, \dots, J'_{\alpha_1 \alpha_2 \dots \alpha_K}$,

$$\eta_i(\alpha) = \sqrt{q_1 - q_0} J_{\alpha_1}^i + \sqrt{q_2 - q_1} J_{\alpha_1 \alpha_2}^i + \dots + \sqrt{q_K - q_{K-1}} J_{\alpha_1 \alpha_2 \dots \alpha_K}^i, \quad (75)$$

$$\mathcal{K}(\alpha) = \sqrt{q_1^2 - q_0^2} J'_{\alpha_1} + \sqrt{q_2^2 - q_1^2} J'_{\alpha_1 \alpha_2} + \dots + \sqrt{q_K^2 - q_{K-1}^2} J'_{\alpha_1 \alpha_2 \dots \alpha_K}. \quad (76)$$

It is immediate to verify that $E(\eta_i(\alpha)\eta_{i'}(\alpha'))$ is zero if $i \neq i'$, while

$$\begin{aligned} E(\eta_i(\alpha)\eta_i(\alpha')) &= 0 \text{ if } \alpha_1 \neq \alpha'_1, \\ &= q_1 \text{ if } \alpha_1 = \alpha'_1, \alpha_2 \neq \alpha'_2, \\ &= q_2 \text{ if } \alpha_1 = \alpha'_1, \alpha_2 = \alpha'_2, \alpha_3 \neq \alpha'_3, \\ &\dots \\ &= 1 \text{ if } \alpha_1 = \alpha'_1, \alpha_2 = \alpha'_2, \dots, \alpha_K = \alpha'_K. \end{aligned} \quad (77)$$

Similarly, we have

$$\begin{aligned} E(\mathcal{K}(\alpha)\mathcal{K}(\alpha')) &= 0 \text{ if } \alpha_1 \neq \alpha'_1, \\ &= q_1^2 \text{ if } \alpha_1 = \alpha'_1, \alpha_2 \neq \alpha'_2, \\ &= q_2^2 \text{ if } \alpha_1 = \alpha'_1, \alpha_2 = \alpha'_2, \alpha_3 \neq \alpha'_3, \\ &\dots \\ &= 1 \text{ if } \alpha_1 = \alpha'_1, \alpha_2 = \alpha'_2, \dots, \alpha_K = \alpha'_K. \end{aligned} \quad (78)$$

This ends the definition of the α system, associated to a given piecewise constant order parameter.

Now, it is simple to verify that (66) (67) hold. Let us consider for example (66). With the α system chosen as before, the repeated application of the stochastic equivalence of (72) and (73) will give rise to a sequence of interchained Gaussian integrations exactly equivalent to those arising from the expression for f , as solution of the equation (60). For (73), there are equivalent considerations.

Therefore we see that the estimate in Theorem 3 is also a consequence of the generalized variational principle.

Up to this point we have seen how to obtain upper bounds. The problem arises whether, as in the ferromagnetic case, we can also get lower bounds,

so to shrink the thermodynamic limit to the value given by the \inf_x in Theorem 3. After a short announcement in [12], Michel Talagrand wrote an extended paper [13], to appear on Annals of Mathematics, where the complete proof of the control of the lower bound is firmly established. We refer to the original paper for the complete details of this remarkable achievement. About the methods, here we only recall that in [11] we have given also the corrections to the bounds appearing in Theorem 3, albeit in a quite complicated form. Talagrand has been able to establish that these corrections do in fact vanish in the thermodynamic limit.

In conclusion, we can establish the following extension of Theorem 1 to spin glasses.

Theorem 4. *For the mean field spin glass model we have*

$$\lim_{N \rightarrow \infty} N^{-1} E \log Z_N(\beta, h, J) = \sup_N N^{-1} E \log Z_N(\beta, h, J) \quad (79)$$

$$= \inf_x (\log 2 + f(0, h; x, \beta) - \frac{\beta^2}{2} \int_0^1 q x(q) dq). \quad (80)$$

7 Diluted models

Diluted models, in a sense, play a role intermediate between the mean field case and the short range case. In fact, while in the mean field model each site is interacting with all other sites, on the other hand, in the diluted model, each site is interacting with only a fixed number of other sites. However, while for the short range models there is a definition of distance among sites, relevant for the interaction, no such definition appears in the diluted models, where all sites are in any case equivalent. From this point of view, the diluted models are structurally similar to the mean field models, and most of the techniques and results explained before can be extended to them.

Let us define a typical diluted model. The quenched noise is described as follows. Let K be a Poisson random variable with parameter αN , where N is the number of sites, and α is a parameter entering the theory, together with the temperature. We consider also a sequence of independent centered random variables J_1, J_2, \dots , and a sequence of discrete independent random variables $i_1, j_1, i_2, j_2, \dots$, uniformly distributed over the set of sites $1, 2, \dots, N$. Then we assume as Hamiltonian

$$H_N(\sigma) = - \sum_{k=0}^K J_k \sigma_{i_k} \sigma_{j_k}. \quad (81)$$

Only the variables σ participate to thermodynamic equilibrium. All noise coming from K, J_k, i_k, j_k is considered quenched, and it is not explicitly indicated in our notation for H .

The role played by Gaussian integration by parts in the Sherrington-Kirkpatrick model, here is assumed by the following elementary derivation formula, holding for Poisson distributions,

$$\begin{aligned} \frac{d}{dt}P(K = k, t\alpha N) &\equiv \frac{d}{dt} \exp(-t\alpha N)(t\alpha N)^k/k! \\ &= \alpha N(P(K = k - 1, t\alpha N) - P(K = k, t\alpha N)). \end{aligned} \quad (82)$$

Then, all machinery of interpolation can be easily extended to the diluted models, as firstly recognized by Franz and Leone in [20].

In this way, the superadditivity property, the thermodynamic limit, and the generalized variational principle can be easily established. We refer to [20], and [15], for a complete treatment.

There is an important open problem here. While in the fully connected case the Poisson probability cascades provide the right auxiliary α systems to be exploited in the variational principle, on the other hand in the diluted case more complicated probability cascades have been proposed, as shown for example in [20], and in [22]. On the other hand, in [15], the very interesting proposal has been made that also in the case of diluted models the Poisson probability cascades play a very important role. Of course here the way how the auxiliary system interact with the original system is different, and involves a multi-overlap structure as explained in [15]. In this way a kind of very deep universality is emerging. Poisson probability cascades are a kind of universal class of auxiliary systems. The different models require different cavity fields ruling the interaction between the original system and the auxiliary system. But further work will be necessary in order to clarify this very important issue. For results about diluted models in the high temperature region, we refer to [23].

8 The short range model and its connections with the mean field version

The investigation of the connections between the short range version of the model and its mean field version are at the beginning. Here we limit ourselves to a synthetic description of what should be done, and to a short presentation of the results obtained so far.

First of all, according to the conventional wisdom, the mean field version should be a kind of limit of the short range model on a lattice in dimension d ,

when $d \rightarrow \infty$, with a proper rescaling of the strength of the Hamiltonian, of the form $d^{-\frac{1}{2}}$. Results of this kind are very well known in the ferromagnetic case, but the present technology of interpolation does not seem sufficient to assure a proof in the spin glass case. So this very basic result is still missing. In analogy with the ferromagnetic case, it would be necessary to arrive at the notion of a critical dimension, beyond which the features of the mean field case still hold, for example in the expression of the critical exponents and in the ultrametric hierarchical structure of the pure phases, or at least for the overlap distributions. For physical dimensions less than the critical one, then the short range model would need corrections with respect to its mean field version. Therefore, this is a completely open problem.

Moreover, always according to the conventional wisdom, the mean field version should be a kind of limit of the short range models, in finite fixed dimensions, as the range of the interaction goes to infinity, with proper rescaling. Important work of Franz and Toninelli shows that this is effectively the case, if a properly defined Kac limit is performed. Here, interpolation methods are effective, and we refer to [24], [25], [26] for full details.

Due to the lack of efficient analytical methods, it is clear that numerical simulations play a very important role in the study of the physical properties emerging from short range spin glass models. In particular, we refer to [27] and [28], for a detailed account about the evidence, coming from theoretical considerations and extensive computer simulations, that some of the more relevant features of the spontaneous replica breaking scheme of the mean field are also present in short range models in three dimensions. Different views are expressed for example in [29], where it is argued that the phase space structure of short range spin glass models is much simpler than that foreseen by the Parisi spontaneous replica symmetry mechanism.

Such very different views, both apparently strongly supported by reasonable theoretical considerations and powerful numerical simulations, are a natural consequences of the extraordinary difficulty of the problem.

It is clear that extensive additional work will be necessary before the clarification of the physical features exhibited by the realistic short range spin glass models.

9 Conclusion and outlook for future developments

As we have seen, in these last few years there has been an impressive progress in the understanding of the mathematical structure of spin glass models,

mainly due to the systematic exploration of comparison and interpolation methods. However many important problems are still open. The most important one is to establish rigorously the full hierarchical ultrametric organization of the overlap distributions, as appears in Parisi theory, and to fully understand the decomposition in pure states of the glassy phase, at low temperatures.

Moreover, it would be important to extend these methods to other important disordered models as for example neural networks. Here the difficulty is that the positivity arguments, so essential in comparison methods, do not seem to emerge naturally inside the structure of the theory.

Finally, the problem of connecting properties of the short range model, with those arising in the mean field case, is still almost completely open.

Acknowledgments

We gratefully acknowledge useful conversations with Michael Aizenman, Pierluigi Contucci, Giorgio Parisi and Michel Talagrand. The strategy explained in this report grew out from a systematic exploration of comparison and interpolation methods, developed in collaboration with Fabio Lucio Toninelli, and Luca De Sanctis.

This work was supported in part by MIUR (Italian Minister of Instruction, University and Research), and by INFN (Italian National Institute for Nuclear Physics).

References

- [1] P. Young, ed., *Spin glasses and random fields*, World Scientific, Singapore, 1987.
- [2] D.L. Stein, *Disordered Systems: Mostly Spin Glasses*, in: *Lectures in the Sciences of Complexity*, ed. D.L. Stein, Addison-Wesley, NY, 1989.
- [3] S.F. Edwards and P.W. Anderson, *Theory of spin glasses*, J. Phys. F: Met. Phys. **5**, 965-974 (1975).
- [4] D. Sherrington and S. Kirkpatrick, *Solvable Model of a Spin-Glass*, Phys. Rev. Lett. **35**, 1792-1796 (1975).
- [5] S. Kirkpatrick and D. Sherrington, *Infinite-ranged models of spin-glasses*, Phys. Rev. **B17**, 4384-4403 (1978).
- [6] M. Mézard, G. Parisi and R. Zecchina, *Analytic and Algorithmic Solution of Random Satisfiability Problems*, Science **297**, 812 (2002).

- [7] M. Mézard, G. Parisi and M. A. Virasoro, *Spin glass theory and beyond*, World Scientific, Singapore, 1987.
- [8] M. Talagrand, *Spin glasses: a challenge for mathematicians. Mean field models and cavity method*, Springer-Verlag, Berlin (2003).
- [9] F. Guerra, *Sum rules for the free energy in the mean field spin glass model*, Fields Institute Communications **30**, 161 (2001).
- [10] F. Guerra and F. L. Toninelli, *The Thermodynamic Limit in Mean Field Spin Glass Models*, Commun. Math. Phys. **230**, 71-79 (2002).
- [11] F. Guerra, *Broken Replica Symmetry Bounds in the Mean Field Spin Glass Model*, Commun. Math. Phys. **233**, 1-12 (2003).
- [12] M. Talagrand, *The Generalized Parisi Formula*, Compte Rendu de l'Académie des Sciences, Paris **337**, 111-114 (2003).
- [13] M. Talagrand, *The Parisi formula*, Annals of Mathematics, to appear (2005).
- [14] M. Aizenman, R. Sims and S. Starr, *Extended variational principle for the Sherrington-Kirkpatrick spin-glass model*, Phys. Rev. **B68**, 214403 (2003).
- [15] L. De Sanctis, *Structural Approachs to Spin Glasses and Optimization Problems*, Ph.D. Thesis, Department of Mathematics, Princeton University, 2005.
- [16] H.E. Stanley, *Introduction to phase transitions and critical phenomena*, Oxford University Press, New York and London, 1971.
- [17] D. Ruelle, *Statistical mechanics. Rigorous results*, W.A. Benjamin Inc., New York, 1969.
- [18] K. Joag-dev, M.D. Perlman and L.D. Pitt, *Association of normal random variables and Slepian's inequality*, Annals of Probability **11**, 451-455 (1983).
- [19] J.-P. Kahane, *Une inégalité du type Slepian and Gordon sur les processus gaussiens*, Israel J. Math. **55**, 109-110 (1986).
- [20] S. Franz and M. Leone, *Replica bounds for optimization problems and diluted spin systems*, J. Stat. Phys. **111**, 535-564 (2003).

- [21] G. Parisi, *A sequence of approximate solutions to the S-K model for spin glasses*, J. Phys. **A13**, L-115 (1980).
- [22] D. Panchenko and M. Talagrand, *Bounds for diluted mean-field spin glass models*, Prob. Theory Related Fields **130**, 319 - 336 (2004).
- [23] F. Guerra and F.L. Toninelli, *The high temperature region of the Viana-Bray diluted spin glass model*, J. Stat. Phys. **115**, 531-555 (2004).
- [24] F. Guerra and F.L. Toninelli, *Some comments on the connection between disordered long range spin glass models and their mean field version*, J. Phys. A: Math. Gen. **36**, 10987-95 (2003)
- [25] S. Franz and F.L. Toninelli, *The Kac limit for finite-range spin glasses*, Phys. Rev. Lett. **92**, 030602 (2004).
- [26] S. Franz and F.L. Toninelli, *Finite-range spin glasses in the Kac limit: free energy and local observables*, J. Phys. A: Math. Gen. **37**, 7433 (2004).
- [27] E. Marinari, G. Parisi and J. J. Ruiz-Lorenzo, *Numerical Simulations of Spin Glass Systems*, pp. 59-98, in [1].
- [28] E. Marinari, G. Parisi, F. Ricci-Tersenghi, J. J. Ruiz-Lorenzo and F. Zuliani, *Replica Symmetry Breaking in Short Range Spin Glasses: a Review of the Theoretical Foundations and of the Numerical Evidence*, J. Stat. Phys. **98**, 973-1074 (2000)
- [29] C. M. Newman and D. L. Stein, *Simplicity of state and overlap structure in finite-volume realistic spin glasses*, Phys. Rev. E **57**, 1356-1366 (1998).