

Linear Response for Granular Fluids

James Dufty and Aparna Baskaran

Department of Physics, University of Florida, Gainesville, FL 32611

J. Javier Brey

Física Teórica, Universidad de Sevilla, Apartado de Correos 1065, E-41080, Sevilla, Spain

(Dated: May 1, 2019)

The linear response of an isolated, homogeneous granular fluid to small spatial perturbations is studied by methods of non-equilibrium statistical mechanics. The long wavelength linear hydrodynamic equations are obtained, with formally exact expressions for the susceptibilities and transport coefficients. The latter are given in equivalent Einstein-Helfand and Green-Kubo forms. The context of these results and their contrast with corresponding results for normal fluids are discussed.

PACS numbers: 45.70.-n, 05.60.-k, 47.10.+g

Granular fluids consist of mesoscopic particles activated by external forces or initial conditions, and appear ubiquitously both in nature and in technological applications. Their flow properties are remarkably similar to those of normal fluids in many cases [1]. This has led to the speculation that the methods of non-equilibrium statistical mechanics developed for normal fluids might be adapted to explore the properties of granular fluids [2, 3]. To date, this has been realized in the examples of numerical molecular dynamics simulation and kinetic theory. However, the extensive formal theoretical tools have been applied to a more limited degree (e.g., impurity diffusion [4] and liquid state structure [5]). In some respects experimental studies of the response functions of interest here are more advanced than those of theory (see, for example, [6]). Here we report an application of linear response methods to obtain the hydrodynamic description of small perturbations in an isolated, homogeneous granular fluid. Although this example is perhaps of limited direct experimental relevance, it provides an instructive controlled test for the extension of these methods to granular fluids, illuminating important differences from normal fluids.

The simplest model of a granular fluid is considered: N smooth, inelastic, hard spheres ($d = 3$) or disks ($d = 2$) of mass m . The only difference from a corresponding model for normal fluids is a loss of energy in each binary collision, characterized by a restitution coefficient $0 < \alpha \leq 1$, with $\alpha = 1$ corresponding to the elastic limit. At the microscopic level, the fluid state is specified by the positions and velocities of all particles, i.e. by a point in the $2dN$ -dimensional phase space $\Gamma \equiv \{\mathbf{q}_1, \dots, \mathbf{q}_N, \mathbf{v}_1, \dots, \mathbf{v}_N\}$. This point evolves according to a deterministic dynamics of free streaming and collisions. The initial macroscopic state is specified by a probability density $\rho(\Gamma, 0)$, characterizing the known (incomplete) initial data. Consequently, all the ingredients required for a statistical mechanical description are available. In particular, the Liouville equation determines the macrostate $\rho(\Gamma, t)$ at all later times. The elements of a hydrodynamic description

then follow from averaging the local number, energy, and momentum densities to obtain the exact macroscopic balance equations. From them, the *phenomenological* hydrodynamic equations are obtained with the additional assumption of constitutive equations for the fluxes, as well as for the source term appearing in the equation for the energy. For systems with small gradients of the hydrodynamic fields, the Navier-Stokes equations for a granular fluid are obtained. Of course, the transport coefficients in these equations are unknown and must be supplied separately from experiment.

For a normal fluid, a formal theoretical description of hydrodynamics has been given by considering the linear response of an equilibrium fluid to small spatial perturbations [7]. The solutions to the linearized hydrodynamic equations, the so-called hydrodynamic modes, can then be identified from the long time, long wavelength behavior of the response functions. As a result, formally exact expressions for the transport coefficients are obtained in terms of time correlation functions. These are the Einstein-Helfand (EH) and Green-Kubo (GK) forms, which have proved useful as the basis for practical (approximate) theoretical predictions of these coefficients. The objective here is to report the application of this linear response approach to granular fluids.

The isolated granular fluid does not support a stationary, homogeneous equilibrium state. Instead, the simplest homogeneous state, known as the homogeneous cooling state (HCS), changes in time due to collisional energy loss. Spatial perturbations of the HCS therefore have a dynamics due to both the perturbation and the background reference state. However, the latter occurs only through the average energy and can be removed by a suitable choice of dimensionless variables. Effectively then, the analysis is similar to that for the equilibrium case, although the reference state is not the Gibbs distribution. The distribution function of the HCS, $\rho_0(\Gamma)$, is an example of a “normal distribution”, in the sense that its time dependence occurs entirely through the temper-

ature,

$$\rho_0(\Gamma, t) = (\ell v_0(t))^{-dN} \rho_0^* (\{\mathbf{q}_{ij}/\ell, \mathbf{v}_i/v_0(t)\}, n_0). \quad (1)$$

Here $\mathbf{q}_{ij} \equiv \mathbf{q}_i - \mathbf{q}_j$ and ℓ is the mean free path. The uniform hydrodynamic fields are the average number density, n_0 , and the temperature, $T_0(t)$, the latter occurring through the thermal velocity $v_0(t) \equiv \sqrt{2T_0(t)/m}$. The granular temperature is defined in terms of the average energy density, e_0 , in the same way as for equilibrium, $e_0 = dn_0 T_0/2$. The HCS distribution is the solution to the Liouville equation in the form [4]

$$\bar{\mathcal{L}}\rho_0 = 0, \quad \bar{\mathcal{L}} \equiv \bar{\mathcal{L}} + \frac{\zeta_0}{2} \sum_{i=1}^N \left(d + \mathbf{v}_i \cdot \frac{\partial}{\partial \mathbf{v}_i} \right), \quad (2)$$

where $\bar{\mathcal{L}}$ is the Liouville operator for hard spheres [2, 3], and $\zeta_0 \equiv -\partial_t \ln T_0$ is the ‘‘cooling rate’’.

To study the response of the HCS to spatial perturbations, an initial *local* HCS, $\rho_l(\Gamma, 0)$, is chosen. This distribution is a functional of specified nonuniform hydrodynamic fields $n(\mathbf{r})$, $T(\mathbf{r})$, and $\mathbf{u}(\mathbf{r})$, characterizing the average number density, energy density, and momentum density for this ensemble. To construct it, consider first the HCS for an inhomogeneous fluid (i.e., the solution to (2) in an external inhomogeneous field). Since the average nonuniform density is a functional of the external field, inverting that relationship gives the HCS distribution as a functional of the density $\rho_0^{(inh)} = (\ell v_0)^{-dN} \rho_l^* (\{\mathbf{q}_{ij}/\ell, \mathbf{v}_i/v_0\} | n)$. The special case in (1) is the value of this functional at the uniform state. Next, the non-uniform flow field and temperature are incorporated through the replacements of \mathbf{v}_i by $\mathbf{v}_i - \mathbf{u}(\mathbf{q}_i)$ and $v_0(T_0)$ by $v_0(\mathbf{q}_i) \equiv v_0[T(\mathbf{q}_i)]$ for each particle in the arguments of $\rho_0^{(inh)}$. The local HCS distribution is then

$$\rho_l(\Gamma) \equiv \left\{ \sum_{i=1}^N [\ell v_0(\mathbf{q}_i)]^{-d} \right\} \rho_l^* \left(\left\{ \frac{\mathbf{q}_{ij}}{\ell}, \frac{\mathbf{v}_i - \mathbf{u}(\mathbf{q}_i)}{v_0(\mathbf{q}_i)} \right\} | n \right). \quad (3)$$

This initial preparation is natural, in the sense that it is expected that locally there is a rapid relaxation of the velocity distribution as each region attempts to approach the HCS at its own values of the hydrodynamic fields. It is the analogue of rapid collisional approach to a local equilibrium state for normal fluids. For smooth spatial variations of small amplitude, to linear order

$$\begin{aligned} \rho(0) &= \rho_l(0) \simeq \rho_0 + \sum_{\beta} \int d\mathbf{r} \left[\frac{\delta \rho_l(0)}{\delta y_{\beta}(\mathbf{r}, 0)} \right]_0 \delta y_{\beta}(\mathbf{r}, 0) \\ &= \rho_0(\Gamma, 0) + \sum_{\beta} \phi_{\beta}(-\mathbf{k}) \delta \tilde{y}_{\beta}(\mathbf{k}, 0), \end{aligned} \quad (4)$$

where the $y_{\beta}(\mathbf{r}, 0)$ are linear combinations of the hydrodynamic fields, as specified below. The second equality follows by considering a single Fourier component for these fields, and it defines $\phi_{\beta}(\mathbf{k})$.

This preparation is also special in the sense that it excites only hydrodynamic modes in the long wavelength limit. This follows from the fact that the y_{β} can be chosen such that $\Phi_{\beta} \equiv \phi_{\beta}(\mathbf{k} = 0)$ are eigenfunctions of the extended Liouville operator $\bar{\mathcal{L}}$ [8]

$$\bar{\mathcal{L}}\Phi_{\beta} = \lambda_{\beta}\Phi_{\beta}, \quad \Phi_{\beta} \equiv \int d\mathbf{r} \left[\frac{\delta \rho_l}{\delta y_{\beta}(\mathbf{r}, 0)} \right]_0, \quad (5)$$

with the eigenvalues $\{\lambda_{\beta}\} = \{0, \zeta_0/2, -\zeta_0/2, \dots, -\zeta_0/2\}$. The specific set of fields in (4) leading to this result are

$$\{y_{\beta}\} = \left\{ \frac{n}{n_0}, \frac{T}{T_0} + 2 \frac{\partial \ln \zeta_0}{\partial \ln n_0} \frac{n}{n_0}, \frac{\hat{\mathbf{k}} \cdot \mathbf{u}}{v_0}, \frac{\mathbf{u}_{\perp}}{v_0} \right\}. \quad (6)$$

The field flow \mathbf{u} has been decomposed into its longitudinal component in the direction of $\hat{\mathbf{k}} = \mathbf{k}/k$, and the remaining $d-1$ transversal components denoted by \mathbf{u}_{\perp} . The eigenvalue $-\zeta_0/2$ is d -fold degenerated. Significantly, the eigenvalues λ_{β} are the same as those of the linearized macroscopic balance equations for these variables in the long wavelength limit. Hence they can be considered as hydrodynamic modes reflected in the microscopic dynamics. These observations are critical to the following linear response analysis. In the elastic limit, the eigenfunctions become linear combinations of the global conserved number, energy, and momentum.

The response of the hydrodynamic fields δy_{β} to the initial perturbation (4) is

$$\delta \tilde{y}_{\beta}^*(\mathbf{k}^*, s) = \sum_{\gamma} C_{\beta\gamma}(\mathbf{k}^*, s) \delta \tilde{y}_{\gamma}^*(\mathbf{k}^*, 0), \quad ds = \frac{v_0}{\ell} dt, \quad (7)$$

$$C_{\beta\gamma}(\mathbf{k}^*, s) \equiv \int d\Gamma^* \delta Y_{\beta}^*(\mathbf{k}^*) e^{-s\bar{\mathcal{L}}^*} \phi_{\gamma}^*(-\mathbf{k}). \quad (8)$$

The dimensionless time s is a measure of the average number of collisions, and Y_{β} is the phase function whose average is y_{β} . Moreover, the asterisks indicate dimensionless variables in the units defined by ℓ , v_0 , and m [4]. Clearly, if the $\delta \tilde{y}_{\beta}^*(\mathbf{k}^*, s)$ obey hydrodynamic equations at long space and time scales, the corresponding excitations must exist in the spectrum of the correlation functions $C_{\beta\gamma}(\mathbf{k}^*, s)$. This suggests defining an exact transport matrix $\mathcal{K}_{\alpha\beta}(\mathbf{k}^*, s)$ by

$$[\partial_s + \mathcal{K}(\mathbf{k}^*, s)] \delta \tilde{y}^*(\mathbf{k}^*, s) = 0, \quad (9)$$

with the identification

$$\mathcal{K}(\mathbf{k}^*, s) = -[\partial_s C(\mathbf{k}^*, s)] C^{-1}(\mathbf{k}^*, s). \quad (10)$$

A matrix notation has been introduced for simplicity. The hydrodynamic equations are then identified from the matrix $\mathcal{K}(\mathbf{k}^*, s)$ for large s and small \mathbf{k}^* . In particular, the linear Navier-Stokes approximation is defined by an expansion of $\mathcal{K}(\mathbf{k}^*, s)$ to order k^{*2} , with the coefficients

evaluated for large s . In general, this procedure does not lead to tractable results since initial perturbations excite microscopic as well as hydrodynamic modes. However, the special initial preparation chosen here is such that only hydrodynamic excitations occur in the long wavelength limit. This follows directly from (5) and (8)

$$C_{\beta\gamma}(0, s) = e^{-s\Lambda_\beta^*} \delta_{\beta\gamma}. \quad (11)$$

As a consequence, the expansion of (10) is assured to start on the hydrodynamic branch. Define

$$C(\mathbf{k}^*, s) = e^{-s\Lambda} + ik^* C^{(1)}(s) + (ik^*)^2 C^{(2)}(s) + \dots, \quad (12)$$

where $\Lambda_{\alpha\beta} = \lambda_\alpha^* \delta_{\alpha\beta}$. Then it is found

$$\mathcal{K}(\mathbf{k}^*, s) = \Lambda + ik^* \mathcal{K}^{(1)}(s) + (ik^*)^2 \mathcal{K}^{(2)}(s) + \dots, \quad (13)$$

with

$$\mathcal{K}^{(1)}(s) = - \left[(\partial_s + \Lambda) C^{(1)}(s) \right] e^{s\Lambda}, \quad (14)$$

$$\mathcal{K}^{(2)}(s) = - \left[(\partial_s + \Lambda) C^{(2)}(s) + \mathcal{K}^{(1)}(s) C^{(1)}(s) \right] e^{s\Lambda}. \quad (15)$$

The transport coefficients and susceptibilities of the phenomenological Navier-Stokes equations can be identified with appropriate matrix elements in these expressions: $\mathcal{K}^{(1)}$ gives the Euler coefficients, $\mathcal{K}^{(2)}$ gives the Navier-Stokes coefficients, and so on. In particular, these expressions provide an extension of the EH representation of transport coefficients to granular fluids. The GK forms follow from these results by partial integration. The detailed expressions and analysis will be given elsewhere.

In this brief presentation, further details will be limited to the case of the transverse flow field response, i.e. $\delta\tilde{y}_\perp^*(\mathbf{k}^*, s)$ associated with one component of \mathbf{u}_\perp in (6). The only non-vanishing element $C_{\alpha\perp}$ is $C_{\perp\perp}$. Also note that $C_{\perp\perp}^{(1)} = 0$ from symmetry. Then, Eq. (9) becomes to order k^2

$$\left[\partial_s - \frac{\zeta_0^*}{2} - k^{*2} \mathcal{K}_{\perp\perp}^{(2)}(s) \right] \delta\tilde{y}_\perp^*(\mathbf{k}^*, s) = 0. \quad (16)$$

If $\mathcal{K}_{\perp\perp}^{(2)}(s)$ has a limit for large s , this becomes the shear diffusion equation, with $\mathcal{K}_{\perp\perp}^{(2)}$ identified in terms of the dimensionless shear viscosity $\eta^* \equiv \eta/mn_0\ell v_0$,

$$\begin{aligned} \eta^* &= - \lim_{s \gg 1} \mathcal{K}_{\perp\perp}^{(2)}(s) = \lim_{s \gg 1} \partial_s \left[C_{\perp\perp}^{(2)}(s) e^{-\frac{s\zeta_0^*}{2}} \right] \\ &= \lim_{s \gg 1} \partial_s \frac{1}{N} \int d\Gamma^* \left[e^s \left(\mathcal{L}^* - \frac{\zeta_0^*}{2} \right) M^* \right] \mathcal{N}^* \rho_0^*, \end{aligned} \quad (17)$$

where $-\mathcal{L}^*$ is the adjoint of $\bar{\mathcal{L}}^*$. The phase function M^* is the space moment of the momentum density

$$M^* = \sum_{i=1}^N (\hat{\mathbf{k}} \cdot \mathbf{q}_i^*) (\hat{\mathbf{e}}_1 \cdot \mathbf{v}_i^*), \quad (18)$$

where $\hat{\mathbf{e}}_1$ is the unit vector in the direction of \mathbf{u}_\perp considered. The function $\mathcal{N}^* \rho_0^* \equiv \phi_\perp^{(1)}$ is given by

$$\begin{aligned} \mathcal{N}^* \rho_0^* &= - \int d\mathbf{r}^* (\hat{\mathbf{k}} \cdot \mathbf{r}^*) \hat{\mathbf{e}}_1 \cdot \left[\frac{\delta \rho_i^*}{\delta \mathbf{u}^*(\mathbf{r}, 0)} \right]_0 \\ &= \sum_{i=1}^N (\hat{\mathbf{k}} \cdot \mathbf{q}_i^*) \hat{\mathbf{e}}_1 \cdot \frac{\partial}{\partial \mathbf{v}_i^*} \rho_0^*. \end{aligned} \quad (19)$$

In the elastic limit and for $\rho_0 \rightarrow \rho_e$, the equilibrium canonical distribution function, the Helfand form for the shear viscosity of a normal fluid is recovered

$$\eta_{elast}^* = - \lim_{s \gg 1} \partial_s \frac{2}{N} \int d\Gamma^* \left(e^{sL^*} M^* \right) M^* \rho_e^*. \quad (20)$$

Both (17) and (20) contain correlation functions involving the momentum density moment M . For a normal fluid, this is an autocorrelation function, while for the granular fluid a new moment \mathcal{N} appears that is determined from the HCS distribution.

These correlation functions must grow as s for long times to obtain a constant viscosity. This is analogous to the definition of the diffusion coefficient for a Brownian particle in terms of its mean square displacement, and for that reason the representation above is referred to as the EH form. An equivalent representation is obtained by carrying out the time derivative, shifting the time dependence, and integrating with respect to time. The result is

$$\begin{aligned} \eta^* &= \frac{1}{N} \int d\Gamma^* F^* \mathcal{N}^* \rho_0^* \\ &+ \lim_{s \ll 1} \int_0^s ds' \frac{1}{N} \int d\Gamma^* F^* e^{-s' \left(\bar{\mathcal{L}}^* + \frac{\zeta_0^*}{2} \right)} \mathcal{F}^* \rho_0^*. \end{aligned} \quad (21)$$

The fluxes are given by

$$F^* = \left(\mathcal{L}^* - \frac{\zeta_0^*}{2} \right) M^*, \quad \mathcal{F}^* \rho_0^* = - \left(\bar{\mathcal{L}}^* + \frac{\zeta_0^*}{2} \right) \mathcal{N}^* \rho_0^*. \quad (22)$$

This is the GK representation. For normal fluids, F^* becomes the volume integrated momentum flux and the second term of (21) becomes the expected time integral of a flux autocorrelation function. As noted below, hard sphere systems entail the additional first term on the right side even for normal fluids. Moreover, as with the EH form, the correlation functions for the granular fluid involve new phase functions defined from the HCS distribution.

The objective here has been to explore the application of linear response methods to the simplest model and state of a granular fluid. The details for all hydrodynamic modes and all transport coefficients will be reported elsewhere. These results illustrate the applicability of such methods and their potential for more complex conditions of experimental interest. They also expose many subtleties in the differences from corresponding results for

normal fluids and provide a warning against a simple translation of normal fluid theory to granular fluids. In closing, some observations are highlighted:

i) The identification of the transport matrices in (13)-(15) constitutes a formal derivation of the linear Navier-Stokes equations. Important limitations on this derivation are placed by restrictions on length and time scales. As for normal fluids, it is expected that the conditions are times long compared to the mean free time ($s \gg 1$) and wavelengths long compared to the mean free path ($k^* = k\ell \ll 1$).

ii) These formal results have no *a priori* limitation on the density or degree of inelasticity. However, peculiarities of the isolated system (cluster instability [10], inelastic collapse [11]) must be confronted at high densities and large inelasticity. These problems can be removed in practice by considering small system sizes and protocols for simulations to avoid dangerous configurations with no physical consequences.

iii) The transport coefficients are given in terms of averages over the HCS reference state, which differs from the equilibrium canonical distribution. Consequently, one of the two phase functions in the EH and GK expressions is different from that for normal fluids.

iv) The generator $\bar{\mathcal{L}}$ for hard sphere dynamics is singular and must be expressed in terms of binary collision operators rather than forces. Important structural changes result in the form of the GK expressions and the associated microscopic fluxes [9]. The GK expressions for both elastic and inelastic hard sphere fluids have a new contribution due to instantaneous collisions, as well as the usual time integral of correlation functions of fluxes. The latter are expressed in terms of the binary collision operators and are different for forward and backward dynamics, both occurring in the GK expressions.

v) The reference HCS is time dependent, due to collisional cooling. A dimensionless form of the equations is required to eliminate this dependence, resulting in a new generator $\bar{\mathcal{L}}^*$ rather than the Liouville operator $\bar{\mathcal{L}}$. In this representation, the HCS is a stationary solution, $\bar{\mathcal{L}}^* \rho_0^* = 0$.

vi) The stationary HCS distribution is singular in the sense that it is restricted to a zero total momentum \mathbf{P} surface, i.e., $\rho_0^* \propto \delta(\mathbf{P}^*)$. Furthermore, the elastic limit is proportional to a delta function in the total energy, since an isolated system is being considered. This constant momentum, constant energy ensemble is often referred to as the “computer” ensemble, as it corresponds to usual conditions in molecular dynamics simulation. For granular fluids, the energy changes, even in isolation, but the constant momentum condition must be retained for consistency with $\bar{\mathcal{L}}^* \rho_0^* = 0$.

vii) The presence of hydrodynamic excitations is expected for a wide class of initial perturbations. However, the choice made here is both physically and mathematically motivated. As the HCS distribution characterizes

the homogeneous state through its dependence on the hydrodynamic fields, it is natural to consider their spatial perturbation through this same dependence. More important for the analysis here is that the resulting perturbation excites *only* hydrodynamic modes in the long wavelength limit. This follows from the property (5), and plays a central role in deriving the explicit forms for the expansion (13). For general initial preparations, the leading term in (13) will not be Λ , but rather some complex time dependent quantity that must be calculated to be Λ at long times.

viii) An earlier derivation of GK expressions leads to different results [12]. The method is formally correct, but does not have the simplifying features of the initial preparation considered here. In addition, some of the special properties of the hard sphere generator were not accounted for in detail, so that different fluxes are obtained and the instantaneous contribution is missed.

The research of J.D. and A.B. was supported in part by the Department of Energy Grant (DE-FG03-98DP00218). J.J.B. acknowledges partial support from the Ministerio de Educación y Ciencia (Spain) through Grant No. BFM2005-01398 (partially financed by FEDER funds). This work also was supported in part by the National Science Foundation under Grant No. PHY99-0794 to the Kavli Institute for Theoretical Physics, UC Santa Barbara.

-
- [1] *Granular Gases*, edited by T. Pöschel and S. Luding (Springer-Verlag, Berlin, 2000); *Granular Gas Dynamics*, edited by T. Pöschel and N. Brilliantov (Springer-Verlag, Berlin, 2003).
 - [2] J. J. Brey, J. W. Dufty, and A. Santos, *J. Stat. Phys.* **87**, 1051 (1997).
 - [3] T. P. C. van Noije and M. H. Ernst, in *Granular Gases*, edited by T. Pöschel and S. Luding (Springer-Verlag, Berlin, 2000).
 - [4] J. Dufty, J. J. Brey, and J. Lutsko, *Phys. Rev. E* **65**, 051303 (2002); J. Dufty and V. Garzó, *J. Stat. Phys.* **105**, 723 (2001).
 - [5] J. Lutsko, *Phys. Rev. E* **63**, 061211 (2001).
 - [6] N. Xu and C. S. O’Hern, *Phys. Rev. Lett.* **94**, 055701 (2005).
 - [7] J.A. McLennan, *Introduction to Nonequilibrium Statistical Mechanics*, (Prentice-Hall, New Jersey, 1989).
 - [8] J. Dufty and A. Baskaran in *Nonlinear Dynamics in Astronomy and Physics*, S. Gottesman ed., *Annals of the New York Academy of Sciences* vol. 1045, 93 (2005); A. Baskaran, J. Brey, and J. Dufty (unpublished).
 - [9] J. Dufty, *Mol. Phys.* **100**, 2331 (2002); J. Dufty and M. H. Ernst, *Mol. Phys.* **102**, 2123 (2004).
 - [10] I. Goldhirsch and G. Zanetti, *Phys. Rev. Lett.* **70**, 1619 (1993).
 - [11] S. McNamara and W.R. Young, *Phys. Rev. E* **53**, 5089 (1996).
 - [12] I. Goldhirsch and T. P. C. van Noije, *Phys. Rev. E* **61**, 3241 (2000).