

Metallic phase in a two-dimensional disordered Fermi system with singular interactions

Victor M. Galitski

Kavli Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106-4030 and
Department of Physics, University of Virginia, Charlottesville, VA 22904-4714

We consider a disordered system of gapless fermions interacting with a singular transverse $(2+1)$ -dimensional gauge field. We study quantum corrections to fermion conductivity and show that they are very different from those in a Fermi liquid with non-singular interactions. In particular, the weak-localization effect is suppressed by magnetic field fluctuations. We argue that these fluctuations can be considered static at time scales of fermionic diffusion. By inducing fluxes through disusive loops that contribute to weak localization, they dephase via the Aharonov-Bohm effect. It is shown that while the flux-flux correlator due to thermal fluctuations of magnetic field is proportional to the area enclosed by the loop, the correlator due to quantum fluctuations is proportional to the perimeter of the loop. The possibility of dephasing due to these quasistatic configurations and the corresponding rates are discussed. We also study interaction induced effects and show that perturbation theory contains infrared divergent terms originating from unscreened magnetic interactions. These singular (Hartree) terms are related to scattering of a fermion off of the static potential created by the other fermions. We show that due to singular small-angle scattering, the corresponding contributions to the density of states and conductivity are very large and positive indicating that the fermion-gauge system remains metallic at low temperatures.

PACS numbers: 71.27.+a, 71.10.Hf, 71.30.+h, 72.15.Rn

I. INTRODUCTION

There are many seemingly unrelated physical systems, whose low-energy description contains fermionic quasi-particles coupled to a fluctuating gauge field. A notable example is a variety of insulating spin liquid phases, which are in principle allowed in strongly interacting electronic systems.^{1,2} These spin liquid states are often described in terms of non-standard quasiparticles coupled to dynamical gauge fields.³ Recent experiments of Kanoda et al.^{4,5} indicate the possibility of a spin-liquid phase in the organic compound $(\text{ET})_2\text{Cu}_2(\text{CN})_3$. Motrunich⁶ and S.-S. Lee and P. Lee⁷ argued that the so-called uniform resonating-valence-bond (URVB) phase is the most likely candidate for the phase discovered in these experiments. This phase is described by a degenerate Fermi gas of spinons interacting with a dynamical $U(1)$ gauge field. There are other physical models, whose low-energy description has an identical or very similar mathematical structure: In particular, the Halperin-Lee-Read theory⁸ of a compressible Fermi-liquid-like state at the filling factor $\nu = 1/2$ is formulated in terms of composite fermions interacting with a dynamical Chern-Simons field. Another example is a "vortex metal" phase suggested in Ref. [9]. To access this phase, one starts with a bosonic theory describing preformed Cooper pairs, and using duality transforms it into a theory of bosonic vortices interacting with a gauge field. The corresponding dual magnetic field represents the (fluctuating) Cooper pair charge density, whose mean value can be very large. If the dual filling factor is $\nu_{\text{dual}} = 1/2$, then vortices may become fermions^{9,10} forming the Halperin-Lee-Read state ("vortex metal"), which is also described by a theory with singular interactions. Remarkably, similar sin-

gular interactions may arise in more conventional systems in the vicinity of quantum phase transitions, where soft bosonic fluctuation modes can mimic longitudinal gauge-field fluctuations. Finally we note that strictly speaking physical electrons in metals and semiconductors should be described in terms of a gauge theory, since they are coupled to a fluctuating $[(3+1)\text{-dimensional}]$ electromagnetic field. The corresponding relativistic effects are typically negligible at all reasonable temperatures, but in the strict zero-temperature limit, they destroy the usual Fermi liquid picture.¹¹ An important difference between the usual quantum electrodynamics and the model studied in this paper is the magnitude of the effects associated with gauge fluctuations: The corresponding gauge-field-fermion coupling in the abelian electrodynamics appearing in the context of quantum magnets³ or the quantized Hall effect⁸ is not small, and thus quantum gauge fluctuations manifest themselves much stronger.

The aforementioned fermion-gauge model has been studied by Nayak and Wilczek¹², by Altshuler et al.¹³, and by Polchinski,¹⁴ who concluded that at least in the limit of small gauge-field-fermion coupling (large- N limit), it is described by a non-Fermi liquid strong-coupling fixed point. This fixed point can be visualized as a collection of patches on the Fermi surface, which do not interact with each other unless the momentum transfer is two Fermi momenta. This phase has a number of other unusual properties: For instance, weak non-singular density-density interactions are irrelevant, regardless the sign, and the physics is determined almost entirely by the interactions between the fermions and singular transverse fields. These results are valid in very clean systems, where the fermion motion is ballistic at low temperatures.

In this paper we investigate the effects of disorder on the properties of the fermion-gauge system and study quantum interference effects. Understanding these effects of disorder is important in attempts to relate theoretical results to real experiments, especially those searching for spin-liquid phases (in particular thermal transport experiments). All real systems contain finite amount of disorder, which may strongly affect their properties at low temperatures.¹⁵ In a usual Fermi-liquid, even weak disorder leads to weak localization¹⁶ and similar interaction-induced effects¹⁷ and eventually to an insulating behavior. These quantum interference effects have been observed in numerous experiments and theoretical results have been quantitatively verified.¹⁷ An important question is whether disorder results in a similar localization transition in fermion-gauge systems. The results presented in this paper suggest that the fermion-gauge system may remain metallic at low temperatures, as long as disorder is relatively weak (dimensionless conductance is large, $\gg 1$). This is due to the existence of unscreened magnetic forces. First, magnetic field fluctuations produce vortices through closed trajectories, which contribute to weak localization. These vortices result in Aharonov-Bohm phase shifts that fermions acquire, while traveling around the diusive loops in the opposite directions. This topological effect leads to a relatively large dephasing rate at finite temperatures $\propto g^2 T$ (where g is the dimensionless gauge-field-fermion coupling constant). We suggest that similar quasistatic configurations due to quantum fluctuations of the magnetic field may, in principle, lead to a similar dephasing effect with the characteristic rate $\propto (g^2 = 2)^{-1}$ (where τ is the scattering time). Second, gauge fluctuations lead to interaction corrections of Ahlshuler-Aronov type. In a fermion gauge system the leading contributions come from the Hartree processes, which correspond to the scattering of fermions off of the static interaction potential, created by the other fermions. Due to the long-range (unscreened) nature of magnetic interactions, the corresponding scattering rate diverges, due to singular small angle scattering. These Hartree contributions to the density of states and conductivity are positive, which may indicate the stability of the metallic phase.

The structure of the paper is as follows:

In Sec. II, we introduce basic notations and describe the model, which consists of a degenerate Fermi gas of particles (that we occasionally call spinons) coupled to a dynamic non-compact $U(1)$ gauge field in the presence of disorder. In Sec. III, we show that the clean strong-coupling Nayak-Wilczek fixed point¹² is unstable against weak disorder, and the system should be described by a diusive theory.

In Sec. IV, we qualitatively study the weak localization correction to conductivity taking into account fluctuating gauge fields. We show that gauge fluctuations are slow compared to fermionic motion, and therefore the magnetic field can be considered static on the time-scales of fermionic diffusion. We explicitly calculate the correla-

tion function of this random magnetic field. We argue that dephasing due to this factor is characterized by the flux correlator of a typical diusive trajectory. At finite temperatures, this correlator is proportional to the area of the path, while at zero temperature the correlator is proportional to the perimeter of the path.

In Sec. IV, we study quantum corrections of Ahlshuler-Aronov type due to the interactions mediated by the transverse gauge field. We find that direct perturbation theory contains infrared divergent diagrams. These singular contributions to the fermionic self-energy, density of states, and conductivity come from the Hartree processes in the diusion channel. The Hartree diagrams and other less divergent exchange diagrams are calculated. It is shown that the correction to conductivity is proportional to a formally infrared divergent integral coming from unscreened magnetic interactions.

II. THE MODEL

In this paper we consider a two-dimensional system of fermions coupled to a fluctuating gauge field in the presence of a disorder potential. The Hamiltonian is

$$H = H_{a,f} + H_{dis} + H_a; \quad (1)$$

where

$$H_{a,f} = \sum_{p,i} f_{p,i}^\dagger E(p) f_{p,i} + \sum_{p,q,i} f_{p+q,i}^\dagger a(q) \frac{\partial E(p)}{\partial p} f_{p,i}; \quad (2)$$

In Eq. (2), f^\dagger and f are the fermion creation and annihilation operators, $E(p)$ is the fermionic spectrum, $i = 1, 2, \dots, N$ is a spin index, H_{dis} is the disorder Hamiltonian, and H_a is the bare Hamiltonian for the gauge field a , which can be written as

$$H_a = \frac{1}{2} \int d^2r \left[g_0^2 e^2(r) + \frac{1}{g_0^2} [r - a(r)]^2 \right]; \quad (3)$$

where a and $e = \frac{i\partial}{\partial a}$ are canonically conjugate quantum operators and g_0 is the bare gauge coupling constant. In the case of the $uRVB$ phase in the tJ -model, this bare coupling is of order¹⁸ $g_0^2 = \frac{1}{s^{(0)}}$, where s is the charge gap for holons and $s^{(0)}$ is the bare phase stiffness.

A clean fermion-gauge system (i.e., $H_{dis} = 0$) has been studied in Refs. [12, 13, 14] in the large- N limit. It was found that in this limit the system is described by a non-Fermi liquid strong coupling fixed point, in which the physics is dominated by singular magnetic interactions. The results of the large- N treatment can be summarized as follows: First, vertex corrections with zero momentum transfer are small, $\propto N^{-1}$. This is due to the fact that gauge bosons are slow compared to fermions, and the analog of Migdal's theorem holds.¹⁹ Due to Migdal's theorem, a self-consistent (Eliashberg-like²⁰) theory can be developed. In this theory, the electronic self-energy

depends only on frequency^{21,22} $\omega / \omega_c^{2=3}$, and is determined by interaction corrections in leading order perturbation theory. Likewise, the effective action for the gauge field is determined by fermionic quasiparticles and can be calculated in leading order perturbation theory. The longitudinal (electric) forces are screened and become effectively short-range. The transverse (magnetic) forces are not screened, and the corresponding gauge-field propagator has the form :

$$K^{(\text{clean})}(\mathbf{l}; \mathbf{q}) = P^{(\text{tr})}(\mathbf{q}) \frac{1}{i\mathbf{l} \cdot (\mathbf{v}\mathbf{q}) + q^2}; \quad (4)$$

where \mathbf{l} and \mathbf{q} are constants and $P^{(\text{tr})}(\mathbf{q}) = \frac{1}{2} q q = q^2$. Even though, the vertices with small momentum transfers are negligible, the $2p_F$ -vertices are singular. This singularity may in principle lead to a spin-density-wave transition^{14,23,24} or a Kohn-Luttinger pairing instability.^{25,26} Considering energy-scales above the corresponding transition temperatures (if any), one can study the following gapless fixed point, discovered in Refs. [12,13]:

$$S = \sum_{\mathbf{p}} \int_{\mathbf{p}} \bar{\psi}_{\mathbf{p}} (\mathbf{l}) + v_F p_z + \frac{p_{jj}^2}{2m} \bar{\psi}_{\mathbf{p}} \psi_{\mathbf{p}} + \int_{\mathbf{q}} a(\mathbf{l}; \mathbf{q}) q^2 + \frac{j^l j}{q} a(\mathbf{l}; \mathbf{q}) + g \sum_{\mathbf{k}} \int_{\mathbf{q}} \bar{\psi}_{\mathbf{k}} \psi_{\mathbf{k}+\mathbf{q}} \bar{\psi}_{\mathbf{p}} v_F(\mathbf{q}) a(\mathbf{l}; \mathbf{q}) + \text{h.c.} \quad (5)$$

This action describes a collection of patches on the Fermi line, which do not interact with each other. In Eq. (5), p_{jj} and p_z correspond to the directions tangent and perpendicular to the Fermi line in the center of the region being considered. One can add different operators to the action and consider their relevance under scaling: $\mathbf{l} \cdot \mathbf{l} \sim l^2$, $p_z \sim l^2$, and $p_{jj} \sim l^2$. The gauge-field-fermion coupling [the last term in Eq. (5)] is exactly marginal and can be considered perturbatively with respect to the Gaussian part. This allows for controlled calculations within the renormalization group scheme. Using this scheme, one can find that non-singular density-density interactions are irrelevant at the tree level, which justifies dropping short-range interactions and screened electric forces.

We are interested in the effects of disorder on the properties of the fermion-gauge system. Let us add the following term to action (5)

$$S_{\text{dis}} = \int d^2 r V(\mathbf{r}) \int dt f(\mathbf{r}; t) f(\mathbf{r}; t);$$

where $V(\mathbf{r})$ is a quenched disorder potential with the correlation function

$$\langle V(\mathbf{r}) V(\mathbf{r}') \rangle = u_0 \delta(\mathbf{r} - \mathbf{r}')$$

Using the replica technique, we perform an averaging over disorder realizations and obtain the following contribution to the action

$$S_{nn^0} = u_0 \int d^2 q O_n(\mathbf{q}) O_{n^0}(-\mathbf{q}); \quad (6)$$

where n and n^0 are the replica indices and $O_n(\mathbf{q}) = \int d\mathbf{p} \bar{\psi}_{\mathbf{p}} \psi_{\mathbf{p}+\mathbf{q}}$. This operator scales as

$$O(s^{2=3} q; s^{1=3} q_{jj}) \sim s^{2=3} O(q); \quad (7)$$

From Eqs. (6) and (7), we conclude that operator (6) is a relevant perturbation, and therefore the Nayak-Wilczek strong coupling phase is unstable against weak quenched disorder, and flows toward a disusive phase.

To access this disusive phase, we will go back to Eq. (1) and include disorder at the very beginning. The strategy is to treat gauge interactions perturbatively with respect to the disusive Fermi liquid fixed point. An important issue is to determine a parameter regime (i.e., temperature and couplings), in which the perturbative description is reliable. Another important problem is to calculate interaction corrections to the properties of the disusive Fermi liquid fixed point in this regime. The main difficulty is due to the fact that we must treat disorder and singular gauge interactions self-consistently.

Our model consists of a disusive Fermi liquid coupled to a dynamic gauge field. The propagator of this gauge field is determined by the fermions. In the disusive regime, Landau damping [the term $j^l j$ in Eq. (4)] is replaced with the dissipative term $j^l j$. The corresponding propagator of the transverse gauge field reads

$$K^{(\text{di})}(\mathbf{l}; \mathbf{q}) = P^{(\text{tr})}(\mathbf{q}) \frac{1}{j^l j + q^2}; \quad (8)$$

where $\chi = (E_F)^{-1}$ is the dimensionless conductance, which is assumed large, $\tau = 1/(2\chi)$ is the scattering time, $\rho = m$ is the density of states, and $\chi = \chi_m$ is a non-universal "Landau diamagnetic susceptibility" of fermions, which depends strongly on the form of the fermionic spectrum, and χ is a numerical coefficient characterizing this non-universal dependence [in a Fermi gas, $\chi_0 = 1/(12)$]. We also assume that the dimensionless fermion-gauge-field coupling constant g is small, which corresponds to the large- N limit with $g \sim N^{-1/2} \sim 1$. In what follows we will treat the dimensionless fermion-gauge-field coupling as a formal small parameter without referring to the large- N approximation. In this paper, we mostly concentrate on the effects due to a singular transverse gauge field, and assume that all the other interactions (including dynamically screened electric forces) are not important. We emphasize that treatment of singular density-density interactions is similar. Such singular longitudinal fluctuations may appear in the vicinity of quantum phase transitions.

III. DEPHASING BY MAGNETIC FIELD FLUCTUATIONS

In this section, we qualitatively discuss the weak localization correction to conductivity in the presence of a fluctuating magnetic field. We assume the validity of the

straightforward, and we obtain the following expression:

$$C^t; (r; r^0) = \frac{1}{2} \int_{r(t)=r}^{\infty} D[r(t)] \exp \left(- \int_{t_1}^{\infty} dt_1 \frac{r^2(t_1)}{4D} \right) \frac{g^2}{2} \int_{t_1}^{\infty} dt_2 \int_{r(t_2)=r}^{\infty} K^-(t_1; t_2) r(t_2); \quad (12)$$

In what follows, we will use the notation S for the last term in Eq. (12). We note that in (12), K^- is the sum of the correlation functions:

$$K^-(t_1; t_2) = K^-(t_1 + t_2; r) + K^-(t_1 - t_2; r) + K^-(t_1 - t_2; r) + K^-(t_2 - t_1; r); \quad (13)$$

with $r = r(t_2) = r(t_1)$.

At this point, one assumes that the typical frequencies are much smaller than temperature, which justifies the classical approximation used. This assumption should be verified a posteriori by comparing the dephasing rate and temperature. The corresponding classical correlation function thus reads (we drop here the terms containing higher powers of \hbar):

$$K^{(class)}(\hbar; q) = \frac{2T}{2\hbar^2 + \hbar^2 q^4} P^{(tr)}(q); \quad (14)$$

To proceed further we make a crucial observation mentioned in the beginning of this section: The field fluctuations are slow compared to fermions. Really, if $\hbar \ll \hbar^2 q^2$ and $q \ll 1 = \frac{1}{D}$, then $\hbar = (\hbar^2 q^2)^{-1/2} \ll 1$ and the static approximation is applicable. In this approximation correlator (14) is replaced by

$$K^{(class; static)}(q) = \frac{2T}{q^2} P^{(tr)}(q); \quad (15)$$

This correlation function describes a static random magnetic field with the correlation function $\langle b(r)b(r^0) \rangle = \langle T \rangle \langle r \rangle^2$. In this case the term $S = 2\hbar \langle \rangle \langle \rangle$ is just the same time flux-flux correlator of the quasistatic random magnetic field, arising from thermal gauge fluctuations. The flux-flux correlator depends on the path and according to (15) is proportional to the area enclosed by the path

$$\langle \rangle \langle \rangle_{classical} = g^2 \frac{T}{A} [];$$

Since the typical area of the trajectory is of order $A[t] = D t$ we find

$$\frac{1}{S} = \frac{g^2 D}{T t}; \quad (16)$$

At this qualitative level, one can identify the dephasing rate as

$$\Gamma(T) = g^2 T; \quad (17)$$

It is possible to calculate the corresponding weakly-temperature dependent coefficient, but this is beyond the scope of the present discussion [according to Wolé, ²⁹ $S = g^2 D = (8\pi) T \ln(\hbar^{-1})$].

There are a few important points, which make dephasing by magnetic field fluctuations drastically different from dephasing by electric field fluctuations: First of all, the physical mechanism involved in the former case is not related to any inelastic scattering and has a purely topological origin. Magnetic field fluctuations are slowed down by fermions and become quasistatic. The fermions see the corresponding random background and this leads to dephasing due to a finite flux through the diusive loops around which the fermions travel. Second, the typical temperature-dependent dephasing rate is proportional to the conductivity $(T) T^{-1} g^2$ [while in the two-dimensional electronic problem, the situation is exactly the opposite, the dephasing rate is proportional to the resistivity $\rho_{Coulomb}(T) T^{-1}$]. We note that if the gauge-field-fermion coupling is not small, then the dephasing rate is much larger than temperature, which invalidates the assumption about only classical fluctuations being important. This observation suggests that modes with frequencies larger than temperature may play an important role.

We now consider quantum fluctuations of the magnetic field. We will show that (i) Quantum fluctuations of magnetic field can also be considered quasistatic at time scales of fermionic diffusion and (ii) Flux-flux correlator due to these quasistatic fluctuations is proportional to the perimeter of the loop.

First, let us calculate the correlation function in real space-time. At zero temperature, this can be done by a straightforward evaluation of the Fourier transform of (11). The result is

$$K(x) = \frac{\hbar}{K^{(1)}(x)} K^{(2)}(x) + K^{(2)}(x) \frac{r \cdot r}{r^2}; \quad (18)$$

where

$$K^{(1)}(x) = \frac{1}{8\pi^2 t} e^{u \cdot Ei(u)} e^{u \cdot Ei(-u)} \quad (19)$$

and

$$K^{(2)}(x) = \frac{1}{4\pi^2 r^2} \left(2 \ln(u) + e^{u \cdot Ei(u)} + e^{u \cdot Ei(-u)} \right); \quad (20)$$

In Eqs. (18), (19), and (20), $x = (t; r)$ is a short-hand notation for the space-time coordinate,

$$u = \frac{r^2}{4(t + \hbar_m^{-1})}; \quad (21)$$

$R_1 = 1;781$ is the Euler's constant, and $Ei(u) = \int_u^\infty (e^x/x)dx$ is the exponential integral function. We have introduced a \high-energy cut-off Λ_m in the definition of the parameter u in Eq. (21). This cut-off is needed to regularize the behavior of the correlation function in the $t \rightarrow 0$ limit. We note that neglecting the term Λ_0^2 in Eq. (11) would lead to a logarithmically divergent equal-time Fourier transform. This divergence is cut-off by the quadratic part Λ_0^2 . We remind that this term comes from the bare action for the gauge field (in the case of the uRVB state in the tJ-model, the constant Λ_0 is of the order of the tunneling amplitude divided by the holon charge gap squared¹⁸). Since the diffusion approximation breaks down at the time-scales smaller than the scattering time τ , we can write the cut-off as

$$\Lambda_m = m \ln \tau = \Lambda_0; \quad (22)$$

We note that results (18), (19), and (20) are exact as long as τ is finite and have logarithmic accuracy if $\tau \rightarrow 0$.

Let us now consider fermions diffusing in the fluctuating background described by Eqs. (18), (19), and (20). Since for a typical fermionic diffusive trajectory $\langle \mathbf{r}(t) | \mathbf{r}(0) \rangle = \frac{D}{t}$, the value of the parameter u [see Eq. (21)] can be estimated as $u = D/(4\tau)^2 \ll 1$. We see that this parameter is large (as long as the system is metallic), and we can use the asymptotic expression for the correlation functions:

$$\lim_{u \rightarrow 1} K(\mathbf{r}; \mathbf{r}) = \frac{1}{2\pi^2 r^2} \ln \left(\frac{\Lambda_m^2 r^2}{4e^2} \frac{r}{r^2} \right) \ln \left(\frac{\Lambda_m^2 r^2}{4} \frac{r}{r^2} \right); \quad (23)$$

where $e = 2;718$ is the base of the natural logarithm. Apparently, Eq. (23) describes a static random magnetic field described by the non-trivial spatial correlation function (23).

Using correlation function (23) one can calculate the ux-ux correlator $h(\mathbf{r})$.

$$h(\mathbf{r}) = g^2 \int \int d\mathbf{r}_1 d\mathbf{r}_2 h_a(\mathbf{r}; \mathbf{r}_1) h_a(\mathbf{r}; \mathbf{r}_2); \quad (24)$$

The exact value of the ux-ux correlator $h(\mathbf{r})$ depends on the geometry of the path (diffusive paths should be averaged over the distances of order mean free path l and viewed as continuous curves). As an example we can consider a circular path S^1 of radius R . We have to calculate integrals of the following type:

$$I_{S^1} = \int_{S^1} \int_{S^1} \frac{d\mathbf{r}_1 d\mathbf{r}_2}{|\mathbf{r}_1 - \mathbf{r}_2|^2};$$

As it stands, the integral is divergent at small distances. If we require that $|\mathbf{r}_1 - \mathbf{r}_2| > r_{min}$, we obtain

$$I_{S^1} = \frac{2\pi R}{r_{min}^2} \ln 2;$$

We see that the leading contribution is proportional to the circumference of the circle, with the second term being a small correction. This result is universal and applies for any loop: Indeed, from Eq. (23) it follows that the main contribution to the correlator always comes from the regions, where points \mathbf{r}_1 and \mathbf{r}_2 are located very close to each other, regardless the form of the trajectory. This is due to the overall factor r^{-2} in Eq. (23). Therefore, for a generic continuous path, the integral is proportional to the perimeter of the path, $P[\gamma]$:

$$I[\gamma] \sim \frac{P[\gamma]}{r_{min}} + \text{geometrical factor} \ln 2; \quad (25)$$

In the context of a diffusive Fermi system, the cut-off distance is of order mean free path $r_{min} = l$, which is the length-scale at which the diffusive description breaks down. Within logarithmic accuracy, we can put $r = l$ in the arguments of the logarithms in Eq. (23). Assuming that $\Lambda_m \gg 1$, we find the ux-ux correlator

$$h(\mathbf{r}) \sim g^2 \frac{\ln^2 \Lambda_m}{1} P[\gamma]; \quad (25)$$

This behavior of the ux-ux correlator can be called a perimeter law. It should be compared with the "area law" behavior of the ux-ux correlator of classical thermal fluctuations (16). The perimeter of a typical fermionic trajectory is $P[\gamma] \sim \sqrt{D/t}$ and therefore the ux-ux correlation function of a trajectory of this size is $h(\mathbf{r}) \sim \frac{g^2 \ln^2 \Lambda_m}{1} \sqrt{D/t}$. It is plausible to assume that these uxes may lead to dephasing with the corresponding time-scale being

$$\tau_1 \sim g^4 \frac{\ln^2 \Lambda_m}{2}; \quad (26)$$

In the usual electronic systems it is known that quantum fluctuations of electric field (i.e., longitudinal modes with $\omega > T$) do not contribute to dephasing in the following sense:²⁷ One can separate different contributions to conductivity by their magnetic field dependence (the external magnetic field is assumed to provide the leading contribution to dephasing with the interaction terms being small corrections). The weak localization term is defined as the leading term in expansion in $1/B$, which depends on the external magnetic field as $1/B^2$. Diagrammatically it is described by the maximally crossed diagram. The perturbative statement about the absence of quantum dephasing is that the first order interaction correction to the weak localization term does not contain contributions from fluctuation modes with $\omega > T$ (or more precisely, this contribution is exponentially small). The corresponding cancellation of the quantum terms is related to the detailed balance in equilibrium. The remaining linear-in- T dephasing term is due to inelastic scattering of electrons; the corresponding rate vanishes at zero temperature.

An interesting (open) question is whether a similar cancellation of "quantum dephasing terms" in the maximally crossed diagram happens in the case of a Fermi gas coupled to a slow fluctuating magnetic field. At the moment, the existence of such terms would not contradict to any known results. As we have seen above, the main contribution to dephasing is not related to any inelastic scattering, and therefore the detailed balance argument is perhaps not relevant here. Second, quantum dephasing expected from the above arguments based on the "perimeter law" (25) should appear only in second order of a perturbation theory and there are no known analytic results in this order. Third, the usual structure of perturbation theory with an external magnetic field providing the leading contribution to the weak localization cut-off is not appropriate in this context, since the main effect, which is based on magnetic field fluctuations, will be smeared out by such a large external field. From the qualitative arguments described in this section, it seems conceivable that quantum fluctuations of magnetic field may still lead to a dephasing effect with the characteristic time-scale given by Eq. (26).

IV. ALTSHULER-ARONOV CORRECTIONS

In this section, we calculate the exchange and Hartree contributions to the self-energy, density of states, and conductivity. We show that the direct leading order perturbation theory is ill-behaved since certain diagrams are infrared divergent. These singular contributions come from the Hartree-type processes. The corresponding corrections to the density of states (39) and conductivity (42) are positive.

A. Interaction corrections to the density of states

In two-dimensional electron liquids, long-range Coulomb interactions produce singular contributions to the density of states and conductivity, which typically have the same sign and order of magnitude that the weak localization correction. These Altshuler-Aronov corrections originate from Coulomb interactions, but in certain regimes (dense electron liquid) do not depend on the strength of these interactions. There is no contradiction here, since in deriving these corrections one assumes that the temperature is smaller than interactions in the following sense: $T = E_F / r_s^2$, with $r_s = (E_F / \epsilon) =$ being the dimensionless conductance and $r_s = e^2 / (\hbar v_F)$. In this parameter regime, the leading correction to the conductivity is universal and negative. This universality is specific to screened Coulomb interactions. In the case of a fermion-gauge system, the interaction-induced corrections are due to gauge fields. The corresponding forces are unscreened, which leads to a different structure of perturbation theory. The existence of corrections of

this type was emphasized by Halperin, Lee, and Read⁸ in the context of the $\nu = 1/2$ quantized Hall state. Also, Khveshchenko³⁰ and Mirlin and Wolf³¹ studied quantum corrections to conductivity, focusing on the exchange (Fock) contributions.

Below we show that the Altshuler-Aronov corrections to the density of states and conductivity are dominated by positive Hartree terms. The corresponding diagrams are found logarithmically divergent. We suggest that the dominance of positive Hartree terms correspond to antilocalization at the one-loop level. The underlying physical argument behind this result is the following: When a fermion travels through a disordered system, it sees a static self-consistent potential created by the other fermions. The fermion scatters off of this static background and this scattering is responsible for Hartree contributions to the density of states and conductivity. We note that the static gauge propagator is proportional to $1/q^2$. This kind of unscreened interaction leads to a divergent single-particle scattering rate (naively, one would expect a power law divergence coming from singular small-angle scattering $\hbar p \rightarrow \hbar p_{\parallel}$), but we show that the divergence is much weaker). Therefore, the effect of scattering of a given fermion off of the static background created by the other fermions lead to singular Hartree terms, which thus dominate the physics. From the theory of electron-electron interactions in disordered conductors, it is known that the Hartree contribution to conductivity is positive for repulsive interactions. In a fermion-gauge system, the magnetic interactions are indeed repulsive and unscreened (at least in the non-compact formulation), and thus the leading order interaction correction to conductivity is expected to be large and positive.

We also note that a similar situation may occur in the vicinity of quantum phase transitions in fermionic systems, where the fermions are coupled to a soft bosonic fluctuation mode, which is in a sense equivalent to a longitudinal singular gauge field. Indeed, the fluctuation propagator in this case is $K_{uc}(\mathbf{q}) /$

$i\omega = (-i\omega + Dq^2) + c^2 / \omega^2$, where c is a constant and ξ is the correlation length. The correlation length diverges at criticality, and the fluctuation mode becomes massless (or unscreened in our language). This should lead to an enhancement of interaction corrections.




FIG. 1: Hartree and Fock contributions to the fermion self-energy. The wavy line corresponds to singular magnetic interactions.

The quantum correction to the fermionic self-energy is described by the diagrams presented in Fig. 1.³² In what

follows we assume a white-noise correlated disorder potential [see Eq. (28)], and therefore the vector vertices are not renormalized by disorder. The basic ingredients needed to calculate the diagrams in Fig. 1 are as follows:¹⁹ The fermionic Matsubara Green's function

$$G_n^{(0)}(p) = \frac{1}{i\omega_p + \text{sgn}\omega = (2)}; \quad (27)$$

where $\omega_p = E(p) - E_F - v_F(p - p^0)$. The impurity propagator (a single dashed line in Fig. 2) reads

$$n(r)u(r^0) = \frac{1}{2} (r - r^0); \quad (28)$$

where $\omega = m = 1$ is the density of states. We note that the impurity lines do not depend on frequency. The particle-




FIG. 2: Diagram series for diffusion; impurity ladder in the particle-hole channel

hole propagator (diffusion) shown in Fig. 2 reads

$$D_{n,n^0}(p; p^0) = \frac{1}{2} + \frac{1}{2} \frac{1}{\omega_j^2} \frac{(i\omega_n)}{\omega_j + D(p - p^0)^2}; \quad (29)$$

where $D = v_F^2 = 2$ is the diffusion coefficient.

The wavy line represents magnetic interactions,

$$K(\omega; Q) = P^{(tr)}(Q) \frac{1}{j\omega + Q^2}; \quad (30)$$

where $\omega = (E_F) = 1$ is the dimensionless conductance and $\omega = \chi_m$ is the diamagnetic susceptibility with $\chi_m = 1$. Finally, the points connecting the wavy lines and the solid fermion lines are vector vertices gv , where $g = 1$ is the dimensionless coupling to the fluctuating gauge field and $v = p/m$ is the fermion velocity.

Using Eqs. (27), (28), (29), and (30), one can calculate the fermionic self-energy to the leading order in the magnetic interaction. The self-energy is the sum of two contributions:

$$\Sigma = \Sigma_H + \Sigma_F;$$

where the first term is the Hartree correction in the diffusion channel (Fig. 1a) and the second term is the Fock correction (Fig. 1b). We do not consider here effects in the Cooper channel (see also Sec. III). The Fock term is relatively small [see Eq. (38) below], since the vector vertices are not renormalized by disorder. Let us study the Hartree term in the diffusion channel. The corresponding self-energy has the form:

$$\Sigma_H(\omega; p) = 2T \sum_{n!}^X \int \frac{d^2q}{(2)^2} G_{n+!}(p+q) D_{n,n+!}^2(p; p+q) B_{n,p}(\omega; q); \quad (31)$$

where 2 is the spin degeneracy factor and B corresponds to the block of four Green's functions (see also Fig. 4c):

$$B_{n,p}(\omega; q) = g^2 \int \frac{d^2p^0}{(2)^2} \frac{d^2p^0}{(2)^2} v^0 v^0 K(\omega; p^0 - p^0) G_n(p^0) G_{n+!}(p^0 + q) G_{n!}(p^0) G_{n+!}(p^0 + q); \quad (32)$$

The crucial point is that the interaction line in this block carries no frequency. The integral above is infrared divergent due to long-range magnetic interactions. At this point, let us introduce a small- Q cut-off Q_{min} as a phenomenological parameter. The value of this parameter can not be determined within the general formulation. We note that we can not replace the magnetic interaction by its average over the Fermi surface, since the Fermi momenta p and p^0 are constrained to be very close to each other due to the dominance of small-angle scattering. Having this in mind, we calculate the B-block (32), which to the leading order is just a constant

$$B = \frac{g^2 v_F^3}{2} L; \quad (33)$$

where $L = \ln(p_F/Q_{min})$ is the logarithmic divergence coming from long-range interactions. Using the self-energy (31) and Eq. (33), we can calculate the density of states,

$$\rho = \frac{2}{\omega} \int \frac{d^2p}{(2)^2} \text{Im} G^2(\omega; p) (i\omega_n - \Sigma; p); \quad (34)$$

where $G(\omega; p)$ and $(i\omega_n - \Sigma; p)$ are the retarded Green's function and self-energy correspondingly. After the analytical continuation in Eq. (34), we obtain the density of states

$$\rho(\omega; T) = \frac{2B}{(2)^2} \text{Im} \int \frac{d^2q}{(2)^2} \frac{1}{(i\omega + Dq^2)^2} \tanh \frac{\omega + !}{2T} + \tanh \frac{!}{2T}; \quad (35)$$

Evaluating the integral at $\omega = 0$ and using the Fermi-gas value of the diamagnetic susceptibility $\chi_m = 1/(2m)$ (the result should be multiplied by χ_m if the susceptibility is different), we obtain the Hartree correction

$$\rho_H(0; T) = \frac{3g^2 L}{2} \ln \frac{1}{T}; \quad (36)$$

where $\omega = m = 1$ and we remind that $L = \int_{Q_{min}}^{R_{PF}} \frac{dQ}{Q} = 1$ is the divergent part coming from long-range magnetic interactions.

The Fock (exchange) correction to the self-energy has the following form:

$$\Sigma_F(\omega; p) = g^2 T \sum_{n!}^X \int \frac{d^2Q}{(2)^2} \frac{p}{m} \frac{p}{m} K(\omega; Q) G_{n+!}(p+Q); \quad (37)$$

It was emphasized earlier that this expression does not contain impurity vertices due to interactions being current-current. It is also clear that this diagram is free of any infrared divergences. The corresponding integrals can be easily calculated and we obtain the following expression for the exchange part of the self-energy (in the Matsubara representation):

$$\Sigma_F(\omega_n) = 4g^2 \omega_n \ln \frac{e}{12 \omega_n} ; \quad (38)$$

where $e = 2.718$ is the base of the natural logarithm [Eq. (38) has better than logarithmic accuracy]. From Eq. (38), it follows that the negative exchange correction to the density of states is at most

$$\Sigma_F(0;T) / g(T) \ln \frac{1}{T}$$

and does not contain the large factor L . Therefore, we conclude that the exchange correction to the density of states is negligible in the low-temperature limit.

A few remarks are in order: In the usual theory of disordered electronic systems it is well known that the Hartree correction is parametrically smaller than the exchange correction in the case of weak long-range interactions. We argue that there is no contradiction between this statement and the above results. In the context of two-dimensional electronic systems, a "long-range interaction" always implies the screened Coulomb interaction $U(0;q) = [q^2(2p_F) + r_s]^{-1}$. In real space, the screened Coulomb interaction falls off as $U_{\text{screened}}(r) = e^2 a_B^2 / r^3$, where a_B is the Bohr's radius. The corresponding Hartree correction is finite, because the interaction between an electron and density fluctuations far from it does not diverge. For the Coulomb interaction, one can not "control" the overall strength and the screening properties independently. The limit of weak screening leads to the small value of the interaction itself. The interaction averaged over the Fermi surface is $\langle U(0;p-p') \rangle = r_s \ln(1/r_s) \approx 1$. Therefore the Hartree terms are small, while the Fock terms remain finite.

In our theory, the fermion-gauge-field coupling g and the "screening length" Q_{min}^{-1} are completely different parameters, and therefore the regime $Q_{\text{min}}^{-1} \gg 1$ does not imply an equally small g . In this limit, the infrared divergence of the Hartree diagram is not compensated by the overall weakness of the interaction.

Summarizing this part, we present the correction to the density of states in leading order perturbation theory:

$$\rho(T) = \frac{g^2 L}{4\pi^3} \ln \frac{1}{T}; \quad (39)$$

where $g \ll 1$ is the fermion-gauge-field coupling, $\chi = m$ with χ being the fermion magnetic susceptibility, and $L = \ln(p_F / Q_{\text{min}})$, with Q_{min}^{-1} being a phenomenological cut-off parameter at this stage. For perturbation theory to be valid, we must require $g^2 L \ln(T) \ll 1$. This is a condition on the coupling g and on temperature.

B. Interaction corrections to conductivity

In a non-interacting system, the conductivity is connected to the density of states via the simple Einstein relation, $\sigma = e^2 D$, where D is the diffusion coefficient. In the presence of interactions, the relation between the density of states and conductivity may be more complicated. In this section, we calculate the Hartree and Fock corrections to conductivity using the usual diagrammatic technique.




FIG. 3: Diagrams for the correction to conductivity. Graphs (a) and (b) are the Hartree contributions; graphs (c) and (d) are the exchange (Fock) contributions.

The Hartree contributions to conductivity are given by diagrams (a) and (b) in Fig. 3.³³ It is possible to reduce the problem of calculating these diagrams to the standard Altshuler-Aronov theory by simply redrawing the graphs as shown in Fig. 4, where the sawtooth line represents the effective interaction (see also Fig. 4c):

$$U(\omega;Q) = \frac{2B(\omega;Q)}{(2\pi)^2} \quad (40)$$

where B is given by Eq. (32) with the fermionic frequencies and momenta set to zero. In leading order perturbation theory, this function can be replaced by a constant (33). Therefore, the effective density-density interaction has essentially zero range, and the usual theory applies. We note that the scheme used here is possible only if one can neglect the contribution coming from the explicit dependence of the gauge interaction on the external frequency ω , which "runs" through the wavy line in Fig. 3b. The corresponding contribution is small if

$$\frac{T}{Q_{\text{min}}^2} \ln \frac{1}{T}$$

or if one is interested in the AC conductivity with frequencies small but finite: $\omega \gg Q_{\text{min}}^2$. Here we assume that one of these conditions applies. In this case, we can use the following formula for the correction to

conductivity

V. CONCLUSION

$$\sigma_H = ie^2 - \frac{1}{2} \int \frac{d^2 Q}{(2\pi)^2} \frac{U D Q^2}{(i + D Q^2)^3}; \quad (41)$$

where the effective interaction U is given by Eq. (40) with $\omega = 0$ and $Q = 0$. We see that the Hartree contribution to conductivity due to the long-range interaction $K(\omega; Q)$ is described by the exchange contribution due to the short-range effective interaction U . Calculating the integral in Eq. (41), we obtain the following expression for the conductivity:




FIG. 4: (a) and (b): Hartree diagrams expressed through the effective interaction U . (c): The effective short-range interaction U [see also Eq. (40)].

$$\sigma_H = \frac{e^2 g^2}{4} \frac{L}{3} \ln \frac{1}{T}; \quad (42)$$

where e is the coupling (charge) to the field with respect to which we are calculating conductivity. Incidentally, we find that the Einstein relation holds for the Hartree contribution to conductivity [see Eq. (39)], $\sigma_H = e^2 \chi_D$.

We now discuss the Fock contribution to the conductivity. It is given by graphs (c) and (d) in Fig. 3. These diagrams were considered in Refs. [30,31]. We rederive the corresponding contribution in Appendix A, confirming the numerical coefficient found in Ref. [31]. The result is

$$\sigma_F = \frac{e^2 g^2}{2} \ln \frac{1}{T}; \quad (43)$$

We see that the Einstein relation does not apply for the Fock contribution. The Fock correction to conductivity is relatively large, even though the Fock correction to the density of states is essentially zero. Nevertheless, the exchange correction (43) is still much smaller than the Hartree term (42) due to the singularity in the factor L .

In this paper we found that in a two-dimensional Fermi system with singular interactions, the dominant quantum corrections to the density of states and conductivity are positive, which suggests that the system remains in a metallic state at low temperatures. By singular interactions we mean long-range interactions $V(\omega; Q)$ for which the integral $L = \int d^2 Q V(0; Q)$ is infrared divergent.

Singular current-current interactions may appear in the context of quantum magnets³ and the quantized Hall effect,⁸ and correspond to the interactions mediated by transverse gauge fields. If the fermion-gauge-field coupling is small (large- N limit), a controllable theory can be developed. In the clean case, the system is in the strong coupling phase discovered in Refs. [12,13,14]. This non-Fermi liquid phase is described by a collection of regions on the Fermi line, which do not interact with each other, unless they are located on opposite sides of the Fermi line. In the latter case, the regions interact strongly, and the strength of these $2p_F$ -interactions is non-universal, in the sense that it depends on the effective gauge coupling. These interactions may lead to instabilities at low temperatures.^{24,26} At high enough temperatures or in the absence of instabilities, the system remains gapless. This gapless non-Fermi liquid phase is unstable against weak disorder. At low but finite temperatures and in the presence of disorder, the fermion-gauge system is described by a diffusive Fermi liquid coupled to a "dissipative" gauge field. This "dissipative" gauge field remains singular (i.e., long-range) in the sense defined above. This transverse gauge field has two important effects on electron transport: (i) First, gauge field fluctuations suppress the weak localization effect by inducing fluxes through closed diffusive trajectories and lead to dephasing topological in origin. At finite temperatures, classical fluctuations of magnetic field lead to a relatively large dephasing rate²⁸ proportional to conductance rather than resistance (c.f., Ref. [17]). It is conceivable that similar quasistatic configurations due to quantum fluctuations of the magnetic field still result in dephasing. The corresponding time-scale (26) should be related to the $\chi_x - \chi_x$ correlator calculated in this paper, which was found to be proportional to the perimeter of the trajectory. (ii) Second, gauge fluctuations result in singular interaction corrections to conductivity. Due to interactions being long-range, the main contributions come from the processes describing scattering of fermions off of the static self-consistent potential created by the other fermions. The corresponding Hartree contribution to conductivity is positive and formally infrared divergent. This strongly suggests that the metallic phase is stable.

We note that somewhat similar singular interactions may appear in electronic systems in the vicinity of quantum critical points. In this case, the role of a singular longitudinal gauge field is played by soft fluctuation modes.³⁴ In the clean case, treatment of these fluctuations

tions is similar to the theory developed in Refs. [12,13,14]. In particular, Eliashberg-type theories developed to describe Fermi systems in the vicinity of quantum critical points^{35,36} are closely related to the strong-coupling theory of Nayak and Wilczek and Altshuler et al. It also seems likely that the non-Fermi liquid phase discovered in the context of gauge theories can be realized in the vicinity of quantum critical points. If the transition happens at zero wave-vector $q_0 = 0$, then a direct mapping exists between the two theories. If the transition happens at a non-zero wave vector³⁵ $q_0 = 2p_F$, then the existence and structure of the non-Fermi liquid phase should depend on whether $z = \arcsin(q_0/2p_F)$ is a rational number or not. If $z = p/q$ is rational, then the system is described by a collection of sets of regions on the Fermi line (q regions in each set), with different sets not interacting with each other, unless the momentum transfer between two sets is exactly two Fermi momenta. These strong-coupling non-Fermi liquid fixed points, if exist, are unstable against weak disorder. In the disordered regime, singular interactions should modify quantum corrections to conductivity. It is possible that in certain cases long-range interactions mediated by soft fluctuations may enhance the Hartree terms compared to the other contributions. In these cases, the competition between negative weak localization and exchange corrections to conductivity and positive Hartree terms would result in a quantum metal-insulator transition.

A serious limitation of the approach used here and in Refs. [12,13,14] is the assumption about a weak gauge-fermion coupling. In the systems of interest, there are no physical reasons for this coupling to be small. We note that it however might be possible to connect the results derived within the large- N approximation to real systems. This can be done by relating the effective coupling strength to observables: In the clean strong coupling phase, the Kohn anomaly in the fermion polarizability can be much stronger^{13,37} than in an ordinary Fermi liquid.³⁸ The corresponding singularity is related to the strength of the gauge-fermion coupling. One can calculate the general form of the dynamic polarizability near $q = 2p_F$,²⁶ which then determines unusual dynamic Friedel oscillations. It is known that in an ordinary Fermi liquid, scattering of electrons off of the dynamic Friedel oscillations leads to a "non-analytic" linear-in- T temperature dependence of thermodynamic properties^{39,40,41,42} and transport properties^{43,44} in the ballistic temperature regime (where "non-analytic" means that these corrections are different from the T^2 -corrections expected in the naive Fermi gas theory). In the fermion-gauge theory, these corrections scale differently with the exponent being related to the effective gauge-fermion coupling.²⁶ Therefore by measuring these non-universal temperature corrections (e.g., corrections to the specific heat, susceptibility, electrical transport, and thermal transport), one in principle can determine the effective gauge coupling. This construction relies on the assumption that the strong-coupling phase survives

at intermediate couplings $g \gg 1$, which is a reasonable conjecture.

Acknowledgments

The author is grateful to Leon Balents, Matthew Fisher, Pavel Kovtun, Anatoly Larkin, Andreas Ludwig, Lesik Motrunich, Gil Refael, and Maxim Vavilov for discussions about this work. This work was supported by the David and Lucile Packard foundation.

APPENDIX A: EXCHANGE CORRECTION TO CONDUCTIVITY

In this Appendix we derive the formula for the exchange correction to conductivity. The leading exchange contribution to the conductivity is coming from diagrams (c) and (d) in Fig. 3. It is not a priori evident that one can use the formulas from the standard Altshuler-Aronov theory of density-density interaction corrections; since in the latter case, the scalar vertices are renormalized by disorder, which produces additional constraints on the frequency summations. We do not have these restrictions. Nevertheless, we show below that the analytical structure of the exchange contributions in our case is essentially identical to the structure of the standard Altshuler-Aronov treatment.

The contributions to the A-eld response tensor from diagrams (c) and (d) are

$$Q^{(c)}(\omega) = e^2 g^2 T \sum_{\mathbf{p}} \frac{d^2 Q}{(2\pi)^2} T_{\omega}(\omega; \mathbf{p}; \mathbf{p} + \mathbf{Q}) K_{\omega}(\omega; \mathbf{Q}) D_{\omega}(\omega; \mathbf{p}; \mathbf{p} + \mathbf{Q}) \quad (A1)$$

and

$$Q^{(d)}(\omega) = e^2 g^2 T \sum_{\mathbf{p}} \frac{d^2 Q}{(2\pi)^2} T_{\omega}(\omega; \mathbf{p}; \mathbf{p} + \mathbf{Q}) K_{\omega}(\omega; \mathbf{Q}) D_{\omega}(\omega; \mathbf{p}; \mathbf{p} + \mathbf{Q}) \quad (A2)$$

where functions T_{ω} correspond to the following block of three fermionic Green's functions

$$T_{\omega}(\omega; \mathbf{p}_1; \mathbf{p}_2) = \frac{d^2 \mathbf{p}}{(2\pi)^2} \frac{\mathbf{p}}{m} \frac{\mathbf{p}}{m} G_{\omega}(\mathbf{p}) G_{\omega}(\mathbf{p} + \mathbf{Q}) G_{\omega}(\mathbf{p} + \mathbf{Q}) \quad (A3)$$

Using the free fermion Green's functions (27), we find:

$$T_{\omega}(\omega; \mathbf{p}_1; \mathbf{p}_2) = 2 \text{sgn}(\omega) \left[\frac{1}{\omega} \left(\frac{1}{\omega} + \frac{1}{\omega} \right) \right] \quad (A4)$$

The last factor is introduced to take into account the constraint on the frequency summation originating from the dispersion (with $\omega = \omega + \omega$). Using Eqs. (A1), (A2), and (A4), we can write the total response tensor due to two

pairs of diagrams of type (c) and (d). This sum has the same analytical structure as the corresponding sum in the usual Altshuler-Aronov theory even though they set of ϵ -functions involved is slightly different.

$$Q^{(c)}(\epsilon) = 2Q^{(c)}(\epsilon) + 2Q^{(d)}(\epsilon) \quad \# \\ = 4DT \sum_{\mathbf{X}^1} F(\epsilon; \mathbf{X}^1) + \sum_{\mathbf{X}^2} F(\epsilon; \mathbf{X}^2); \quad (A5)$$

where

$$F(\epsilon; \mathbf{X}) = \frac{1}{(2\pi)^2} \frac{d^2 Q}{j + |\mathbf{X}| + DQ^2} \frac{1}{j + |\mathbf{X}| + DQ^2} : \quad (A6)$$

Equations (A5) and (A6) were first derived in Ref. [31]

by Mirlin and Wole. Using the standard analytical continuation procedure (see Appendix 4. of Ref. [17] for details), they found the correction to the DC conductivity as

$$\sigma_F = \frac{e^2 g^2 D}{2} \text{Im} \int_0^{\infty} d\epsilon \frac{\epsilon}{\epsilon^2 + D} \coth \frac{\epsilon}{2T} F(0; j, j, \epsilon) : \quad (A7)$$

At low temperatures, this leads to the following result:

$$\sigma_F = \frac{e^2 g^2 \ln 2}{2} \ln \frac{1}{T} : \quad (A8)$$

-
- ¹ P. A. Lee, N. Nagaosa, and X.-G. Wen, cond-mat/0410445.
 - ² X.-G. Wen, Phys. Rev. B 65, 165113 (2002).
 - ³ L. B. Ioffe and A. I. Larkin, Phys. Rev. B 39, 8988 (1989).
 - ⁴ Y. Shimizu, K. Miyagawa, K. Kanoda, M. Maesato, and G. Saito, Phys. Rev. Lett. 91, 107001 (2003).
 - ⁵ Y. Kurosaki, Y. Shimizu, K. Miyagawa, K. Kanoda, and G. Saito, cond-mat/0504273.
 - ⁶ O. I. Motrunich, Phys. Rev. B 72, 045105 (2005).
 - ⁷ S.-S. Lee and P. A. Lee, cond-mat/0502139.
 - ⁸ B. I. Halperin, P. A. Lee, and N. Read, Phys. Rev. B 47, 7312 (1993).
 - ⁹ V. M. Galitski, G. Refael, M. P. A. Fisher, and T. Senthil, Phys. Rev. Lett. 95 077002 (2005)].
 - ¹⁰ M. V. Feigel'man, V. B. Geshkenbein, L. B. Ioffe, and A. I. Larkin, Phys. Rev. B 48, 16641 (1993).
 - ¹¹ M. Y. Reizer, Phys. Rev. B 39, 1602 (1989); *ibid.* 44, 5476 (1991); *ibid.* 40, 11571 (1989).
 - ¹² C. Nayak and F. Wilczek, Nucl. Phys. B 417, 359 (1994).
 - ¹³ B. L. Altshuler, L. B. Ioffe, and A. J. M. ILLIS, Phys. Rev. B 50, 14048 (1994).
 - ¹⁴ J. Polchinski, Nuclear Physics B 422, 617 (1994).
 - ¹⁵ D. Belitz and T. R. Kirkpatrick, Rev. Mod. Phys. 66, 261 (1994).
 - ¹⁶ E. Abrahams, P. W. Anderson, D. C. Licciardello, and T. V. Ramakrishnan, Phys. Rev. Lett. 42, 673 (1979); L. P. Gor'kov, A. I. Larkin, and D. E. Khmel'nitskii, JETP Lett. 30, 228 (1979) [Pis'ma Zh. Eksp. Teor. Fiz. 30, 248 (1979)].
 - ¹⁷ B. L. Altshuler and A. G. Aronov, "Electron-electron interactions in disordered systems," ed. by M. Pollak and A. L. Efros, pp. 1-151, In: Modern Problems in Condensed Matter Physics, North Holland (1985).
 - ¹⁸ N. Nagaosa and P. A. Lee, Phys. Rev. B 61, 9166 (2000).
 - ¹⁹ A. A. Abrikosov, L. P. Gor'kov, and E. D. Zyaloshinskii, "Methods of Quantum Field Theory in Statistical Physics," Englewood Cliffs, NJ: Prentice-Hall, (1963).
 - ²⁰ G. M. Eliashberg, Sov. Phys. JETP 11, 696 (1960) [Zh. Eksp. Teor. Fiz. 38, 966 (1960)].
 - ²¹ P. A. Lee, Phys. Rev. Lett. 63, 680 (1989).
 - ²² Y. B. Kim and P. A. Lee, Phys. Rev. B 54, 2715-2717 (1996).
 - ²³ B. L. Altshuler, L. B. Ioffe, and A. J. M. ILLIS, Phys. Rev. B 52, 5563 (1995).
 - ²⁴ B. L. Altshuler, L. B. Ioffe, A. I. Larkin, and A. J. M. ILLIS, Phys. Rev. B 52, 4607 (1995).
 - ²⁵ W. Kohn and J. M. Luttinger, Phys. Rev. Lett. 15, 524 (1965).
 - ²⁶ V. M. Galitski, to be published.
 - ²⁷ I. L. Aleiner, B. L. Altshuler, and M. E. Gershenson, Waves in Random Media 9, 201 (1999).
 - ²⁸ A. G. Aronov and P. Wole, Phys. Rev. B 50, 16574 (1994).
 - ²⁹ P. Wole, Foundations of Physics 30, 2125 (2000).
 - ³⁰ D. V. Khveshchenko, Phys. Rev. Lett. 77, 362 (1996).
 - ³¹ A. D. Mirlin and P. Wole, Phys. Rev. B 55, 5141 (1997).
 - ³² P. Lee and T. V. Ramakrishnan, Rev. Mod. Phys. 57, 287 (1985).
 - ³³ H. Fukuyama, J. Phys. Soc. Jpn. 48, 2169 (1980); *ibid.* 50, 649 (1981).
 - ³⁴ W. Metzner, D. Rohe, and S. Andergassen, Phys. Rev. Lett. 91, 066402 (2003).
 - ³⁵ A. V. Chubukov, V. M. Galitski, and V. M. Yakovenko, Phys. Rev. Lett. 94, 046404 (2005).
 - ³⁶ A. V. Chubukov, Phys. Rev. B 71, 245123 (2005).
 - ³⁷ L. B. Ioffe and A. J. M. ILLIS, Physics { Uspekhi 41, 595 (1998).
 - ³⁸ F. Stern, Phys. Rev. Lett. 18, 546-548 (1967).
 - ³⁹ V. M. Galitski and S. D. Sarm, Phys. Rev. B 70, 035111 (2004).
 - ⁴⁰ A. V. Chubukov and D. L. Maslov, Phys. Rev. B 68, 155113 (2003).
 - ⁴¹ A. V. Chubukov, D. L. Maslov, S. Gangadharan, and L. I. Glazman, Phys. Rev. B 71, 205112 (2005).
 - ⁴² V. M. Galitski, A. V. Chubukov, and S. D. Sarm, Phys. Rev. B 71, 201302 (2005).
 - ⁴³ G. Zala, B. N. Narozhny, and I. L. Aleiner, Phys. Rev. B 64, 214204 (2001); *ibid.* 65, 180202 (2002).
 - ⁴⁴ S. D. Sarm, Phys. Rev. B 33, 5401 (1986); A. Gold and V. T. Dolgoplov, *ibid.* 33, 1076 (1986); S. Das Sarm and E. H. Hwang, Phys. Rev. Lett. 83, 164 (1999).