

A Large Atom Number Metastable Helium Bose-Einstein Condensate

A. S. Tychkov, T. Jelten, J. M. M. Namara, P. J. J. Tol, N. Herschbach, W. Hogervorst, and W. Vassen
 Laser Centre Vrije Universiteit, De Boelelaan 1081, 1081 HV Amsterdam, The Netherlands
 (dated: December 17, 2021)

We have produced a Bose-Einstein condensate of metastable helium ($^4\text{He}^*$) containing over $1 \cdot 10^7$ atoms, which is more than an order of magnitude higher than previously achieved. The improved starting conditions for evaporative cooling are obtained by applying one-dimensional Doppler cooling inside a magnetic trap. The same technique is successfully used to cool the spin-polarized fermionic isotope (^3He), for which thermalizing collisions are highly suppressed. We detect the atomic cloud applying different techniques: absorption imaging to monitor ballistic expansion and microchannel plate detectors for time-of-flight experiments as well as to measure the ions produced during the formation and decay of the condensate.

Ten years after the first experimental realization of Bose-Einstein condensation (BEC) in dilute atomic gases [1] the field of degenerate quantum gases has developed into a major area of research. For most elements it has not yet been possible to produce Bose-Einstein condensates containing large numbers of atoms. Only for the one-electron atoms hydrogen, sodium, and rubidium condensates with more than 10^6 atoms have been realized [2]. Large condensates provide better signal-to-noise ratio and are especially useful for atom optics and atom laser applications as well as for sympathetic cooling experiments. In this realm metastable atoms are particularly interesting as they provide a high detection efficiency due to their high internal energy. Absorption imaging is the technique most frequently applied to detect and measure BEC. Only in the case of metastable helium (2^3S_1 state) has detection of the condensate been performed using a microchannel plate (MCP) detector [3]. The same detector is also used to measure ions produced by Penning ionization, while absorption imaging is not used. In a recent experiment, after pioneering work with metastable neon [4], Schellekens et al. used a position sensitive MCP detector to observe the Hanbury Brown and Twiss effect both above and below the BEC threshold [5]. In the second experiment in which BEC was realized with He^* [6], optical detection was used and up to $6 \cdot 10^5$ atoms could be condensed. Recently, this group reported a high precision measurement of the metastable helium scattering length a , which was performed using a two-photon dark resonance [7]. The value $a = 7.512 \text{ nm}$ is favorable for experiments with the fermionic isotope ^3He . It ensures a stable ultracold $^4\text{He}/^3\text{He}$ boson-fermion mixture, as the inter-isotope scattering length will be large and positive [8]. Large clouds of ^4He down to BEC provide an efficient reservoir for sympathetic cooling and will facilitate the observation of degenerate ^3He clouds with large numbers of atoms and phase separation.

In this letter we present an experiment that combines the various detection methods used previously [3, 6] and describe the realization of a BEC of ^4He containing more than $1 \cdot 10^7$ atoms. This large improvement is primarily due to the application of one-

dimensional Doppler cooling inside the magnetic trap rather than three-dimensional Doppler cooling prior to magnetic trapping. This leads to lower temperatures and higher densities at the start of the evaporative cooling ramp. Doppler cooling of polarized atoms was originally proposed for atomic hydrogen [9], and has recently been demonstrated for optically dense samples of magnetically trapped chromium atoms [10]. Compared to the schemes we investigated previously [11, 12], this configuration is more efficient and simple. Comprised of a single retro-reflected laser beam, this robust Doppler-cooling scheme allows us to reach a temperature of three times the Doppler limit in the magnetic trap, while practically no atoms are lost in the cooling process.

The experimental setup is an extended and improved version of our previous setup [11]. In short, we start with a beam of metastable atoms produced by a liquid-nitrogen cooled dc-discharge source. The atomic beam is collimated, deflected and slowed applying laser beams resonant with the $2^3S_1 \rightarrow 2^3P_2$ transition at 1083 nm. Typically $2 \cdot 10^9$ atoms are loaded into a magneto-optical trap (MOT) at a temperature of 1 mK. Since our previous experiments [11] we have installed a new ultrahigh vacuum (UHV) chamber and magnetic trap. The coils for our cloverleaf magnetic trap are placed in water-cooled plastic boxes and positioned in re-entrant windows. Inside the UHV chamber two MCP detectors (Hamamatsu F4655) and an rf-coil are mounted. The first MCP detector is positioned 10 cm from the trap center and attracts all positively charged ions produced by two-body collisions: $\text{He}^* + \text{He}^* \rightarrow \text{He}^+ + \text{He}(1^1S) + e$ (or $\text{He}^* + \text{He}^* \rightarrow \text{He}_2^+ + e$). This Penning ionization process is the primary loss mechanism in cold clouds of He^* . A second identical MCP detector with a grounded grid in front is positioned 17 cm below the trap center and detects with high efficiency all neutral He^* atoms that fall upon it. At the MOT temperature of 1 mK this is only a fraction ($\sim 10^{-3}$) of the total number of atoms, due to the small solid angle. However, at decreasing temperatures the detected fraction increases as a result of slower expansion; a condensate released from our magnetic trap falls almost completely on the MCP.




FIG. 1: Time-of-flight signals of ^4He atoms released from the magnetic trap, with and without one-dimensional Doppler cooling. The MOT contained $N = 2.4 \times 10^9$ atoms at a temperature $T = 0.8 \text{ mK}$ in this particular experiment. The apparent signal increase after Doppler cooling is due to the increased fraction of atoms that is detected at lower temperature. The line is a fit assuming a Maxwell-Boltzmann velocity distribution; the number of data points (dots) is reduced by a factor of 50 for clarity.

This detector is mounted on a translation stage and can be displaced horizontally to allow a vertical laser beam to pass through the trap center for absorption imaging. Absorption imaging of the MOT cloud determines the number of atoms in the MOT with an accuracy of about 20% [11]. This is used to calibrate our He^* MCP detector. Thus the time-of-flight (TOF) signal of atoms released from the trap allows extraction of the temperature as well as the number of atoms in the cloud. When the MOT is loaded we switch off all currents and laser beams. In a weak applied magnetic field we spin-polarize the cloud (all atoms into $M = +1$) and switch on the currents of the cloverleaf magnetic trap, which has a trap depth of about 7 mK. Typically we transfer about 60% of the atoms from the MOT into the magnetic trap in this procedure. The small losses are probably due to the 2 ms delay between the magneto-optical and magnetic trapping. This delay is limited by the current rise time of our power supply. We operate the cloverleaf trap at a bias magnetic field $B_0 = 20 \text{ G}$ to suppress excitation of depolarizing transitions in the subsequent one-dimensional Doppler cooling stage.

One-dimensional Doppler cooling starts at the same time as the cloverleaf trap is switched on. It is implemented by retro-reflecting a weak circularly polarized laser beam along the (horizontal) symmetry axis of the magnetic field. During the cooling pulse the axial temperature decreases reducing the axial size and increasing the optical thickness. Cooling in the radial directions relies on the reabsorption of spontaneously emitted red-detuned photons by the optically thick cloud. Other possible energy redistribution mechanisms are collisional thermalization and anharmonic mixing. While

the collision rate increases from 1.5 s^{-1} to 20 s^{-1} during Doppler cooling, anharmonic mixing is negligible in our trap. With a laser detuning of one natural linewidth below the resonance frequency for an atom at rest in the center of the trapping magnetic field and an intensity of $10^{-3} I_{\text{sat}}$ ($I_{\text{sat}} = 0.17 \text{ mW/cm}^2$), optimum cooling is realized in 2 seconds. In a different realisation 10^8 ^3He atoms were loaded into the magnetic trap and cooled using the same technique. For a spin-polarized fermionic isotope s-wave collisions are forbidden, while the contribution of the higher-order partial waves is highly suppressed in this temperature range. We observed a temperature decrease from 1 mK to 0.3 mK. This proves that cooling in three dimensions relies upon the reabsorption of the red-detuned photons scattered by atoms in the cloud.

Figure 1 shows two TOF traces, illustrating the effect of one-dimensional Doppler cooling in our cloverleaf magnetic trap. Fits of a Maxwell-Boltzmann velocity distribution show that we typically trap $N = 1 \times 10^9$ atoms, which are cooled to a temperature $T = 0.13 \text{ mK}$. This implies a temperature reduction by a factor of 8 and an increase in phase-space density by a factor of 30. In previous experiments [11, 12] we applied three-dimensional Doppler cooling or a two-color magneto-optical trap to improve the starting conditions for evaporative cooling. One-dimensional Doppler cooling provides lower temperatures, higher phase-space density and is much easier to implement. The lifetime of the atoms in the magnetic trap is about 3 minutes, limited by collisions with background gas. To compress the cloud we reduce the bias field in the trap center to 3 G in 2.5 seconds, which increases the temperature to 0.2 mK. The parameters of our magnetic trap then are modest: the axial and radial trap frequencies are 47 Hz and 237 Hz respectively. Higher frequencies are possible but are not required to achieve BEC.

We cool the gas by rf-induced evaporative cooling in 12 s to BEC. Shorter ramps, down to 2 s, also produce a condensate, albeit with fewer atoms. Our procedure is similar to previous experiments on evaporative cooling in our group [13]. We apply a single ramp, starting at an rf frequency of 50 MHz. The frequency decreases exponentially to zero but the ramp is terminated at around 8.4 MHz. This end frequency varies over 0.05 MHz as a result of small day-to-day variations in B_0 . The most sensitive method to detect BEC is TOF analysis of the expanding condensate on the He^* MCP detector when the trap is switched off. In contrast to the Orsay experiments [3, 5], all atoms stay in the $M = +1$ sublevel during the trap switch-off. We use the TOF signal to determine the number of atoms in the condensate as well as in the thermal cloud. A typical TOF spectrum is shown in Fig. 2a. This spectrum is obtained in a single shot and is a textbook example of the inverted parabola indicating the presence of a Bose-Einstein condensate on top




FIG. 2: Observation of BEC, (a) on the He⁺ MCP detector, mounted 17 cm below the trap; the condensed atoms are visible on top of a broader thermal cloud (see dashed-dotted line), (b) on a CCD camera, after an expansion time of 19 ms, (c) on an ion detector; the condensate starts to grow at $t = -0.9$ s (at $t = 0$ the trap is switched off).

of a broad thermal distribution. From the thermal distribution we deduce that BEC is typically reached at a temperature of 2 to 3 K. Applying the MCP calibration, the area under the fitted curve determines the number of thermal atoms that have hit the detector. This number is, however, only a small fraction of the total number of thermal atoms: the detector has a sensitive diameter of 1.45 cm, while a thermal cloud at temperatures around 1 K has a diameter of several centimeters. As the condensate has a much smaller size it will fall almost completely on the detector allowing us to measure the number of condensed atoms N_0 . However, the radial size of the condensate at the MCP detector surface is close to its diameter. A small residual magnetic field gradient may therefore already push the atoms so that some miss the MCP. In order to determine the number of condensed atoms accurately we therefore have applied a small mag-

netic field pulse to push the atoms on the MCP.

When the MCP detector is shifted horizontally, we can detect the condensate on a CCD camera. We apply a weak ($I = 10^{-1} I_{\text{sat}}$), 50 μ s long on-resonance laser pulse (at 1083 nm) to image the shadow of the atoms on a cooled CCD camera (Hamamatsu C4880-21). Figure 2b shows a typical absorption image after an expansion time of 19 ms. As expected, the condensate expands non-isotropically while the thermal cloud shows an isotropic expansion. Combining both techniques, expansion can be monitored in all three dimensions of space: the image shows the horizontal xy-plane and the MCP signal the vertical z-direction. The absorption images also provide details on the direction and velocity of the center of mass movement. This information is used to adjust the position of the MCP in the horizontal plane.

In the MOT, due to a two-body loss rate constant of $5 \cdot 10^{-9} \text{ cm}^3/\text{s}$ [11], the atomic cloud generates 10^{10} ions per second. This large rate hampers ion counting at this stage and therefore the ion current is measured at low MCP gain. In a spin-polarized cloud the two-body loss rate is as low as $10^{-14} \text{ cm}^3/\text{s}$ [7] and therefore ion counting is feasible in a magnetic trap. For this purpose we increase the MCP voltage in the magnetic trapping phase and count the ions produced during evaporative cooling. The ion signal increases although the number of trapped atoms decreases. This is due to the increase in density. When BEC sets in, a sharp increase is expected, indicating the formation of a dense cloud in the centre of the trap [3]. This is demonstrated in Fig. 2c, which shows the growth of the condensate as well as its decay. The decay in this experiment was primarily due to collisions with fast but still trapped atoms, as no rf knife was used. We also applied an rf knife in some of our experiments to measure the lifetime of a pure condensate and found a lifetime of a few seconds.

It has been particularly difficult to extract the number of condensed atoms. There are two main reasons for this. First, it is difficult to calibrate the He⁺ MCP detector for a large flux of atoms. When the number of He atoms that hits one channel of the He⁺ MCP detector per unit of time is too large, its response is no longer linear. This occurs for our MCP only when detecting a falling condensate; thermal atoms, both above and below the BEC threshold, spread out over a sufficiently large part of the MCP detector and over a sufficiently long time that linearity is preserved down to the lowest temperatures. A condensate expands to a dimension of 0.15×2 cm in the plane of the detector and 2 cm in vertical direction, covering only a small fraction of the MCP surface in less than 10 ms. We have determined the fitted number of condensed atoms as a function of the voltage over the detector (using the calibration of gain vs. voltage) and found that the extracted number increased for decreasing voltage. Based on these observations we conclude that we condense at least $1 \cdot 10^7$ atoms; probably this number is

a factor of 2 higher. Second, calibration of the number of atoms by absorption imaging could not be realized. The number of atoms is systematically underestimated up to a factor of 10 compared with the MCP integral method. We attribute this discrepancy to a combination of small effects including remnant magnetic fields, lensing, CCD dark current (background) fluctuations and low detection efficiency at 1083 nm.

As an independent check of the calibration of the number of atoms we studied the size of the condensate as a function of expansion time. We examined this expansion both with the He MCP detector, measuring the radial (vertical) expansion after 190 ms, and by absorption imaging, measuring the radial expansion and axial expansion in the horizontal plane. From the measured expansion in three dimensions the release energy is extracted. This release energy equals, if $N_0 \gg 1$, the interaction energy of the condensate in the magnetic trap. From the interaction energy the number of condensed atoms can be calculated, when the trap frequencies and the scattering length are known [14]. When we adopt $a = 7.512 \text{ nm}$ [7], our expansion provides $N_0 = 7.5 \cdot 10^7$. However, this number must be too high as we do not fulfil

the assumption that the trap should be switched off faster than the inverse of the largest trap frequency, i.e. 3 ms. If we assume that extra energy is added during the switch-off of the trap, the extracted value for N_0 is closer to the value deduced from the integrated TOF signal. We have tried to reach the diabatic regime by decompression of the trapped cloud before releasing the atoms. For this purpose we reduced both trap frequencies to 30 Hz and could then switch off the trap fast enough. However, this strongly increased the problem of MCP saturation as the expansion was slowed down considerably. Moreover, the absorption imaging showed a slightly anisotropic expansion of the condensate whereas an isotropic expansion is expected at equal radial and axial trap frequency. It could only be made isotropic by introducing a delay of 2 ms between switching off the axial and radial confinement. It is questionable whether under these conditions the expansion is fully undisturbed. Nevertheless, from the width of the TOF signal on the MCP detector after expansion from the decompressed trap we can deduce $N_0 = (1 \text{ to } 2) \cdot 10^7$, which is consistent with the number obtained from the integrated TOF signal.

To summarize, we have realized a condensate of ^4He with more than $1 \cdot 10^7$ atoms and detected it measuring the ion production rate in situ, observing its ballistic expansion by absorption imaging and by recording the time-of-flight signal on an MCP detector. The main ingredient that made this large atom number possible is one-dimensional Doppler cooling in a magnetic trap. We demonstrated that this technique can also be applied to cool spin-polarized helium fermions, where the Pauli principle forbids s-wave collisions. Combining both isotopes in one setup may open up a new field of quantum degenerate boson-fermion mixtures of metastable atoms.

We thank Jacques Bouma for technical support. This work was supported by the 'Cold Atoms' program of the Dutch Foundation for Fundamental Research on Matter (FOM) and by the European Union (ESF BEC 2000+ / QUDED IS program and 'Cold Quantum Gases' network).

Electronic address: w.vassen@few.vu.nl

- [1] M. H. Anderson et al., *Science* 269, 198 (1995); K. B. Davis et al., *Phys. Rev. Lett.* 75, 3969 (1995).
- [2] E. W. Streed et al., *arXiv: cond-mat/0507348* (2005).
- [3] A. Robert et al., *Science* 292, 461 (2001).
- [4] M. Yashida and F. Shimizu, *Phys. Rev. Lett.* 77, 3090 (1996).
- [5] M. Schellekens et al., *arXiv: cond-mat/0508466* (2005), accepted for publication in *Science*.
- [6] F. Pereira Dos Santos et al., *Phys. Rev. Lett.* 86, 3459 (2001).
- [7] S. Møller et al., *arXiv: cond-mat/0509286* (2005).
- [8] R. J. W. Stas, J. M. M. Châmarra, W. Hogervorst, and W. Vassen, *Phys. Rev. Lett.* 93, 053001 (2004).
- [9] T. H. J. van der Meer et al., *J. Opt. Soc. Am. B* 6, 2235 (1989).
- [10] P. O. Schmidt et al., *J. Opt. Soc. Am. B* 20, 203 (2003).
- [11] N. Herschbach et al., *J. Opt. Soc. Am. B* 20, 203 (2003); P. J. J. Tol et al., *Phys. Rev. A* 60, R761 (1999).
- [12] A. S. Tychkov et al., *Phys. Rev. A* 69, 055401 (2004).
- [13] P. J. J. Tol, W. Hogervorst, and W. Vassen, *Phys. Rev. A* 70, 013404 (2004).
- [14] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, *Rev. Mod. Phys.* 71, 463 (1999); M. J. Holland, D. Jin, M. L. Chiofalo, and J. Cooper, *Phys. Rev. Lett.* 78, 3801 (1997).