

κ -generalization of Stirling approximation and multinomial coefficients

Tatsuaki Wada*

*Department of Electrical and Electronic Engineering,
Ibaraki University, Hitachi, Ibaraki, 316-8511, Japan*

Hiroki Suyari†

*Department of Information and Image Sciences,
Chiba University, Chiba, 263-8522, Japan*

(Dated: October 2, 2018)

Abstract

Stirling approximation of the factorials and multinomial coefficients are generalized based on the one-parameter (κ) deformed functions introduced by Kaniadakis [Phys. Rev. E **66** (2002) 056125]. We have obtained the relation between the κ -generalized multinomial coefficients and the κ -entropy by introducing a new κ -product operation.

PACS numbers: 05.20.-y, 05.90.+m

Keywords: κ -entropy, κ -product, Stirling approximation, multinomial coefficients

*Electronic address: wada@ee.ibaraki.ac.jp

†Electronic address: suyari@faculty.chiba-u.jp, suyari@ieee.org

I. INTRODUCTION

As is well-known in the fields of statistical mechanics, Ludwig Boltzmann clarified the concept of entropy. He considered a macroscopic system consists of a large number of particles. Each particle is assumed to be in one of the energy level E_i ($i = 1, \dots, k$) and the number of particles in the level E_i is n_i . The total number of particles is then $n = \sum_{i=1}^k n_i$ and the total energy of a macrostate is $E = \sum_i n_i E_i$. The number of ways of arranging n particles in k energy levels such that each level has n_1, n_2, \dots, n_k particles are the multinomial coefficients

$$\left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right] \equiv \frac{n!}{n_1! \cdots n_k!}, \quad (1)$$

which is proportional to the probability of the macrostates if all microscopic configurations are assumed to be equally likely. In the thermodynamic limit n increasing to infinity, we consider the relative number $p_i \equiv n_i/n$ as the probability of the particle occupation of a certain energy level E_i . In order to find the most probable macrostate, Eq. (1) is maximized under a certain constraint. The Stirling approximation of the factorials leads to

$$\ln \left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right] \approx n S^{\text{BGS}} \left(\frac{n_1}{n}, \dots, \frac{n_k}{n} \right), \quad (2)$$

where

$$S^{\text{BGS}} \left(\frac{n_1}{n}, \dots, \frac{n_k}{n} \right) \equiv - \sum_{i=1}^k p_i \ln(p_i), \quad (3)$$

is the Boltzmann-Gibbs-Shannon (BGS) entropy.

Quite recently one of the authors (H.S.) has shown one-parameter generalizations of Gauss' law of error [1], Stirling approximation [2], and multinomial coefficients [3] based on the Tsallis q -deformed functions and associated multiplication operation (q -product) [4, 5]. These mathematical structures are quite fundamental for the basis of any generalization of statistical physics as in the standard statistical physics. In particular Ref. [3] has shown that the one-to-one correspondence between the q -multinomial coefficient and Tsallis q -entropy [6, 7, 8], i.e., the one-parameter (q) generalization of Eq. (2).

On the other hand, Kaniadakis [9, 10, 11] has proposed the κ -generalized statistical mechanics based on the κ -deformed function, which is another type of one-parameter deformations for the exponential and logarithmic functions. Based on the κ -deformed functions and

the associated product operation (κ -product) we have already shown the κ -generalization of Gauss' law of error [12].

In this work we show the κ -generalizations of the Stirling approximation and the relation between the κ -multinomial coefficient and κ -entropy. In the next section the κ -factorial is introduced based on the κ -product. Then the κ -generalization of the Stirling approximation is obtained. We see that the naive approach fails in order to relate the κ -multinomial coefficients with the κ -entropy. In order to overcome this difficulty, we introduce a new kind of κ -product in section 3 and show the explicit relation between the κ -multinomial coefficients and the κ -entropy. The final section is devoted to our conclusions.

II. κ -STIRLING APPROXIMATION

Let us begin with the brief review of the κ -factorial and its Stirling approximation. The κ -generalized statistics [9, 10, 11] is based on the κ -entropy

$$S_\kappa \equiv - \sum_i p_i \ln_{\{\kappa\}}(p_i), \quad (4)$$

where κ is a real parameter in the range $(-1, 1)$, and the κ -logarithmic function is defined by

$$\ln_{\{\kappa\}}(x) \equiv \frac{x^\kappa - x^{-\kappa}}{2\kappa}. \quad (5)$$

The inverse function is the κ -exponential function defined by

$$\exp_{\{\kappa\}}(x) \equiv \left[\sqrt{1 + \kappa^2 x^2} + \kappa x \right]^{\frac{1}{\kappa}}. \quad (6)$$

In the limit of $\kappa \rightarrow 0$ the both κ -logarithmic and κ -exponential functions reduce to the standard logarithmic and exponential functions, respectively. We thus see that S_κ reduces to the BGS entropy of Eq. (3) in the limit of $\kappa \rightarrow 0$.

Based on the above κ -deformed functions, the κ -product is defined by

$$\begin{aligned} x \otimes_\kappa y &\equiv \exp_{\{\kappa\}} \left[\ln_{\{\kappa\}}(x) + \ln_{\{\kappa\}}(y) \right] \\ &= \left[\left(\frac{x^\kappa - x^{-\kappa}}{2} \right) + \left(\frac{y^\kappa - y^{-\kappa}}{2} \right) + \sqrt{1 + \left\{ \left(\frac{x^\kappa - x^{-\kappa}}{2} \right) + \left(\frac{y^\kappa - y^{-\kappa}}{2} \right) \right\}^2} \right]^{\frac{1}{\kappa}}, \quad (7) \end{aligned}$$

which reduces to the standard product $x \cdot y$ in the limit of $\kappa \rightarrow 0$. Similarly the κ -division is defined by

$$\begin{aligned} x \otimes_{\kappa} y &\equiv \exp_{\{\kappa\}} [\ln_{\{\kappa\}}(x) - \ln_{\{\kappa\}}(y)] \\ &= \left[\left(\frac{x^{\kappa} - x^{-\kappa}}{2} \right) - \left(\frac{y^{\kappa} - y^{-\kappa}}{2} \right) + \sqrt{1 + \left\{ \left(\frac{x^{\kappa} - x^{-\kappa}}{2} \right) - \left(\frac{y^{\kappa} - y^{-\kappa}}{2} \right) \right\}^2} \right]^{\frac{1}{\kappa}}, \end{aligned} \quad (8)$$

which reduces to the standard division x/y in the limit of $\kappa \rightarrow 0$. By utilizing this κ -product, the κ -factorial $n!_{\kappa}$ with $n \in \mathbb{N}$ is defined by

$$\begin{aligned} n!_{\kappa} &\equiv 1 \otimes_{\kappa} 2 \otimes_{\kappa} \cdots \otimes_{\kappa} n \\ &= \left[\sum_{k=1}^n \left(\frac{k^{\kappa} - k^{-\kappa}}{2} \right) + \sqrt{\left\{ \sum_{k=1}^n \left(\frac{k^{\kappa} - k^{-\kappa}}{2} \right) \right\}^2 + 1} \right]^{\frac{1}{\kappa}} = \exp_{\{\kappa\}} \left[\sum_{k=1}^n \ln_{\{\kappa\}}(k) \right]. \end{aligned} \quad (9)$$

Now we come to the Stirling approximation of the κ -factorials. For sufficient large n , the summation is well approximated with the integral as follows.

$$\ln_{\{\kappa\}}(n!_{\kappa}) = \sum_{k=1}^n \ln_{\{\kappa\}}(k) \approx \int_0^n dx \ln_{\{\kappa\}}(x) = \frac{n^{1+\kappa}}{2\kappa(1+\kappa)} - \frac{n^{1-\kappa}}{2\kappa(1-\kappa)}. \quad (10)$$

Clearly this reduces to the standard Stirling approximation as

$$\lim_{\kappa \rightarrow 0} \ln_{\{\kappa\}}(n!_{\kappa}) \approx \lim_{\kappa \rightarrow 0} \frac{n}{1-\kappa^2} \left\{ \ln_{\{\kappa\}}(n) - \frac{n^{\kappa} + n^{-\kappa}}{2} \right\} = n(\ln n - 1). \quad (11)$$

Next the κ -multinomial coefficient is defined by utilizing the κ -product and κ -division as follows

$$\left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right]_{\kappa} \equiv n!_{\kappa} \otimes_{\kappa} \left(n_1!_{\kappa} \otimes_{\kappa} \cdots \otimes_{\kappa} n_k!_{\kappa} \right), \quad (12)$$

where we assume

$$n = \sum_{i=1}^k n_i. \quad (13)$$

In the limit of $\kappa \rightarrow 0$, Eq. (12) reduces to the standard multinomial coefficient of Eq. (1).

Let us try to relate the κ -multinomial coefficients with the κ -entropy. Taking the κ -logarithm of Eq. (12) and applying the κ -Stirling approximation leads to

$$\ln_{\{\kappa\}} \left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right]_{\kappa} = \ln_{\{\kappa\}}(n!_{\kappa}) - \sum_{i=1}^k \ln_{\{\kappa\}}(n_i!_{\kappa}), \quad (14)$$

we then obtain

$$\ln_{\{\kappa\}} \left[\begin{array}{c} n \\ n_1 \cdots n_k \end{array} \right]_{\kappa} \approx \frac{n^{1+\kappa}}{2\kappa(1+\kappa)} \left\{ 1 - \sum_{i=1}^k \left(\frac{n_i}{n} \right)^{1+\kappa} \right\} + \frac{n^{1-\kappa}}{2\kappa(1-\kappa)} \left\{ \sum_{i=1}^k \left(\frac{n_i}{n} \right)^{1-\kappa} - 1 \right\}. \quad (15)$$

From this relation, we see that the above naive approach fails. Since the r.h.s. consists of the two terms with different factors (one is proportional to $n^{1+\kappa}/(1+\kappa)$ and the other is proportional to $n^{1-\kappa}/(1-\kappa)$), this cannot be proportional to the κ -entropy, which can be written by

$$\begin{aligned} S_{\kappa} \left(\frac{n_1}{n}, \dots, \frac{n_k}{n} \right) &\equiv - \sum_{i=1}^k \left(\frac{n_i}{n} \right) \ln_{\{\kappa\}} \left(\frac{n_i}{n} \right) \\ &= \frac{1}{2\kappa} \left\{ 1 - \sum_{i=1}^k \left(\frac{n_i}{n} \right)^{1+\kappa} \right\} + \frac{1}{2\kappa} \left\{ \sum_{i=1}^k \left(\frac{n_i}{n} \right)^{1-\kappa} - 1 \right\}. \end{aligned} \quad (16)$$

III. INTRODUCING A NEW κ -PRODUCT

In order to overcome the above difficulty, we introduce another κ -generalization of the products based on the following function defined by

$$u_{\{\kappa\}}(x) \equiv \frac{x^{\kappa} + x^{-\kappa}}{2} = \cosh(\kappa \ln(x)). \quad (17)$$

We here call it the κ -generalized unit function since $\lim_{\kappa \rightarrow 0} u_{\{\kappa\}}(x) = 1$. The basic properties of $u_{\{\kappa\}}(x)$ are as follows.

$$u_{\{\kappa\}}(x) = u_{\{\kappa\}}\left(\frac{1}{x}\right), \quad (18)$$

$$u_{\{\kappa\}}(x) \geq 1, \text{ because } u_{\{\kappa\}}(x) - 1 = \frac{(x^{\frac{\kappa}{2}} - x^{-\frac{\kappa}{2}})^2}{2} \geq 0. \quad (19)$$

The inverse function of $u_{\{\kappa\}}(x)$ can be defined by

$$u_{\{\kappa\}}^{-1}(x) \equiv \left[\sqrt{x^2 - 1} + x \right]^{\frac{1}{\kappa}}, \quad (x \geq 1). \quad (20)$$

In Ref. [13], the canonical partition function associated with the κ -entropy is obtained in terms of this $u_{\{\kappa\}}$ function. Note that

$$\ln_{\{\kappa\}}(x) = \frac{x^{\kappa} - x^{-\kappa}}{2\kappa} = \frac{1}{\kappa} \sinh(\kappa \ln(x)), \quad (21)$$

the two kinds of the κ -deformed functions ($\ln_{\{\kappa\}}(x)$ and $\exp_{\{\kappa\}}(x)$; $u_{\{\kappa\}}(x)$ and $u_{\{\kappa\}}^{-1}(x)$) are thus associated each other. This can be seen from the following relations

$$\sqrt{1 + \kappa^2 \ln_{\{\kappa\}}^2(x)} = u_{\{\kappa\}}(x), \quad (22)$$

$$u_{\{\kappa\}}^{-1}\left(\sqrt{1 + \kappa^2 x^2}\right) = \exp_{\{\kappa\}}(x), \quad \text{for } x \geq 0. \quad (23)$$

Now by utilizing these functions, a new κ -product is defined by

$$\begin{aligned} x \odot_{\kappa} y &\equiv u_{\{\kappa\}}^{-1}\left[u_{\{\kappa\}}(x) + u_{\{\kappa\}}(y)\right] \\ &= \left[\left(\frac{x^{\kappa} + x^{-\kappa}}{2}\right) + \left(\frac{y^{\kappa} + y^{-\kappa}}{2}\right) + \sqrt{\left(\frac{x^{\kappa} + x^{-\kappa}}{2} + \frac{y^{\kappa} + y^{-\kappa}}{2}\right)^2 - 1} \right]^{\frac{1}{\kappa}} \end{aligned} \quad (24)$$

Similarly the corresponding κ -division is defined by

$$\begin{aligned} x \oplus_{\kappa} y &\equiv u_{\{\kappa\}}^{-1}\left[u_{\{\kappa\}}(x) - u_{\{\kappa\}}(y)\right] \\ &= \left[\left(\frac{x^{\kappa} + x^{-\kappa}}{2}\right) - \left(\frac{y^{\kappa} + y^{-\kappa}}{2}\right) + \sqrt{\left(\frac{x^{\kappa} + x^{-\kappa}}{2} - \frac{y^{\kappa} + y^{-\kappa}}{2}\right)^2 - 1} \right]^{\frac{1}{\kappa}}, \end{aligned} \quad (25)$$

where $\left(\frac{x^{\kappa} + x^{-\kappa}}{2}\right) - \left(\frac{y^{\kappa} + y^{-\kappa}}{2}\right) \geq 1$.

Note that $\lim_{\kappa \rightarrow 0} x \odot_{\kappa} y = \infty$, consequently there is no corresponding operation in the standard case of $\kappa = 0$. However unless $\kappa = 0$ this new product satisfies

$$x \odot_{\kappa} y = y \odot_{\kappa} x, \quad \text{comutativity} \quad (26)$$

$$(x \odot_{\kappa} y) \odot_{\kappa} z = x \odot_{\kappa} (y \odot_{\kappa} z). \quad \text{associativity} \quad (27)$$

There exists no unit element within a real number. Therefore real numbers and this product consist a semigroup!

Note also that $x \oplus_{\kappa} x \neq 1$ because $u_{\{\kappa\}}^{-1}(0)$ does not exist by the definition Eq. (20). However we see that the following identity holds.

$$x \odot_{\kappa} y \oplus_{\kappa} x = y, \quad (\kappa \neq 0). \quad (28)$$

Using the new κ -product, the associated κ -factorial can be introduced as

$$\begin{aligned} n!^{\kappa} &\equiv 1 \odot_{\kappa} 2 \odot_{\kappa} \cdots \odot_{\kappa} n \\ &= \left[\sum_{k=1}^n \left(\frac{k^{\kappa} + k^{-\kappa}}{2}\right) + \sqrt{\left\{ \sum_{k=1}^n \left(\frac{k^{\kappa} + k^{-\kappa}}{2}\right) \right\}^2 - 1} \right]^{\frac{1}{\kappa}}. \end{aligned} \quad (29)$$

Similar to Eq. (10), the κ -Stirling approximation can be obtained as

$$\begin{aligned} u_{\{\kappa\}}(n!^\kappa) &= \sum_{k=1}^n u_{\{\kappa\}}(k) \approx \int_0^n dx u_{\{\kappa\}}(x) \\ &= \frac{n^{1+\kappa}}{2(1+\kappa)} + \frac{n^{1-\kappa}}{2(1-\kappa)}. \end{aligned} \quad (30)$$

Also the corresponding κ -multinomial coefficient can be defined by

$$\left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right]^\kappa \equiv n!^\kappa \underset{\kappa}{\oplus} \left(n_1!^\kappa \underset{\kappa}{\odot} \cdots \underset{\kappa}{\odot} n_k!^\kappa \right). \quad (31)$$

Applying the above Stirling approximation to Eq. (31) we obtain

$$u_{\{\kappa\}} \left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right]^\kappa \approx \frac{n^{1+\kappa}}{2(1+\kappa)} \left\{ 1 - \sum_{i=1}^k \left(\frac{n_i}{n} \right)^{1+\kappa} \right\} - \frac{n^{1-\kappa}}{2(1-\kappa)} \left\{ \sum_{i=1}^k \left(\frac{n_i}{n} \right)^{1-\kappa} - 1 \right\}. \quad (32)$$

This is the complementary relation to Eq. (15), and by combining Eqs (15) and (32) we have

$$\ln_{\{\kappa\}} \left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right]_\kappa \pm \frac{1}{\kappa} u_{\{\kappa\}} \left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right]^\kappa \approx \pm \frac{n^{1\pm\kappa}}{\kappa(1\pm\kappa)} \left\{ 1 - \sum_{i=1}^k \left(\frac{n_i}{n} \right)^{1\pm\kappa} \right\}, \quad (33)$$

We thus obtain the final result

$$\begin{aligned} &\left(\frac{1-\kappa}{2n^{1-\kappa}} \right) \left(\ln_{\{\kappa\}} \left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right]_\kappa - \frac{1}{\kappa} u_{\{\kappa\}} \left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right]^\kappa \right) \\ &\quad + \left(\frac{1+\kappa}{2n^{1+\kappa}} \right) \left(\ln_{\{\kappa\}} \left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right]_\kappa + \frac{1}{\kappa} u_{\{\kappa\}} \left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right]^\kappa \right) \\ &= \frac{1}{2\kappa} \left\{ 1 - \sum_{i=1}^k \left(\frac{n_i}{n} \right)^{1+\kappa} \right\} + \frac{1}{2\kappa} \left\{ \sum_{i=1}^k \left(\frac{n_i}{n} \right)^{1-\kappa} - 1 \right\} = S_\kappa \left(\frac{n_1}{n}, \dots, \frac{n_k}{n} \right). \end{aligned} \quad (34)$$

Note that since

$$\lim_{\kappa \rightarrow 0} \frac{1}{\kappa} u_{\{\kappa\}} \left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right]^\kappa = 0, \quad (35)$$

as shown in the Appendix, Eq. (34) reduces to the standard case Eq. (2) in the limit of $\kappa \rightarrow 0$.

IV. CONCLUSION

We have generalized Stirling approximation of the factorials and multinomial coefficients based on the κ -deformed function introduced by Kaniadakis, which is different one-parameter generalization from the q -deformed functions by Tsallis. In order to relate the κ -generalized multinomial coefficients to the κ -entropy, we showed the naive approach, which is similar to that [3] for the q -generalization, failed. In order to overcome this difficulty, we have introduced a new kind of the κ -product operations, which never exist in the standard case of $\kappa = 0$, and have obtained the relation between the κ -generalized multinomial coefficients and the κ -entropy.

The final result Eq. (34) clearly states that the maximizing κ -entropy is equivalent to the maximizing the l.h.s. of Eq. (34) as same as in the standard case of Eq. (2). A next step in a future work is thus to clarify what microscopic states are described by the both κ -multinomial coefficients Eqs. (12) and (31).

V. APPENDIX

We here show the proof of Eq. (35).

$$\frac{1}{\kappa} u_{\{\kappa\}} \left[\begin{matrix} n \\ n_1 \cdots n_k \end{matrix} \right]^\kappa = \frac{1}{\kappa} \left(\sum_{\ell=1}^n u_{\{\kappa\}}(\ell) - \sum_{i=1}^k \sum_{j=1}^{n_i} u_{\{\kappa\}}(j) \right), \quad (36)$$

In the limit of $\kappa \rightarrow 0$ the numerator reduces to $n - \sum_{i=1}^k n_i$ since $\lim_{\kappa \rightarrow 0} u_{\{\kappa\}} = 1$. Consequently both the denominator κ and the numerator become null. Then applying l'Hopital's rule and using the relation

$$\frac{d}{d\kappa} u_{\{\kappa\}}(x) = \ln x \left(\frac{x^\kappa - x^{-\kappa}}{2} \right), \quad (37)$$

we obtain Eq. (35).

-
- [1] H. Suyari and M. Tsukada, IEEE Trans. Inform. Theory, **51** (2005) 753-757.
 - [2] H. Suyari, "q-Stirling's formula in Tsallis statistics", arXiv: cond-mat/0401541.
 - [3] H. Suyari, "Mathematical structure derived from the q-multinomial coefficient in Tsallis statistics", arXiv: cond-mat/0401546.

- [4] L. Nivanen, A. Le Mehaute, and Q. A. Wang, Rep. Math. Phys., **52** (2003) 437-444.
- [5] E. P. Borges, Physica A, **340** (2004) 95-101.
- [6] C. Tsallis, Possible generalization of Boltzmann-Gibbs statistics, J. Stat. Phys. **52** (1988) 479-487.
- [7] C. Tsallis, R.S. Mendes, A.R. Plastino, Physica A **261** (1998) 534-554.
- [8] M. Gell-Mann, C. Tsallis, *Nonextensive Entropy: Interdisciplinary Applications* (Oxford University Press, Oxford 2004).
- [9] G. Kaniadakis and A.M. Scarfone, Physica A, **305** (2002) 69.
- [10] G. Kaniadakis, Phys. Rev. E **66** (2002) 056125.
- [11] G. Kaniadakis, Phys. Rev. E **72** (2005) 036108.
- [12] T. Wada and H. Suyari, “ κ -generalization of Gauss’ law of error”, Phys. Lett. A (2005) in press.
- [13] A.M. Scarfone and T. Wada, “Canonical partition function for anomalous systems described by the κ -entropy”, arXiv: cond-mat/0509364