

The generalized equation of motion for the orbital dynamics in the presence of current

Chyh-Hong Chen¹ and Naoto Nagaosa^{1,2,3y}¹ERATO-SSS, Department of Applied Physics, University of Tokyo, Bunkyo-ku, Tokyo 113-8656, Japan.²Correlated Electron Research Center (CERC), National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba Central 3, Tsukuba 305-8562, Japan⁴CREST, Japan Science and Technology Corporation (JST), Saitama, 332-0012, Japan

The orbital dynamics induced by the charge current is studied theoretically. The equation of motion for the isospin vector T^A in the $SU(N)$ case is derived in the presence of the current, and is applied to the cases of e_g ($N=2$) and t_{2g} ($N=3$) orbitals. In spite of the anisotropic transfer integrals between orbitals, the dynamics is found to be isotropic for e_g -orbitals similarly to the spin case, while it is anisotropic for t_{2g} -orbitals. The implication of this result to the current driven orbital domain wall motion is discussed.

I. INTRODUCTION

The dynamics of the spins in the presence of the current is an issue of recent intensive interests^{1,2,3,4,5,6,7,8,9,10,11,12,13}. Especially it has been predicted theoretically^{3,4,5,10,11,12,13} and experimentally confirmed that the magnetic domain wall (DW) motion is driven by the spin polarized current in the metallic ferromagnets⁸ and also magnetic semiconductors⁹. The basic mechanism of this current driven dynamics is the spin torque due to the current, which is shown to be related to the Berry phase⁵. In addition to the spin, there is another internal degree of freedom, i.e., orbital in the strongly correlated electronic systems. Therefore it is natural and interesting to ask what is the current driven dynamics of the orbital, which we shall address in this paper. In the transition metal oxides, there often occurs the orbital ordering concomitant with the spin ordering. Especially it is related to the colossal magnetoresistance (CMR)¹⁴ in manganese oxides. One can control this ordering by magnetic/electric field, and/or the light radiation. Even the current driven spin/orbital order melting has been observed. On the other hand, an orbital liquid phase has been proposed for the ferromagnetic metallic state of manganites¹⁵. The effect of the current on this orbital liquid is also an interesting issue.

There are a few essential differences between spin and orbital. For the doubly degenerate e_g orbitals, the $SU(2)$ pseudospin can be defined analogous to the spin. However, the rotational symmetry is usually broken in this pseudospin space since the electronic transfer integral depends on the pair of the orbitals before and after the hopping, and also the spatial anisotropy affects as the pseudo-magnetic field. For t_{2g} system, there are three degenerate orbitals, and hence we should define the $SU(3)$ Gell-Mann matrices to represent its orbital state. There is also an anisotropy in this 8 dimensional order parameter due to the same reason as mentioned above.

In this paper, we derive the generic equation of motion of the $SU(N)$ internal degrees of freedom in the presence of the orbital current. $N=2$ corresponds to the e_g orbitals and $N=3$ to the t_{2g} orbitals. Especially, the anisotropy

of the DW dynamics is addressed. Surprisingly, there is no anisotropy for the e_g case while there is for the t_{2g} case. Based on this equation of motion, we study the DW dynamics between different orbital orderings. The effect of the current on the orbital liquid is also mentioned.

II. CP^{N-1} FORMALISM

In this section, we are only interested in the orbital dynamics. To be general, we consider a system with N electronic orbital degeneracy. The model that we investigate is very similar to the lattice CP^{N-1} sigma model with anisotropic coupling between the nearest-neighbor spinors. In contrary to the prediction in the band theory, the system is an insulator when it is 1/N filled, namely one electron per unit cell. Because of the strong on-site repulsive interaction, the double occupancy is not allowed. Therefore, it costs high energy for electrons to move, and the spin degrees of freedom is quenched to form some spin ordering. Due to the complicated spin, orbital, and charge interplay scheme, it is convenient to use the slave-fermion method in which we express the electron as $d_i = h_i^y z_i^{(t)} z_i^{(s)}$ where h_i^y , $z_i^{(s)}$, $z_i^{(t)}$ are baptized as holon, spinon, and pseudo spinon (for the orbital) respectively, and the index i denotes the position¹⁵. If holes are introduced into the system, their mobility leads to frustrate the spin ordering and thus leads to a new phase which might possess finite conductivity. Therefore, we consider the following effective Lagrangian

$$L = i \sum_i \left(1 - h_i h_i \right) z_i z_i + \sum_{\langle ij \rangle} \left(t_{ij} h_i h_j z_i z_j + c.c. \right) \quad (1)$$

where z is short for the spinor $z^{(t)}$. The t_{ij} in Eq.(1) is the transfer integral which is in general anisotropic because of the symmetry of the orbitals. Therefore, the system does not have $SU(N)$ symmetry in general. To introduce the current, we consider the following mean field:

$$\langle h_i h_j \rangle = x e^{i \phi_{ij}} \quad (2)$$

where x denotes the doping concentration, and ϕ_{ij} is the bond current with the relation $\phi_{ij} = -\phi_{ji}$. Then, the

Lagrangian can be written as

$$L = i\sum_i \tilde{z}_i^\dagger \dot{z}_i + \sum_{\langle ij \rangle} (t_{ij} x e^{i\phi_{ij}} z_i z_j^\dagger + c.c.) \quad (3)$$

Note that the constraint $\sum_i \tilde{z}_i^\dagger \tilde{z}_i = 1$ is imposed, and the lowest energy state within this constraint is realized in the ground state. The most common state is the orbital ordered state, which is described as the Bose condensation of z , i.e., $\langle z_i \rangle \neq 0$. In the present language, it corresponds to the gauge symmetry breaking. On the other hand, when the quantum and/or thermal fluctuation is enhanced by frustration etc., the orbital could remain disordered, i.e., the orbital liquid state¹⁵. Then the Lagrangian Eq.(3) describes the liquid state without the gauge symmetry breaking. In this case, the gauge transformation

$$\begin{aligned} z_i &\rightarrow e^{i\phi_i} z_i \\ \tilde{z}_i &\rightarrow e^{i\phi_i} \tilde{z}_i \end{aligned} \quad (4)$$

is allowed. Given $\phi_{ij} = \phi_i - \phi_j$, where $\phi_{ij} = \phi_i - \phi_j$, the local gauge transformation in Eq.(4) with $\phi_i = \phi_j$ corresponding to the simple shifts of the momentum from k to $k + \phi_j$. Therefore, the presence of current does not affect the state significantly since the effect is canceled by the gauge transformation. On the other hand, if z_i represent an orbital ordering, the current couples to the first order derivative of z in the continuum limit. Define $a = iz\partial z$, the second term in Eq.(3) can be written as $\sum_j a$ in the continuum limit. Namely, the current couples to the Berry's phase connection induced by the electron hopping. Therefore, we expect some non-trivial effect similar to the spin case.

To derive the equation of motion for the orbital moments, we use the SU(N) formalism. Introducing the SU(N) structure factors

$$[A; B] = if_{ABC} C \quad (5)$$

$$f[A; B]g = d_{ABC} C + g_A{}^{AB} \quad (6)$$

where A are the general SU(N) Gell-Mann matrices, $[]$ are the commutators, and f, g are the anti-commutators. The f_{ABC} is a totally-anti-symmetric tensor, and d_{ABC} is a totally-symmetric tensor. Let us express t_{ij} in this basis

$$t_{ij} = t_{ij}^0 1 + t_{ij}^A A \quad (7)$$

We will only consider the nearest-neighbor hopping: $t_{ij}^A = t_{\langle ij \rangle}^A$. In the rest of the paper, $t_{i,i+\hat{k}}^A$ will be written as t_k^A , so does \tilde{z}_{ij} . Define the CP^{N-1} superspin vector as

$$T^A(i) = z_i^\dagger A z_i \quad (8)$$

The equation of motion of $T^A(i)$ given by Eq.(3) can be obtained as¹⁶.

$$\begin{aligned} \dot{T}^A(i) = & \frac{xa}{(1-x)\sim} [2\cos k f_{ABC} t_k^B T^C(i) \\ & 2\sin k t_k^0 T^A(i) - \sin k t_k^B d_{ABC} T^C(i) \\ & + f_{ABC} t_k^B \sin k \frac{[k]}{C}(i)] \end{aligned} \quad (9)$$

where the dummy k is summed over x, y , and z direction, a is the lattice constant, and the orbital current $t_c(i)$ is given by

$$t_c(i) = i(\tilde{z}_i^\dagger C z_i - z_i^\dagger C \tilde{z}_i); \quad (10)$$

which is the second order in \tilde{z} . Up to the first order in \tilde{z} , the first term in the right hand side of the Eq.(9) is zero provided that

$$t_x^A + t_y^A + t_z^A = 0 \quad (11)$$

which is true for most of the systems that we are interested in. Consequently, the dominant terms in Eq.(9) will be the second and the third ones, which can be simplified as

$$\begin{aligned} \dot{T}^A(i) = & \frac{xa}{(1-x)\sim} [2\sin k t_x^0 T^A(i) \\ & + \sin k t_k^B d_{ABC} T^C(i)]; \end{aligned} \quad (12)$$

which is one of the main results in this paper. Using Eq.(12), we will discuss the orbital DW motion in the e_g and the t_{2g} systems.

Here some remarks are in order on the mean field approximation Eq.(2) for the Lagrangian Eq.(1). First, it is noted that the generalized Landau-Lifshitz equation obtained by Bazaliy et. al⁵ can be reproduced in the present mean field treatment when applied to the spin problem. As is known, however, there are two mechanisms of current-induced domain wall motion in ferromagnets¹¹. One is the transfer of the spin torque and the other is the momentum transfer. The latter is due to the backward scattering of the electrons by the domain wall. Our present mean field treatment and that in Bazaliy's paper⁵ take the former spin torque effect correctly, while the latter momentum transfer effect is dropped, since the scattering of electrons is not taken into account. However, the latter effect is usually small because the width of the domain wall is thicker than the lattice constant, and we can safely neglect it.

III. $N = 2, e_g$ SYSTEM

First, we consider the application on the (La,Sr)MnO₃ (113 or 214) system¹⁴. Without losing the generality, we show the result on the 113 system. The other one can be obtained in a similar way.

In LaMnO₃, the valence of Mn ion is Mn^{3+} with the electronic configuration $(t_{2g})^3(e_g)^1$. By doping with Sr, one hole is introduced to Mn^{3+} and make it to be Mn^{4+} . The transfer integral between Mn ions depends on the Mn 3d and O 2p orbitals. After integrating over the oxygen p orbitals, the effective hopping between the Mn d orbitals can be obtained. If we denote the up state as $d_{3z^2-r^2}$

and the down state as $d_{x^2-y^2}$, t_{ij} have the following form

$$\begin{aligned} t_x &= t_0 \begin{pmatrix} \frac{1}{3} & \frac{p}{4} \\ \frac{p}{4} & \frac{1}{3} \end{pmatrix} = t_0 \left(\frac{1}{2} + \frac{p}{4} x + \frac{1}{4} z \right) \\ t_y &= t_0 \begin{pmatrix} \frac{1}{3} & \frac{p}{4} \\ \frac{p}{4} & \frac{1}{3} \end{pmatrix} = t_0 \left(\frac{1}{2} + \frac{p}{4} x + \frac{1}{4} z \right) \\ t_z &= t_0 \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = t_0 \left(\frac{1}{2} + \frac{1}{2} z \right) \end{aligned} \quad (13)$$

where $\hat{\tau}^i$ are the Pauli matrices. For $N=2$, the pseudospin moment has the $O(3)$ symmetry given by $T^A = z^A z$. The consequent equations of motion is given by

$$\left(\frac{\partial}{\partial t} + \frac{x}{1-x} \frac{a}{x} t_0 \sim \right) T^A(i) = 0 \quad (14)$$

where \sim is $(x; y; z)$, which can be related to the orbital current as $\mathbf{v}_o = \frac{x}{1-x} \frac{a}{x} t_0 \sim$. Taking the continuum limit, Eq.(14) becomes

$$\left(\frac{\partial}{\partial t} - \mathbf{v}_o \cdot \nabla \right) T^i(\mathbf{r}) = 0 \quad (15)$$

which suggests the solution to be the form $T^i(\mathbf{r} + \mathbf{v}_o t)$. The result is similar to spin case. While the spin domain wall moves opposite to the spin current¹³, in our case, the orbital domain wall also moves opposite to the orbital current.

We can estimate the order of magnitude of the critical current to drive the orbital DW. The lattice constant a is about 3Å. The transfer integral constant t_0 is around 2eV in the LSMO system estimated from the photoemission measurement. If we set v around 1m/s, the critical current can be estimated as $e a^3 \sim 6 \times 10^9$ A/m², which is roughly the same as the order of magnitude of that to drive the spin domain wall.

It should be noted that the current only couples to the first order derivative of z . The double exchange term which is given by the second order derivative is not shown in the equation of motion. However, the double exchange term plays a role to stabilize the DW configuration before the current is switched on.

There are two orbitals degenerate to $d_{x^2-y^2}$, which are $d_{y^2-z^2}$ and $d_{z^2-x^2}$. Similarly, $d_{3y^2-r^2}$ and $d_{3x^2-r^2}$ are degenerate to $d_{3z^2-r^2}$. In the Manganite system, $d_{3z^2-r^2}$ and $d_{x^2-y^2}$ may have different energy due to slight structural distortion in the unit cell. Therefore, in most cases, domain walls are the type of those which separate two degenerate domains. For example, let's consider the orbital domain wall separating $3y^2-r^2$ and $3x^2-r^2$. In Fig.1, the domain wall sits at $y=0$, and suppose the current is along the positive x direction. $3y^2-r^2$ and $3x^2-r^2$ orbitals are described by the spinors $(1=2; \frac{1}{3}=2)$ and $(1=2; \frac{1}{3}=2)$ respectively. The configuration given in Fig.1 is described by the spinor field $(\cos(\mathbf{x}_1); \sin(\mathbf{x}_1))$ where $(\mathbf{x}_1) = 2=3(2 \cot^{-1} e^{x_1/w} + 1)$, where w is the width of the domain wall.

Moreover, due to the special properties of $SU(2)$ algebra, the equation of motion is isotropic regardless how anisotropic the transfer integral is. Therefore, the motion of orbital domain wall is undistorted as in the spin case.




FIG. 1: (Color online) An example of the domain structure in the e_g system. The domain boundary lies on the y -axis.

IV. $N=3$, t_{2g} SYSTEM

Let us consider the t_{2g} systems, for example, in the Vanadate or Titanate systems. The t_{2g} orbitals contain three orbits d_{xy} , d_{yz} , and d_{zx} . The hopping integral t_{ij} between the Ti^{3+} sites or the V^{3+} ones is given as

$$\begin{aligned} t_x &= t_0 \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} = t_0 \left(\frac{2}{3} + \frac{1}{2} x + \frac{1}{2} z \right) \\ t_y &= t_0 \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = t_0 \left(\frac{2}{3} + \frac{1}{2} x + \frac{1}{2} z \right) \\ t_z &= t_0 \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} = t_0 \left(\frac{2}{3} + \frac{1}{2} x + \frac{1}{2} z \right) \end{aligned} \quad (16)$$

where $\hat{\tau}^A$ are $SU(3)$ Gell-Mann matrices with the normalization condition $\text{Tr}(\hat{\tau}^A \hat{\tau}^B) = 2 \delta^{AB}$. The super-spin $\hat{\mathbf{T}}^A$ given by $z^A z$ is normalized because $zz = 1$ and $\hat{\mathbf{T}}^A \hat{\mathbf{T}}^A = 2 \frac{2}{3}$.

The equation of motion in this case will be anisotropic because d_{ABC} is non-trivial in the $SU(3)$ case. It is inspiring to work on an example to see how it goes. Let's consider the d_{xy} d_{yz} orbital DW shown in the Fig.2. Because of the orbital symmetry, such DW can only be stabilized along the y direction. Similarly, the d_{xy} d_{zx} DW can only be stabilized along the x direction and so on. Therefore, the effect of current is anisotropic. In the

hand, when the classical fluctuation of the orbital plays the dominant role for the orbital disordering, the short range orbital order can be regarded as the distribution of the domain walls, which shows the translational motion due to the current as discussed in this paper. Note however that the current does not induce any anisotropy in the orbital pseudospin space in the e_g case.

Orbital degrees of freedom is not preserved in the vacuum or the usual metals, where the correlation effect is not important, in sharp contrast to the spin. Therefore the orbital quantum number can not be transmitted along the long distance and the pseudospin valve phenomenon is unlikely in the orbital case.

VI. ACKNOWLEDGEMENT

We are grateful for the stimulating discussions with Y. Tokura, Y. Oginoto, and S. Murakami. CHC thanks the Fellowship from the ERATO Tokura Super Spin Structure project. This work is supported by the NAREGI Grant, Grant-in-Aids from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

APPENDIX A: e_g ORBITALS

There are two class of e_g orbitals, i.e., one is $3x^2 - r^2$, $3y^2 - r^2$, $3z^2 - r^2$, and the other is $x^2 - y^2$, $y^2 - z^2$, and $z^2 - x^2$. They can be described by a two-component spinor. If we use $(1;0)$ to describe $3z^2 - r^2$ and $(0;1)$ for $x^2 - y^2$, the other degenerate orbitals are given by

$$\begin{aligned} 3y^2 - r^2 &: (1=2; \bar{P}_{3=2}) \\ 3x^2 - r^2 &: (1=2; \bar{P}_{3=2}) \\ y^2 - z^2 &: (\bar{P}_{3=2}; 1=2) \\ z^2 - x^2 &: (\bar{P}_{3=2}; 1=2) \end{aligned} \quad (A1)$$

Their pseudospin moments given by the transformation $T^A = \frac{1}{2} \bar{z} z$ are shown as the following

$$\begin{aligned} T_{3x^2 - r^2} &= (\bar{P}_{3=2}; 0; 1=2) \\ T_{3y^2 - r^2} &= (\bar{P}_{3=2}; 0; 1=2) \\ T_{3z^2 - r^2} &= (0; 0; 1) \\ T_{x^2 - y^2} &= (0; 0; 1) \\ T_{y^2 - z^2} &= (\bar{P}_{3=2}; 0; 1=2) \\ T_{z^2 - x^2} &= (\bar{P}_{3=2}; 0; 1=2) \end{aligned} \quad (A2)$$

APPENDIX B: GELL-MANN MATRICES

The Gell-Mann matrices used in the text are given explicitly in the following:

$$\begin{aligned} \lambda_1 &= \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}; \quad \lambda_2 = \frac{1}{2} \begin{pmatrix} 0 & i & 0 \\ -i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}; \\ \lambda_3 &= \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}; \quad \lambda_4 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}; \\ \lambda_5 &= \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}; \quad \lambda_6 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}; \\ \lambda_7 &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & i \\ 0 & i & 0 \end{pmatrix}; \quad \lambda_8 = \frac{1}{3} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}; \end{aligned} \quad (B1)$$

Electronic address: chem@appitu-tokyo.ac.jp

^y Electronic address: nagaosa@appitu-tokyo.ac.jp

¹ M. N. Baibich, J. M. Broto, A. Fert, F. Nguyen Van Dau, F. Petro, P. Etienne, G. Creuzet, A. Friederich, and J. Chazelas, Phys. Rev. Lett. 61, 2472 (1988).

² B. Dieny, V. S. Speriosu, S. M. Etin, S. S. P. Parkin, B. A. Gumey, P. Baumgart, and D. R. Wilhoit, J. Appl. Phys. 69, 4774 (1991).

³ J. C. Slonkowski, J. Magn. Mater. 159, L1 (1996).

⁴ L. Berger, Phys. Rev. B 54, 9353 (1996).

⁵ Ya. B. Bazaliy, B. A. Jones, and S.-C. Zhang, Phys. Rev. B 57, R3213 (1998).

⁶ J. J. Versluijs, M. A. Bari, and M. D. Coey, Phys. Rev. Lett. 87, 026601 (2001).

⁷ D. A. Allwood, Gang Xiong, M. D. Cooke, C. C. Faulkner, D. Atkinson, N. Vemier, and R. P. Cowburn, Science 296, 2003 (2002).

⁸ A. Yamaguchi, T. Ono, S. Nasu, K. Miyake, K. Mibu, T. Shinjo, Phys. Rev. Lett. 92, 077205 (2004).

⁹ M. Yamaguchi, D. Chiba, F. Matsukura, H. Ohno, Nature 428, 539 (2004).

¹⁰ S. Zhang, P. M. Levy, and A. Fert, Phys. Rev. Lett. 88, 236601 (2002).

¹¹ G. Tatara and H. Kohno, Phys. Rev. Lett. 92, 086601 (2004).

¹² S. Zhang, and Z. Li, Phys. Rev. Lett. 93, 127204 (2004).

¹³ S. E. Barnes and S. M. Aekawa, Phys. Rev. Lett. 95, 107204 (2005)

¹⁴ For a review see Y. Tokura and N. Nagaosa, Science 288, 462 (2000).

¹⁵ S. Ishihara, M. Yamaguchi, and N. Nagaosa, Phys. Rev. B 56, 686 (1997).

¹⁶ We also apply the chain rule on the difference operator. It differs from the differential operator by the second order difference. Since the width of the domain wall is much larger than the atomic scale, our analysis applies.