

# Note on cond-mat/0510119: Jarzynski equation for adiabatically stretched rotor

Markus Bier

Max-Planck-Institut für Metallforschung, Heisenbergstraße 3, 70569 Stuttgart, Germany, and  
Institut für Theoretische und Angewandte Physik,  
Universität Stuttgart, Pfaffenwaldring 57, 70569 Stuttgart, Germany  
(Dated: October 10, 2005)

In a recent article (cond-mat/0510119) it has been argued that the Jarzynski equation is violated for adiabatic stretching processes of a three dimensional rotor system. Here we want to show that the reasoning is not correct. Rather, the Jarzynski equation is fulfilled for this adiabatically stretched rotor.

Consider a particle of mass  $m$  which can freely move on a sphere around the origin with temporal varying radius  $r(t)$  changing according to some prescribed protocol. The Hamiltonian in terms of spherical coordinates  $\theta, \phi$  and the conjugate momenta  $p_\theta, p_\phi$  is

$$H(\theta, \phi; p_\theta, p_\phi; t) = \frac{1}{2} \frac{p_\theta^2}{r(t)^2} + \frac{p_\phi^2}{2 \sin^2 \theta} + U(r(t)); \quad (1)$$

where  $U$  is an external potential which depends on the sphere radius  $r(t)$  but not on the coordinates  $\theta$  or  $\phi$ . If the system is in contact with a heat bath of temperature  $T$  and if the sphere radius is constant,  $r(t) = R$ , a simple calculation yields the canonical partition function

$$Z(\theta, \phi; R) = \frac{4\pi R^2}{\Lambda^3} \exp(-\beta U(R)); \quad (2)$$

Therefore, the free energy difference  $F(\theta, \phi; R_1; R_0)$  between the equilibrium states for sphere radii  $R_1$  and  $R_0$  fulfills

$$\begin{aligned} & \exp(-\beta F(\theta, \phi; R_1; R_0)) \\ &= \frac{Z(\theta, \phi; R_1)}{Z(\theta, \phi; R_0)} \\ &= \frac{R_1^2}{R_0^2} \exp(-\beta (U(R_1) - U(R_0))); \end{aligned} \quad (3)$$

which agrees with Ref. [1], Eq. (4).

Now we consider the following stretching process: For  $t < 0$ , let the system be in equilibrium with a heat bath of temperature  $T$  while the sphere radius  $r(t)$  is held fixed at  $R_0$ . At  $t = 0$ , the system is decoupled from the heat bath. In the time interval  $t \in [0; t_s]$ , the sphere radius is changed from  $r(0) = R_0$  to  $r(t_s) = R_1$ . At  $t = t_s$ , the system is again coupled to the heat bath of temperature  $T$  and the system is equilibrated while keeping the sphere radius at  $R_1$ . After equilibration, the free energy has changed by  $F(\theta, \phi; R_1; R_0)$ . On the other hand, by changing the sphere radius from  $R_0$  to  $R_1$ , one has to perform the work  $W$  to the system. In general,  $W$  depends on the protocol  $r$  and the microstate of the system at  $t = 0$ . Remarkably, for a fixed protocol  $r$ , the

Jarzynski equation [2] asserts

$$\langle \exp(-\beta W) \rangle_i = \exp(-\beta F(\theta, \phi; R_1; R_0)); \quad (4)$$

where  $\langle \cdot \rangle_i$  denotes the average over the canonically distributed microstates at  $t = 0$ .

It is argued in Ref. [1] that was " $W = U(R_1) - U(R_0)$ " which, in conjunction with Eq. (3), would violate the Jarzynski equation (4). In the following, we want to show that this expression for the work  $W$  is not correct. Rather the Jarzynski equation (4) is fulfilled for the system under consideration.

From Eq. (1) one deduces the Hamiltonian equations

$$\begin{aligned} \dot{\theta} &= \frac{p_\theta}{r(t)^2} \\ \dot{\phi} &= \frac{p_\phi}{r(t)^2 \sin^2 \theta} \\ \dot{p}_\theta &= -\frac{\cos \theta p_\phi^2}{r(t)^2 \sin^3 \theta} \\ \dot{p}_\phi &= 0; \end{aligned} \quad (5)$$

The time dependence of the kinetic energy

$$T = \frac{1}{2} \frac{p_\theta^2}{r(t)^2} + \frac{p_\phi^2}{2 \sin^2 \theta} \quad (6)$$

under the Hamiltonian dynamics Eq. (5) is given by

$$\begin{aligned} \frac{dT}{dt} &= -\frac{\dot{r}(t)}{r(t)^3} p_\theta^2 + \frac{p_\phi^2}{\sin^3 \theta} \\ &\quad + \frac{1}{r(t)^2} p_\theta p_\phi + \frac{p_\phi p_\theta}{\sin^2 \theta} \frac{\cos \theta p_\phi^2}{\sin^3 \theta} \\ &= -\frac{\dot{r}(t)}{r(t)^3} p_\theta^2 + \frac{p_\phi^2}{\sin^3 \theta} \\ &= -2 \frac{\dot{r}(t)}{r(t)} T; \end{aligned} \quad (7)$$

which yields

$$T(r(t)) = \frac{\text{const}}{r(t)^2}; \quad (8)$$

The work  $W$  performed on the system by stretching the sphere radius of the rotor from  $R_0$  to  $R_1$  is given by

$$\begin{aligned} W &= (T(R_1) + U(R_1)) - (T(R_0) + U(R_0)) \\ &= T(R_0) \left( \frac{R_0^2}{R_1^2} - 1 \right) + U(R_1) - U(R_0). \end{aligned} \quad (9)$$

This expression for the work  $W$  disagrees with that given in Ref. [1].

For the rotor initially in contact with a heat bath of temperature  $T_0$ ,  $T(R_0)$  is a Boltzmann distributed stochastic variable with probability density  $\exp(-T(R_0)/T_0)$ . Therefore, one finds

$$\begin{aligned} & \langle \exp(-W/T_0) \rangle \\ &= \int_0^\infty dT(R_0) \exp(-T(R_0)/T_0) \\ & \quad \exp\left(-T(R_0) \left( \frac{R_0^2}{R_1^2} - 1 \right) - (U(R_1) - U(R_0))\right) \\ &= \exp(-U(R_1)/T_0) \int_0^\infty dT(R_0) \exp\left(-\frac{R_0^2}{R_1^2} T(R_0)\right) \\ &= \exp(-U(R_1)/T_0) \frac{R_1^2}{R_0^2} : \end{aligned} \quad (10)$$

As the rightmost expressions in Eqs. (3) and (10) are identical, the validity of the Jarzynski equation (4) for the adiabatically stretched rotor is proved.

---

Electronic address: bier@uidsm pi-stuttgart.mpg.de

[1] J. Sung, cond-mat/0510119 (2005).

[2] C. Jarzynski, Phys. Rev. Lett. 78, 2690 (1997).