

# Dissociative attachment of an electron to a molecule: kinetic theory

A. Rossani and A.M. Scarfone<sup>y</sup>

Istituto Nazionale di Fisica della Materia (CNR-INFM), Dipartimento di Fisica,  
Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy.

(Dated: March 23, 2022)


Test particles interact with a medium by means of a bimolecular reversible chemical reaction. Two species are assumed to be much more numerous so that they are distributed according to fixed distributions: Maxwellians and Dirac's deltas. Equilibrium and its stability are investigated in the first case. For the second case, a system is constructed, in view of an approximate solution.

PACS numbers: 05.20.Dd; 51.10.+y; 51.50.+w; 82.20.-w

Keywords: Kinetic and transport theory; Boltzmann equation; Chemical reaction

## I. INTRODUCTION

Among the elementary processes of collision between ions and molecules, the dissociative attachment of an electron to a molecule



plays a role in the physics of weakly ionized gases [1, 2]. However, a full kinetic study is lacking. In particular, when the electric field is not vanishing, equilibrium solutions are not available for the distribution functions.

In the present paper we shall assume that particles 2 and 4 are much more numerous and can be treated as a neutral background, while particles 1 and 3 have the same (negative) charge  $e$  and are subjected to a constant electric field  $E$ . (Hereinafter, particles  $P$ ,  $AB$ ,  $A$ , and  $PB$  will be labeled by the subscripts 1, 2, 3, 4. Each of these particles is endowed with mass  $m_i$  and internal energy of chemical link  $E_{i1}$ .)

First of all we recall the full Boltzmann equations for particles 1 and 3. The exact form of the collision integrals is shown. The weak form of the kinetic equations is constructed, in order to investigate conservation laws and equilibrium. Two cases are considered:

(1) particles 2 and 4 are distributed according to Maxwellians without drift velocity. Under this assumption, equilibrium and its stability are investigated when  $E = 0$ . Firstly we show how to construct the equilibrium solutions. Successively, we show the existence of a Lyapunov functional for the present problem. A physical counterpart of these mathematical results is discussed.

(2)  $B$  is heavier than  $P$  and  $A$ . In the limit  $m_B \rightarrow 1$  particles 2 and 4 can be considered to be distributed according to Dirac's deltas [3]. As a consequence, the Boltzmann equations are modified. Moreover, if the electric field is small, we introduce a first order spherical harmonic expansion for the distribution functions of particles 1 and 3. The result is a system of differential-  
nite

difference equations for the problem. By taking advantage of the smallness of  $E$ , a first order solution can be constructed for the current functions.

## II. BOLTZMANN EQUATIONS FOR PARTICLES 1 AND 3

The nonlinear integrodifferential Boltzmann equations governing the evolution of the distribution function for the reacting particles 1 and 3 reads as follows [4]:

$$\frac{\partial}{\partial t} + \mathbf{v} \cdot \frac{\partial}{\partial \mathbf{x}} + e \frac{E}{m_1} \frac{\partial}{\partial v} f_1 = J_1[f] + Q_1[f]; \quad (2)$$

where  $J_i[f]$  and  $Q_i[f]$  are the chemical and elastic collision integrals with  $f = f_1; f_2; f_3; f_4$ .

The chemical collision integrals are given by

$$J_i[f] = \int \int K_i[f] d\mathbf{w} d\mathbf{n}^0; \quad (3)$$

where, for  $i = 1$ , we have

$$K_{1\mathbf{w}}[f] = \int \int \int \mathcal{G}_{12}^{34}(\mathbf{g}; \mathbf{n}^0) \frac{m_1 m_2}{m_3 m_4} f_3(\mathbf{v}_{12}^{34}) f_4(\mathbf{w}_{12}^{34}) f_1(\mathbf{v}) f_2(\mathbf{w}) d\mathbf{v} d\mathbf{w} d\mathbf{n}^0; \quad (4)$$

being  $\chi(x)$  the Heaviside step function whereas, for  $i = 3$ , we have

$$K_{3\mathbf{w}}[f] = \int \int \mathcal{G}_{34}^{12}(\mathbf{g}; \mathbf{n}^0) \frac{m_3 m_4}{m_1 m_2} f_1(\mathbf{v}_{34}^{12}) f_2(\mathbf{w}_{34}^{12}) f_3(\mathbf{v}) f_4(\mathbf{w}) d\mathbf{v} d\mathbf{w} d\mathbf{n}^0; \quad (5)$$

where

$$\mathbf{v}_{12}^{34} = \frac{1}{M} (m_1 \mathbf{v} + m_2 \mathbf{w} + m_4 \mathbf{g}_{12}^{34} n^0); \quad (6)$$

$$\mathbf{w}_{12}^{34} = \frac{1}{M} (m_1 \mathbf{v} + m_2 \mathbf{w} - m_3 \mathbf{g}_{12}^{34} n^0); \quad (7)$$

$$\mathbf{v}_{34}^{12} = \frac{1}{M} (m_3 \mathbf{v} + m_4 \mathbf{w} + m_2 \mathbf{g}_{34}^{12} n^0); \quad (8)$$

$$\mathbf{w}_{34}^{12} = \frac{1}{M} (m_3 \mathbf{v} + m_4 \mathbf{w} - m_1 \mathbf{g}_{34}^{12} n^0); \quad (9)$$

<sup>E</sup>lectronic address: alberto.rossani@polito.it

<sup>y</sup>Electronic address: antonio.scarfone@polito.it

with  $M = m_1 + m_2 = m_3 + m_4$ . In Eqs. (4) and (5) we have introduced the differential collision frequencies of the forward and backward reaction  $g_{12}^{34}(g; n, n^0)$  and  $g_{34}^{12}(g; n, n^0)$ , where  $g = |v - w|$  whilst  $n$  and  $n^0$  are the unit vectors of the relative velocities before and after collision, respectively. We observe that the following microreversibility condition holds

$$(m_1 m_2)^2 g_{12}^{34}(g; n, n^0) = (m_3 m_4)^2 g_{34}^{12}(g; n, n^0) (g^2 - 34); \quad (10)$$

where

$$g_{kl}^{ij} = \frac{r}{m_i m_j} \frac{m_k m_l}{(g^2 - ij)}; \quad (11)$$

with  $ij = 2M - E = m_i m_j$  and  $E = E_3 + E_4 - E_1$ .  $E_2 > 0$  is the molecular heat of reaction.

Differently, the elastic collision integrals are given by

$$Q_i[f] = \int R_i[f] dw dn^0; \quad (12)$$

where

$$R_i[f] = \sum_{i'=2,4} \int_{i'}^{h_i} g_{i'}^{i'}(g; n, n^0) f_i(v_{i'}^{i'}) f_i(w_{i'}^{i'}) f_i(v) f_i(w); \quad (13)$$

and

$$v_{i'}^{i'} = \frac{1}{m_i + m_{i'}} (m_i v + m_{i'} w + m_{i'} g n^0); \quad (14)$$

$$w_{i'}^{i'} = \frac{1}{m_i + m_{i'}} (m_i v + m_{i'} w - m_{i'} g n^0); \quad (15)$$

The weak form of the kinetic equations for  $i=1$  and  $3$  is obtained by multiplication times a pair of sufficiently smooth functions  $\phi_1(v)$  and  $\phi_3(v)$ , respectively, integration over  $v$ , and by summing [5]:

$$\begin{aligned} \int \frac{\partial f_1}{\partial t} \phi_1(v) dv + \int \frac{\partial f_3}{\partial t} \phi_3(v) dv = & \int K_1[f] \phi_1(v) \phi_3(v_{12}^{34}) dv dw dn^0 \\ & + \frac{1}{2} \int (R_1[f] \phi_1(v) \phi_1(v_{12}^{12}) + R_3[f] \phi_3(v) \phi_3(v_{34}^{34})) dv dw dn^0; \quad (16) \end{aligned}$$

Observe that for  $\phi_1(v) R_3(v) = 1$  we get  $dn_1 = dt + dn_3 = dt = 0$ , with  $n_i = \int f_i dv$ , that is the total number of test particles is conserved.

with  $i' = 2, 4$  and  $i = 1, 3$  fixed.

From the weak form of the Boltzmann equation, by setting

### III. CASE (1)

We assume particles 2 and 4 much more numerous, so that they can be treated as an equilibrium background at a fixed temperature  $T$  [6]:

$$f_{i'} = m_{i'}^3 \exp\left(-\frac{E_{i'}}{2} m_{i'} v^2\right); \quad (17)$$

$$\phi_1 = \ln f_1 \exp\left(-\frac{E_2}{2} m_2 v^2\right); \quad (18)$$

$$\phi_3 = \ln f_3 \exp\left(-\frac{E_4}{2} m_4 v^2\right); \quad (19)$$

where  $f_{i'} = f_{i'} m_{i'}^3$ , we obtain

$$\begin{aligned} D = & \int g_{12}^{34}(g; n, n^0) (m_1 m_2)^3 \ln \frac{\tilde{f}_3(v_{12}^{34}) \tilde{f}_4(w_{12}^{34})}{\tilde{f}_1(v) \tilde{f}_2(w)} [\tilde{f}_1(v) \tilde{f}_2(w) - \tilde{f}_3(v_{12}^{34}) \tilde{f}_4(w_{12}^{34})] dv dw dn^0 \\ & + \frac{1}{2} \sum_{i'} \int (m_i m_{i'})^3 g_{i'}^{i'}(g; n, n^0) \ln \frac{\tilde{f}_i(v_{i'}^{i'}) \tilde{f}_{i'}(w_{i'}^{i'})}{\tilde{f}_i(v) \tilde{f}_{i'}(w)} [\tilde{f}_i(v) \tilde{f}_{i'}(w) - \tilde{f}_i(v_{i'}^{i'}) \tilde{f}_{i'}(w_{i'}^{i'})] dv dw dn^0 = 0; \quad (20) \end{aligned}$$

( $i=1$  and  $3$ ;  $i'=2$  and  $4$ ), where  $D$  is the left hand side of (16). Based on these results, by standard methods of

kinetic theory [4], we have

Proposition 1. The equilibrium condition

$$\frac{\partial f_1}{\partial t} = \frac{\partial f_3}{\partial t} = 0; \quad (21)$$

is equivalent to

$$f_1(v_{12}^{i'}) f_3(w_{12}^{i'}) = f_1(v) f_3(w); \quad (22)$$

$$f_3(v_{12}^{34}) f_4(w_{12}^{34}) = f_1(v) f_2(w); \quad (23)$$

with  $i = 1, 3$  and  $i' = 2, 4$

From the first equation we get

$$f_i = \exp \left( -\frac{E_i}{T} - \frac{1}{2} m_i v^2 \right); \quad (24)$$

while the second one gives  $n_1 + n_2 = n_3 + n_4$ :

In order to investigate the stability of such equilibrium solution, we introduce the following functional:

$$L = H + \int_{\mathbb{R}^3} f_1 \left( E_1 + \frac{1}{2} m_1 v^2 \right) dv + \int_{\mathbb{R}^3} f_3 \left( E_3 + \frac{1}{2} m_3 v^2 \right) dv; \quad (25)$$

where  $H = \int_{\mathbb{R}^3} H dv$ , being  $H = H_1 + H_3$  and  $\partial H_i = \partial f_i = \ln f_i$ .

Proposition 2.  $L$  is a Lyapunov functional for the present problem

Proof. First of all, from (16) and (20) we verify that

$$\frac{dL}{dt} = D \leq 0; \quad (26)$$

Moreover, by introducing the first Taylor expansion of  $H$  around the equilibrium

$$\begin{aligned} \hat{H} dv &= \int_{\mathbb{R}^3} H + \frac{\partial H}{\partial f_1} (f_1 - f_1^0) + \frac{\partial H}{\partial f_3} (f_3 - f_3^0) dv \\ &= \int_{\mathbb{R}^3} H + \int_{\mathbb{R}^3} \left( E_1 + \frac{1}{2} m_1 v^2 \right) (f_1 - f_1^0) + \int_{\mathbb{R}^3} \left( E_3 + \frac{1}{2} m_3 v^2 \right) (f_3 - f_3^0) dv; \end{aligned} \quad (27)$$

where  $f_i^0$  means "at equilibrium", from Eqs. (25) and (27) we obtain

$$\begin{aligned} L - L^0 &= \int_{\mathbb{R}^3} H - H^0 dv \\ &+ \int_{\mathbb{R}^3} \left( E_1 + \frac{1}{2} m_1 v^2 \right) (f_1 - f_1^0) dv \\ &+ \int_{\mathbb{R}^3} \left( E_3 + \frac{1}{2} m_3 v^2 \right) (f_3 - f_3^0) dv \\ &= \int_{\mathbb{R}^3} (H - \hat{H}) dv; \end{aligned} \quad (28)$$

Due to the convexity of  $H$  we can conclude that  $L - L^0 \geq 0$ . The inequality

$$\frac{dL}{dt} \leq 0; \quad (29)$$

can be interpreted on a physical ground. In fact, by introducing the entropy  $S = - \int_{\mathbb{R}^3} H dv$ , we get the following thermodynamic inequality:

$$dS \geq \frac{1}{T} (dE - p_1 dn_1 - p_3 dn_3); \quad (30)$$

where  $E = \int_{\mathbb{R}^3} f_1 (E_1 + m_1 v^2/2) dv + \int_{\mathbb{R}^3} f_3 (E_3 + m_3 v^2/2) dv$  is the total energy density of test particles. With respect

to Clausius inequality, we observe an additional term due to the fact that the medium of field particles 2 and 4 not only provides heat to the gas of test particles 1 and 3 but also modifies its composition.

#### IV. CASE (2)

In the limit  $m_B \rightarrow 1$  we have

$$\frac{m_1 m_2}{m_3 m_4} \rightarrow \frac{m_1}{m_3}; \quad (31)$$

and the following relations hold

$$\begin{aligned} g_{12}^{34} &\rightarrow v = \frac{m_1}{m_3} v^2 + v_3; \\ g_{34}^{12} &\rightarrow v^+ = \frac{m_3}{m_1} v^2 + v_1; \\ v_{12}^{34} &\rightarrow w + v n^0; \quad v_{34}^{12} \rightarrow w + v^+ n^0; \\ v_{12}^{i'} &\rightarrow w + v n^0; \quad w_{12}^{i'} \rightarrow w - v n^0; \\ w_{34}^{12} &\rightarrow w; \quad w_{12}^{34} \rightarrow w; \end{aligned} \quad (32)$$

where  $i = 2, 4$  and  $i' = 1, 3$ . Moreover we can pose  $n^0 = n$ ,  $n^0 \rightarrow 0$  and  $g \rightarrow v$ .

By taking into account that  $f_i(w) = N_i(w)$  for  $i = 2$  and  $4$ , the integrals  $J_i[f]$  and  $Q_i[f]$  read now

$$J_1[f] = \int_0^Z (v^2 - 1) \frac{34}{12} (v; 0) \frac{m_1}{m_3} N_4 f_3(v; 0) - N_2 f_1(v) dv; \quad (33)$$

$$J_3[f] = \int_0^Z \frac{m_1}{m_3} \frac{v^+}{v} \frac{34}{12} (v^+; 0) \frac{m_3}{m_1} N_2 f_1(v^+; 0) - N_4 f_3(v) dv; \quad (34)$$

$$Q_1[f] = \int_0^Z N_2 \frac{12}{12} (v; 0) + N_4 \frac{14}{14} (v; 0) f_1(v; 0) - f_1(v) dv; \quad (35)$$

$$Q_3[f] = \int_0^Z N_2 \frac{32}{32} (v; 0) + N_4 \frac{34}{34} (v; 0) f_3(v; 0) - f_4(v) dv; \quad (36)$$

Equilibrium and its stability for the present problem are investigated in [5]. Our purpose here is to construct model equations suitable for an approximate solution. As usual in the physics of weakly ionized gases [1], if both the spatial gradients and the electric field are small we may resort to a first order spherical harmonic expansion of  $f_i(v)$ :

$$f_i(v) = N_i(v) + J_i(v); \quad (37)$$

where

$$N_i(v) = \frac{1}{4} \int_0^Z f_i(v) dv; \quad (38)$$

and

$$J_i(v) = \frac{3}{4} \int_0^Z f_i(v) dv; \quad (39)$$

By projecting over 1 and 3 we get, after some manipulations, the following system for the new unknown functions  $F_i(\tau) = N_i(v)$  and  $G_i(\tau) = J_i(v)$ :

$$\frac{\partial F_1(\tau)}{\partial \tau} + \frac{P}{3} r - G_1(\tau) \frac{eE}{m_1} \frac{2}{3} \frac{\partial}{\partial \tau} [G_1(\tau)] = (v^2 - 1) \frac{34}{12(0)}(\tau) \frac{m_1}{m_3} N_4 F_3 - N_2 F_1(\tau); \quad (40)$$

$$\frac{\partial F_3(\tau)}{\partial \tau} + \frac{P}{3} r - G_3(\tau) \frac{eE}{m_3} \frac{2}{3} \frac{\partial}{\partial \tau} [G_3(\tau)] = \frac{m_1}{m_3} \frac{34}{12(0)} + \frac{m_3}{m_1} N_2 F_1 + N_4 F_3(\tau); \quad (41)$$

$$\frac{\partial G_1(\tau)}{\partial \tau} + \frac{P}{3} r F_1(\tau) - \frac{2eE}{m_1} \frac{P}{3} \frac{\partial F_1(\tau)}{\partial \tau} = (v^2 - 1) \frac{34}{12(1)}(\tau) \frac{m_1}{m_3} N_4 G_3 - N_2 G_1(\tau) \frac{34}{12(0)}(\tau) + G_1(\tau) G_1(\tau); \quad (42)$$

$$\frac{\partial G_3(\tau)}{\partial \tau} + \frac{P}{3} r F_3(\tau) - \frac{2eE}{m_3} \frac{P}{3} \frac{\partial F_3(\tau)}{\partial \tau} = \frac{m_1}{m_3} \frac{34}{12(1)} + \frac{m_3}{m_1} N_2 G_1 + N_4 G_3(\tau) \frac{34}{12(0)} + G_3(\tau) G_3(\tau); \quad (43)$$

where we have posed  $\tau = (v^2 - 1)$  and

$$i(\tau) = N_2 \frac{12}{12(\tau)}(\tau) + N_4 \frac{14}{14(\tau)}(\tau); \quad (44)$$

being  $\lim_{ij(\tau)}(\tau) = \lim_{ij(0)}(\tau) - \lim_{ij(1)}(\tau)$  and

$$\lim_{ij(k)}(\tau) = 2 \int_0^Z + 1 \lim_{ij}(\tau; \tau) d\tau; \quad (45)$$

with  $k = 1$  and  $2$ .

Consider now the stationary space-homogeneous equa-

tions We observe that for  $E = 0$  the following equilibrium solutions hold:

$$F_i = C_i \exp \left( \frac{m_i}{2T} \right); \quad (46)$$

where

$$\frac{C_3 N_4}{C_1 N_2} = \frac{m_3}{m_1}^3 \exp \left( \frac{E}{T} \right); \quad (47)$$

(mass action law). Finally, we observe that the electric

Since  $F_i^{(0)}$  are already known, by solving equations (40) and (41), we can obtain the expression of  $G_i^{(1)}$ , in the case of isotropic reaction collision frequency  $\frac{34}{12(0)}$ :

$$G_1^{(1)} = \frac{eE C_1}{T} \frac{P - \exp \left( \frac{m_1}{2T} \right)}{\left( \frac{1}{1} \right) N_2 \frac{34}{12(0)} \left( \frac{1}{1} \right) + \frac{1}{1} \left( \frac{1}{1} \right)}; \quad (49)$$

$$G_3^{(1)} = \frac{eE C_3}{T} \frac{\exp \left( \frac{m_3}{2T} \right)}{N_4 \frac{m_1}{m_3}^2 \frac{34}{12(0)} (v^+) v^+ + P - \frac{1}{3} \left( \frac{1}{1} \right)}; \quad (50)$$




FIG. 1: Plot of the current function  $\mathcal{G}_1^{(1)}(\xi)$  (full line) for the charge particles P and  $\mathcal{G}_3^{(1)}(\xi)$  (dotted line) for the charge particles A, in arbitrary units.

field must be a small quantity,  $E = \epsilon$ , so that we can expand  $F_i$  and  $G_i$  as follows

$$F_i = F_i^{(0)} + F_i^{(1)} + \dots; \quad G_i = G_i^{(1)} + \dots; \quad (48)$$

where  $G_i^{(1)} = G_i^{(1)} \mathbf{e}$ , with  $\mathbf{e}$  the unit vector of  $E$ . In figure 1 we depict, in arbitrary unity, the plots of  $\mathcal{G}_1^{(1)}(\xi)$  (full line) and  $\mathcal{G}_3^{(1)}(\xi)$  (dotted line). Since the forward equation has a threshold for  $\xi = 1$ , the collision frequency of particles 1 suddenly increases, and a discontinuity in the relevant plot of  $\mathcal{G}_1^{(1)}(\xi)$  occurs.

## V. CONCLUSIONS

Two linear Boltzmann models have been constructed for test particles reacting with a medium of numerous field particles. In the first case the field particles are distributed according to Maxwellians with vanishing drift velocity. Theorems on equilibrium and its stability are given, as well as their connection with thermodynamics. In the second case we consider particles B heavier than particles P and A. By means of first order spherical harmonic expansion, four equations can be constructed, suitable for an approximate solution.

- 
- [1] E.H. Holt and R.E. Haskell, Foundations of Plasma Dynamics, (Macmillan, New York, 1965).
  - [2] B.M. Smirnov, Physics of Weakly Ionized gases (Mir Publishers, Moscow, 1981).
  - [3] C.R. Garibotti and G. Spiga, J. Phys. A: Math. Gen. 27, 2709 (1994).

- [4] A. Rossani and G. Spiga, Phys. A 252, 563 (1999).
- [5] M. Groppi and A. Rossani, AIP Conf. Proc. 585, 567 (2001).
- [6] M. Groppi, A. Rossani, and G. Spiga, Trans. Theor. Stat. Phys. 32, 567 (2003).