

First-principles approach to Non-Collinear Magnetism : towards Spin Dynamics

S. Sharma^{1,2,5}, J. K. Dewhurst^{2,3}, C. Ambrosch-Draef^{2,4}, S. Kurth⁵, N.Helbig^{1,5}, S. Pittalis⁵, S. Shallcross⁶, L. Nordstrom⁷, and E. K. U. Gross⁵¹ Fritz Haber Institute of the Max Planck Society, Faradayweg 4-6, D-14195 Berlin, Germany.² Institut für Physik, Karl-Franzens-Universität Graz, Universitätsplatz 5, A-8010 Graz, Austria.³ School of Chemistry, The University of Edinburgh, Edinburgh EH9 3JJ⁴ Department Materials Physics, University of Leoben, Erzerherzog Johann - Strasse 3, A-8700 Leoben⁵ Institut für Theoretische Physik, Freie Universität Berlin, Am Mühlentempel 14, D-14195 Berlin, Germany.⁶ Department of Physics, Technical University of Denmark DK-2800 Lyngby, Denmark.⁷ Department of Physics, Uppsala University, Box 530, 751 21 Uppsala, Sweden.

(dated: January 28, 2022)

A description of non-collinear magnetism in the framework of spin-density functional theory is presented for the exact exchange energy functional which depends explicitly on two-component spinor orbitals. The equations for the effective Kohn-Sham scalar potential and magnetic field are derived within the optimized effective potential (OEP) framework. With the example of a magnetically frustrated Cr monolayer it is shown that the resulting magnetization density exhibits much more non-collinear structure than standard calculations. Furthermore, a time-dependent generalization of the non-collinear OEP method is well suited for an ab-initio description of spin dynamics. We also show that the magnetic moments of solids Fe, Co and Ni are well reproduced.

PACS numbers: 71.15Mb, 71.10.-w, 71.22.+i

The extension of the original density functional theory (DFT) Hohenberg-Kohn-Sham approach to the case of spin polarized systems was given, under the name spin DFT (SDFT) more than three decades ago [1]. While this formulation was for arbitrary directions of the magnetization vector field, even today most applications are based on a restricted collinear version. This has the advantage of computational simplicity: one then works with two separate KS equations, one yielding the spin-up orbitals the other the spin-down orbitals, whereas the general formulation involves Pauli spinors. Nevertheless, there exists a wealth of non-collinearity in nature. To give only a few examples it is widely seen in molecular magnets, exchange frustrated solids (Fe, spin glasses), and all magnets at finite temperatures.

Crucial for practical calculations using SDFT is the approximation made for the exchange-correlation (xc) energy functional. The Local Spin Density Approximation (LSDA) and the Generalized Gradient Approximations (GGAs) are currently the most popular ones. These have been developed for collinear magnetism, and their use in non-collinear situations relies on the magnetization, $m(\mathbf{r})$, and exchange correlation magnetic field, $B_{xc}(\mathbf{r})$, being made collinear in a local reference frame at each point in space [2]. This is only possible with purely local functionals like LSDA [3, 4], though it has been used under additional approximations for gradient functionals as well [5]. Such approximations (that lead to locally collinear magnetization and xc magnetic field) cause $m(\mathbf{r}) \cdot B_{xc}(\mathbf{r})$ to vanish everywhere in space. As noted recently, this fact renders the adiabatic time dependent extension of these functionals improper [6] for the study

of spin dynamics, because in the absence of external magnetic fields and within adiabatic approximation, the local torque on the spins ($m(\mathbf{r};t) \cdot B_{xc}(\mathbf{r};t)$) vanishes [7]. This is a serious limitation since the dynamics of the spin degree of freedom is responsible for a number of important phenomena such as spin injection, the dynamics of Bloch walls, spin wave excitations [8], and spin filtering, mechanisms crucial for recent developments in spintronics [9]. The search for approximate xc functionals which depend on all three components of the spin magnetization m beyond the form of the locally collinear LSDA has remained a major challenge in the description of non-collinear magnetism.

In recent years, an alternative route to the construction of approximate xc functionals has enjoyed increasing interest. These involve functionals depending explicitly on the single-particle KS orbitals which, through the KS single-particle equation, are implicit functionals of the density [10]. Technically, one needs to employ the Optimized Effective Potential (OEP) [11] method to compute the local xc potential. The simplest orbital-dependent approximation to the xc energy is the EXact EXchange (EXX) functional which is the Fock exchange energy but evaluated with KS orbitals (i.e. orbitals coming from a local potential). A number of successful EXX calculations have been reported for semiconductors [12, 13, 14] and magnetic metals [15]. However, for magnetic systems again the collinear formalism has been employed.

In this Letter we extend the OEP formalism for SDFT to non-collinear magnetic systems. Most importantly, we do not rely on a condition of local collinearity and treat the wavefunctions as Pauli spinors for high lying and

Dirac spinors for deep lying (3 Ha below the Fermi level) electrons. Using the EXX functional, we demonstrate with the example of an unsupported Cr(111) monolayer, that (i) the magnetization and B_{xc} are generally not locally parallel in contrast to what has been assumed in all calculations to date and (ii) that the non-collinearity is much more pronounced than found with the LSDA functional. Against popular belief [16], we find that this non-collinearity is not restricted to just the interstitial region but spreads all the way to the atom center. With the examples of bulk Fe, Co and Ni we further show that our formalism can also be effectively used for collinear magnets.

To derive the OEP equations in the general non-collinear case, we start with the Kohn-Sham (KS) equation

for two-component spinors ψ_i , which has the form of a Pauli equation. For non-interacting electrons moving in an effective scalar potential v_s and a magnetic vector field B_s it reads as (atomic units are used throughout)

$$\frac{1}{2} \nabla^2 \psi_i + v_s(r) \psi_i - B_s(r) \sigma \psi_i = \epsilon_i \psi_i \quad (1)$$

This equation can be derived by minimizing the total energy which, in SDFT, is given as a functional of the density $n(r) = \sum_i^{\text{occ}} |\psi_i(r)|^2$ and the magnetization density $m(r) = \sum_i^{\text{occ}} \psi_i^\dagger(r) \sigma \psi_i(r)$. For a given external scalar potential v_{ext} and magnetic field B_{ext} this total energy reads

$$\begin{aligned} E[\psi; m] &= T_s[\psi; m] + \int d^3r n(r) v_{\text{ext}}(r) + \int d^3r m(r) B_{\text{ext}}(r) + U[\psi] + E_{xc}[\psi; m] \\ &= \sum_i^{\text{occ}} \int d^3r \psi_i^\dagger(r) v_{xc}(r) \psi_i(r) + \int d^3r m(r) B_{xc}(r) + U[\psi] + E_{xc}[\psi; m]; \end{aligned} \quad (2)$$

where $U[\psi] = \frac{1}{2} \int d^3r \int d^3r' n(r) n(r') / |r - r'|$ is the Hartree energy. The xc potential and xc magnetic field are given by

$$v_{xc}(r) = \frac{E_{xc}[\psi; m]}{n(r)} \quad \text{and} \quad B_{xc}(r) = \frac{E_{xc}[\psi; m]}{m(r)}; \quad (3)$$

respectively. The exact functional form of $E_{xc}[\psi; m]$ is unknown and has to be approximated in practice.

Assuming that the densities $(\psi; m)$ are non-interacting $(v; B)$ -representable one may, equivalently, minimize the total-energy functional (2) over the effective scalar potential and magnetic field. Thus the conditions

$$\frac{\delta E[\psi; m]}{\delta v_s(r)} \Big|_{B_s} = 0 \quad \text{and} \quad \frac{\delta E[\psi; m]}{\delta B_s(r)} \Big|_{v_s} = 0 \quad (4)$$

must be satisfied.

If the functional derivatives in Eq. 4 are evaluated for an xc functional that depends explicitly on the KS spinors, one obtains the natural extension of the OEP equations to non-collinear magnetism. By the usage of spinor valued wavefunctions we can stay within a single global reference frame, in contrast to the case where functionals originally designed for collinear magnetism are used in a non-collinear context by introducing a local reference frame at each point in space. The most commonly used orbital functional is the EXX energy given

by

$$E_x^{\text{EXX}}[\psi; m] = \frac{1}{2} \sum_{i,j}^{\text{occ}} \int d^3r \int d^3r' \frac{\psi_i^\dagger(r) \psi_j(r) \psi_j^\dagger(r') \psi_i(r')}{|r - r'|} \quad (5)$$

where the label occ indicates that the summation runs only over occupied states. In the following we restrict ourselves to an exchange-only treatment although generalization to other orbital functionals is straightforward.

For the energy functional Eq.(2) using the EXX approximation to E_{xc} one obtains the following coupled integrodifferential equations for the exchange potential and magnetic field

$$\begin{aligned} R_v(r) &= \frac{E[\psi; m]}{v_s(r)} \Big|_{B_s} \\ &= \sum_i^{\text{occ}} \sum_j^{\text{occ}} \frac{\psi_i^\dagger(r) \psi_j(r)}{\psi_i^\dagger(r) \psi_j(r)} + \text{c.c.} = 0 \end{aligned} \quad (6)$$

and

$$\begin{aligned} R_B(r) &= \frac{E[\psi; m]}{B_s(r)} \Big|_{v_s} \\ &= \sum_i^{\text{occ}} \sum_j^{\text{occ}} \frac{m_{ij}(r)}{\psi_i^\dagger(r) \psi_j(r)} + \text{c.c.} = 0; \end{aligned} \quad (7)$$

where $\psi_{ij}(r) = \psi_i^\dagger(r) \psi_j(r)$, $m_{ij}(r) = \sum_i^{\text{occ}} \psi_i^\dagger(r) \sigma \psi_j(r)$ and j runs only over the unoccupied states. The matrix

is given by

$$V_{ij}^{NL} = \int_V \int_V \frac{v_i(\mathbf{r}) v_j(\mathbf{r})}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r} d\mathbf{r}'; \quad (8)$$

where

$$V_{ij}^{NL} = \sum_k \frac{v_i(\mathbf{r}) v_k(\mathbf{r}) v_k(\mathbf{r}') v_j(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r} d\mathbf{r}'; \quad (9)$$

are the non-local matrix elements of the Coulomb interaction between states i and j .

To ensure that our numerical analysis be as accurate as possible, we use the full-potential linearized augmented plane wave (FP-LAPW) method [17] implemented within the EXCITING code [18]. Here the single electron potential is calculated exactly without any shape approximation and the space is divided into muffin-tin (MT) regions, where atomic orbitals are used as a basis and interstitial region, where plane waves are used as a basis. The deep lying core states (3 Ha below the Fermi level) are treated as Dirac spinors and valence states as Pauli spinors. More importantly the magnetization density and exchange magnetic field are both treated as unconstrained vector fields throughout space. In our implementation of the OEP method the exchange fields are iteratively updated by subtracting the residue functions R_v and R_B from the exchange fields. In other words, if i is the iteration number then

$$\begin{aligned} v_x^i(\mathbf{r}) &= v_x^{i-1}(\mathbf{r}) - R_v^i(\mathbf{r}); \\ B_x^i(\mathbf{r}) &= B_x^{i-1}(\mathbf{r}) - R_B^i(\mathbf{r}) \end{aligned} \quad (10)$$

is repeated until convergence is reached, with R_v^i and R_B^i calculated by inserting v_x^{i-1} and B_x^{i-1} into Eqs. (6) and (7). i is then fixed chosen in such a manner as to achieve a speedy convergence. In the collinear case this method is similar to the one previously suggested in the Ref. 19.

In order to explore the impact of treating non-collinear magnetism in the way outlined above we compare our approach with the standard LSDA functional using the example of an unsupported Cr (111) monolayer. We set the lattice parameter of the Cr monolayer to that of the Ag (111) surface. The result is a topologically frustrated anti-ferromagnet, known from LSDA calculations to exist as a non-collinear Neel state with the net magnetization direction of the three non-equivalent atoms pointing at 120° to each other. In Fig. 1 we show the magnetization density and B field for both the LSDA and EXX functionals. Both find, as they must, the non-collinear Neel state, and in fact the EXX and LSDA MT averaged moments are similar, being 2.60 μ_B and 2.0 μ_B , respectively. The details of the xc density and field however are very different with the EXX functional producing a lot more structure, in contrast to its fairly homogeneous LSDA counterpart. In the past, the LSDA results (of




FIG. 1: Fully non-collinear magnetization density and B field obtained using the LSDA and exchange-only EXX functionals for an unsupported Cr monolayer in Neel state. Arrows indicate the direction and information about the magnitude (in atomic units) is given in the colour bar.

the kind shown in Fig. 1), which show almost no non-collinearity in the MT region, led to the conclusion that it is sufficient to treat only the interstitial region as non-collinear [16]. The present work shows that orbital functionals such as EXX are more sensitive to the atomic shell structure and this sensitivity also manifests itself in the magnetization density and exchange B field. This is clear from the flower petal like structure visible in the magnitude of EXX density and B field. The Neel walls are also much narrower in the EXX case. Adding LSDA correlations to the EXX functional does not significantly change these results. A striking feature of the EXX B field is that, unlike its LSDA counterpart, it is not locally parallel to the magnetization density.




FIG. 2: $m(\mathbf{r})$ $B_x(\mathbf{r})$ for an unsupported Cr monolayer, in the same plane as Fig. 1, obtained using the EXX functional. Arrows indicate the direction and information about the magnitude (in atomic units) is given in the colour bar.

Another appealing property of the EXX functional

- [18] J. K. Dewhurst, S. Sharma, and C. Ambrosch-Daxl (2004), URL <http://exciting.sourceforge.net>.
- [19] S. Kummel and J. P. Perdew, Phys. Rev. B 68, 035103 (2003).
- [20] T. Kotani and H. Akai, Physica B 237, 332 (1997).