

Definition and relevance of nonequilibrium intensive thermodynamic parameters

Eric Bertin,¹ Olivier Dauchot,² and Michel Droz¹

¹Department of Theoretical Physics, University of Geneva, CH-1211 Geneva 4, Switzerland

²SPEC, CEA Saclay, F-91191 Gif-sur-Yvette Cedex, France

(Dated: April 14, 2024)

We show that intensive thermodynamic parameters associated to additive conserved quantities can be naturally defined from a statistical approach in far-from-equilibrium steady-state systems, under few assumptions, and without any detailed balance requirement. It may apply, e.g., to dissipative systems like granular gases where volume or mass is still conserved, or to systems with periodic boundary conditions where fluxes of conserved quantities are present. We emphasize the usefulness of this concept to characterize the coexistence of different nonequilibrium phases, and discuss the influence of the contact between two different systems, in relation with measurement issues.

PACS numbers: 05.20.-y, 05.70.Ln, 05.10.Lc

One of the essential features of intensive thermodynamic parameters (ITPs), like temperature, pressure or chemical potential, is that these parameters take equal values in two different systems that can exchange the conjugated extensive quantity (energy, volume, or particles) once equilibrium is reached. This is a strong property, since it holds even if the systems put into contact have very different microscopic dynamics. This equilibration between different systems makes ITPs a powerful concept to describe the influence of the environment or phase coexistence, and is essential to allow, say, the temperature of a system to be measured with a thermometer.

Several approaches have been developed in nonequilibrium statistical physics to generalize the notion of ITP, usually focusing on temperature. For aging systems, an effective temperature may be defined from the long-time slope of the fluctuation-dissipation relation (FDR) [1, 2, 3]. For systems weakly driven into a stationary nonequilibrium state, ITPs are defined locally as in equilibrium [4]. On the other side, phenomenological extensions to far-from-equilibrium regimes [5] seem to lack microscopic foundations. For such systems s , statistical temperatures have been proposed in specific models, from FDR [6], maximum entropy principle [7], or generalized microcanonical approaches [8]. Yet, most of these studies do not discuss the relevance and generality of the so-defined temperature.

In this Letter, we propose a general definition of ITPs conjugated to additive conserved quantities in nonequilibrium systems, valid under few assumptions. This definition, which does not require any detailed balance relation to be fulfilled, is illustrated on a stochastic model in which a dissipated energy is coupled to a globally conserved mass. The usefulness of nonequilibrium ITPs to characterize the coexistence of nonequilibrium phases is outlined, on the example of a two-species mass transport model. Finally, we discuss the problem of the contact between two nonequilibrium systems with different microscopic dynamics, as well as the relation with the large deviation functional approach [9].

Framework and hypotheses. Starting with a general formulation of the problem, we consider the steady state of a macroscopic nonequilibrium system for which some additive quantities Q_k , $k = 1; \dots$, are globally conserved by the dynamics. On general grounds, the stationary probability to be in a microstate γ , with probability weight f_γ , is of the form

$$P_\gamma = \frac{f_\gamma}{Z(Q_1; \dots; Q_N)} \prod_{k=1}^N \delta(Q_k - Q_{k,\gamma}) \quad (1)$$

where the product of delta functions accounts for the conservation laws, and $Z(Q_1; \dots; Q_N)$ is a normalization factor, named hereafter a "partition function". Splitting the system into two subsystems S_1 and S_2 , so that $\gamma = \gamma_1; \gamma_2$, we introduce the probability $(Q_{11}; \dots; Q_{1N}; Q_{21}; \dots; Q_{2N})$ that conserved quantities have values Q_{k1} in S_1 , given their total values Q_k

$$(f_{Q_{k1}} g f_{Q_k}) = \sum_{\gamma_1; \gamma_2} P_{\gamma_1; \gamma_2} \prod_{k=1}^N \delta(Q_{k1} - Q_{k1,\gamma_1}) \delta(Q_k - Q_{k,\gamma_2}) \quad (2)$$

The key assumption in the following derivation is that this conditional probability satisfies an asymptotic factorization property, namely

$$\ln (f_{Q_{k1}} g f_{Q_k}) = \ln Z_1(f_{Q_{k1}} g) + \ln Z_2(f_{Q_k - Q_{k1}} g) - \ln Z(f_{Q_k} g) + o_N(f_{Q_{k1}} g; f_{Q_k} g) \quad (3)$$

where $Z(f_{Q_k} g)$ refers to subsystem S , when isolated, and $o_N(f_{Q_{k1}} g; f_{Q_k} g)$ becomes negligible with respect to $\ln Z$ in the large N limit, N being the number of degrees of freedom. Although not obvious in nonequilibrium systems where long-range correlations might develop, this factorization property actually holds in some rather large classes of models. It is satisfied, for instance, in lattice models like the ZRP [10], the ARAP [11], or more general mass transport models [12, 13], where the probability weight f factorizes as a product of one-site weights $f = \prod_{i=1}^N f_{i; \gamma_i}$, with $f_i = f_{i; \gamma_i}$. In this case, $o_N = 0$ in Eq. (3). Besides, this asymptotic factorization property

also holds in one-dimensional models with conservation laws, for which the stationary probability is given by a matrix product ansatz with finite-dimensional matrices: $f = \text{Tr} \prod_{i=1}^N M_i$, where M_i is a matrix associated to the local state Q_i (for nonperiodic systems, one rather has $f = \langle \mathbf{w} | \prod_{i=1}^N M_i | \mathbf{v} \rangle$, with vectors \mathbf{w} and \mathbf{v} determined by the boundary conditions). This approach has been used, e.g., in the context of the ASEP model, and can lead to finite-dimensional [14, 15] or infinite-dimensional matrices [15, 16]. We shall simply sketch here the argument leading to Eq. (3), and defer a more detailed presentation to a later publication [17]. For simplicity, we consider a generic model with local variables q_i and one conserved quantity $Q = \sum_{i=1}^N q_i$. The stationary distribution $P(fq_i g)$ reads

$$P(fq_i g) = \frac{1}{Z(Q)} \text{Tr} \prod_{i=1}^N M_i(q_i) \prod_{i=1}^N q_i^{f_i} Q^{g_i}; \quad (4)$$

where $M_i(q)$ is a finite-dimensional matrix. Introducing the matrix $R(Q)$ such that $Z(Q) = \text{Tr} R(Q)$, one has

$$(Q_1 | Q) = \frac{1}{Z(Q)} \text{Tr} [R_1(Q_1) R_2(Q - Q_1)] \quad (5)$$

Denoting by $\hat{R}(s)$ and $\hat{M}(s)$ the Laplace transforms of $R(Q)$ and $M(q)$, one has $\hat{R}(s) = \hat{M}(s)^N$. If $\hat{M}(s)$ is invertible, $\hat{M}(s) = \exp(B(s))$, so that $\hat{R}(s) = \exp(NB(s))$. $B(s)$ may be decomposed as $B(s) = D(s) + L(s)$ where $D(s)$ is diagonalizable, $L(s)$ is nilpotent (i.e. $L(s)^{p(s)} = 0$), and $D(s)$ commutes with $L(s)$. It follows that $\hat{R}(s) = \exp(ND(s)) \exp(NL(s))$, where $\exp(NL(s))$ is a polynomial in $NL(s)$: the dominant contribution to $\hat{R}(s)$ is proportional to $N^{p(s)-1} \exp(N\lambda_1(s))$, with $\lambda_1(s)$ the largest eigenvalue of $D(s)$. If $p(s)$ is bounded when $s \rightarrow 0$, one can show performing an inverse Laplace transform that the asymptotic factorization property (3) holds [17].

For infinite matrix representations, this argument may break down as the eigenvalues of $D(s)$ might not be bounded. From a physical viewpoint, it is reasonable to think that the factorization property (3) is related to the presence of a finite correlation length in the system, as large subsystems then become essentially independent. Both properties are expected to hold for finite matrices [15]. Yet, for infinite matrices, whether these two properties are actually related or not remains an open issue.

Definition of nonequilibrium ITPs. { To define a nonequilibrium ITP, we compute, guided by equilibrium procedures, the derivative of $\ln(fQ_{k1} g fQ_{k2} g)$ with respect to Q_{k1} at the most probable value Q_{k1} . Equating this derivative to zero yields

$$\frac{\partial \ln Z_1}{\partial Q_{k1} - Q_{k1}} = \frac{\partial \ln Z_2}{\partial Q_{k2} - Q_k - Q_{k1}} \quad (6)$$

Thus it is natural to define the ITP π_k of the system as

$$\pi_k = \frac{\partial \ln Z}{\partial Q_k} \quad (7)$$

since this parameter, when defined within subsystems that can exchange the quantity Q_k , takes equal values in both subsystems once the steady-state is reached, as seen from Eq. (6) {see [8] for a more detailed discussion on a specific example. In the following, this property is loosely called the "equilibration" of subsystems, although we deal with nonequilibrium situations.

The above partition into two subsystems is also useful when one subsystem is small, but still macroscopic. The effect of the rest of the system, acting as a reservoir of Q_k , may then be encoded into the parameters π_k . Integrating the distribution (1) over the degrees of freedom of the reservoir leads for the subsystem to

$$P = \frac{f}{Z(\pi_1; \dots; \pi_N)} \exp \left(\sum_{k=1}^N \pi_k Q_k \right) \quad (8)$$

which may be called a "grand-canonical" distribution. Note also that $\ln Z$ may be interpreted as a nonequilibrium thermodynamic potential, and that, as in the equilibrium formalism, the cumulants of Q_k are given by the successive derivatives of $\ln Z$ with respect to π_k [17].

Illustration on a dissipative model. { Let us illustrate on a simple nonequilibrium model the definition (7) of the ITP. We consider a one-dimensional model with two different physical quantities, say, an "energy" $E = \sum_{i=1}^N \epsilon_i$ and a "mass" $M = \sum_{i=1}^N m_i$. Energy is flowing through the boundaries, and is partially dissipated in the bulk, whereas mass is conserved and cannot flow through the boundaries. The continuous time dynamics is defined as follows. An amount of energy ϵ may be either injected at site $i=1$ with rate $J(\epsilon)$, moved from site i to $i+1$ with rate $\epsilon'_1(\epsilon; m_i)$ (if $i=N$, the energy flows out of the system), or dissipated on site i with rate $\epsilon(\epsilon; m_i)$. Besides, an amount of mass m may be transported from site i to $j=i+1$ (except for $i=1$ or N , where $j=2$ or $N-1$ respectively) with rate $\epsilon'_2(\epsilon; m_i)$, and with equal probability for both target sites. To our knowledge, this model, which generalizes the cascade model introduced in [18], was not considered previously in the literature. In order to allow for a factorized distribution, the rate functions are chosen as follows

$$J(\epsilon) = v_1(\epsilon) e^{b\epsilon} \quad (9)$$

$$\epsilon'_1(\epsilon; m) = v_1(\epsilon) \frac{g(\epsilon; m)}{g(\epsilon; m)} \quad (10)$$

$$\epsilon'_2(\epsilon; m) = v_2(\epsilon) \frac{g(\epsilon; m)}{g(\epsilon; m)} \quad (11)$$

$$\epsilon(\epsilon; m) = (e^{a\epsilon} - 1) v_1(\epsilon) \frac{g(\epsilon; m)}{g(\epsilon; m)} \quad (12)$$

with positive functions $v_1(\epsilon)$, $v_2(\epsilon)$, $g(\epsilon; m)$, and two positive parameters a and b . Clearly, due to the presence of flux and dissipation, the above dynamics cannot satisfy

detailed balance. The steady-state distribution reads

$$P(f_j; m_j g) = \frac{\prod_{j=1}^N f_j(f_j; m_j)}{Z(M)} \exp\left(-\sum_{j=1}^N m_j M_A\right) \quad (13)$$

with $f_j(f_j; m_j) = g(f_j; m_j) \exp[(a_j N + b)f_j]$ on site j . Let us choose for the function $g(f; m)$ the simple form $g(f; m) = \exp(-mf)$ with $a > 1$ and $b > 0$. The partition function can be computed in the large N limit, and one finds $Z(M) = A M^{(1-1/N)}$ with A independent of M . This gives for the ITP $\mu = (1-1/N)M$ with $\mu = M/N$, showing that μ depends, through N , on the coupling to the nonconserved energy.

Relevance of ITPs for phase coexistence. Let us now discuss the case where two systems with the same microscopic dynamics, but different macroscopic states, can exchange some globally conserved quantities. A well-studied example of such a situation is the condensation transition observed in ZRP [10, 19, 20, 21], or more generally mass transport models [13]. This transition occurs when the overall density exceeds a critical value ρ_{crit} , so that a finite fraction of the total mass condense onto a given site. In standard approaches [10], the critical density is obtained by looking for the convergence radius of a formal grand-canonical partition function [22]. Details about the condensed phase need to be studied in the canonical ensemble where the total mass is fixed [13].

In the following, we interpret the condensation as a phase coexistence and show that enforcing the equality of the ITP defined in Eq. (7) (here, a chemical potential) in the two phases leads to a natural quantitative description of the condensation. To this aim, we compute separately the chemical potential of the condensate, taken as isolated, and the equation of state of the fluid phase. Let us illustrate this point on a concrete example. We consider a variant of the model defined by Eqs. (9) to (12), defined on a periodic lattice, where \mathcal{J} is now a conserved quantity (that is, $\mathcal{J}(\cdot) = 0$ and $\mathcal{J}(\cdot; m) = 0$), so that we shall replace the notations \mathcal{J}_i and m_i with m_{1i} and m_{2i} respectively (both quantities are now "masses"). The continuous time dynamics proceeds by transferring a mass ± 1 or ± 2 from a random site i to $i+1$, with rates $\gamma_1(\pm 1; m_{1i}; m_{2i})$ or $\gamma_2(\pm 2; m_{1i}; m_{2i})$ given in Eqs. (10) and (11) respectively. In dimension $d > 1$, the mass ± 1 or ± 2 may also be transferred in any direction transverse to the axis, with equal probability. The steady-state distribution in arbitrary dimension d is given by

$$P(f_{m_{ki}} g) = \frac{\prod_{i=1}^N g(m_{1i}; m_{2i})}{Z(M_1; M_2)} \exp\left(-\sum_{k=1}^d \sum_{i=1}^N m_{ki} M_k\right) \quad (14)$$

Guided by the two-species ZRP studied in [21], we choose $g(m_1; m_2) = \exp(-m_2 m_1)$, with $a > 0$. Assuming that a condensate containing large amounts M_{1c} and M_{2c} forms, the partition function of the isolated condensate

is $Z_c(M_{1c}; M_{2c}) = g(M_{1c}; M_{2c})$, and the chemical potentials $\mu_k = -\partial \ln Z_c / \partial M_{kc}$ read

$$\mu_1 = \frac{M_{2c}}{M_{1c} + 1}; \quad \mu_2 = \frac{1}{M_{1c}} \quad (15)$$

Thus in the thermodynamic limit, $\mu_2 = 0$. Enforcing the equality of the chemical potentials in both phases $\mu_k = \mu_{kc}$, the densities ρ_k in the fluid phase are computed in the grand-canonical ensemble, yielding $\rho_1 = (1 + \mu_1) = 1$ and $\rho_2 = (1 + 2\mu_1) = [1 + (1 + \mu_1)]$. Hence $\rho_1 > 0$, and $M_{1c} = M_{2c}^{1/(1+\mu_1)}$. The condensate M_{1c} is subextensive, and $\rho_1 = 1$. It follows that ρ_1 , and thus ρ_2 , are fixed by the overall density ρ . A condensate forms if $\rho > \rho_2^{\text{crit}}$, meaning that ρ_2^{crit} is the critical density ρ_2^{crit} . One has $M_{2c} = N(\rho_2 - \rho_2^{\text{crit}})$, and Eq. (15) for μ_1 gives

$$M_{1c} = \frac{1}{1 + \mu_1} N(\rho_2 - \rho_2^{\text{crit}})^{\frac{1}{1 + \mu_1}} \quad (16)$$

Accordingly, the framework of ITPs provides a simple description of the condensation in terms of the (out-of-equilibrium) "equilibration" of two coexisting phases.

Contact of systems with different dynamics. In this part, we consider the issue of the equalization of ITPs when two systems with different microscopic dynamics are put into contact. This is precisely the type of problem encountered when one wishes to define a "thermometer", which in the present more general nonequilibrium context, we shall call "ITP-meter". For such an "ITP-meter", one requires two essential properties. First, its ITP must equalize with that of the system over which the measure is performed, without perturbing this system. Second, one needs to know the equation of state of the "ITP-meter" to relate its ITP to a directly measurable quantity. This is a highly non-trivial problem, which depends on the microscopic dynamics of each system, but also on the dynamics at the contact. Let us imagine that two different systems S_1 and S_2 , that separately conserve the same physical quantity $Q = \sum_i q_i$, are put into contact. The dynamics at the contact indeed imposes the distribution (Q_1, Q_2) for the random partition of Q into Q_1 and $Q_2 = Q - Q_1$ over the two systems. The probability distribution of the whole system then reads, assuming for simplicity that the probability weights of each system strictly factorize (the index i refers to S_1)

$$P(f_{q_i} g) = \frac{\prod_{i_1, i_2} Q_{i_1, i_2} f_{q_{i_1}} f_{q_{i_2}}}{Z_1(Q_1) Z_2(Q - Q_1)} \prod_i q_i Q \quad (17)$$

with $Q_1 = \sum_{i \in S_1} q_i$. The ITPs of the two systems equalize only if $(Q_1, Q_2) = Z_1(Q_1) Z_2(Q - Q_1) = Z(Q)$. This relation is indeed satisfied in equilibrium due to the detailed balance relation, if the Hamiltonian is additive [17]. Out of equilibrium, the above factorization and the

equalization of the ITPs does not hold in general, but are recovered inside some classes of nonequilibrium systems.

To illustrate this point, let us consider two single conserved mass transport models S_1 and S_2 with site-independent transport rates $v_{1,2}(j) = v(j)f_{1,2}(j) = f_{1,2}(j)$, with different $f_1(j)$ and $f_2(j)$, but the same $v(j)$. The dynamics at the contact is defined as follows: a mass leaving a boundary site may be transferred to one of the neighboring sites or, with the same probability, to the other system. The global probability weights factorize (as $v(j)$ is site-independent [12, 17]), and the ITPs of the two systems equalize. The key point is that the equalization of the ITPs occurs for two different microscopic dynamics ($f_1(j) \neq f_2(j)$), but belonging to a given class (same $v(j)$). Indeed, if $v_1(j) \neq v_2(j)$, such an equalization does not hold. This result suggests the existence of classes of systems which mutually "equilibrate", so that any member of a class, with known equation of state, may be used as an "ITP-meter" for the other members, under suitable size ratios.

Discussion. We have proposed a general definition of ITPs associated to additive conserved quantities in nonequilibrium steady states, under the asymptotic factorization assumption (3). This property holds in particular if the stationary distribution is factorized or, for one-dimensional systems, if it can be computed from a matrix product ansatz with finite-size matrices. Its applicability to more general situations remains an important open issue. Indeed, it would be essential to have a criterion allowing to determine experimentally or numerically whether the factorization property holds, without having to know the stationary probability distribution explicitly. Note that in equilibrium, the same difficulty arises in principle, but it reduces there to the "additivity" of the hamiltonian (short-range interactions).

Besides, we note that if (Q_1, Q) may be written as $(Q_1, Q) = \exp[NG(\cdot; \cdot)]$, with $Q_1 = Q|_{N_1}$ and $Q = Q|_N$, Eq. (3) corresponds to an additivity property for $G(\cdot; \cdot)$, which somehow resembles the additivity principle for the large deviation functional $F(f(x)|g)$ of the density profile $\rho(x)$ in the open ASEP model [9]. As the physically relevant states minimize $G(\cdot; \cdot)$ with respect to Q_1 in the thermodynamic limit, $G(\cdot; \cdot)$ appears as a "free energy" function [23]. In analogy with the ASEP where $F(f(x)|g)$ is a nonlocal functional of $\rho(x)$, $G(\cdot; \cdot)$ is a nonlocal function as it depends on the global density ρ , due to the conservation law. Still, the physical origin of this nonlocality may be different in the two problems. Also, the quantity Q is conserved here, whereas particles are exchanged with boundary reservoirs in the ASEP.

From Eq. (6), one sees that the equality of ITPs between subsystems is related to the minimization of a nonlocal "free energy" function. Note however that this is already true in equilibrium systems. The specificity of nonequilibrium dynamics shows up mostly when the sys-

tems into contact have different microscopic dynamics. In this case, the ITPs of the two systems may not equalize, due to the essential role played by the dynamics at the contact (a role hidden at equilibrium due to detailed balance). Yet, there may exist classes of nonequilibrium systems that mutually "equilibrate".

Acknowledgements. This work has been partly supported by the Swiss National Science Foundation.

-
- [1] L. F. Cugliandolo, J. Kurchan, and L. Peliti, Phys. Rev. E 55, 3898 (1997).
 - [2] J. Kurchan, J. Phys.: Cond. Mat. 12, 6611 (2000); Nature 433, 222 (2005).
 - [3] A. Garriga and F. Ritort, Phys. Rev. E 72, 031505 (2005); F. Ritort, e-print cond-mat/0509488.
 - [4] S. de Groot and P. Mazur, Non-equilibrium Thermodynamics, (North-Holland, Amsterdam, 1962).
 - [5] J. Casas-Vazquez and D. Jou, Rep. Prog. Phys. 66, 1937 (2003).
 - [6] A. Puglisi, A. Baldassarri, and V. Loreto, Phys. Rev. E 66, 061305 (2002); A. Barrat, V. Loreto, and A. Puglisi, Physica A 334, 513 (2004).
 - [7] B. N. Miller and P. M. Larson, Phys. Rev. A 20, 1717 (1979); F. Sastre, I. Domic, and H. Chate, Phys. Rev. Lett. 91, 267205 (2003).
 - [8] E. Bertin, O. Dauchot, and M. Droz, Phys. Rev. Lett. 93, 230601 (2004); Phys. Rev. E 71, 046140 (2005).
 - [9] B. Derrida, J. L. Lebowitz and E. R. Speer, Phys. Rev. Lett. 89, 030601 (2002); J. Stat. Phys. 110, 775 (2003).
 - [10] M. R. Evans and T. Hanney, J. Phys. A 38, R195 (2005).
 - [11] R. Rajesh and S. N. Majumdar, J. Stat. Phys. 99, 943 (2000); F. Ziesler and A. Schadschneider, J. Stat. Phys. 106, 173 (2002).
 - [12] M. R. Evans, S. N. Majumdar and R. K. P. Zia, J. Phys. A 37, L275 (2004).
 - [13] S. N. Majumdar, M. R. Evans, and R. K. P. Zia, Phys. Rev. Lett. 94, 180601 (2005).
 - [14] F. H. L. Essler and V. Rittenberg, J. Phys. A 29, 3375 (1996); K. Mallick and S. Sandow, J. Phys. A 30, 4513 (1997).
 - [15] P. F. Amdt, T. Heinzel, and V. Rittenberg, J. Stat. Phys. 97, 1 (1999).
 - [16] B. Derrida, M. R. Evans, V. Hakim, and V. Pasquier, J. Phys. A 26, 1493 (1993).
 - [17] E. Bertin, O. Dauchot, and M. Droz, to be published.
 - [18] E. Bertin, Phys. Rev. Lett. 95, 170601 (2005).
 - [19] S. Grosskinski, G. M. Schutz, and H. Spohn, J. Stat. Phys. 113, 389 (2003).
 - [20] C. Godreche, J. Phys. A 36, 6313 (2003).
 - [21] T. Hanney and M. R. Evans, Phys. Rev. E 69, 016107 (2004); S. Grosskinski and T. Hanney, Phys. Rev. E 72, 016129 (2005).
 - [22] By "formal", we mean that it is defined by analogy with equilibrium as a superposition of different canonical partitions functions, without being clearly associated with a well-defined physical ensemble (see, e.g., [15]).
 - [23] Introducing a partition into a large number of subsystems, one may also construct in this way a large deviation functional for the density profile $\rho(x)$.