

# On the Bose-Einstein Condensation of Magnons in $\text{Cs}_2\text{CuCl}_4$

S.E. Sebastian<sup>1</sup>, V.S. Zapf<sup>2</sup>, N.H. Harrison<sup>2</sup>, C.D. Batista<sup>3</sup>, P.A. Sharma<sup>2</sup>, M. Jain<sup>2</sup>, I.R. Fisher<sup>1</sup>, A. Lacerda<sup>2</sup>

<sup>1</sup>Geballe Laboratory for Advanced Materials and Department of Applied Physics, Stanford University, Stanford, CA 94305

<sup>2</sup>MS-T-NHMF, Los Alamos National Laboratory, Los Alamos, NM 87545 and

<sup>3</sup>Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545

(Dated: April 14, 2024)

In a recent paper [1], Radu et al. report experimental results they claim to support Bose-Einstein condensation (BEC) of magnons in  $\text{Cs}_2\text{CuCl}_4$ . It is true that an experimentally measured critical power law scaling exponent in agreement with the BEC universality class would support the realization of a BEC in magnetic systems that order as a canted antiferromagnet. It can be shown, however, that the claim of Radu et al. is overstated in this instance, because their determination of the critical exponent relies on a model-dependent theoretical approximation to the critical field  $H_{c1}$  for which the associated errors are neglected. We show that when these errors are included, the uncertainty in the obtained exponent is so large that the available experimental data cannot be used to differentiate between contending universality classes.

A two parameter fit to only a few data points delineating the critical ordering temperature ( $T_c$ ) versus magnetic field ( $H$ ) in the vicinity of the quantum critical point (QCP), to the power law

$$T_c \propto (H - H_{c1})^{\frac{1}{\nu}}$$

with both  $H_{c1}$  and the critical exponent  $\nu$  varying has been shown to be unreliable [2, 3]. An independent experimental determination of  $H_{c1}$  is therefore required to obtain an accurate estimate of  $\nu$ . Given that neutron scattering measurements on  $\text{Cs}_2\text{CuCl}_4$  presented in Ref. [4] have provided such a determination, yielding  $H_{c1} = 8.44 \pm 0.01$  T, this would be an appropriate value to use in the fit to Eqn. (1). Radu et al. instead use a value of  $H_{c1}' = 8.51$  T in their fit to Eqn. (1), calculated using an approximate theoretical Hamiltonian, that is subsequently assumed to have zero error in their analysis. This assumption has two principal inaccuracies. The first is that the model Hamiltonian neglects higher order interactions, thereby introducing an unknown systematic error in  $H_{c1}'$ . The second is that the exchange couplings used in its computation have significant experimental uncertainty, introducing a large error in  $H_{c1}'$ . We obtain  $H_{c1}' = 8.51 \pm 0.12$  T on using the published errors in the exchange interactions [4].

Fig. 1 shows fits of Eqn. (1) to the experimentally measured phase boundary data points using both the experimental value of  $H_{c1} = 8.44 \pm 0.01$  T of Coldea et al. [4] and the theoretical estimate of  $H_{c1}' = 8.51 \pm 0.12$  T, yielding  $\nu = 2.8 \pm 0.4$  and  $\nu = 1.5 \pm 0.9$  respectively, on considering the dominant contribution to the error:

$$\nu = \frac{d}{dH} \left( \frac{1}{T_c} \right)_{H_{c1}} H_{c1}. \text{ The single most important factor}$$




FIG. 1: Points on the ordering phase boundary from the experimental data in [1]. The solid line represents the best fit to using the experimentally measured value of  $H_{c1} = 8.44$  T from [4]. The dashed line represents the best fit to using the theoretical estimate of  $H_{c1}' = 8.51$  T as per the analysis technique used in [1]. The inset shows the variation in the  $\nu$  value of  $\nu$  with the value of  $H_{c1}$ .

responsible for the very large error of  $\sim 60\%$  in the case of the latter as compared to the error of  $\sim 14\%$  in the former fit, is the extreme sensitivity of the fit to the theoretical estimate of the critical field  $H_{c1}'$ , as depicted graphically in the inset to Fig. 1.

Given the substantial uncertainty in the value of  $\nu$  that is obtained from a rigorous analysis, it is clear that the available experimental data do not favor the 3d BEC universality class ( $\nu = 1.5$ ) over other possibilities, including the 3d Ising universality class ( $\nu = 2$ ).

- [1] T. Radu et al., Phys. Rev. Lett. 95, 127202 (2005).
- [2] O. Nohadani et al., Phys. Rev. B. 69, 220402(R) (2004).
- [3] S.E. Sebastian et al., Phys. Rev. B. 72, 100404(R) (2005).
- [4] R. Coldea et al., Phys. Rev. Lett. 88, 137203 (2002).
- [5] M. Y. Veillette et al., Phys. Rev. B 71, 214426 (2005).
- [6] S. Sachdev Quantum Phase Transitions (Cambridge University Press, 1999).