

## On the conservation of spin currents in spin-orbit coupled systems

R. Shen,<sup>1,2</sup> Yan Chen,<sup>1</sup> and Z. D. Wang<sup>1,2</sup><sup>1</sup>Department of Physics, The University of Hong Kong, Pokfulam Road, Hong Kong, China<sup>2</sup>National Laboratory of Solid State Microstructures and Department of Physics, Nanjing University, Nanjing, 210093 China

Applying the Gordon-decom position-like technique, the convective spin current (CSC) is extracted from the total angular-momentum current. The CSC describes the transport properties of the electron spin and is conserved in the relativistic quantum mechanics approach where the spin-orbit coupling has been intrinsically taken into account. Interestingly, in the presence of external electromagnetic field, the component of the convective spin along the field remains still conserved. This conserved CSC is also derived for the first time in the nonrelativistic limit using the Foldy-Wouthuysen transformation.

PACS numbers: 72.25.-b, 03.65.-w, 71.70.Ej

Spintronics is a fast developing field in condensed matter physics, where the research on the spin transport and manipulation in solid state systems appears to be a central subject.<sup>1,2</sup> Typically, the spin transport is strongly affected by a coupling of spin and orbital degrees of freedom in semiconductors. Of particular interest is the spin Hall effect in a spin-orbit coupled system. The spin Hall effect means that a transverse spin current can be induced by applying an electric field. Earlier studies focused on the extrinsic effects due to the spin dependent scattering.<sup>3,4,5</sup> Recently, an intrinsic spin Hall effect (ISHE), which is resulted from the spin-orbit coupled bands and may exist in the clean systems, has attracted considerable attentions.<sup>6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28</sup> On the theoretical side, the ISHE has been predicted in the p-doped semiconductors described by the Luttinger-like model<sup>6</sup> and in the two-dimensional electron gas described by the Rashba-like model<sup>7</sup>. The ISHE in the two-dimensional hole gas<sup>20</sup> and the impurity scattering effect on the ISHE<sup>15,16,17,18,19,20,21</sup> have also been analyzed. On the experimental side, the spin accumulation related to the spin Hall effect has been observed in the n-type bulk semiconductor<sup>30</sup> and in the two-dimensional hole gas.<sup>31</sup>

The spin current is an important issue in discussing the spin Hall effect and other spin transport phenomena. In spite of the extensive investigation on the ISHE, there is still a fundamental problem that what is the proper definition of the spin current operator in the spin-orbit coupled system. The conventional definition of the spin current operator is the anticommutator of the Pauli spin matrix and the velocity of the electron.<sup>6,7</sup> The electron spin is not a conserved quantity in the spin-orbit coupled system. As a result, there is a source term, namely spin torque, in the continuity-like equation related to the conventional spin current.<sup>25,26,27</sup> The spin torque represents the spin precession caused by the external field and the spin-orbit coupling. The presence of the nonzero source term in the continuity-like equation makes the conventional spin current an ill-defined quantity to describe the transport of spin. On the other hand, the non-conserved part of the conventional spin current has

very fast dynamics and can not contribute to the low frequency response.<sup>24</sup> Therefore, it is essential to define a conserved effective spin current from a first principle for studying the spin transport properties in the spin-orbit coupled system.

There are already some efforts focusing on the understanding of the nonconservation of the conventional spin current.<sup>22,23,24,25,26,27,28,29</sup> In the p-doped semiconductor described by the Luttinger-like model, Murakami et al.<sup>24</sup> have proposed a conserved effective spin current of the spin  $S=2$  hole by modifying both the spin density and the spin current operator. By rewriting the spin torque as the divergence of a torque dipole density, another approach to the conserved effective spin current was also proposed<sup>29,30,31</sup> where the spin density operator remains unchanged and only the spin current operator is modified. In this paper, motivated by the fact that the spin-orbit coupling in the semiconductors comes from the relativistic correction of the electron motion, we develop a conserved effective spin current in the relativistic quantum mechanics approach where the spin-orbit coupling has been intrinsically taken into account. For a relativistic electron, the spin is not conserved, but the total angular-momentum (TAM) consisting of both spin and orbit angular-momentum is a conserved quantity in the absence of external fields. Therefore, there is the conserved TAM current according to the Noether's theorem. Under the Gordon-decom position-like scheme, the TAM current can be divided into two parts: the convective TAM current and the internal one. It is found that the spin part of the convective TAM current, namely the convective spin current (CSC) satisfies the continuity equation. The CSC is not only conserved but also closely related to the spin flux of the electron moving in the spatial space and therefore is an appropriate candidate for the spin current operator in discussing the spin transport properties. In the presence of external electromagnetic field, neither the TAM nor the CSC is conserved in a general case because of the mutual exchange of the angular-momentum between the electron and the external field. However, more intriguingly, we find that if both the electric field and the magnetic field are along the same direction, namely the z-direction, the z-components of

the CSC and the TAM still remain conserved. Remarkably, this is just the case in the Rashba-like spin-orbit coupled system. As a result, the CSC is also likely robust subject to the external field in some sense. Finally, by using the Foldy-Wouthuysen transformation, the conserved CSC is derived in the nonrelativistic (NR) limit, which has a potential application in spintronics.

We start with the brief review on the Gordon decomposition of the probability current of the relativistic electron. For the electron of the mass  $m$  traveling in the three-dimensional free space, the four-component wave function of the electron satisfies the Dirac equation

$$i\hbar \frac{\partial}{\partial t} \psi = c \boldsymbol{\alpha} \cdot \mathbf{p} \psi + m c^2 \beta \psi; \quad (1)$$

where the momentum  $\mathbf{p} = -i\hbar \nabla$ , the matrices  $\boldsymbol{\alpha} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ ,  $\beta = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ ,  $\boldsymbol{\sigma}$  is the Pauli spin matrix and  $\mathbb{1}$  is the  $2 \times 2$  unity matrix. The invariance of the Dirac Eq. (1) under the U(1) gauge transformation results in the continuity equation for the probability current

$$\frac{\partial}{\partial t} \rho + \nabla \cdot \mathbf{j} = 0; \quad (2)$$

where the probability density  $\rho = \psi^\dagger \psi$  and the current  $\mathbf{j} = c \psi^\dagger \boldsymbol{\alpha} \psi$ . The Dirac Eq. (1) can be rewritten as

$$i\hbar \frac{\partial}{\partial t} \psi = c \boldsymbol{\alpha} \cdot \nabla \psi + m c^2 \beta \psi; \quad (3)$$

Substituting Eq. (3) into the definitions of  $\rho$  and  $\mathbf{j}$ , one finds that the density and current can be divided into the convective and internal parts, known as the Gordon decomposition,<sup>32,33</sup> in the form

$$\rho = \rho_c + \rho_i; \quad \mathbf{j} = \mathbf{j}_c + \mathbf{j}_i; \quad (4)$$

$$\begin{aligned} \rho_c &= \frac{i\hbar}{2m c^2} \left( \psi^\dagger \frac{\partial}{\partial t} \psi - \frac{\partial}{\partial t} \psi^\dagger \psi \right); \\ \mathbf{j}_c &= \frac{\hbar}{2m i} \left( \psi^\dagger \nabla \psi - \nabla \psi^\dagger \psi \right); \end{aligned} \quad (5)$$

$$\rho_i = \psi^\dagger \beta \psi; \quad \mathbf{j}_i = c \psi^\dagger \boldsymbol{\alpha} \beta \psi + \frac{\partial \mathbf{P}}{\partial t}; \quad (6)$$

where

$$\mathbf{M} = \frac{\hbar}{2m c} \nabla \times \mathbf{j}; \quad \mathbf{P} = \frac{\hbar}{2m c} \nabla (\rho_i); \quad (7)$$

and the matrix  $\boldsymbol{\alpha} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  as well as  $\beta = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ . The internal parts of the density and current are closely related to the internal state of the electron. For the electron of charge  $q$ ,  $q\mathbf{M}$  is essentially the magnetic moment density of the electron arising from its spin and  $q\mathbf{P}$  is the electric polarization belonging to the spin. The convective parts of the density and current are determined by the rate of change of  $\psi$  in space and time and closely related

to the motion of the electron in the spatial space. Notice that the convective and internal parts of the density are separately conserved. For the NR limit,  $\rho_i$  is zero up to the zeroth order in  $(1/c)$ ,  $\mathbf{j}_i$  is only a curl term and has nothing to do with the electron accumulation, and the convective parts of the current and the density are reduced to the usual form in the NR quantum mechanics,

$$\rho_{c,NR} = \psi^\dagger \psi; \quad \mathbf{j}_{c,NR} = \frac{\hbar}{2m i} \nabla \psi^\dagger \psi - \psi^\dagger \nabla \psi; \quad (8)$$

with the two-component wave function of the electron.

Next, we turn to the spin current. According to the Noether's theorem, the invariance of the Dirac Eq. (1) under the Lorentz rotation leads to the continuity equation for the TAM current<sup>34</sup>

$$\frac{\partial}{\partial t} \mathbf{r} \cdot \mathbf{j} = 0; \quad (9)$$

where the density of the  $\mathbf{r}$ -component of the TAM is

$$\mathbf{r} \cdot \mathbf{j} = \frac{1}{2} \psi^\dagger \mathbf{J} \psi + \hbar \mathbf{x}; \quad (10)$$

and the current of the  $\mathbf{r}$ -component of the TAM is

$$\mathbf{j} = \frac{1}{2} c \psi^\dagger \mathbf{J} \psi + \hbar \mathbf{x}; \quad (11)$$

where the  $\mathbf{r}$ -component of the TAM operator  $\mathbf{J} = (\hbar/2) (\boldsymbol{\sigma} + \mathbf{r} \times \mathbf{p})$  with the position operator  $\mathbf{r}$  and the index  $\mathbf{x} = x; y; z$ . The TAM contains two parts, the spin part  $(\hbar/2) \boldsymbol{\sigma}$  and the orbit part  $\mathbf{r} \times \mathbf{p}$ . For a relativistic electron, unlike the TAM, neither the spin density nor the orbit angular-momentum density is conserved alone. Similar to the probability current, the TAM current can also be divided into the convective and internal parts<sup>35</sup>

$$\begin{aligned} \mathbf{r} \cdot \mathbf{j}_c &= \frac{1}{2} \frac{i\hbar}{2m c^2} \left( \psi^\dagger \frac{\partial}{\partial t} \mathbf{J} \psi - \frac{\partial}{\partial t} \psi^\dagger \mathbf{J} \psi \right) + \hbar \mathbf{x}; \\ \mathbf{j}_c &= \frac{1}{2} \frac{\hbar}{2m i} \left( \psi^\dagger \nabla (\mathbf{J} \psi) - \nabla \psi^\dagger \mathbf{J} \psi \right) + \hbar \mathbf{x}; \end{aligned} \quad (12)$$

$$\mathbf{r} \cdot \mathbf{j}_i = \mathbf{r} \cdot \mathbf{P}; \quad \mathbf{j}_i = c \mathbf{r} \times \mathbf{M} + \frac{\partial \mathbf{P}}{\partial t}; \quad (13)$$

where

$$\begin{aligned} \mathbf{M} &= \frac{1}{2} \frac{\hbar}{2m c} \nabla \times \mathbf{J} + \hbar \mathbf{x}; \\ \mathbf{P} &= \frac{1}{2} \frac{\hbar}{2m c} \nabla (\mathbf{r} \cdot \mathbf{j}_i) + \hbar \mathbf{x}; \end{aligned} \quad (14)$$

The convective and internal parts of the TAM current have the same structure as those of the probability current. For the NR limit,  $\rho_i$  is zero up to the zeroth order in  $(1/c)$ ,  $\mathbf{j}_i$  is only a curl term and has nothing to do with the accumulation of the TAM. Let's consider the spin

part of the convective TAM current, namely the CSC, with the density for the spin-component

$$s_c^- = \frac{\hbar^2}{4m c^2} - \frac{\partial}{\partial t} + \hbar \mathbf{z}; \quad (15)$$

and the current

$$\mathbf{j}_c^- = \frac{\hbar}{2m i} - \frac{\hbar}{2} (\mathbf{r} \cdot) \mathbf{r} - \frac{\hbar}{2} : \quad (16)$$

The CSC is determined by the rate of change of in space and time and ought to be closely related to the spin flux of the electron moving in the spatial space. Moreover, noting that the solution of the Dirac Eq. (1) is also a solution of the Klein-Gordon equation, one can easily find that the CSC satisfies the continuity equation

$$\frac{\partial s_c^-}{\partial t} + \mathbf{r} \cdot \mathbf{j}_c^- = 0; \quad (17)$$

despite that the bare spin is not a conserved quantity for the relativistic electron. Therefore, we argue that the CSC is just the correct effective spin current for discussing the spin transport properties of the relativistic electron.

The results we present above are valid for the electrons moving in the free space. The influence of the external electromagnetic field can be taken into account by making the transformation

$$\mathbf{p}' = \mathbf{p} - \frac{q}{c} \mathbf{A}; \quad \hbar \frac{\partial}{\partial t}' = \hbar \frac{\partial}{\partial t} - qV \quad (18)$$

with the charge of the electron  $q$ , the scalar potential  $V$  and the vector potential  $\mathbf{A}$ . The density and current of the convective spin in the external electromagnetic field are written as

$$s_c^- = \frac{\hbar^2}{4m c^2} - \frac{\partial}{\partial t} + \hbar \mathbf{z}; \quad \frac{\hbar}{2m c^2} qV - ; \quad (19)$$

$$\mathbf{j}_c^- = \frac{\hbar^2}{4m i} - (\mathbf{r} \cdot) \hbar \mathbf{z}; \quad \frac{q\hbar \mathbf{A}}{2m c} - : \quad (20)$$

In the external electromagnetic field, neither the TAM nor the CSC is conserved generally because of the mutual exchange of the angular momentum between the electron and the external field. The continuity-like equation of the CSC reads

$$\frac{\partial s_c^-}{\partial t} + \mathbf{r} \cdot \mathbf{j}_c^- = - (\mathbf{B} \cdot \mathbf{i} \cdot \mathbf{E}) ; \quad (21)$$

where the Bohr Magneton  $\mu_B = q\hbar/2m c$ , the magnetic field  $\mathbf{B} = \mathbf{r} \cdot \mathbf{A}$ , and the electric field  $\mathbf{E} = -(\partial \mathbf{A} / \partial t) - \mathbf{r} V$ . Although the CSC is not conserved, the source and drain of the CSC in Eq. (21) are only the external fields, while those of the bare spin include the contributions from both the external fields and the orbit angular momentum. Therefore, the CSC is still more

suitable than the bare spin current for discussing the spin transport of the relativistic electron in the external fields. For a particular case that both the electric and magnetic fields are along the same direction, namely the  $z$  direction, one can find from Eq. (21) that the  $z$ -component of the convective spin is still conserved. This particular case is relevant to the two-dimensional electron gas with the Rashba interaction where the electron is free in the  $x$ - $y$  plane and the confinement potential is only along the  $z$  direction. The  $z$ -component of the spin is also of the most interest in the Rashba system. In the following text we are restricted to this particular case, where the CSC is still conserved and can be served as the proper quantity for discussing the spin transport of the relativistic electron.

In order to apply the idea of the CSC into the spintronics, the NR limit of the CSC operator is needed. The Hamiltonian of the relativistic electron in the external electromagnetic field reads

$$H = c \quad + qV + m c^2 ; \quad (22)$$

The NR limit of the Hamiltonian (22) can be made by the Foldy-Wouthuysen transformation.<sup>34,36</sup> Up to the  $(1=c^2)$  order, the NR Hamiltonian of the electron is given as

$$\begin{aligned} H_{NR} = & \frac{1}{2m} p'^2 + qV - \frac{q\hbar}{8m^2 c^2} \mathbf{r} \cdot \mathbf{E} \\ & - \frac{q\hbar}{8m^2 c^2} (\mathbf{E} \cdot \mathbf{E}) \\ & + \frac{1}{8m^3 c^2} p'^4 + \frac{\mu_B^2}{2m c^2} \mathbf{E}^2 - \mathbf{B}^2 \\ & + \frac{\mu_B}{4m^2 c^2} (\mathbf{B} \cdot \mathbf{B})^2 + \mathbf{B}^2 (\mathbf{B} \cdot \mathbf{B}) ; \end{aligned} \quad (23)$$

and the two-component wave function, i.e., the large component of the four-component spinor in the Foldy-Wouthuysen representation corresponding the positive energy solutions, satisfies the Schrodinger equation  $\hbar (\partial / \partial t) = H_{NR}$ . The Schrodinger equation and the Hamiltonian (23) are the starting point for discussing the motion of the electron in the NR quantum mechanics including some relativistic corrections up to the  $(1=c^2)$  order. We recall the physical meanings of each term in Eq. (23). The first three terms are the NR kinetic energy, the potential energy, and the Zeeman energy arising from the electron spin, respectively. The fourth and fifth terms are the so-called Darwin term and the spin-orbit interaction, respectively. All other terms are of even higher orders.

With the help of the Hamiltonian (22) and the Dirac equation, the convective spin density in Eq. (19) can be rewritten as  $s_c^{-z} = \hat{y} \hat{s}_c^{-z} + \hbar \mathbf{z} : = 2$ , with the operator

$$\hat{s}_c^{-z} = \frac{\hbar}{2} z - \frac{\hbar}{2m c} (\quad) z ; \quad (24)$$

One can easily verify that  $\hat{s}_c^{-z}$  is an Hermitian operator and commutes with the Hamiltonian (22) and therefore is a conserved quantity. The difference between the

convective spin and the bare spin is the second term in Eq. (24) which is a higher order relativistic correction term coupling the large and small components of the four-component spinor. The NR limit of the convective spin density operator up to the  $(1=c^2)$  order, the same order as that of the Hamiltonian, is obtained by the same unitary transformations applied to the Hamiltonian (22) in the Foldy-Wouthuysen scheme. After the Foldy-Wouthuysen transformation, the convective spin density is also reduced to an even operator as the Hamiltonian does, which does not couple the large and small components of. The NR convective spin density operator acting on the two-component spinor is obtained as

$$\begin{aligned}\hat{C}_{C,NR}^{S-Z} &= \frac{\hbar}{2} z + \frac{i\hbar}{4m^2 c^2} (\hat{p}_x \hat{p}_x + \hat{p}_y \hat{p}_y) z \\ &= \frac{\hbar}{2} z + \frac{\hbar}{4m^2 c^2} \hat{C};\end{aligned}\quad (25)$$

with the operator  $\hat{C} = \frac{2}{x} + \frac{2}{y} z (\hat{p}_x \hat{p}_x + \hat{p}_y \hat{p}_y) z$ . With the help of the operator (25) the convective spin density is written as  $\hat{C}_{C,NR}^{S-Z} = \hat{C}_{C,NR}^{S-Z} + \hbar z = 2$ . Due to the unitarity of the transformation, the NR convective spin density operator Eq. (25) commutes with the Hamiltonian (23) and remains as the conserved quantity up to the  $(1=c^2)$  order. The difference between the convective spin and the bare spin is the second term in Eq. (25) which is of the order of  $(1=c^2)$  and is very small. However, the spin-orbit interaction is also originated from the relativistic correction of the electron motion up to the  $(1=c^2)$  order. Therefore, the second term in Eq. (25) can not be neglected when discussing the spin current in the spin-orbit coupled system.

Although there are many terms in the NR Hamiltonian (23), only the spin-orbit term is of particular interest on the topic of the spin Hall effect. Therefore, we remove the other relativistic corrections and rewrite the effective Hamiltonian in the form

$$H_{SO} = \frac{1}{2m} p^2 + qV + \frac{q\hbar E}{4m^2 c^2} (\hat{p}_x \hat{p}_x + \hat{p}_y \hat{p}_y) z; \quad (26)$$

It should also be noted that, in going from Eq. (23) to Eq. (26), it has been assumed that the electric field is static, i.e.,  $E = -\nabla V$ , and the external fields are along the  $z$ -direction. It can be easily verified that  $\hat{C}_{C,NR}^{S-Z}$  commutes with  $H_{SO}$  and therefore served as a conserved quantity to calculate the spin flux in the spin-orbit coupled system. With the help of Eqs. (25), (26), and the Schrodinger equation  $\hat{H} \psi = E \psi$ , the continuity equation for the CSC in the NR limit is obtained in the form

$$\frac{\partial \hat{C}_{C,NR}^{S-Z}}{\partial t} + \nabla \cdot \hat{J}_{C,NR}^{S-Z} = 0; \quad (27)$$

with the current

$$\hat{J}_{C,NR}^{S-Z} = \frac{1}{4m} \hbar \nabla \cdot \hat{C}_{C,NR}^{S-Z} + (\hat{p}_x \hat{p}_x + \hat{p}_y \hat{p}_y) z + \hbar z; \quad (28)$$

Eqs. (25), (27), and (28) constitute the main results of this paper. Comparing to the conventional spin current,<sup>7</sup> the CSC (28) has an additional correction term

$$\hat{J}_{corr} = \frac{\hbar}{16m^3 c^2} \nabla \cdot \hat{C} + (\hat{p}_x \hat{p}_x + \hat{p}_y \hat{p}_y) z + \hbar z; \quad (29)$$

which comes from the second term in Eq. (25). One finds that both the spin density and current have been modified to construct the conserved CSC. The modification, the second term in Eq. (25) and Eq. (29), is a kind of the relativistic correction resulting from the coupling between the large and small components of the four-component spinor. Those corrections are of the  $(1=c^2)$  order and are often small enough to be neglected. However, the spin-orbit interaction is also a relativistic correction of the  $(1=c^2)$  order. Therefore, the CSC ought to be used instead of the bare spin current in the spin-orbit coupled system.

Although the Hamiltonian (26) looks similar to the Rashba model<sup>37,38</sup> in the semiconductor heterostructure, a realistic situation is quite complicated. The Rashba coefficient may not be directly determined by the external electric field, but the asymmetry in the band structure parameters of the heterostructure like the effective mass.<sup>2,39</sup> The discussion in this paper is restricted to the case that the electron moves in a uniform space. Therefore, it is interesting and desirable to apply the above idea of the conserved CSC to the semiconductor heterostructures or quantum wires in future, though is highly non-trivial.

In summary, the CSC has been extracted from the TAM current by the Gordon-like decomposition in the relativistic quantum mechanics approach. The NR limit of the CSC is also derived. The CSC describes the transport properties of the electron spin and is conserved in the spin-orbit coupled system. It is also shown that the CSC remains robust in the external electromagnetic field if both the electric field and the magnetic field are along the same direction. As a result, the CSC is a good substitution for the bare spin current in studying the spin transport phenomena in spin-orbit coupled systems.

We thank J.C.Y. Teo and D.Y. Xing for valuable discussions. This work was supported by the State Key Program for Basic Research of China, the National Natural Science Foundation of China under Grant No. 10504011, the RGC funding of Hong Kong, and Seed Funding for Basic Research of the University of Hong Kong.

<sup>1</sup> S. A. Wolf, D. D. Awschalom, R. A. Buhrman, J. M. Daughton, S. von Molnar, M. L. Roukes, A. Y. Chtchelkanova, and D. M. Treger, Science 294, 1488 (2001).

- <sup>2</sup> I. Zutic, J. Fabian, and S. Das Sarma, *Rev. Mod. Phys.* **76**, 323 (2004).
- <sup>3</sup> M. I. Dyakonov and V. I. Perel, *Sov. Phys. JETP* **13**, 467 (1971).
- <sup>4</sup> J. E. Hirsch, *Phys. Rev. Lett.* **83**, 1834 (1999).
- <sup>5</sup> S. Zhang, *Phys. Rev. Lett.* **85**, 393 (2000).
- <sup>6</sup> S. Murakami, N. Nagaosa, and S. C. Zhang, *Science* **301**, 1348 (2003).
- <sup>7</sup> J. Sinova, D. Culcer, Q. Niu, N. A. Sinitsyn, T. Jungwirth, and A. H. MacDonald, *Phys. Rev. Lett.* **92**, 126603 (2004).
- <sup>8</sup> E. I. Rashba, *Phys. Rev. B* **70**, 161201(R) (2004); 201309(R) (2004).
- <sup>9</sup> J. P. Hu, B. A. Bernevig, and C. J. Wu, *Int. J. Mod. Phys. B* **17**, 5991 (2003).
- <sup>10</sup> J. Schliemann and D. Loss, *Phys. Rev. B* **69**, 165315 (2004); **71**, 085308 (2005).
- <sup>11</sup> N. A. Sinitsyn, E. M. Hankiewicz, W. Teizer, and J. Sinova, *Phys. Rev. B* **70**, 081312(R) (2004).
- <sup>12</sup> B. A. Bernevig, *Phys. Rev. B* **71**, 073201 (2005).
- <sup>13</sup> S. Q. Shen, M. Ma, X. C. Xie, and F. C. Zhang, *Phys. Rev. Lett.* **92**, 256603 (2004).
- <sup>14</sup> G. Y. Guo, Y. Yao, and Q. Niu, *Phys. Rev. Lett.* **94**, 226601 (2005).
- <sup>15</sup> J. I. Inoue, G. E. W. Bauer, and L. W. Molenkamp, *Phys. Rev. B* **70**, 041303(R) (2004).
- <sup>16</sup> E. G. Mishchenko, A. V. Shytov, and B. I. Halperin, *Phys. Rev. Lett.* **93**, 226602 (2004).
- <sup>17</sup> O. V. Dimitrova, *Phys. Rev. B* **71**, 245327 (2005).
- <sup>18</sup> O. Chalaev and D. Loss, *Phys. Rev. B* **71**, 245318 (2005).
- <sup>19</sup> S. Murakami, *Phys. Rev. B* **69**, 241202(R) (2004).
- <sup>20</sup> B. A. Bernevig and S. C. Zhang, *Phys. Rev. Lett.* **95**, 016801 (2005).
- <sup>21</sup> N. Sugimoto, S. Onoda, S. Murakami, and N. Nagaosa, *arXiv cond-mat/0503475* (2005).
- <sup>22</sup> S. Zhang and Z. Yang, *Phys. Rev. Lett.* **94**, 066602 (2005).
- <sup>23</sup> D. Culcer, J. Sinova, N. A. Sinitsyn, T. Jungwirth, A. H. MacDonald, and Q. Niu, *Phys. Rev. Lett.* **93**, 046602 (2004).
- <sup>24</sup> S. Murakami, N. Nagaosa, and S. C. Zhang, *Phys. Rev. B* **69**, 235206 (2004).
- <sup>25</sup> Q. F. Sun and X. C. Xie, *arXiv cond-mat/0502317* (2005).
- <sup>26</sup> P. Q. Jin, Y. Q. Li, and F. C. Zhang, *arXiv cond-mat/0502231* (2005).
- <sup>27</sup> P. Zhang, J. R. Shi, D. Xiao, and Q. Niu, *arXiv cond-mat/0503505* (2005).
- <sup>28</sup> J. Wang, B. G. Wang, W. Ren, and H. Guo, *arXiv cond-mat/0507159* (2005).
- <sup>29</sup> Y. Wang, K. Xia, Z. B. Su, and Z. Ma, *arXiv cond-mat/0508340* (2005).
- <sup>30</sup> Y. K. Kato, R. C. Myers, A. C. Gossard, and D. D. Awschalom, *Science* **306**, 1910 (2004).
- <sup>31</sup> J. Wunderlich, B. Kestner, J. Sinova, and T. Jungwirth, *Phys. Rev. Lett.* **94**, 047204 (2005).
- <sup>32</sup> W. Gordon, *Z. Phys.* **50**, 630 (1928).
- <sup>33</sup> G. Baym, *Lectures on Quantum Mechanics* (Benjamin, New York, 1969).
- <sup>34</sup> F. Schwabl, *Advanced Quantum Mechanics*, translated by R. Hilton and A. Lahee (Springer, Berlin, 2004).
- <sup>35</sup> F. W. Hehl, A. Macias, E. W. Mielke, Y. N. Obukhov, *arXiv gr-qc/9706009* (1997).
- <sup>36</sup> L. L. Foldy and S. A. Wouthuysen, *Phys. Rev.* **78**, 29 (1950).
- <sup>37</sup> E. I. Rashba, *Sov. Phys. Solid State* **2**, 1109 (1960).
- <sup>38</sup> Y. A. Bychkov and E. I. Rashba, *J. Phys. C* **17**, 6039 (1984).
- <sup>39</sup> P. Pfeiffer and W. Zawadzki, *Phys. Rev. B* **52**, 14332(R) (1995); **59**, 5312(R) (1999).