

Pomeranchuk and Topological Fermi Surface Instabilities from Central Interactions

J. Quintanilla

ISIS Spallation Facility, CCLRC Rutherford Appleton Laboratory, Chilton, Didcot, OX11 0QX, U.K.

A. J. Schofield

School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham, B15 2TT, U.K.

We address at the mean field level the emergence of a Pomeranchuk instability in a uniform Fermi liquid with central particle-particle interactions. We find that Pomeranchuk instabilities with all symmetries except $l=1$ can take place if the interaction is repulsive and has a finite range r_0 of the order of the inter-particle distance. We demonstrate this by solving the mean field equations analytically for an explicit model interaction, as well as numerical results for more realistic potentials. We find in addition to the Pomeranchuk instability other, subtler phase transitions in which the Fermi surface changes topology without rotational symmetry-breaking. We argue that such interaction-driven topological transitions may be as generic to such systems as the Pomeranchuk instability.

I. INTRODUCTION

Experimental evidence of "hidden" phases of itinerant electron systems^{1,2,3} and the prospect of realizing novel conditions in layered heterostructures and ultra-cold gases have led to increased efforts to identify unconventional phase transitions and predict their manifestations. To give three examples: the Fulde-Ferrell-Larkin-Ovchinnikov state has been proposed in organic superconductors⁴, superconductor-ferromagnet heterostructures⁵ and in imbalanced mixtures of ultracold atoms^{6,7}; a supersolid phase is a possibility in Bose gases loaded on optical lattices⁸; and a "d-density wave" may be realized in ladder compounds^{10,11} and possibly "hide" in the phase diagram of cuprate superconductors⁹, where other hidden order parameters have been proposed^{47,48}.

In this context there has been a surge of interest in the Pomeranchuk Instability (PI)¹². Through it a Fermi liquid may enter a "nematic" state characterized by a deformed Fermi surface. It has been argued that such an instability may take place in quantum Hall systems^{13,14} and in the metamagnets $\text{Sr}_2\text{Ru}_2\text{O}_7$ ³ and URu_2Si_2 ¹⁵. Moreover there is evidence that the Hubbard model has a phase with a distorted Fermi surface^{16,17,18,19,20,21} and the Emery model of a CuO_2 plane has been shown to have a nematic ground state in the strong coupling limit²².

More generally the PI is an interesting candidate unconventional phase transition on account of its subtlety. Thus, considerable effort is going into characterizing it theoretically^{23,24,25,26,27,28,29,30} on the basis of phenomenological models featuring anisotropic effective interactions. This approach is proving very successful in establishing some generic features of the phase diagram^{23,24,25} and describing collective excitations and quantum critical fluctuations^{26,27,28,29,30}. On the other hand it sidelines the question of how the anisotropy emerges in the first place²⁶ and what other³¹, perhaps even subtler instabilities may generically arise in such contexts. It is these questions that we address here.

In this paper we present a mean field (MF) theory of the PI in a three-dimensional, uniform fermion liquid with a central effective interaction potential $V(r)$. The authors of Ref. 26 have pointed out that such interaction may lead to a PI. Here we show that the emergence of the anisotropic state from a Galilean invariant fluid requires repulsion with an intermediate range of the order of the inter-particle distance. This is confirmed by explicit calculation for a model interaction potential for which the theory can be solved analytically. However we also find that the intermediate-range repulsion leads, quite generally, to a different instability in which there is no symmetry breaking but the topology of the Fermi surface changes. We discuss the nature of this subtler quantum phase transition. A few instances of the two distinct types of Fermi surface shape instabilities that we find are pictured in Fig. 1. These two types of instability compete and we show that this conclusion is robust when we consider more realistic finite range interactions.

II. MEAN FIELD THEORY

To motivate a microscopic theory of the PI we start by recalling the original, phenomenological theory due to Pomeranchuk¹². Like him, we start with an unpolarized Fermi sphere and consider an infinitesimal change of the occupation numbers, $N_k \rightarrow N_k + \delta N_k$; arising from an angle-dependent modulation of the Fermi vector, $k_F \rightarrow k_F + \delta k_F(\hat{k})$, $k_F \rightarrow k_F + \delta k_F(\hat{k})$ in the symmetric or antisymmetric spin channel, respectively⁴⁹. We then use Landau's expression for the corresponding change in the ground state energy: $E \rightarrow E + \delta E$; with

$$E = \sum_k \epsilon(k) N_k + \frac{1}{2} \sum_{k,k^0} f^s(k;k^0) N_k N_{k^0} + f^a(k;k^0) S_k \cdot S_{k^0} \quad (1)$$

the phenomenological functions $\epsilon(k)$ (which here may depend on the spin) and $f^{s;a}(k; k^0)$ are now derived from our microscopic parameters via Eq. (8) and

$$f^{s;a}(k; k^0) = \frac{1}{2} V - \frac{1}{2} V(k - k^0); \quad (12)$$

where $\epsilon = 1; 0$ in the $s; a$ channels, respectively.

Let us pause briefly to note the following subtlety. Eqs. (9,10), from which all the subsequent results follow, could have been derived by minimizing the energy of the trial state given in Eq. (11). However note this only determines the Fermi surface, but it under-determines the dispersion relation $\epsilon(k)$. The justification of the particular form given in Eq. (8) thus relies on the assumption that the low-lying excitations correspond to rearrangements of the electrons in momentum space, whose energy is given by Eq. (1) [or, equivalently, that the equilibrium state at finite temperatures can be adequately described by the mean field Hamiltonian of Eq. (5)]. This is necessary to justify the language we use below, e.g. in defining the Fermi velocity in terms of $\epsilon(k)$. We stress, however, that the results themselves refer only to the equilibrium shape of the Fermi surface in the $l=0$ ground state and are therefore more general, and independent of the meaning assigned to $\epsilon(k)$.

To study the PI in our microscopic model we postulate an unpolarized, spherical Fermi surface

$$N_{k; \sigma} = (k_F - k)^2; \quad (13)$$

completely described by the Fermi vector $k_F > 0$ [Fig. 1 (a)], and use the above equations to determine whether the system has a PI. In the state described by Eq. (13), the electron dispersion relation of Eq. (8) is given by

$$\epsilon(k) = \frac{\hbar^2 k^2}{2m} - \frac{k_F^2}{2} + \frac{2k_F^2}{0} \int_0^1 dr r^2 V(r) j_1(k_F r) j_0(k - k_F) j_0(k_F r); \quad (14)$$

This yields the following expression for the Fermi velocity:

$$\begin{aligned} v_F &= \frac{\hbar}{m} k_F + \frac{2k_F^2}{m} \int_0^1 dr r^2 V(r) j_1(k_F r)^2 \\ &= \frac{\hbar}{m} k_F + \frac{P_F^2}{(2\hbar)^3} V_1; \end{aligned} \quad (15)$$

where in the second line we have expressed v_F in terms of one of the coupling constants defined by Eq. (18), below. Note that the state described by Eq. (13) requires $v_F > 0$. Together Eqs. (12) and (15) give the Landau parameters in Eq. (3),

$$F_1^{a;s} = \frac{21 + \frac{1}{2} \frac{k_F^2}{\hbar^2 v_F}}{(2\hbar)^3 \hbar v_F} \approx 0.8 \frac{V}{V_1}; \quad (16)$$

in terms of the microscopic parameters of the model. This, in turn, allows us to express the PI equations as

$$V_1 \approx 0.8 \frac{V}{V_1} > \frac{(2\hbar)^3}{P_F^2} v_F; \quad = \frac{(2\hbar)^3}{m P_F} + V_1 \quad (17)$$

where the strength of the interaction potential in a given angular momentum channel $l = 0; 1; 2; \dots$ is given by

$$V_l = \frac{4\pi}{0} \int_0^1 dr r^2 V(r) j_l(k_F r)^2; \quad (18)$$

For antisymmetric instabilities, $\epsilon = 0$, Eq. (17) takes the following, explicit form:

$$\frac{4\pi}{0} \int_0^1 dr r^2 V(r) j_1(k_F r)^2 > \frac{2\hbar^2}{m k_F}; \quad (19)$$

Eq. (17) is our microscopic expression of the PI condition of Eq. (2). It is valid, within our MF ansatz of Eq. (5), for any system whose Hamiltonian has the form given by Eq. (4). From it we can derive a series of conclusions concerning a Pomerenchuk instability in an isotropic system with central interactions:

1. Purely attractive interactions can only lead to the gas-liquid transition ($l = 0; \epsilon = 1$)⁵¹; conversely, purely repulsive interactions can only lead to the Stoner or PI.
2. There are no $l = 1$ Pomerenchuk instabilities. This is the type of PI^{15,31,36} where rotational symmetry-breaking is achieved by displacing the Fermi surface so as to set up a charge (s) or spin (a) current. This is quite a general consequence of the well known relation between the effective mass and the Landau parameter, F_1^s , in a Galilean invariant system, which is captured by Eqs. (15) and (16). On the other hand for spin-dependent interactions (or in lattice systems), not considered here, we may have $F_1^s \neq F_1^a$ and then the instabilities considered in Refs.^{15,31,36} could be realized.
3. The PI for $l = 2$ is degenerate in the spin channel. These instabilities break rotational symmetry by changing the shape of the Fermi surface, without generating any currents of charge or spin [see Fig. 1 (b,c)]. Our result implies that, at the instability, it does not matter whether the lobes of the spin-up and spin-down Fermi surface point in the same direction⁵². Note this is quite different from the situation at $l = 0$ (see point 1, above).
4. Finally, from Eq. (18) we can also deduce that $V_1 - V_l$ cannot be large and positive, as required by Eq. (17), if the repulsive part of the interaction is of very short range $r_0 \ll k_F^{-1}$: In effect, $r^2 j_1(k_F r)^2 \sim r^{2(l+1)}$ for $r \ll k_F^{-1}$ so for such short-ranged interactions Eq. (18) gives $V_l \sim \int_0^{r_0} dr r^{2(l+1)} \sim r_0^{2l+3}$ whence for small r_0 Eq. (17) can only be satisfied for $l = 0$. The extreme case of this is the repulsive contact potential $V(\vec{r}) = \delta(\vec{r})$; for which $r_0 = 0$ and $V_1 = 4\pi \int_0^1 dr r^2 \delta(r) = 0$. For this potential, our theory leads only to the Stoner instability.




Figure 2: (color online) Phase diagram showing zero-temperature instabilities of the types depicted in Fig. 1 for the repulsive delta-shell model. The Stoner instability occurs above the solid line. The other instabilities described in Fig. 1 occur above the long dash line (b), medium dash line (c), short dash line (d), long dash-dot line (e) and short dash-dot line (f). The colors correspond with Fig. 1

III. DELTA-SHELL MODEL

From point 4 we conclude that repulsive interactions with range at least of the order of the Fermi wavelength, $r_0 \propto k_F^{-1}$; are necessary for the PI in an isotropic system with isotropic interactions. We investigate this further by choosing a specific form of the central interaction potential, namely the \delta-shell" potential:

$$V(\mathbf{r}) = g \delta(\mathbf{r} - r_0) \quad (20)$$

This is an idealization of an interaction with a very sharp peak at a particular inter-particle distance, $r = r_0$: The \coupling constant" g has dimensions of energy \times length and represents the product of the height and width of the potential barrier.

For $g < 0$; this interaction potential can lead to superconductivity with unconventional pairing^{37,38}. Likewise, we expect that for $g > 0$ it will lead to a PI. Indeed Eq. (18) gives

$$V_1 = (4\pi)^2 g r_0^2 j_1(k_F r_0)^2 \quad (21)$$

so, depending on the value of $k_F r_0$; any value of l may become dominant.

The particularly simple form of the interaction potential in Eq. (20) allows us to write the key expressions in our theory of the PI analytically. In particular Eq. (15) giving the Fermi velocity on the Fermi sphere reads

$$v_F = \frac{\hbar}{m} k_F + g \frac{2(k_F r_0)^2}{\hbar} j_1(k_F r_0)^2 \quad (22)$$

Together with Eq. (21) and $V = g 4\pi r_0^2$ these equations allow us to write the following, simple form of the Stoner

and PI equations:

$$j_1(k_F r_0)^2 - j_1(k_F r_0)^2 > \frac{\hbar^2}{2m r_0^2 g k_F}; l = 0; 1; 2; \dots \quad (23)$$

If the interaction is very strong, $g k_F \hbar^2 = 2m r_0^2$; this gives a sequence of fixed phase boundaries at $j_1(k_F r_0) = j_1(k_F r_0)$: In the opposite limit of very weak interaction the unpolarized Fermi sphere is, as expected, stable.

The solid, long dash and medium dash lines of Fig. 2 are the phase diagram obtained by solving Eq. (23) for $l = 0, 1, 2$. Note that there are effectively only two, dimensionless parameters in the theory^{37,38}: the \effective range" $k_F r_0$ and \coupling constant" $g = r_0 \epsilon_0$ (where $\epsilon_0 = \hbar^2 / 2m r_0^2$). The first of these parameters is the range of the interaction measured in units of $l = k_F$. The second is the product of the width of the potential barrier measured in units of that range and its height measured in units of the corresponding \localization energy" ϵ_0 . As expected, for small range, $k_F r_0 \lesssim 3$; we only find the Stoner instability. For longer ranges or, equivalently, higher densities, and sufficiently large values of the dimensionless coupling constant (which we note that depends not only on g but also on m and r_0), the PI can take place.

IV. TOPOLOGICAL TRANSITIONS

In addition to the anticipated PI, our analytic treatment of the delta-shell model also reveals a competing class of Fermi surface instabilities which, unlike the PI, occur without symmetry breaking. Consider the electron dispersion relation in the isotropic state, Eq. (14). For the delta-shell potential it takes the form

$$\epsilon(k) = \frac{\hbar^2}{2m} k^2 - \epsilon_F + \frac{g}{r_0} \frac{2k_F^2 r_0^2}{j_1(k_F r_0)} j_1(k r_0) j_1(k_F r_0) \quad (24)$$

This is plotted in Fig. 3 for three different values of $k_F r_0$. The free-electron dispersion relation is modified by an oscillatory term due to electron-electron interactions. The period of the oscillations is r_0^{-1} . For small $g = r_0 \epsilon_0$ the effect of these is the usual renormalization of the effective mass $m^* = \hbar^2 / v_F$, which follows from Eq. (22). However at large $g = r_0 \epsilon_0$ the effect of interaction on this \bare" dispersion relation cannot be described simply as a renormalization of m . In fact it can lead to a dramatic change of the state of the system as the amplitude of the oscillations becomes large enough that either

1. the dispersion relation dips below the Fermi level somewhere outside the Fermi sphere [Fig. 3 (a)],
2. it goes above the Fermi level at the center of the Fermi sphere [Fig. 3 (b)] or
3. it peaks above the Fermi level at some intermediate k , $0 < k < k_F$ [Fig. 3 (c)].




Figure 3: Electron dispersion relation for the "delta-shell" model in the state with an unpolarized Fermi sphere: (a) $k_F r_0 = 4$, $g/r_0 \epsilon_0 = 0$ (solid); 20 (long dash); 97.66 (short dash); 150 (dotted); (b) $k_F r_0 = 6$, $g/r_0 \epsilon_0 = 0$; 2; 8.84; 16 (same order); (c) $k_F r_0 = 9$, $g/r_0 \epsilon_0 = 0$; 20; 41.40; 60.

In either case, Eq. (10) no longer reduces to Eq. (13). Instead, either a thin shell of occupied states forms outside the Fermi sphere [Fig. 1 (d)], or states inside the Fermi sphere become vacated [Fig. 1 (e,f)]. The associated instabilities are quite distinct from the Stoner and PI, as the change N_k ; although infinitesimal, takes place away from the Fermi surface. Instead, they are continuous phase transitions in which no symmetry is broken, but the topology of the Fermi surface changes. In that sense they are more reminiscent of the Lifshitz




Figure 4: Evolution of the Fermi vector as a function of the dimensionless coupling constant $g/r_0 \epsilon_0$ for fixed particle density $N = 9 r_0^{-3}$. The plot shows both the "original" Fermi vector k_F and the "emerging" Fermi vector k_F' , namely the radius of the sphere of empty states depicted in Fig. 1 (e), as the system enters from below the corresponding dome in the phase diagram (see Fig. 2).

transition^{39,40}. There is, however, a crucial difference, namely that the present phase transitions are driven by electron-electron interactions, which induce the fermions to "migrate" to other regions of reciprocal space, rather than by the band structure. On the basis of this one would expect the present instabilities to have a much stronger thermodynamic signature. For example, the transitions illustrated in Fig. 1 (d) and (f), with their underlying dispersions of Fig. 3 (a) and (c) respectively, result in the appearance of entire new Fermi surface sheets with finite k_F . This will lead in mean-field theory to a discontinuous jump in the density of states and hence in C/T .

A general framework to understand such topological quantum phase transitions has been put forward in Ref. 41. In this formalism, the Fermi surface is a vortex "loop" in a four-dimensional space and the phase transitions we have just described correspond to the nucleation of new loops. One can also view these instabilities as generalizations to dimension larger than one of the phenomenon of "quantum Hall edge reconstruction". The latter can be described as the emergence, due to interactions, of new Fermi points in the one-dimensional chiral Fermi liquid on the edge of a quantum Hall system^{42,43,44,45}. Indeed in the instabilities described here always one of the new Fermi surfaces has negative Fermi velocity – analogous to the creation of left-moving quasiparticles in a right-moving chiral Fermi liquid⁵³. Yang and Sachdev⁴⁶ have recently described the quantum critical fluctuations for a phase transition of type 1, above [Figs. 1 (d) and 3 (a)].

It is important to note that the plots in Fig. 3 correspond to evaluating Eq. (24) at a fixed value of k_F . For a fixed number of particles, N , such solutions are valid only up to the instability, as beyond it they would violate Luttinger's theorem. To describe the migration of elec-




Figure 5: (color online) Phase diagram showing zero-temperature instabilities of the unpolarized, spherical Fermi surface for the "hard-core" model. The coupling constant is measured in units of the Fermi energy $\epsilon_F = \hbar^2 k_F^2 / 2m$. Key as in Fig. 2.

trons in reciprocal space mentioned above, which happens beyond the instability, it is necessary to determine k_F self-consistently by requiring that the total number of particles be fixed. For example, for an instability of the type 2, above Figs. 3 (b) and 1 (e)], this means that

$$\frac{N}{\Omega} = 2 \frac{1}{(2\pi)^3} \frac{4}{3} k_F^3 - k_F^0{}^3; \quad (25)$$

where k_F^0 is the "emerging" Fermi vector at the center of the Fermi sphere, determined by $\epsilon(k_F^0) = 0$. This is demonstrated in Fig. 4. Note that the way k_F^0 grows as a function of coupling suggests thinking of this quantity as a sort of topological "order parameter".

It is evident from Fig. 3 that the new sheet of electron-like or hole-like Fermi surfaces are initially formed by localized states, with $v_F = 0$: As we progress into the new state, v_F becomes finite. Conversely, if we run the process backward, the effective masses on the additional Fermi surfaces diverge (except for the hole-like Fermi surface in Fig. 1 (e), for which the Fermi vector goes to zero at the same time as the Fermi velocity).

The short dash, long dash-dot and short dash-dot lines on Fig. 2 show the boundaries of these "Fermi surface topology transitions". Notably, except for a range of densities near where the $l=1$ PI would have been, the unpolarized Fermi sphere is only stable at small coupling

V. HARD-CORE MODEL

Our results suggest that the essential ingredient of the PI in the present context, namely the finite range $r_0 \propto k_F^{-1}$, also leads to the interaction-driven Lifshitz transition. To probe the generality of this observation, we have repeated the calculation (this time by evaluating

the mean-field equations (14) and (17) numerically) for a repulsive "hard core" potential

$$V(\mathbf{r}) = V(r_0 - |\mathbf{r}|); \quad (26)$$

We have again found that, for $r_0 \propto k_F^{-1}$; there are, in addition to the Stoner instability, PI with $l=2;4;6;\dots$. Moreover we also find an interaction-driven Lifshitz transition which, for certain ranges of values of $k_F r_0$, takes place before the Stoner or PI set in. A phase diagram is presented in Fig. 5.

Unlike the delta-shell potential, for the hard core potential the PI domes are contained within the Stoner ones, i.e. the PI can only take place on a polarized Fermi surface, or for spinless fermions. However note that there are regions of the phase diagram where the boundary of the $l=2$ instability nearly overlaps with that for $l=0$, indicating that the two transitions happen almost simultaneously.

The plot does not show all the instabilities: the $l > 2$ PI take place at higher values of $k_F r_0$ than those shown. There are also other domes of topological instability, though for this potential all of them are of the type in Fig. 1 (e).

Our results for the hard-core potential not only support our identification of a sharp feature of $V(r)$ at $r_0 \propto k_F^{-1}$ as the crucial ingredient for a PI, but suggest as well that, in such situations, the interaction-driven Lifshitz transition is at least as likely to occur as the Stoner or PI.

We conclude this section by noting that a similar analysis using the screened Coulomb interaction

$$V(\mathbf{r}) = \frac{e^2}{4\pi\epsilon_0} \frac{e^{-|\mathbf{r}|/r_0}}{|\mathbf{r}|} \quad (27)$$

does not reveal either PI or instabilities of the Fermi surface topology — only the Stoner instability is realized, and that only if we allow r_0 to deviate from its Thomas-Fermi value.

VI. CONCLUSIONS

In summary we have studied, at the mean-field level, PI of a uniform Fermi liquid with central fermion-fermion interactions. We find that PI of different symmetries may emerge from repulsive interactions of sufficiently long, but finite range $r_0 \propto k_F^{-1}$ (with the interesting exception of the $l=1$ PI which never takes place). We have confirmed this by solving the theory analytically for an explicit form of the interaction potential featuring repulsion at a particular distance r_0 . Surprisingly we have found that, in addition to the PI, there is also a new type of Fermi surface instability: the interaction-driven Lifshitz transition. This topological phase transition is even subtler than the PI and seems to be generically associated with the class of models leading to the PI. Further support for this picture is provided by analysis of an additional

model, featuring hard-core repulsion. On the other hand the screened Coulomb interaction does not lead to these effects suggesting that a sharp feature (either a spike of repulsion or a sudden drop) must be present at the distance r_0 .

Unlike the Lifshitz transition, the new quantum phase transition that we have described is fundamentally driven by interactions. Thus one would expect it to have a stronger thermodynamic signature. It will be of great interest, in the near future, to establish this signature and the properties of the novel state of matter this phase transition may lead to.

Acknowledgments

We acknowledge discussions with J.M.F. Gunn, C. Hooley, M. Haque, B. J. Powell, S. Simon, J.F. Chalker, N.I. Gidopoulos, M.W. Long, G. Volovik, A.H. Castro-Neto, E. Fradkin, S. Ramos and W.J.L. Buyers. JQ thanks the University of Birmingham for hospitality and acknowledges financial support by the Leverhulme Trust and an Atlas fellowship awarded by CCLRC in association with St. Catherine's College, Oxford.

- ¹ J.L. Tallon et al, Phys. Stat. Sol. (b) 215, 531 (1999).
- ² T.T.M. Palstra et al, Phys. Rev. Lett. 55, 2727 (1985); M.B. Walker et al, Phys. Rev. Lett. 71, 002630 (1993); K.H. Kim et al, Phys. Rev. Lett. 91, 256401 (2003).
- ³ S.A. Grigera et al, Science 306, 1154 (2004).
- ⁴ L. Balicas et al, Phys. Rev. Lett. 87, 067002 (2001).
- ⁵ J. F. Annett, M. Krawiec and B. L. Gyor'y, cond-mat/0510591.
- ⁶ T.M. Izushima et al, Phys. Rev. Lett. 94, 060404 (2005).
- ⁷ Kun Yang, Phys. Rev. Lett. 95, 218903 (2005).
- ⁸ K. Goral, L. Santos and M. Lewenstein, Phys. Rev. Lett. 88, 170406 (2002).
- ⁹ S. Chakravarty, R.B. Laughlin, D.K. Morr and C. Nayak, Phys. Rev. B 63, 094503 (2001).
- ¹⁰ U. Schollwöck et al, Phys. Rev. Lett. 90, 186401 (2003).
- ¹¹ S. Lee, J.B. Marston and J.O. Fjærestad, Phys. Rev. B 72, 075126 (2005).
- ¹² I. A. Pomeranchuk, JETP 35, 524-525 (1958).
- ¹³ E. Fradkin and S.A. Kivelson, Phys. Rev. B 59, 8065 (1999).
- ¹⁴ E. Fradkin, S.A. Kivelson, E. Manousakis and K. Nho, Phys. Rev. Lett. 84, 1982 (2000).
- ¹⁵ C.M. Vamä and Lijun Zhu, cond-mat/0502344.
- ¹⁶ C.J. Halboth and W. Metzner, Phys. Rev. Lett. 85, 5162 (2000).
- ¹⁷ V. Hankevych, I. Grote and F. Wegner, Phys. Rev. B 66, 094516 (2002).
- ¹⁸ W. Metzner, D. Rohe and S. Andergassen, Phys. Rev. Lett. 91, 066402 (2003).
- ¹⁹ A. Neumayr and W. Metzner, Phys. Rev. B 67, 035112 (2003).
- ²⁰ E.C. Carter and A.J. Schofield, Phys. Rev. B 70, 045107 (2004).
- ²¹ The PImay also be favored by non-linearities in the electron dispersion relation: D.G. Barci and L.E. Oxman, Phys. Rev. B 67, 205108 (2003).
- ²² S.A. Kivelson, E. Fradkin and T.H. Geballe, Phys. Rev. B 69, 144505 (2004).
- ²³ Hae-Young Kee and Yong Baek Kim, Phys. Rev. B 71, 184402 (2005).
- ²⁴ I. Khavkine, Chung-Hou Chung, V. Oganesyan and Hae-Young Kee, Phys. Rev. B 70, 155110 (2004).
- ²⁵ H. Yamase, V. Oganesyan and W. Metzner, Phys. Rev. B 72, 35114 (2005).
- ²⁶ V. Oganesyan, S.A. Kivelson and E. Fradkin, Phys. Rev. B 64, 195109 (2001).
- ²⁷ Hae-Young Kee, Phys. Rev. B 67, 073105 (2003).
- ²⁸ J. Nilsson and A.H. Castro Neto, Phys. Rev. B 72, 195104 (2005).
- ²⁹ L. Dell'Anna and W. Metzner, cond-mat/0507532.
- ³⁰ M. J. Lawler et al, cond-mat/0508747.
- ³¹ C.M. Vamä, Phil. Mag. 85, 1657-1666 (2005).
- ³² A.J. Leggett, Rev. Mod. Phys. 47, 331 (1975).
- ³³ S.A. Kivelson, E. Fradkin and V.J. Emery, Nature 393, 550 (1998).
- ³⁴ P. Nozières and S. Schmitt-Rink, J. Low. Temp. Phys. 59, 195 (1985).
- ³⁵ P. Nozières, J. Phys. I France 2, 443-458 (1992).
- ³⁶ Congjun Wu and Shou-Cheng Zhang, Phys. Rev. Lett. 93, 36403 (2004).
- ³⁷ J. Quintanilla and B.L. Gyor'y, Physica B 284-288, 421-422 (2000).
- ³⁸ J. Quintanilla, B.L. Gyor'y, J. F. Annett and J. P. Wallington, Phys. Rev. B 66, 214526 (2002).
- ³⁹ I.M. Lifshitz, Zh. Eksp. Teor. Fiz. 38, 1569 (1960) [Sov. Phys. JETP 11, 1130 (1960)].
- ⁴⁰ A.A. Abrikosov, "Fundamentals of the Theory of Metals" (Elsevier, New York, 1988).
- ⁴¹ G. Volovik, cond-mat/0601103.
- ⁴² A.H. MacDonald, S.R. Eric Yang and M.D. Johnson, Aust. J. Phys. 46, 345 (1993).
- ⁴³ C. de C. Chamon and X.G. Wen, Phys. Rev. B 49, 8227 (1994).
- ⁴⁴ Xin Wan, Kun Yang and E.H. Rezayi, Phys. Rev. Lett. 88, 056802 (2002).
- ⁴⁵ A.H. Castro Neto, F. Guinea and N.M.R. Peres, Phys. Rev. B 73, 205408 (2006).
- ⁴⁶ Kun Yang and Subir Sachdev, Phys. Rev. Lett. 96, 187001 (2006).
- ⁴⁷ C.M. Vamä, Phys. Rev. B 55, 14554 (1997); Phys. Rev. Lett. 83, 3538 (1999).
- ⁴⁸ B. Fauque, Y. Sidis, V. Hinkov, S. Pailhes, C.T. Lin, X. Chaud, and P. Bourges, Phys. Rev. Lett. 96, 197001 (2006).
- ⁴⁹ We assume a k -independent quantization axis for the spin and choose this to be in the z direction. One could envisage richer states in which such axis cannot be defined, for example analogous to the Balian-Werthamer superconducting state³². A state of this type has been discussed in Ref. 36.
- ⁵⁰ We follow the notations of Ref. 32.
- ⁵¹ Our M F Hamiltonian H_0 lacks any term hybridizing par-

ticles with holes and therefore cannot describe the key instability present for purely attractive interactions, namely superconductivity. When this is included, all other instabilities are precluded^{34,35}.

⁵² Inside the region of instability, higher order terms in the free energy decide between these configurations.

⁵³ A similar phase transition has also been described in S. A. Artamonov, V. R. Shaginyan and Yu. G. Pogorelov, JETP Lett. 68, 942 (1998). It has also been suggested that the opening of a full gap to single-particle excitations on approach to the Bose-Einstein condensation limit in a d-

wave superconductor should be regarded as an interaction-driven Lifshitz transition: S. S. Botelho and C. A. R. Sade Melo, Phys. Rev. B. 71, 134507 (2005). A change of topology of the Fermi surface has also been noticed in the RVB state on a frustrated square lattice [J. Powell, private communication], for fermions on a lattice with boson-controlled hopping [D. M. Edwards, Physica B 378-380, 133-134 (2006)] and in the pseudogap region of the cuprate phase diagram [C. M. Varma and Lijun Zhu, cond-mat/0607777].