

Quantum magnetism with multicomponent dipolar molecules in an optical lattice

Ryan Barnett¹, Dmitry Petrov^{1,2}, Mikhail Lukin¹, and Eugene Demler¹

¹Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA

²ITAMP, Harvard-Smithsonian Center for Astrophysics, Cambridge, Massachusetts 02138, USA
(Dated: March 23, 2024)

We consider bosonic dipolar molecules in an optical lattice prepared in a mixture of different rotational states. The $1/r^3$ interaction between molecules for this system is produced by exchanging a quantum of angular momentum between two molecules. We show that the Mott states of such systems have a large variety of quantum phases characterized by dipolar orderings including a state with ordering wave vector that can be changed by tilting the lattice. As the Mott insulating phase is melted, we also describe several exotic superfluid phases that will occur.

In ultracold physics, systems with long-range dipolar interactions have recently attracted considerable attention both theoretically and experimentally (for a recent review of ultracold dipolar molecules see [1] and references therein). For atoms, dipolar interactions come from their magnetic moments and become important for large electronic spin [2]. Recent experiments demonstrated the relevance of such dipolar interactions for the expansion of Cr atoms from the BEC state [3]. On the other hand, for heteronuclear molecules, dipolar interactions arise from their electric dipole moments. Recent experiments have succeeded in trapping and cooling several types of heteronuclear molecules [1, 4, 5, 6]. In a state with a well-defined angular momentum, molecules do not have a dipole moment. However, when an external electric field is used to polarize the molecules, dipole moments can be induced. There has been considerable theoretical effort to study the resulting dipole interactions and many-body physics associated with such systems [8, 9, 10, 11, 12].

In this Letter, we consider an alternative mechanism for obtaining the $1/r^3$ dipolar interactions, and the important concomitant directional character. Namely, we investigate a mixture of heteronuclear dipolar molecules in the lowest ($N = 0$) and the first excited ($N = 1$) rotational states. For such a system, the origin of the long-range interaction is the exchange of angular momentum quanta between molecules. We demonstrate that when loaded into an optical lattice, such mixtures can realize various kinds of non-trivial effective dipolar spin systems with anisotropic, long-range interactions. Several approaches for realizing spin systems using cold atoms have been discussed before, including bosonic mixtures in optical lattices in the Mott state [13, 14, 15, 16], interacting fermions in special lattices [17], and trapped ions interacting with lasers [18]. The system we consider has the practical advantages of the high energy scale for spin-dependent phenomena (set by dipolar interactions) and the new physics associated with the long-ranged nature of the dipole interactions. Experimental realization of the system will give insight into several open questions in condensed matter physics including competition between ferro and antiferroelectric orders in crystals [19, 20] and systems with frustrated spin interactions [21].

Consider the system that contains bosonic molecules in the lowest ($N = 0; N_z = 0$) and first excited ($N = 1, N_z = 1; 0; 1$) rotational states where we let s^y and $t^y_{1;0;1}$ create these respective states. We will often use the change of basis $t^y_x = (t^y_1 + t^y_{-1})/\sqrt{2}$, $t^y_y = i(t^y_1 - t^y_{-1})/\sqrt{2}$, and $t^y_z = t^y_0$. To describe molecules in an optical lattice we use the one-band Hubbard type effective model

$$H = H_{\text{kin}} + H_{\text{Hub}} + H_{\text{dip}}: \quad (1)$$

The first term on the right hand side of (1) is the kinetic energy from nearest-neighbor hopping $H_{\text{kin}} = \sum_{\langle ij \rangle} J_{ij} s^y_i s^y_j + t^y_i t^y_j + \text{h.c.}$. Operators s^y_i and t^y_i create molecules on site i (here and after the summation over repeating indices $i = x, y, z$ is implied). The last term in (1) describes the dipolar interaction between molecules from different sites

$$H_{\text{dip}} = \frac{1}{2} \sum_{i \neq j} \frac{d_i d_j}{R_{ij}^3} \frac{3\mathbf{d}_i \cdot \mathbf{e}_{ij} d_j \cdot \mathbf{e}_{ij}}{R_{ij}^3} \quad (2)$$

where R_{ij} are lattice vectors, \mathbf{e}_{ij} is the i -component of the unit vector along R_{ij} , and parameter d equals $2d^2/3$, where d is the value of the dipole moment associated with the $N = 0 \rightarrow N = 1$ transition, and d_i is the dipole moment operator at site i . The i -component of the operator d_i is written as

$$d_i = s^y_i t^y_i e^{2iB_e t} + t^y_i s^y_i e^{2iB_e t}, \quad (3)$$

where we absorbed the energy difference between the rotational levels $E_{N=1} - E_{N=0} = 2B_e$ into the time dependence of the t operators. Since the rotational constant B_e is considerably larger than any other energy scale in the system, we assume that the terms in (2) that oscillate at frequencies $\pm 4B_e$ average to zero. This forces the number of molecules in the $N = 0$ and $N = 1$ states to be independently conserved. Then (2) reduces to

$$H_{\text{dip}} = \frac{1}{2} \sum_{i \neq j} \frac{(s^y_i t^y_j s^y_j t^y_i + \text{h.c.})}{R_{ij}^3} \frac{3\mathbf{e}_{ij} \cdot \mathbf{e}_{ij}}{R_{ij}^3}: \quad (4)$$

The second term on the right hand side in (1) is the Hubbard on-site interaction. For two molecules in

the absence of an external electric field, the long-range part of their interaction potential is dominated by the van der Waals tail $C_6=R^{-6}$ originating from second order terms in the dipole-dipole interaction operator [22]. For polar molecules with large static rotational polarizabilities one can estimate $C_6 \approx d^4/6B_e$. For the RbCs molecule ($d = 0.5$ a.u., $B_e = 7.7 \cdot 10^8$ a.u.) we have $C_6 \approx 1.5 \cdot 10^5$ a.u. For molecules with smaller dipole moments and larger rotational constants like, for example, CO ($d = 0.043$, $B_e = 9.0 \cdot 10^6$), the van der Waals interaction is comparable in magnitude to interatomic forces. In any case the range of the potential, which scales as $R_e = (\mu C_6)^{1/4}$, is not much different from typical ranges of interatomic potentials (for RbCs $R_e \approx 400$ a.u.). First order terms in the dipole-dipole operator are also absent for two molecules with $N = 1$. In this case, apart from a weak quadrupole-quadrupole contribution proportional to R^{-5} , the long-range part of the intermolecular potential is given by the van der Waals interaction with a comparable C_6 coefficient. Thus, the interactions between molecules with the same N are all short ranged and, in an ultracold system, can be modeled by contact potentials. Then, averaging them over the Gaussian on-site wave functions gives the Hubbard on-site interaction.

The interaction between s and t molecules (without loss of generality we consider \mathbf{z} here) is similar to the resonant interaction of an electronically excited atom and a ground state atom. For even partial waves the intermolecular potential is asymptotically given by $W_z(R) = (1 - 3\cos^2\theta_z)2R^{-3}$, where θ_z is the angle between \mathbf{R} and the z -axis. We consider the weakly interacting regime where the characteristic energy scale of this interaction, $E = l_0^{-3}$, is smaller than the Bloch band separation. Here l_0 is the oscillator length of the on-site harmonic confinement. Then, the two-body problem in a harmonic potential can be solved in the mean-field approximation by using the pseudopotential approach (see [23] and references therein). Due to the anisotropy of $W_z(R)$ the corresponding on-site interaction energy, V_z , can be tuned at will by changing the aspect ratio of the on-site confinement [23].

We arrive at the following expression for H_{Hub} :

$$H_{\text{Hub}} = \sum_i \frac{U}{2} n_{si}(n_{si} - 1) + \frac{U}{2} n_{ti}(n_{ti} - 1) + \sum_i U n_{ti} n_{ti} + V n_{si} n_{ti} : \quad (5)$$

It is easy to see that (5) holds for arbitrary filling factors as long as the on-site density profiles remain Gaussian. However, for the purpose of this paper it is sufficient to consider on average one or two molecules per site. The ten coupling constants in Eq. (5) are difficult to control independently. However, in many cases not all of them are relevant. For a mixture of s and t_z molecules the

relevant coupling constants are U , U_z , and V_z . These are tunable through trap aspect ratio and/or Feshbach resonance. When considering this system we will take $U = U_z$ and $V = V_z$. Another example is the Mott insulating state with one molecule per site (all kinds of molecules are allowed) when the interaction energies are much larger than J . Then, the particular values of the on-site coupling constants are not important (for mechanical stability it is sufficient that they are positive) and the state of the system is found by minimizing the intersite dipolar interactions.

We now discuss the resulting phase diagram, first focusing our attention on the Mott insulating state with one molecule per site. We take the variational wave function

$$|j_M\rangle = \prod_i \left(\cos(\theta_i) s_i^y + \sin(\theta_i) t_i^y \right) |j\rangle_i \quad (6)$$

where θ_i describes the fraction of the molecules excited into $N = 1$ states and \mathbf{t}_i is a normalized complex vector $\mathbf{t}_i \cdot \mathbf{t}_i = 1$. Here, θ_i is the variational parameter which describes the direction the dipole moment points on site i . This allows us to construct variational states that benefit maximally from dipolar interactions. In all cases discussed below we verified the absence of phase separation by checking the eigenvalues of the compressibility matrix for s and t bosons [24]. Taking the expectation value of the dipole operator (3) with our variational wave function (6) we obtain $\langle d_i \rangle = \sin(2\theta_i) j \cdot j \cos(\theta_i - 2B_e t)$ where we have written $\mathbf{t}_i = j \cdot \mathbf{t}_i$. Upon taking the expectation value of dipole Hamiltonian (4), we find for the dipolar energy

$$E_{\text{dip}} = \frac{\sin^2(2\theta)}{8} \sum_{i \neq j} \frac{(\mathbf{t}_i \cdot \mathbf{t}_j + \cos\theta_i \cos\theta_j) (3\mathbf{e}_{ij} \cdot \mathbf{e}_{ij})}{R_i R_j^3} : \quad (7)$$

When minimizing the energy in (7) it is important to keep track of the conservation laws that may be present for certain experimental geometries and on the initial preparation of the system. We will now consider several examples of ordering in the Mott insulating state. Although the dipole interaction in all cases is described by (7) we will see that different preparation leads to very different types of order. Though all discussion in this work will be restricted to 2d, we emphasize that there are nontrivial results in the Mott insulating phase for the 1d and 3d cases as well. As the first example, we consider the square lattice in the xy -plane defined by vectors $\mathbf{a}_1 = \hat{x}$ and $\mathbf{a}_2 = \hat{y}$. Due to cross-terms such as $s^\dagger t_x^\dagger s t_y$ in the dipolar Hamiltonian (4), we see that t_x molecules can be converted to t_y molecules and vice-versa. Thus, N_{t_x} and N_{t_y} are not conserved quantities, and, consequently, the only conserved quantities are N_s and N_{t_z} . Now consider preparing this system in a mixture of $N = 0$

and $N_x = 1; N_z = 1$ states. Then after the system relaxes, taking the constraints into account, we must have $\langle N_{S_i} \rangle = N \cos^2(\alpha)$, $\langle N_{t_x i} \rangle + \langle N_{t_y i} \rangle = N \sin^2(\alpha)$, and $\langle N_{t_z i} \rangle = 0$. This gives the constraints on the variational wave function $\langle t_z \rangle = 0$ and $\langle j_x^2 \rangle + \langle j_y^2 \rangle = 1$. We see that the dipoles are allowed to rotate freely in the xy -plane. For this case, the dipoles will choose to point head-to-tail in the direction of one of the bonds, while alternating in the other. Thus, it is straightforward to see that this gives the ordering wave vector $q = (0; 0)$ with $\langle t_x \rangle = e^{i(q \cdot R_i + \phi_0)}$ and $\langle t_y \rangle = \langle t_z \rangle = 0$ where ϕ_0 is an arbitrary phase corresponding to a change of phase of the time dependent oscillations of the dipolar momentum.

We point out that this configuration is degenerate to the one with dipoles pointing head-to-tail in the y -direction.

As the next example in two dimensions, we take the same lattice as in the previous example, but prepare the system in a mixture of $N_x = 0$ and $N_x = 1; N_z = 0$ states. Recalling that for this geometry, both N_S and N_{t_z} are conserved quantities, we find the constraint on the variational wave function $\langle t_x \rangle = \langle t_y \rangle = 0$ and $\langle t_z \rangle = e^{i\phi}$. With this constraint, the dipole interaction energy is

$$E_{\text{dip}} = \frac{\sin^2(2\alpha)}{4} \sum_{i \neq j} \frac{\cos(\phi_i - \phi_j)}{R_{ij}^3} \quad (8)$$

Here, the dipoles are confined to point in the z -direction, and therefore cannot point head-to-tail. This gives antiferromagnetic ordering in all directions, $q = (\pi; \pi; 0)$, with $\langle t_z \rangle = e^{i(q \cdot R_i + \phi_0)}$ where ϕ_0 is an arbitrary phase.




FIG. 1: The ordering wave vector as the lattice is tilted by angle α . As described in the text, for this situation a magnetic field is used to break the degeneracy between $(N_x = 1, N_z = 1; 0; 1)$ states.

For the final example for the Mott insulating state, we consider a lattice in the xz -plane given by $a_1 = \cos(\alpha)\hat{x} + \sin(\alpha)\hat{z}$ and $a_2 = -\sin(\alpha)\hat{x} + \cos(\alpha)\hat{z}$. In addition, we consider breaking the degeneracy of the $(N_x = 1, N_z = 1; 0; 1)$ states with an external static magnetic

field in the z -direction which will introduce the term proportional to $B L_z$ into our hamiltonian. Preparing the system in a superposition of $N_x = 0$ and $N_x = 1; N_z = 1$ states, we note that because of this degeneracy breaking, there will be no mixing between other angular momentum states. That is, we can completely neglect the $t_{1,0}$ states. This will give $\langle t_x \rangle = \langle t_y \rangle = e^{i\phi} = \frac{1}{\sqrt{2}}$ and $\langle t_z \rangle = 0$ which will confine our dipoles to rotate in the xy -plane as: $\langle t_i(t) \rangle = d_0 \cos(\phi_i - 2B_e t)\hat{x} + d_0 \sin(\phi_i - 2B_e t)\hat{y}$. The dipolar energy of this system is therefore

$$E_{\text{dip}} = \frac{\sin^2(\alpha)}{8} \sum_{i \neq j} \frac{\cos(\phi_i - \phi_j) (1 - \frac{3}{2} e_{ijx}^2)}{R_{ij}^3} \quad (9)$$

We use the ansatz $\phi_i = q \cdot R_i + \phi_0$ to find the minimum of this dipolar energy for a particular lattice defined by the angle α , and the results are summarized in Fig. 1.

We now consider melting the Mott insulator, and entering the superfluid (SF) state. An interesting question to consider is what happens to the ordering wave vector as the Mott insulating state is melted? For instance, deep in the superfluid phase, we will have $q = 0$ which is favorable for Bose-Einstein condensation while we saw that antiferromagnetic ordering is typically favored in the Mott insulating state by dipolar interactions. One possibility is that the wave vector interpolates smoothly between these two extremes as the hopping J increases. Another possibility is that the molecules in the s and t states phase-separate. We will show below that both scenarios are possible depending on on-site energy parameters in our original hamiltonian. For simplicity, we restrict our attention to the third example we discussed above for the Mott insulating state which was a two dimensional lattice in the xy plane prepared with z polarized light. For further simplicity, we take $\langle N_{S_i} \rangle = \langle N_{t_z i} \rangle = N/2$. As we saw before, we can neglect populating the t_x and t_y states, and this phase has antiferromagnetic $q = (\pi; \pi; 0)$ order in the Mott insulating phase.

Allowing for noninteger occupation per site motivates the variational wave function

$$|\psi\rangle = \sum_i \sum_{n=0}^Y \sum_{\alpha=0}^X \frac{(a_i^\alpha)^n}{n!} |i\rangle \quad (10)$$

where $a_i^\alpha = \cos(\alpha) e^{i\phi_s \cdot R_i} s_i^\alpha + \sin(\alpha) e^{i\phi_t \cdot R_i} t_{iz}^\alpha$ and normalization requires $\sum_n \langle j_n^2 \rangle = 1$ (compare with (6)). As before, this wave function maximizes the dipole energy for a given site which is energetically favorable. We can now use a canonical transformation to write our original hamiltonian in terms of the boson operators a_i^α (defined above) and $b_i^\alpha = \sin(\alpha) e^{i\phi_s \cdot R_i} s_i^\alpha + \cos(\alpha) e^{i\phi_t \cdot R_i} t_{iz}^\alpha$ (a new variable resulting from the transformation), and drop the terms which give zero when evaluated using the above variational wave function (10). This leads to the following single-site mean field hamiltonian

$$H_{MF} = 2J \sum_{\langle i,j \rangle} \left[\cos^4(\frac{q}{2}) + \sin^4(\frac{q}{2}) + 2\cos^2(\frac{q}{2})\sin^2(\frac{q}{2})\cos(q) \right] a^\dagger_i a_j + a^\dagger_i a_j a^\dagger_j a_i + a^\dagger_i a_j a^\dagger_j a_i + a^\dagger_i a_j a^\dagger_j a_i \quad (11)$$

$$+ \frac{1}{4} \sin^2(2) (2n_a n_{a_i} - n_a^2) \sum_{\mathbf{R}_i \neq 0} \frac{\cos(\mathbf{q} \cdot \mathbf{R}_i)}{R_i^3} + \frac{1}{2} U n_a (n_a - 1) + \frac{1}{4} (V - U) \sin^2(2) n_a (n_a - 1)$$




FIG. 2: The phase diagram for $V = U$ (left) and $V = 0$ (right). For both cases, the dipolar interaction strength was fixed at $\epsilon = U=5$. Shown are the antiferromagnetic MI states with one and two bosons per site labeled MI1 and MI2. SF1 and SF2 correspond to superfluid states with partial and complete phase separation (described in text). SF3 is a superfluid phase with no phase separation which has an ordering wave vector that interpolates between the Mott insulating and deep superfluid regime.

where $n_a = a^\dagger a$ and we have already performed the minimization over the center of mass momentum $\mathbf{p} = (\mathbf{p}_t + \mathbf{p}_s)/2$. The ground state of this hamiltonian for fixed (relative concentrations) and $\mathbf{q} = \mathbf{p} - \mathbf{p}_s$ (relative momentum) can be determined self-consistently in a_i and b_{a_i} through iteration numerically. The general approach will then be to minimize these ground state energies over $\mathbf{q}_{k,y} \in [0, \pi]$ and $\mathbf{p} \in [0, \pi/2]$. When the minimum occurs for $\epsilon = 4$, phase separation will occur.

The resulting phase diagrams are shown in Fig. 2. The Mott insulating phases are antiferromagnetically aligned and were discussed in the previous section. SF1 corresponds to partial phase separation where part of the lattice will have a larger concentration of s molecules while the other part will have a higher concentration of t molecules. Recall that phase separation will occur for when $\epsilon = 4$ since we initially prepare the system to have equal populations of molecules in the s and t_z states. The region with more s molecules will have $(\mathbf{p}_s)_{x,y} = 0$ and $(\mathbf{p}_t)_{x,y} = \pi$. This will allow the more populated s species to benefit maximally from BEC which prefers zero wave vector while still giving $\mathbf{q}_{k,y} = \pi$ which is preferred for the dipole interaction. The similar situation holds for the region of the lattice with a higher concentration of t_z molecules. SF2 corresponds to the case where the s and t_z molecules completely phase separate. Since the dipole interaction is negligible for this case, we will have

$(\mathbf{p}_s)_{x,y} = (\mathbf{p}_t)_{x,y} = 0$ which will favor BEC. Finally, SF3 corresponds to the case mentioned above where the wave vector \mathbf{q} interpolates between the deep superfluid and Mott insulating states ($0 < \mathbf{q}_{k,y} < \pi$) for which no phase separation occurs ($\epsilon = 4$).

In conclusion, we have shown that polar molecules prepared in a mixture of two rotational states can exhibit long-ranged dipolar interactions in the absence of an external electric field. We have described several novel Mott insulating and superfluid phases that can be realized as a result of such an interaction. Such states can be detected by Bragg scattering or by time-of-flight expansion [25].

This work was supported by the NSF grant DMR-0132874 and career program, the Harvard-MIT CUA, and the Harvard-Smithsonian ITAMP. We thank B. Halperin, R. Klemens, A. Polkovnikov, D.W. Wang, and P. Zoller for useful discussions.

Note added: When this manuscript was close to completion we became aware of a paper considering a similar system [12].

-
- [1] J. Doyle, B. Friedrich, R. V. Klemens, and F. Masnou-Seeuws, Eur. Phys. J. D 31, 149 (2004).
 - [2] K. Goral, K. Rzaewski, and T. Pfau, Phys. Rev. A 61, 051601 (2000).
 - [3] J. Stuhler et. al, Phys. Rev. Lett. 95, 150406 (2005).
 - [4] C. A. Stan et. al, Phys. Rev. Lett. 93, 143001 (2004).
 - [5] S. Inouye et. al, Phys. Rev. Lett. 93, 183201 (2004).
 - [6] J. M. Sage, S. Sainis, T. Bergeman, and D. DeMille, Phys. Rev. Lett. 94, 203001 (2005).
 - [7] S. Yi and L. You, Phys. Rev. Lett. 92, 193201 (2004).
 - [8] L. Santos, G. V. Shlyapnikov, P. Zoller, and M. Lewenstein, Phys. Rev. Lett. 85, 1791 (2000).
 - [9] D. DeMille, Phys. Rev. Lett. 88, 067901 (2002).
 - [10] K. Goral, L. Santos, and M. Lewenstein, Phys. Rev. Lett. 88, 170406 (2002).
 - [11] M. G. Moore and H. R. Sadeghpour, Phys. Rev. A 67, 041603 (2003).
 - [12] A. Micheli, G. Brennen, and P. Zoller, quant-ph/0512222.
 - [13] O. Mandel et. al, Nature 425, 937 (2003).
 - [14] L. M. Duan, E. Demler, and M. Lukin, Phys. Rev. Lett. 91, 090402 (2003).
 - [15] A. Kuklov, N. Prokofev, and B. Svistunov, Phys. Rev. Lett. 92, 050402 (2004).
 - [16] A. Isacsson, M.-C. Cha, K. Sengupta, and S. M. Girvin, Phys. Rev. B 72, 184507 (2005).
 - [17] B. Damaskiet. al, Phys. Rev. Lett. 95, 060403 (2005).

- [18] X.-L. Deng, D. Porras, and J. I. Cirac, cond-mat/0509197.
- [19] J. A. Sauer, Phys. Rev. 57, 142 (1940).
- [20] J. M. Luttinger and L. Tisza, Phys. Rev. 70, 954 (1946).
- [21] R. Moessner and S. L. Sondhi, Phys. Rev. B 68, 054405 (2003).
- [22] The rotational relaxation can be greatly reduced in an optical lattice, similar to the inelastic losses in the atomic case. It can be completely eliminated in the 1 atom Mott case and suppressed in a deep lattice.
- [23] S. Yi and L. You, Phys. Rev. Lett. 92, 193201 (2004).
- [24] See, for instance, chapter 1 and 2 of K. Huang, Statistical Mechanics (John Wiley and Sons, 1987).
- [25] E. Altman, E. Demler, and M. Lukin, Phys. Rev. A 70, 013603 (2004).