## Spin Structure Factor of the Frustrated Quantum M agnet Cs<sub>2</sub>CuCl<sub>4</sub>

Denis Dalidovich, Rastko Sknepnek, A. John Berlinsky, Junhua Zhang, and Catherine Kallin

Department of Physics and Astronomy, M cM aster University, H am ilton, Ontario, Canada L8S 4M 1 (Dated: December 29, 2021)

The ground state properties and neutron structure factor for the two-dimensional antiferrom agnet on the triangular lattice, with uni-directional anisotropy in the nearest-neighbor exchange couplings and a weak D zyaloshinskii-M oriya (DM) interaction, are studied. This H am iltonian has been used to interpret neutron scattering measurements on the spin 1=2 spiral spin-density-wave system,  $C s_2 C u C l_4$ ,  $R \cdot C oldea, et al., P hys. R ev. B 68, 134424 (2003)]. C alculations are performed using a 1=S expansion, taking into account interactions between spin-waves. The ground state energy, the shift of the ordering wave-vector, Q, and the local magnetization are all calculated to order 1=S<sup>2</sup>. The neutron structure factor, obtained using anharm onic spin-wave G reen's functions to order 1=S, is shown to be in reasonable agreement with published neutron data, provided that slightly di erent parameters are used for the exchange and DM interactions than those inferred from measurements in high magnetic eld.$ 

PACS numbers: 75.10 Jm , 75.25.+ z, 75.30 D s, 75.40 G b

#### I. IN TRODUCTION

There is enormous interest in the condensed matter community in the possibility of observing fractionalized quasiparticles in two or three dimensional electronic system s.<sup>1,2,3</sup> Fractionalization is the rule, rather than the exception, in one dimensional metals where electronic spin and charge propagate independently.<sup>4</sup> Fractionally charged quasiparticles have also been found in the two dim ensional electron gas in high m agnetic elds where the fractional quantum Halle ect is observed.<sup>5</sup> On the theoretical side, fractionalization has recently been argued to exist in frustrated two-dimensional quantum model system s,<sup>6,7</sup> although no corresponding experim ental system s have yet been con med. A set of long-wavelength theories have been developed that describe the properties of putative fractionalized magnets.<sup>8,9,10</sup> Subsequently, the theoreticalm odeling of possible fractionalized phases was extended to a wide variety of strongly correlated electron system s.<sup>11,12,13</sup> The discovery of fractionalization in real two or three dimensional systems in zero magnetic eld would provide a striking example of the emergence of new kinds of particles from the collective behavior of strongly interacting electrons.<sup>1</sup>

As a result of this intense interest, the recent claim of the observation of spinons, neutral, spin 1/2 quasiparticles, in the insulating antiferrom agnet  $C_{s_2}C_{UC}L_{l_1}$  has attracted considerable attention.<sup>14,15</sup>  $C_{s_2}C_{UC}L_{l_2}$  consists of nearly uncoupled layers of coupled chains. The chains are staggered, so that each spin interacts equally with two spins on each of two neighboring chains, with an antiferrom agnetic coupling,  $J^0$ , about 1/3 the strength of the intrachain antiferrom agnetic nearest neighbor coupling, J. The layer thus resembles a triangular lattice, with strong interactions along one direction and weaker interactions in the transverse directions. These intralayer couplings, J and  $J^0$  assisted by a much weaker interlayer coupling  $J^{00}$ , stabilize an incommensurate spin density wave (SDW) ground state with a wave vector oriented along the chain direction. A weak D zyaloshinskii-M oriya (DM) interaction D is believed to orient the spins in the 2D layer.

C oblea and cow orkers<sup>14,15</sup> used neutron scattering to study spin-wave excitations from this ordered ground state. W hile C oblea et al. did observe excitations sim – ilar to what is expected for spin-waves, they also saw broad features which they interpreted as excitations of pairs of spinons. Their idea is that the higher energy spin excitations resem ble excitations from a nearby-lying spin-liquid state that supports fractionalized excitations. The low-dimensional nature of the system together with the low spin are argued to place this system close to a quantum critical point separating the ordered SDW state from a 2D spin liquid state. This conjecture has stim ulated a number of theoretical studies of possible 2D spin-liquid states that could account for the observed behavior.<sup>16,17,18,19,20,21,22</sup>

In this paper, we pursue a di erent interpretation of the neutron data, namely that low spin and quasi-onedimensionality, along with non-collinear SDW order, all give rise to substantial anharm onic interactions that couple one- and two-spinwave excitations from the ordered state. The coupling to two spinwave states also yields broad spectra as seen in the neutron data. A similar conclusion has been reached by Veillette et al.<sup>23</sup> who also noted that the low momentum resolution of the neutron data is another signi cant factor in the observed breadth of the spectra. We nd that anharm onic e ects lead to signi cant broadening of the neutron spectra but are sensitive to anisotropies, such as the DM interaction, which suppress quantum uctuations. We show in particular that, if such anisotropies are weak, then quantum uctuations, described in terms of a 1=S expansion about the mean eld SDW state, reduce the local moment and lead to substantial renorm alizations and broadening of the excitation spectra.

The rest of this paper is organized as follows. In Sec.  ${\rm I\!I}$  we describe the H am iltonian and the SDW state of  $C_{2}C_{1}UC_{1}$ . In Sec. III we show how anharmonic e ects m odify the ground state energy and the wave vector Q of the SDW to order  $1=S^2$ , and we also review the calculation of the anham onic one-spin-wave G reen's functions to order 1=S. In Section IV we calculate the sublattice magnetization M to order 1=S<sup>2</sup> as a function of the ratio  $J^0=J$  and strength of DM interaction D = J, showing that the renorm alized value of M depends sensitively on the ratio D = J for sm all D = J. In Sec. V we exam ine expressions for the anharm onic spin-wave energies and dam ping to order 1=S. For D = 0 we study how the Goldstone mode at Q is preserved and show how the preservation of this zero energy mode can be used to de ne a set of renorm alized coupling constants. Section VI reviews the calculation of the neutron structure factor and presents a detailed comparison to the inelastic data of Coldea et al. for speci c wave vectors where two-m agnon scattering is important. In Sec. VII we discuss the implications of our theoretical calculations in interpreting existing data on C s<sub>2</sub>C uC l<sub>4</sub>. W e also m ake suggestions for possible new experim ents.

### II. SPIN DENSITY WAVE STATE FOR C $s_2 C \, u C \, l_4$

The spin density wave state of  $C s_2 C u C l_4$  can be described by the simple, model H am iltonian for a set of decoupled layers:

$$H = \int_{R}^{X} JS_{R} \qquad S_{R+1+2} + J^{0}S_{R} \qquad (S_{R+1} + S_{R+2})$$

$$R \qquad (1)^{n}D \qquad S_{R} \qquad (S_{R+1} + S_{R+2}); \qquad (1)$$

where J is the nearest neighbor coupling constant between  $S = 1=2 \text{ Cu}^{2+}$  spins along chains in one direction, which we take to be the x-direction in a triangular lattice, while  $J^0$ , is the coupling constant along the other two principal directions in each layer as illustrated in Fig.1. Both J and  $J^0$  are antiferrom agnetic, and  $J > J^0$ . The last term in Eq. (1) with D = (0;D;0) describes the D zyaloshinskii-M oriya interaction that alternates in sign between even and odd layers labeled by index n.

Values of the coupling constants, determ ined from measurem ents in high magnetic  $elds_r^{24}$  are:

$$J = 0:374 \text{ meV}; J^0 = 0:128 \text{ meV} J=3;$$
 (2)

$$D = 0.02 \text{ m eV};$$
 (3)

leading to a sm all N eel tem perature  $T_N$  of less than 1K in the presence of the interlayer coupling  $J^{(0)}$ . The interlayer coupling is su ciently sm all  $(J^{(0)}=J=0.045)^{24}$  that




FIG.1: Exchange couplings between the di erent sites of the triangular lattice within a single layer.

we neglect it in our calculations performed at T = 0. The classical ground state for this H am iltonian is a SDW whose wave vector Q<sub>0</sub> is along the strong-coupling direction and has the value

$$Q_0 = 2 (0:5+);$$
 (4)

where experimentally, = 0.030(2) is found for C  $s_2$  C uC  $l_4$  ,  $^{24}$  so that Q  $_0$  is close to .

Thism eans that individual chains are ordered approxim ately antiferrom agnetically, while the spins in adjacent chains, which are slipped by half a lattice spacing along the chain direction, are approximately orthogonal.

To further illustrate this point, consider the calculation of the classical ground state energy of the SDW state, that can be easily determ ined by introducing a local reference frame, so that at every site the averaged spin is directed along the axis:

$$S_{R}^{x} = S_{R} \cos(Q \quad R) \quad \underset{R}{S} \sin(Q \quad R);$$
$$S_{R}^{z} = S_{R} \sin(Q \quad R) + \underset{R}{S} \cos(Q \quad R);$$
$$S_{R}^{y} = S_{R} : \qquad (5)$$

In Eqs. (5), Q is the wave vector of the spiral structure, which at the classical level we write as  $(Q_0;0;0)$  where  $Q_0$  changes sign from layer to layer.  $Q_0$  is determined from minimization of the classical ground state energy with  $S_R = S_R = 0$  for all R. The classical ground state energy (per spin) is given by

$$S^{2}E_{G}^{(0)}(Q) = S^{2}J_{Q}^{T};$$
 (6)

$$J_{O}^{T} = J_{Q} \quad \text{iD}_{Q}; \qquad (7)$$

where

$$J_{k} = J \cos k_{x} + 2J^{0} \cos \frac{k_{x}}{2} \cos \frac{p - 3k_{y}}{2};$$
 (8)

and

$$D_{k} = 2iD \sin \frac{k_{x}}{2} \cos \frac{p - 3k_{y}}{2}$$
(9)

are the Fourier transforms of the exchange and DM interactions of the Ham iltonian respectively.

For D = 0, in Eq. (4) is given by

$$= \frac{1}{2} \arcsin \frac{J^0}{2J} = 0.0547;$$
(10)

while for D = 0.02 meV, the value of  $Q_0$  that minimizes the classical ground state energy corresponds to

= 0:0533 for the experimental values of J and  $J^0$ . The two terms in Eq. (8), represent the intrachain and interchain energies respectively. Substituting the actual value of  $Q_0$  for a given  $J^0=J$  and D = 0, we not that

$$S^{2}E_{G}^{(0)}(Q_{0}) = JS^{2} 1 + \frac{J^{2}}{2J^{2}}$$
: (11)

Thus the contribution to the ground state energy from the interchain coupling is of order  $(J^0)^2 = J^2$  compared to the intrachain contribution. Quantum uctuations reduce the electiveness of interchain coupling further by renormalizing the SDW wave vector to a value even closer to (corresponding to a smaller elective  $J^0=J$ ). At the same time, quantum uctuations lower the total energy. Thus if one were to model the elects of quantum uctuations in terms of elective coupling constants, J and  $J^0$ , the result would be  $J^0=J' < J^0=J$  and J' > J. As is shown in the next section and in Appendix A, the value of obtained at the classical level, is in fact considerably renormalized by quantum uctuations.

### III. 1/S EXPANSION

To address the physics determ ined by quantum uc-tuations, we employ the well-known Holstein-Prim ako transform ation for the spin operators:<sup>25</sup>

$$S_{R} = S_{R} \quad iS_{R} \quad \frac{p}{2S}a_{R}^{Y} \quad 1 \quad \frac{1}{4S}a_{R}^{Y}a_{R} ;$$
 (12)

$$S_{R}^{+} = S_{R}^{+} + iS_{R}^{-} \frac{p}{2S} \frac{1}{4S} a_{R}^{Y} a_{R}^{-} a_{R}^{-};$$
 (13)

$$S_{R} = S \quad a_{R}^{Y} a_{R} :$$
 (14)

These are written in a form with the square root expanded to rst order in 1=(2S). The magnon operators  $a_R^{\gamma}$ ,  $a_R$  describe excitations around the spiral ground state and obey Bose statistics. Substituting the transformations (12)-(14) into Eq. (1), we obtain the Ham iltonian for interacting magnons,

$$H = S^{2}E_{G}^{(0)}(Q) + (H^{(2)} + H^{(3)} + H^{(4)}); \quad (15)$$

where in the Fourier transform ed representation

$$H^{(2)} = 2S X^{k} A_{k} a_{k}^{y} a_{k} \frac{B_{k}}{2} (a_{k}^{y} a_{k}^{y} + a_{k} a_{k}) : (16)$$

The functions  $A_k$  and  $B_k$  are expressed through  $J_k$  as

$$A_{k} = \frac{1}{4} J_{Q+k}^{T} + J_{Q-k}^{T} + \frac{J_{k}}{2} J_{Q}^{T}; \qquad (17)$$

$$B_{k} = \frac{J_{k}}{2} - \frac{1}{4} J_{Q+k}^{T} + J_{Q-k}^{T} :$$
 (18)

The interactions between m agnons are described by the last two terms H  $^{(3)}$  and H  $^{(4)}$ . The three-m agnon term can be written in the form

$$H^{(3)} = \frac{i}{2}^{r} \frac{S}{2N} \frac{X}{1;2;3} (C_{1} + C_{2}) (a_{3}^{y} a_{2} a_{1}) a_{1}^{y} a_{2}^{y} a_{3}) (a_{2}^{y} a_{2} a_{1})$$
(19)

that contains the vertex

$$C_{k} = J_{Q+k}^{T} \quad J_{Q-k}^{T} ; \qquad (20)$$

which is antisymmetric with respect to the transform ation k ! k. For brevity, in longer expressions we use the convention that  $1 = k_1$ , etc.

The four-magnon term can be conveniently written as:

$$H^{(4)} = \frac{1}{4N} X_{1;2;3;4} [(A_{1 3} + A_{1 4} + A_{2 3} + A_{2 4}) (B_{1 3} + B_{1 4} + B_{2 3} + B_{2 4}) (A_{1} + A_{2} + A_{3} + A_{4})]$$

$$(21)$$

$$a_{1}^{Y} a_{2}^{Y} a_{3} a_{4 1+2;3+4} + \frac{2}{3} (B_{1} + B_{2} + B_{3}) a_{1}^{Y} a_{2}^{Y} a_{3}^{Y} a_{4} + a_{4}^{Y} a_{3} a_{2} a_{1 1+2+3;4};$$

and contains only symmetric vertices. It is important to emphasize that H  $^{(3)}$  comes from the coupling between the operators S<sub>R</sub> and S<sub>R</sub> ,which arises for non-collinear ordered states. This term plays a crucial role in calculations of the structure factor as well as in renorm alization of the energy spectrum .

To proceed, one needs to diagonalize the quadratic part of the H am iltonian Eq. (16) by m eans of the B ogoliubov

transform ation:

$$a_{k} = u_{k}c_{k} + v_{k}c_{k}^{y}; a_{k}^{y} = u_{k}c_{k}^{y} + v_{k}c_{k};$$
 (22)

$$u_{k} = \frac{r}{\frac{A_{k} + \mathbf{w}_{k}}{2\mathbf{w}_{k}}}; \quad v_{k} = sgnB_{k} \frac{r}{\frac{A_{k} - \mathbf{w}_{k}}{2\mathbf{w}_{k}}};$$
 (23)

The energy spectrum in the above equations is given by:

$$\mathbf{"}_{k} = \frac{\mathbf{A}_{k}^{2} + \mathbf{B}_{k}^{2}}{r + \frac{\mathbf{A}_{k}^{2} + \mathbf{B}_{k}^{2}}{(\mathbf{J}_{k} + \mathbf{J}_{Q}^{T})^{T} (\mathbf{J}_{Q}^{T} + \mathbf{J}_{Q}^{T} + \mathbf{J}_{Q}^{T})^{T}} = \frac{\mathbf{A}_{k}^{T}}{(\mathbf{J}_{k} + \mathbf{J}_{Q}^{T})^{T} (\mathbf{J}_{Q}^{T} + \mathbf{J}_{Q}^{T} + \mathbf{J}_{Q}^{T})^{T}} = 2 \mathbf{J}_{Q}^{T} : (24)$$

For D = 0, the m agnon spectrum has zeros at k = 0and k = Q in the two-dimensionalBrillouin zone, while a non-zero D leads to the appearance of a nite gap of order DJ around the Q-points. A fler diagonalization, H<sup>(2)</sup> takes the form :

$$H^{(2)} = SE_{G}^{(1)}(Q) + \frac{2S}{N} \sum_{k}^{X} \mathbf{w}_{k} c_{k}^{y} c_{k}; \qquad (25)$$

$$E_{G}^{(1)}(Q) = J_{Q}^{T} + \frac{1}{N}_{k}^{X}$$
"k: (26)

The contribution  $SE_{G}^{(1)}(Q)$  gives the leading 1=(2S) correction to the classical ground state energy Eq. (6). M inimizing the sum  $S^{2}E_{G}^{(0)}(Q) + SE_{G}^{(1)}(Q)$ , we nd the quantum correction to the classical value  $Q_{0}$ . To order 1=S the shifted wave vector is Q = (Q;0;0)

$$Q = Q_0 + \frac{Q^{(1)}}{2S};$$
 (27)

$$Q^{(1)} = \frac{\left(\frac{\theta^2 J_Q^T}{\theta Q^2}\right)^{\# 1}}{\left(\frac{\theta Q^2}{\theta Q^2}\right)^{\# 1}} \frac{1}{N} \frac{X}{k} \frac{(A_k + B_k)}{"_k} \frac{(\theta J_Q^T + k)}{\theta Q} \frac{(28)}{Q_0}$$

In the absence of the DM interaction,

$$Q^{(1)}=2 = 0:0324;$$
 (29)

while for D = 0.02 m eV,

$$Q^{(1)}=2 = 0:0228:$$
 (30)

We see that inclusion of the D zyaloshinskii-M oriya interaction suppresses the renorm alization of Q towards  $\ .$ 

It is possible to go further and also calculate the correction to the ground state energy and the ordering wave vector Q that is of order  $1=(2S)^2$ . The details of those calculations are presented in Appendix A. The results indicate that quantum corrections computed order by order in 1=(2S) are signi cant for the values of J and  $J^0$  given by Eqs. (2). To further explore this issue, one can compare the classical ground state energy Eq. (6) to its leading renorm alization due to quantum uctuations  $SE_G^{(1)}(Q_0)$ . We nd, for D = 0

$$S^{2}E_{G}^{(0)}(Q_{0})=J = 0.265; SE_{G}^{(1)}(Q_{0})=J = 0.157;(31)$$

and, for D = 0.02 m eV,

$$S^{2}E_{G}^{(0)}(Q_{0})=J=0.291; SE_{G}^{(1)}(Q_{0})=J=0.138:(32)$$

We see that the leading quantum correction lowers the ground state energy by about 50%, giving reasonable agreement with a recent experimental determination of this quantity, based on susceptibility measurements,<sup>26</sup> which yields a ground state energy slightly above 0:5J. We show in Appendix A that for D = 0 the next order in 1=(2S) correction reduces the ground state energy by a further 10 percent below the sum of the values given in Eq. (31), bringing the total within a few percent of the experimental value.

To calculate physical observables, we need the G reen's functions for the magnon operators  $a_k$ . They can be conveniently written in the form of a 2 2 matrix, indicating the presence of norm al and anom alous parts. At zero tem perature, the de nition reads

$$i\hat{G}(k;!) = \int_{1}^{Z_{1}} dt e^{i!t} \hat{T} a_{k}(t) h_{k}(0)a_{k}(0);$$
(33)

where  $\hat{T}$  is the time ordering operator, and the average is taken with respect to the ground state. The inverse of the matrix for the bare G reen's functions has the form<sup>29</sup>

$$\hat{G}^{(0) 1}(k;!) = \begin{array}{c} ! & 2SA_{k} + i & 2SB_{k} \\ & 2SB_{k} & ! & 2SA_{k} + i \end{array} ; \qquad (34)$$

corresponding to the G reen's function

$$\hat{G}^{(0)}(k;!) = \frac{1}{(! !_{k} + i)(! + !_{k} i)} \quad \begin{array}{c} ! + 2SA_{k} & i & 2SB_{k} \\ 2SB_{k} & ! + 2SA_{k} i \end{array} ; \quad (35)$$

;

where  $!_k = 2S "_k$ . The self-energy (k;!) determ ining the exact G reen's function is also a 2 2 m atrix, that can be parametrized as

$$\hat{k}_{k}(k; !) = \begin{array}{c} X (k; !) + Y (k; !) & Z (k; !) \\ Z (k; !) & X (k; !) & Y (k; !) \end{array} ;$$

$$(36)$$

and which satis es the Dyson equation

$$\hat{\mathbf{G}}^{-1}(\mathbf{k};!) = \hat{\mathbf{G}}^{(0)-1}(\mathbf{k};!) \qquad (37)$$

The self-energy to order 1= (2S) consists of two parts

$$(k;!) = (4) (k) + (3) (k;!):$$
 (38)

The contribution  $^{(4)}$  (k) is frequency independent and can be calculated simply by the Hartree-Fock decoupling of the quartic term in the Ham iltonian Eq. (21).

$$X^{(4)}(k) = A_{k} + \frac{2S}{N} \frac{X}{q} \frac{1}{!q} \frac{1}{2} B_{k} + B_{q} B_{q}$$
$$+ (A_{k} q B_{k} q A_{q} A_{q} A_{k}) A_{q}; (39)$$

$$Y^{(4)}(k) = B_{k} + \frac{2S}{N} \frac{X}{q} \frac{1}{!_{q}} (B_{k} + B_{q}) A_{q} + A_{k q} B_{k q} A_{q} \frac{1}{2} A_{k} B_{q}; (40)$$

$$Z^{(4)}(\mathbf{k}) = 0$$
: (41)

 $^{(3)}$  (k;!) is the contribution arising from the threem agnon interactions, H  $^{(3)}$ . Its non-zero in aginary part determ ines the m agnon dam ping to rst order in 1=(2S). The corresponding components are most easily calculated by transform ing Eq. (19) to the q<sub>k</sub>-operator basis using the B ogoliubov transform ations, Eqs. (22).

$$X^{(3)}(k;!) = \frac{S}{16N} \frac{X^{n}}{q} [_{1}(q;k q)]^{2} + [_{2}(q;k q)]^{0} \frac{1}{!_{q} + !_{kq} ! i} + \frac{1}{!_{q} + !_{kq} + ! i}$$

$$Z^{(3)}(k;!) = \frac{S}{16N} \frac{X^{n}}{q} \left[ \frac{1}{q}; k q \right]^{2} \left[ \frac{2}{q}; k q \right]^{2} \frac{1}{\left[ \frac{1}{q} + \frac{1}{kq} + \frac{1}{q} + \frac{$$

$$Y^{(3)}(k;!) = \frac{S}{16N} \prod_{q}^{X} f_{2_{1}}(q;k_{q}) \sum_{2_{1}}^{Q}(q;k_{q})g_{\frac{1}{!_{q}+!_{k_{q}}}!i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!_{k_{q}}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!_{q}+!}i_{\frac{1}{!}i_{\frac{1}$$

#### IV. SUBLATTICE MAGNETIZATION

In this section we consider the staggered magnetization for a range of values of  $J^0=J$  and D=J and show that quantum uctuations lead to a considerable renorm alization of the average value of the local moment. From the operator de nition Eq. (14) we can write the staggered magnetization as

$$M = S + \frac{1}{N} \frac{X}{k} \frac{Z}{2i} G_{11} (k;!) e^{i!0}; \quad (44)$$

where the exact G reen's function  $G_{11}$  (k;!) is the corresponding element of the full matrix  $\hat{G}$  (k;!) and 0 is a negative in nitesimal. To lowest order in 1=(2S), one can use  $G_{11}^{(0)}$  (k;!) of Eq. (35), to nd the rst order correction to the S = 1=2 value

$$M^{(1)} = \frac{1}{2N} \frac{X}{k} \frac{A_k}{k} 1 :$$
(45)

For the next order correction in 1=(2S), there are two contributions. The rst one, denoted as M  $_{\rm I}^{(2)}$ , arises from renorm alization by quantum uctuations of the ordering wave vector Q<sub>0</sub>. This correction is obtainable by

substituting the renorm alized value Q given by Eq. (27) into the form ula above and expanding using Eq. (45) up to rst order in 1=(2S).

The second contribution is determ ined by the self-energy corrections of order 1=(2S) to the G reen's function itself and is given by

$$M_{I}^{(2)} = \frac{Q^{(1)}}{2S} \frac{1}{N} \frac{X}{k} \frac{B_{k}(A_{k} + B_{k})}{4^{n_{k}^{3}}} \frac{@J_{Q+k}^{T}}{@Q} (46)$$

$$M_{II}^{(2)} = \frac{1}{N} \frac{X}{k} \frac{Z}{2i} e^{i!0} \hat{G}_{11}(k;!); \qquad (47)$$

where

$$G_{11}(k;!) = \hat{G}^{(0)}(k;!) \hat{K};!) \hat{G}^{(0)}(k;!) = \frac{1}{11} = \frac{1}{11}$$

$$k_{11}(k;!) \hat{G}^{(0)}_{11}(k;!) + 2Z(k;!) \hat{G}^{(0)}_{11}(k;!) \hat{G}^{(0)}_{12}(k;!) + k_{11}(k;!) \hat{G}^{(0)}_{$$

This correction, as a result of integration over !, will contain two parts, so that

$$M_{II}^{(2)} = M_{a}^{(2)} + M_{b}^{(2)} :$$
(49)

O ne part com es from taking the residues at the poles of self-energies, while the other results from the double-poles of the products of two G reen's functions. Using Eqs. (39)-(42), we obtain the rst contribution

$$M_{a}^{(2)} = \frac{S}{8N} \frac{X}{_{k,q}} \left( \frac{\left[ 1 (q;k q) \right]^{2}}{2} \frac{h}{G_{11}^{(0)}} (k; _{k,q}) + G_{12}^{(0)} (k; _{k,q})^{2} + \frac{\left[ 2 (q;k q) \right]^{2}}{2} \right)^{1} \right)^{1} \left( \frac{h}{2} (q;k q)^{2} - \frac{h}{2} \right)$$

$$_{k,q} = !_{q} !_{k q}$$
(51)

The contribution arising from the residues at the double-poles of the products of two G reen's functions in Eqs. (47)-(48), is straightforward to calculate as well. A fler some algebra, we nd that

$$M_{b}^{(2)} = \frac{1}{2SN} \frac{X}{k} \frac{B_{k}}{4_{k}^{*}} \times (k;!_{k}) + Z(k;!_{k}) \times (k;!_{k}) \times (k;!_{k})$$

where  $!_{k} = 2S''_{k}$ , and

$$X^{0}(k;!_{k});Y^{0}(k;!_{k});Z^{0}(k;!_{k}) = \frac{@X(k;!)}{@!};\frac{@Y(k;!)}{@!};\frac{@Z(k;!)}{@!}:=!_{k}:$$
(53)

It is clear from the expressions above that both  ${\rm ~M}_{\rm ~T}$ and M  $_{II}^{(2)}$  contain the overall prefactor 1=(2S). In Fig. 2, we plot the total sublattice magnetization

$$M = S + M^{(1)} + M^{(2)}_{I} + M^{(2)}_{II}$$
 (54)

as a function of D zyaloshinskii-M oriya interaction D, for coupling constants J and  $J^0$  as given by Eqs. (2). The

ratios of each correction to the classical magnetization S = 1=2, M = S, are also plotted, as explained in the caption of Fig. 2.0 ne can see that the renorm alization of the local magnetization by quantum uctuations to second order in 1=(2S) is considerable for all values of D between 0 and 0:025 m eV, although increasing D clearly suppresses uctuations making the system more classical. The gure also shows a calculated sublattice magnetiza-




FIG.2: The total value of sublattice m agnetization M , Eq. (54), calculated up to the second order in 1=(2S) as a function of D zyaloshinskii-M oriya interaction D, is shown by the solid line. The dashed, dotted and dashed-dotted lines are the results for the quantum corrections M  $^{(1)}$ , M  $^{(2)}_{I}$  and M  $^{(2)}_{I}$ , respectively, divided by S = 1=2.

tion that is negative for D between 0 and D 0:003

7

m eV, signaling the melting of the assumed spiral ordered ground state by quantum uctuations when D is too small.

Experimentally, a value of M = S = 0.25 was inferred by Coldea and co-workers<sup>31</sup> from elastic neutron scattering, and this value was used in the analysis of the most recent neutron data.<sup>14</sup> Such a small renormalization of the local moment is incompatible with the theory presented here. It is also unlikely to be explained by the addition of the very small expected interlayer exchange coupling  $J^{00}=J=0.045.^{14,24}$  It is indeed puzzling that such a small moment reduction, which is less than for the spin 1=2 H eisenberg antiferrom agnet on a two-dimensional square lattice, would occur in this system which is both frustrated and close to the one-dimensional lim it.

### V. ENERGY SPECTRUM

The renorm alized m agnon energy spectrum  $k_k = 2S u_k$  can be found from the poles of the exact G reen's function, determ ined by<sup>29</sup>

$$\begin{array}{cccc} n & 0 & n & h & \text{io} \\ \text{Re } \det \hat{G}^{1}(\mathbf{k}; \mathbf{k}_{k}) &= \text{Re } \det \hat{G}^{(0) 1}(\mathbf{k}; \mathbf{k}_{k}) & \hat{}(\mathbf{k}; \mathbf{k}_{k}) &= 0; \end{array}$$
 (55)

which leads to the self-consistency equation:

$$\mathbf{t}_{k} = \mathbf{ff2S} (\mathbf{A}_{k} + \mathbf{B}_{k}) + \mathbf{Re} [\mathbf{X} (\mathbf{k}; \mathbf{t}_{k}) \quad \mathbf{Z} (\mathbf{k}; \mathbf{t}_{k})]\mathbf{g} \mathbf{f2S} (\mathbf{A}_{k} \quad \mathbf{B}_{k}) + \mathbf{Re} [\mathbf{X} (\mathbf{k}; \mathbf{t}_{k}) + \mathbf{Z} (\mathbf{k}; \mathbf{t}_{k})]\mathbf{g}$$

$$\mathbf{fIm} [\mathbf{X} (\mathbf{k}; \mathbf{t}_{k}) + \mathbf{Z} (\mathbf{k}; \mathbf{t}_{k})]\mathbf{g} \mathbf{fIm} [\mathbf{X} (\mathbf{k}; \mathbf{t}_{k}) \quad \mathbf{Z} (\mathbf{k}; \mathbf{t}_{k})]\mathbf{g} + [\mathbf{Im} \mathbf{Y} (\mathbf{k}; \mathbf{t}_{k})]^{2} + \mathbf{ReY} (\mathbf{k}; \mathbf{t}_{k}):$$
(56)

Near these poles, if the imaginary part of the self-energy is small, the shape of the one-m agnon spectrum is approximately Lorentzian and has the form  $^{29}$ 

$$h \qquad i$$

$$det \hat{G}^{1}(k;!)' \quad 2!_{k} z_{k} (! : t_{k}) + i \mathbb{I} m \quad det \hat{G}^{1}(k; t_{k}); \qquad (57)$$

where

$$2!_{k} z_{k} = \frac{0}{0!} \operatorname{Re}^{h} \operatorname{det} \widehat{\mathcal{G}}^{-1} (k; !) : (58)$$

The imaginary part of Eq. (57) determines the inverse life-time of the quasiparticles. To calculate the half-width at half-maximum of the peak, we use the parameter

$$_{k}' \frac{\text{Im} [\det \hat{G}^{-1}(k; \boldsymbol{t}_{k})]}{2!_{k} z_{k}}; \qquad (59)$$

Equation (56), although form ally exact, in plicitly includes contributions of all orders in 1=S. Furtherm ore, quantities such as the term s 2S ( $A_k = B_k$ ) that are of the leading order in S, should be corrected for the rst order shift of the ordering wave vector from  $Q_0$  to Q, given by Eqs. (27)-(28). A lthough the self-energies are only calculated to order 1=S, taking the square root of a sum of squares of self-energies evaluated at renorm alized frequencies, e ectively m ixes in higher order corrections. A lso, particularly for the case D = 0, there is cancellation am ong term s of order 1=S, leaving a result, at the shifted Q-vector, which is the square root of residual 1=S<sup>2</sup> term s, which m ay be negative.

A simple way to preserve physical behavior at the Goldstone wave vector Q for D = 0, is to evaluate the

8

self-energies using renorm alized coupling constants, for which the classical ordering wave vector is Q, that is

$$Q = + 2 \arcsin \frac{J^0}{2J} : \qquad (60)$$

The e ect of this renorm alization, which determ ines the ratio  $J^0=J$  and represents a higher order in 1=S correction to the self-energies, is that the self-energies will vanish at the renorm alized Q. A further renorm alization of J by itself is de ned by the condition that the coe cient of the linear term in k Q j inside the square root vanishes for k = Q, so that the energy dispersion relation around this point is linear in k Q j. The procedure of expanding the self-energies around Q is tedious but com pletely equivalent to that described elsew here.<sup>29,30</sup> O m itting the interm ediate steps, we obtain the condition

$$\frac{\partial J_k}{\partial k}_{k=Q} = \frac{1}{2SN} \frac{X}{p} \frac{(A_p + B_p)}{p} \frac{\partial J_{p+Q}}{\partial p}; \quad (61)$$

where the bars indicate that the renorm alized values J and  $J^0$  should be used in the right hand side of this equation. One can easily check that, within this approach, we retain consistency up to lowest order in 1=2S. Indeed, expanding the derivative  $@J_k = @k$  around  $Q_0$ , we recover Eq. (28) for the correction to the ordering wave vector. U sing Eqs. (2), we not from Eqs. (60)-(61) that

$$J = 0.472 \text{ meV}; J^0 = 0.066 \text{ meV}:$$
 (62)

W e see that in this approach,  $J^0$  is renorm alized down from its bare value, while J increases, as discussed in Sec. II.

In spite of the fact that we expect the system to be unstable for D = 0 and  $J^0 = J = 1=3$ , it is of pedagogical interest to exam ine the e ects of anham onicity and the shift of the G oldstone m ode for this case. The right hand panel of Fig. 3 shows the results for the renorm alized energy spectrum and its comparison to the linear spinwave dispersion along the (1;0) direction for D = 0. The inverse lifetime of quasiparticles Eq. (59) is also shown. The excitations are seen to be heavily overdam ped close to the (10) point. A lso kinks are seen at various places in the renorm alized curves, apparently at points where oneto-two m agnon decay channels turn on. A lthough not shown in Fig. 3, we also nd that the band width of the dispersion along (1;0)! (1;1) is renorm alized dow nw ard by about a factor of 3 due to anham onic e ects.

In the left hand panel of Fig. 3, we plot the renormalized energy spectrum for  $D = 0.02 \text{ meV}^{23}$  In the presence of this DM interaction, it su ces to use the bare values for the coupling constants in expressions for self-energies, since the nite gap in the spectrum does not lead to any problem atic behavior near Q when D is not too small. The e ects of quantum uctuations are very sensitive to the presence of a non-zero D. Increasing the value of D reduces the interaction-induced dam ping of quasiparticles, making them more stable. Quantum




FIG. 3: (Color Online) The two panels present the results for the bare and renormalized energy spectra for the cases of D = 0 and D = 0.02 meV. Spin-wave energies are shown together with the inverse lifetime  $_{k}$ . The path in the twodimensionalBrillouin zone sweeps from (00) point to (10).  $k_{x}$ ,  $k_{y}$  are measured in units of 2 and 2 = 3 respectively.

uctuations, in addition to shifting Q towards , reduce the size of the gap at this point. However, the e ect of suppression of the gap towards zero, is relatively sm all for D = 0.02 m eV.

# VI. SCATTERING INTENSITY

In this Section, the dynam ical properties of an antiferrom agnet described by the model of Eq. (1) are calculated within the fram ework of the 1=(2S) expansion. We are speci cally interested in answering the question of whether anharm onic spin waves can account for the main features seen in inelastic neutron scattering measurements on  $C s_2 C u C l_4$ .<sup>14</sup> We begin with expressions for the structure factor components that enter the form ula for the inelastic neutron scattering cross section. These results were obtained earlier by O hyam a and Shiba,<sup>29</sup> and we sim ply quote them here.

Neutron scattering spectra of spin excitations in m agnetic solids may be expressed in terms of the Fourier transformed real-time dynamical correlation function, i;k = (x;y;z)

$$S^{ik}(k;!) = \frac{1}{2} \int_{1}^{Z_{1}} dt K^{K}_{R}(0)S^{k}_{R}(t)ie^{i(!t k R)}$$
(63)

For the spiral spin density wave state,  $S^{xx}(k;!) = S^{zz}(k;!)$ . The complete expression for the inelastic, differential scattering cross-section, including polarization factors, is given by<sup>27</sup>

$$I(k;!) = p_{x}S^{xx}(k;!) + p_{y}S^{yy}(k;!);$$
(64)

$$p_x = 1 + \cos^2_{k}; \quad p_y = \sin^2_{k};$$
 (65)

where k is the angle between the scattering wave vector k and the axis perpendicular to the plane of the spins.

We begin by calculating the corresponding timeordered spin-spin correlation functions in the rotated coordinate system

$$iF (k;!) = dte^{i!t}hf^{S}_{k}(0)S_{k}(t)i; \quad (66)$$

$$S^{YY}(k;!) = \frac{1}{2} \operatorname{Im} F(k;!);$$
 (67)

$$S^{xx}(k;!) = S^{zz}(k;!) = \frac{1}{-1} Im^{+}(k+Q;!) + (k-Q;!);$$
 (68)

$$S^{xz}(k;!) = S^{zx}(k;!) = \frac{1}{2} \operatorname{Im}^{+}(k+Q;!) \quad (k \quad Q;!); \quad (69)$$

with,

$$(k;!) = \frac{1}{4} F (k;!) + F (k;!) iF (k;!) F (k;!) :$$
 (70)

W e treat the interactions between the m agnons as a perturbation in the form ally sm all parameter 1=(2S), and use the standard  $\hat{S}$ -m atrix expansion while calculating the averages of the m agnon operators  $a_k$  (t).<sup>28</sup> A s a result,

$$F (k;!) = \frac{S}{2}c_{y}^{2} [G_{11}(k;!) + G_{22}(k;!) 2G_{12}(k;!)];$$

$$F (k;!) = \frac{S}{2}c_{x}^{2} [G_{11}(k;!) + G_{22}(k;!) + 2G_{12}(k;!)];$$
(71)

where the G reen's function are determ ined from Eq. (37) with the self-energies given by Eqs. (39)-(42), and, to the relevant order in 1=(2S),

$$c_{y} = 1 \frac{1}{4SN} X 2v_{k}^{2} u_{k}v_{k} ;$$

$$c_{x} = 1 \frac{1}{4SN} X 2v_{k}^{2} + u_{k}v_{k} ;$$
(72)

The term that m ixes the transverse and longitudinal uctuations has the form :

$$i F (k;!) F (k;!) = C_{x} P^{(1)}(k;!) [G_{11}(k;!) + G_{22}(k;!) 2G_{12}(k;!)] + P^{(2)}(k;!) [G_{11}(k;!) - G_{22}(k;!)] :$$
(73)

with the functions  $P^{(1;2)}(k;!)$  de ned as

$$P^{(1)}(k;!) = \frac{S}{4N} \bigvee_{q}^{(1)}(q;k q) (u_{q}v_{k} q + v_{q}u_{k} q) \frac{1}{!_{q} + !_{k} q} + \frac{1}{!_{q} + !_{k} q + ! i};$$

$$P^{(2)}(k;!) = \frac{S}{4N} \bigvee_{q}^{(2)}(q;k q) (u_{q}v_{k} q + v_{q}u_{k} q) \frac{1}{!_{q} + !_{k} q} \frac{1}{!_{q} + !_{k} q + ! i};$$
(74)

As before,  $!_k = 2S''_k$ . The longitudinal correlations expanded in inverse powers of 2S can be expressed as a sum of two contributions

$$F (k;!) = F_0 (k;!) + F_1 (k;!);$$
(75)

where

$$F_{0}(k;!) = \frac{1}{2N} \left( u_{q}v_{k} + v_{q}u_{k} \right)^{2} \frac{1}{!_{q} + !_{k} + !_{q} ! i} + \frac{1}{!_{q} + !_{k} + !_{q} ! i} ;$$
(76)

$$F_{1} (k;!) = \frac{1}{2S} P^{(1)}(k;!) {}^{2} [G_{11}(k;!) + G_{22}(k;!) + 2G_{12}(k;!)] + P^{(2)}(k;!) {}^{2} G_{11}(k;!) + G_{22}(k;!) + 2G_{12}(k;!)] + 2P^{(1)}(k;!) P^{(2)}(k;!) [G_{11}(k;!) - G_{22}(k;!)]$$
(77)

Note that the term  $F_1$  (k;!) is form ally of order  $1=(2S)^2$  compared to the leading one magnon contributions Eq. (71). However, it is kept because it contains the self-energy corrections to Eq. (76) describing the two magnon continuum.

U sing the set of formulas above, we have calculated the scattering intensities as a function of frequency for various trajectories in k and ! space, corresponding to the m easured spectra.<sup>14</sup> In generalour results agree with those of Veillette et al.<sup>23</sup> except for sm all di erences due to their use of self-consistent energy denom inators in the self-energies and our use of shifted Q values in the anharm onic G reen's functions. Veillette et al. restricted their attention to values of J, J<sup>0</sup> and D given by Eqs. (2)-(3). In the discussion that follows, we will also consider the consequences of varying the D parameter, which both shifts certain m odes and also controls the size of quantum uctuations, as well as varying the overall energy scale, J.

W e focus our attention on two trajectories in the (k;!)space whose param eterizations correspond to the scans G and J of Ref. 14,

G: 
$$k_x = 0.5; k_y = 1.53 \quad 0.32! \quad 0.1!^2$$
  
J:  $k_y = 0.47; k_y = 1.0 \quad 0.45!$ : (78)

In Eq. (78),  $k_x$  and  $k_y$  are written in the units of 2 and  $2 = \frac{1}{3}$  respectively, while the energy is measured in m eV. Since the momentum transfer lies in the x-z plane,  $p_x =$  $p_v = 1$  for both of those scans. It is important also to take into account the nite energy and momentum resolution of the scattered neutrons. The nite energy resolution 1 0:016 m eV leads to a sm all Lorentzian broadening of the one-m agnon peaks. How ever, the e ect of sm earing due to the nite momentum resolution k=2 = 0.085is much more pronounced and considerably broadens the structure of the G and J scans. W hat is special about the G and J scans is that they both exhibit sharp low energy m odes with energies around 0.1 m eV. A s we shall see, the energies of these peaks are quite sensitive to the exact value of the DM interaction D.

In Fig. 4, we plot the results of scattering intensity calculations for scan G, a data set which was analyzed in detail by Coldea et al.<sup>14</sup> In particular, it was claim ed that the high-energy tail in this scan has too much intensity, by nearly an order ofm agnitude, to be explained by linear spin-wave theory. Our calculations were done using the bare couplings J and  $J^0$ , and the D zyaloshinskii-M oriya interaction was set equal to D = 0.02 meV, the value obtained from experiments in high magnetic eld.<sup>24</sup> The theory predicts three peaks, one corresponding to the principal mode "<sub>k</sub> and the other two, re ecting the




FIG.4: (ColorOnline) Intensity of the scattered neutrons for scan G in Eq. (78), in the presence of D zyaloshinskii-M oria interaction D = 0.02 meV. The energy and momentum resolutions are taken to be ! = 0.016 meV and k = 0.085 respectively. The thick solid line is the total scattering intensity, Eq. (64). O ther lines represent the various contributions appearing in Eqs. (67)-(71).

presence of the shifted secondary spin waves at " $_{0}$  k. For the non-renorm alized couplings J and  $J^0$ , Eq. (24) for the bare spectrum predicts that the principal peak should be at 0:22 meV while the secondary peaks are very close to each other and located approxim ately at energy 0:28 m eV. Inclusion of self-energies to order 1= (2S) renorm alizes the position of the main peak down to 0:18 m eV. The principal one-m agnon peak should lie below a two-m agnon scattering continuum starting exactly at the location of the principal peak. How ever, the presence of self-energies in the G reen's functions, leads to a shift in positions of their poles, and to a nite gap between the principal peak and the start of the two magnon scattering continuum. This result is an artifact of truncating the perturbation expansion at a nite order, as discussed by Veillette et al.<sup>23</sup> W e also observe that the two secondary peaks are so close to each other that it is di cult to distinguish them, once the nite frequency and m om entum resolution is fully incorporated into the calculations. Experim entally, the main peak is observed at a som ew hat sm aller energy of 0:107 m eV, while the secondary ones are located around 0:25 m eV.<sup>14</sup> Fig. 4 also shows the sign and magnitude of the various contributions to the total intensity. It is seen that the two-m agnon part, integrated over the full range of frequencies, carries a considerable




FIG.5: (Color Online) Intensity of the scattered neutrons corresponding to scan G, for the value of D zyaloshinskii-M oria interaction D = 0.01 m eV. Upper panel: The total structure factor with the energy (m om entum) resolution E = 0.016 m eV ( k = 0.085). Inset: Experim ental data from Ref. 14. Lower panel: D i erent components which contribute to the structure factor (see Eqs. (67)-(71) in the text).

overallweight com pared to the one-m agnon contribution. At the same time, the term mixing the transverse and longitudinal uctuations can have both signs, and being sm all, does not in uence the intensity pro levery much.

W hy is the two-m agnon intensity in our calculation so much larger, relative to the principal peak than in the analysis of C oldea et al.? The reason is the factor  $c_y^2$  which renorm alizes the out-of-plane scattering and which is basically due to the reduction of the localm om ent by quantum uctuations. As noted above, theory predicts a much smaller localm om ent than the surprisingly large value obtained from experim ent.<sup>31</sup> A coording to our calculations, the moment reduction increases with decreasing D , and the ordered moment vanishes for D < 0:003 m eV.

In Fig. 5, we plot the theoretical result for scan G, but this time with a reduced DM interaction D = 0.01 m eV.




FIG. 6: (Color Online) Intensity of the scattered neutrons corresponding to scan J, Eq. (78), in the presence of D zyaloshinskii-M oria interaction D = 0.02 meV. The di erent components are displayed as in Figs. 4, 5

The comparison of our calculations with the experim entalm easurem ents reported in Ref. 14, is displayed in the upperpanel. W e see that for this value of D, the position of the principal magnon peak is shifted towards smaller energies. The gap between the main peak and the energy where the continuum appears, is also decreased; there is further broadening of the principal peak and m ore intensity in the two-magnon continuum. It appears that the value D = 0:01 m eV provides a better explanation of the form of the total scattering intensity for scan G than D = 0.02 meV. Comparison of the plots for a range of values of D, shows that the scattering intensity is very sensitive to the strength of the DM interaction. Note that the DM interaction was not included in the analysis of Ref. 14.] Finally, we observe that even this smaller value of D = 0.01 meV does not account for all of the high energy scattering which seems to extend up to 1 m eV, suggesting additional correlations at high energy which are not contained in the theory.

Next, we exam ine scan J which probes the vicinity of the ordering wave vector.<sup>14</sup> Inelastic scattering at this point should show a primary peak at the wave vector Q and two secondary modes, corresponding to mok.T mentum transfer  $k_J$  Q. The primary mode, which is prom inent in the theoretical spectrum, directly m easures the strength of the DM interaction, since its energy is proportional to DJ. The experim ental data for point J, shown only in the inset of Fig. 5G in the paper of Coldea and  $co-workers_{r}^{14}$  do not extend to low enough energy to display this peak. Nevertheless the position of the peak, at an energy of 0.10(1) m eV, is listed in Table 1 of that paper. This value of the mode at the J point is noticeably smaller than anham onic spin-wave theory would predict for D = 0.02 m eV, suggesting that the e ective value of the DM interaction constant D is smaller than 0.02 meV. Since the theory, for D = 0.02 meV, gives a peak at 0.15 meV, a t to the measured peak at 0.1 meV would suggest a value of D in the range of 0.008 to 0.01 meV. Of the two secondary peaks at the J point, one has an energy close to zero, corresponding to the shifted zone-center mode, while the other lies at higher energy, just above 0.3 meV.

#### VII. DISCUSSION AND CONCLUSIONS

The main motivation for the work presented in this paper was to see if a relatively conventional anharm onic spin-wave calculation could account for neutron scattering measurements on  $Cs_2CuCl_4$ . Our conclusion is a quali ed \yes".

W ithout any calculations, one knows that the observed excitations are from an ordered state exhibiting magnetic B ragg peaks, and hence that the low est-lying excitations must be spin waves. However, both linear and anharm onic spin-wave theory suggest a substantial moment reduction for a broad range around the expected values of D =J and  $J^0=J$ . Further experiments would be useful to verify the amount by which the moment is reduced, since the present result<sup>31</sup> is inconsistent both with anharm onic spin-waves and with close proximity to a spin-liquid state.

At low energies we nd that we can t the sharp spinwave features of the data, albeit with a som ewhat smaller value of the DM interaction constant, D, than was obtained from high eld measurements of a fully polarized state. We also nd from theoretical calculations that, without a su ciently large DM interaction, long range order is destroyed by quantum uctuations. However, the value of D that best ts theory to experiment is well within the range in which order is stable, at least for T = 0. Additional experiments, including measurements of the renormalized D in zero eld, might clarify this point. (We note in passing that a value of J about 30-40% larger than given by Eq. (2), for xed  $J^0=J$ , would provide a much improved t to the higher energy onespin-wave features.)

O ur theory yields a considerable am ount of continuous two-spin-wave scattering. However the am ount of this scattering, although readily measurable, is somewhat less than is observed experimentally. Furthermore, calculation of the relative size of one- and two-spin-wave features is complicated by their sensitivity to the size of D. That is, a smaller D suppresses one-spin-wave scattering and makes two-spin-wave continuous scattering relatively more conspicuous. Quantum uctuations also broaden the one-spin-wave features via lifetime e ects. However, it is di cult to assess, from the data alone, how much of the observed broadening of the principal peaks is due to lifetimes and how much is due to low momentum resolution as was emphasized by Veillette et al.<sup>23</sup> The theory suggests though that experiments with higher momentum resolution are likely to reveal considerably sharper low energy spin-wave peaks.

W ith regard to the shape of the broad scattering at higher energies, we are inclined toward a picture of scattering by quasi-one-dimensional spinons. W ith a bare interchain coupling of 0.128 meV, which we know to be considerably renormalized downward by quantum uctuations, one m ight expect that, at energies above about 0.3 meV, the system would behave like a collection of uncoupled chains. In this sense we would expect the high-energy response of the system to be better described by one-dimensional spinons than by a low-order spin-wave perturbation calculation such as the one that we have studied.

## A cknow ledgm ents

This project has been supported by the Natural Sciences and Engineering Research Council (NSERC) of Canada and by the Canadian Institute for A dvanced Research (CIAR). The authors gratefully acknow ledge valuable discussions and correspondence with Martin Veillette, who also provided access to a particularly useful com puter program. Som e of the research for this paper was done at the A spen Center for Physics.

# APPENDIX A:1=(2S)<sup>2</sup> CORRECTION TO THE GROUND STATE ENERGY AND ORDERING WAVE VECTOR

The ground state energy correction at the order 1=(2S)<sup>2</sup>, denoted here as  $E_{\rm G}^{(2)}$  (Q), consists of two parts,

$$E_{G}^{(2)}(Q) = E_{3}(Q) + E_{4}(Q):$$
 (A1)

 $E_4$  (Q) is obtained simply by substituting the Bogoliubov transformation Eq. (22) into H<sup>(4)</sup> and perform – ing the normal ordering in the terms containing two pairs of quasiparticle operators c and c<sup>V</sup>. In this Appendix, we present the results of calculations in which the D zyaloshinskii-M oria interaction D = 0, so that

$$E_{4} (Q) = \frac{1}{2N} X_{k,p}^{R} + \frac{1}{N} \frac{1}{N} X_{p}^{R} \frac{A_{p}}{P} + \frac{A_{k}}{2} + \frac{1}{N^{2}} X_{k,p}^{R} (A_{k,p} - B_{k,p}) \frac{A_{k}A_{p} + B_{k}B_{p}}{4^{n}_{k} T_{p}} : (A2)$$

 $E_3$  (Q) ensues due to the key role of the cubic interactions (19) in the total H am iltonian. It is technically advantageous rst to express (19) in terms of the quasiparticle c-operators, so that

$$H^{(3)} = \frac{i}{2} \begin{bmatrix} \frac{1}{2} & \frac{1}{2N} \\ \frac{1}{1;2;3} & \frac{1}{3} \\ + \\ 2(1;2;3) & c_3^y c_2 c_1 \\ \frac{1}{2} & c_1^y c_2^y c_3^y \\ + \\ 2(1;2;3) & c_3^y c_2 c_1 \\ \frac{1}{2} & c_1^y c_2^y c_3 \\ \frac{1}{2} & c$$

+ 
$$(C_2 C_3)(u_1v_2u_3 + v_1u_2v_3)$$
: (A5)

At T = 0, the only contribution to  $E_3$  (Q) is given by the 'sunrise' diagram containing two vertices  $_1$  (q;p; q p). Using the form ulas (23), we arrive at the following expression for  $E_3$ :

$$(1;2;3) = (C_{1} + C_{2}) (u_{1}u_{2}v_{3} + v_{1}v_{2}u_{3})$$

$$+ (C_{1} + C_{3}) (u_{1}u_{3}v_{2} + v_{1}v_{3}u_{2})$$

$$+ (C_{2} + C_{3}) (u_{2}u_{3}v_{1} + v_{2}v_{3}u_{1}); (A 4)$$

$$_{2} (1;2;3) = (C_{1} + C_{2}) (u_{1}u_{2}u_{3} + v_{1}v_{2}v_{3})$$

$$+ (C_{1} - C_{3}) (v_{1}u_{2}u_{3} + u_{1}v_{2}v_{3})$$

$$E_{3}(Q) = \frac{1}{24N^{2}} \frac{X}{p_{jq}} \frac{\left[\frac{1}{q}(q;p;q-p)\right]^{2}}{p_{p}(q+q+p_{p+q})} = \frac{1}{16N^{2}} \frac{X}{p_{jq}} \frac{1}{(p_{p}+q+q+p_{p+q})} C_{p+q}^{2} \frac{A_{p+q}+B_{p+q}}{p_{p+q}} \frac{A_{p}A_{q}+B_{p}B_{q}}{p_{p}q} = 1$$

$$+ C_{p}C_{q} \frac{(A_{p+q}+B_{p+q})(A_{p}+B_{p})(A_{q}+B_{q})}{p_{p+q}q^{2}p} + \frac{A_{p+q}}{p_{p+q}} \frac{B_{p+q}}{p_{p+q}} \frac{A_{p}+B_{p}}{p_{p}} \frac{A_{q}+B_{q}}{q} : (A6)$$

Numerical calculations show that for the interaction constants given by Eqs. (2),

$$E_3 (Q_0) = J = 0.0425;$$
 (A 7)

$$E_4 (Q_0) = J = 0.0103;$$
 (A 8)

which are quite sm all. The  $1=(2S)^2$  correction to  $Q_0$  can be computed by substituting the expansion

$$Q = Q_0 + \frac{Q^{(1)}}{2S} + \frac{Q^{(2)}}{(2S)^2}$$
 (A 9)

into the series for the ground state energy

$$E_{G} (Q) = S^{2} J_{Q} + SE_{G}^{(1)} (Q) + E_{G}^{(2)} (Q); \quad (A10)$$

and expanding its minimum in powers of 1=(2S). As a result,

$$Q^{(2)} = \frac{Q^2 J_Q}{Q Q^2} \frac{1}{2} \frac{1}{2} \frac{Q^3 J_Q}{Q Q^3} (Q^{(1)})^2$$

$$+2\frac{@^{2}E_{G}^{(1)}(Q)}{@Q^{2}} (Q^{(1)}) + 4\frac{@E_{G}^{(2)}(Q)}{@Q} (A 11)$$

The calculations of derivatives over Q in expressions entering Eqs. (26), (A6), (A2) are lengthy but straightforward. Om itting the corresponding cum bersom e nal expressions, we state here that as a result of num erical integration:

$$Q^{(2)}=2$$
 0:011: (A 12)

The value of the ground state energy at its m in imum (S = 1=2)

$$E_{m in} = J = \frac{1}{J} S^{2} J_{Q} + S E_{G}^{(1)} (Q) + E_{G}^{(2)} (Q) - \frac{1}{8} \frac{(Q^{2} J_{Q})}{(Q^{2})^{2}} (Q^{(1)})^{2} = 0.459:$$
(A13)

Note that the last term in Eq. (A13) is always negative and arises solely due to the shift in the ordering wave vector  $Q^{(1)}$ . Numerically however, this term is tiny, even compared to the second order correction  $E_{G}^{(2)}(Q_{0})$  that itself gives a small contribution to the total energy Eq. (A13).

<sup>1</sup> See T. Senthil, cond-m at/0411275, and references therein. <sup>2</sup> P.W. Anderson, Mater. Res. Bull. 8, 153 (1973); P. Fazekas

and P.W .Anderson, Philos.M ag. 30, 23 (1974).

- <sup>3</sup> S.A.K ivelson, D.S.R okhsar, and J.P.Sethna, Phys.Rev. B 35, 8865 (1987).
- <sup>4</sup> A. O. Gogolin, A. A. Nersesyan and A. M. Tsvelik, Bosonization in Strongly Correlated Systems (Cambridge University Press, Cambridge, 1999).
- <sup>5</sup> See, e.g., Perspectives in Quantum Hall E ects: Novel Quantum Liquids in Low-D imensional Sem iconductor Structures, Ed.by S.D as Sarm a and A.Pinzcuk (W iley, New York, 1996).
- <sup>6</sup> R.Moessner and S.L.Sondhi, Phys. Rev. Lett. 86, 1881 (2001).
- $^7\,$  C .N ayak and K .Shtengel, Phys.R ev.B 64,064422 (2001).
- <sup>8</sup> L.Balents, M.P.A.Fisher, and C.Nayak, Int.J.Mod. Phys.B 12, 1033 (1998).
- <sup>9</sup> L.Balents, M.P.A.Fisher, and C.Nayak, Phys.Rev.B 60, 1654 (1999).
- <sup>10</sup> L.Balents, M.P.A.Fisher, and C.Nayak, Phys.Rev.B 61 6307 (2000).
- <sup>11</sup> E.Dem ler, C.Nayak, H.-Y.Kee, Y.B.K in , and T.Senthil, Phys. Rev. B 65, 155103 (2002).
- <sup>12</sup> T. Senthil and O. M otrunich, Phys. Rev. B 66, 205104 (2002); O. I. M otrunich and T. Senthil, Phys. Rev. Lett. 89, 277004 (2002).
- <sup>13</sup> L.Balents, M PA.Fisher and SM.Girvin, Phys.Rev.B 65, 224412 (2002).
- <sup>14</sup> R.Cokkea, D.A.Tennant and Z.Tykzynski, Phys.Rev. B 68, 134424 (2003).
- <sup>15</sup> R. Coklea, D. A. Tennant and Z. Tylczynski, Phys. Rev. Lett. 86, 1335 (2001).
- <sup>16</sup> Z.W eihong, R.H.M cK enzie and R.R.P. Singh, Phys. Rev.B 59, 14367 (1999).

- <sup>17</sup> W. Zheng, J.O. F jaerestad, R.R.P. Singh, R.H.McKenzie, and R.Coldea, cond-m at/0506400.
- <sup>18</sup> C.H.Chung, J.B.M arston and R.H.M cK enzie, J.Phys.: Condens. M atter 13, 5159 (2001); S.Takei, C.H.Chung and Y.B.K im, Phys. Rev. B 70, 104402 (2004).
- $^{19}\,$  Y .Zhou and X .G .W en, cond-m at/0210662.
- <sup>20</sup> C.-H. Chung, K. Voelker, and Y. B. Kim, Phys. Rev. B 68,094412 (2003).
- <sup>21</sup> J. Merino, Ross H. McKenzie, J. B. Marston and C. H. Chung, J. Phys.: Condens. Matter 11, 2965 (1999).
- <sup>22</sup> J. Alicea, O. I. Motrunich, M. P. A. Fisher, cond-m at/0512427.
- <sup>23</sup> M.Y.Veillette, A.J.A.Jam es, and F.H.L.Essler, Phys. Rev.B 72, 134429, (2005).
- <sup>24</sup> R. Coldea, D. A. Tennant, K. Habicht, P. Smeibidl, C. Wolters and Z. Tylczynski, Phys. Rev. Lett. 88, 137203 (2002).
- $^{25}$  T.Holstein and H.Prim ako ,Phys.Rev.58,1098 (1940).
- <sup>26</sup> Y.Tokiwa, T.Radu, R.Coldea, H.W ilhelm, Z.Tylczynski, and F.Steglich, cond-m at/0601272 (2006).
- <sup>27</sup> S.W. Lovesey, in Theory of Neutron Scattering from Condensed M atter, (C larendon, O xford, England, 1987).
- <sup>28</sup> See, e.g., A. A. A brikosov, L.P.G orkov & I.E.D zyaloshinski. M ethods of Q uantum Field Theory in Statistical Physics, (D over, N ew York, 1975)
- <sup>29</sup> T. O hyam a and H. Shiba, J. Phys. Soc. Jpn. 62, 3277 (1993).
- <sup>30</sup> A.V. Chubukov, S. Sachdev, and T. Senthil, J. Phys.: Condens.M atter 6, 8891 (1994).
- <sup>31</sup> R.Coldea, D.A.Tennant, R.A.Cowley, D.F.M.dMorrow, B.Dorner, and Z.Tylczynski, J.Phys.: Condens. M atter 8, 7473 (1996).