

Thermodynamically activated vortex-dipole formation in a two-dimensional Bose-Einstein condensate

Daniel Schumayer¹, and David A. W. Hutchinson^{1,2}

¹The Jack Dodd Centre for Photonics and Ultra-Cold Atoms,
Department of Physics, University of Otago, P. O. Box 56, Dunedin, New Zealand

²Laboratoire Kastler Brossel, Ecole Normale Supérieure,
24 rue Lhomond, 75231 Paris Cedex 05, France

(dated: March 23, 2024)

Three distinct types of behaviour have recently been identified in the two-dimensional trapped bosonic gas, namely; a phase coherent Bose-Einstein condensate (BEC), a Berezinskii-Kosterlitz-Thouless-type (BKT) superfluid and normal gas phases in order of increasing temperature. In the BKT phase the system favours the formation of vortex-antivortex pairs, since the free energy is lowered by this topological defect. We provide a simple estimate of the free energy of a dilute Bose gas with and without such vortex dipole excitations and show how this varies with particle number and temperature. In this way we can estimate the temperature for cross-over from the coherent BEC to the (only) locally ordered BKT-like phase by identifying when vortex dipole excitations proliferate. Our results are in qualitative agreement with recent, numerically intensive, classical field simulations.

PACS numbers: 03.75.Hh, 03.75.Lm

Keywords: Bose-Einstein condensation, Berezinskii-Kosterlitz-Thouless phase transition, vortex-dipole

Since the experimental realisation of Bose-Einstein condensation (BEC) [1], tremendous advances have been made in exploring the physics of ultracold atomic and molecular gases. It is now possible to simultaneously prepare multiple condensates and collide them [2, 3] or even slice a condensate into numerous pieces using optical standing-waves [4]. Such an optical potential, in addition to the normal harmonic confinement, allows one to create a highly anisotropic trap and in this way it is possible to enter the quasi-two- [4, 5, 6, 7] or one-dimensional [5] regimes.

In lower dimensions the properties of quantum systems are radically altered. Unlike in the three-dimensional (3D) case, the different density of states means condensation cannot occur either in two- (2D) or one-dimensional (1D) homogeneous ideal Bose gases in the thermodynamic limit at finite temperature [8, 9].

The application, however, of a trapping potential introduces a low-momentum cutoff which suppresses long-wavelength fluctuations and allows the possibility of BEC [10, 11] below a certain critical temperature, which for the ideal 2D case we denote T_0^{2D} .

In the homogeneous 2D system, although BEC is absent at finite temperature, superfluidity can exist with algebraically decaying order [12, 13, 14] in an interacting but spatially uniform Bose gas. This is the Berezinskii-Kosterlitz-Thouless (BKT) phase in which the gas has local order but no global phase coherence, with no uniform order parameter. A local order parameter can be defined and is known as the quasi-condensate [15].

A similar scenario persists in the presence of external

confinement. Three distinct regions can be identified in a two-dimensional trapped interacting bosonic system. At temperatures much below T_0^{2D} a phase coherent BEC appears to exist [16, 17]. At higher temperatures the coherence length decreases to length scales smaller than the system size [18] and bound vortex-antivortex pairs (vortex-dipoles) begin to form [19] indicating a BKT-like phase. Eventually, around T_0^{2D} , the vortex-dipoles unbind and the correlation length decays exponentially. This is the analogue of the BKT phase transition to the thermal gas [20, 21].

A phase diagram and approximate critical temperature for the BEC to BKT cross-over were derived numerically in [19] assuming that the vortex-dipole is always at the centre of the trap, i.e. placed at the highest density. Here, we use simple analytic arguments, without this restriction allowing the vortex-dipole to be placed anywhere inside the Thomas-Fermi radius, to obtain a very simple picture of the vortex dipole proliferation in the gas. Our approach obviates the need for any detailed numerical calculations and qualitatively agrees with the more sophisticated approaches, including the full classical field simulations [20].

In our model, we minimize the Helmholtz free-energy, $F = U - TS$, where U denotes the total internal energy and S the entropy of the state, and the system has constant volume V and an absolute temperature T . We partition the gas into two subsystems, namely, a condensate and a thermal part, each with its own internal energy and entropy contribution, (E_c, E_t) and $(S_c$ and $S_t)$, respectively. In the condensate part we allow for a vortex-dipole excitation, i.e. a bound vortex and antivortex pair. It suffices to consider only the effect of one pair in the condensate with the additional configurational entropy of the vortex dipole denoted as S_v .

Electronic address: dschumayer@physics.otago.ac.nz

The important contributions come from the condensate, with the thermal component only producing global energy shifts. Accordingly, E_t and S_t are only roughly approximated. The free energy of the system can now be written as $F = U - TS = (E_c + E_t) - T(S_c + S_v + S_t)$.

The model system consists of N interacting bosonic particles of mass m confined by an external non-rotating harmonic potential characterized by the $\omega_x, \omega_y, \omega_z$ trap frequencies. The actual values of the physical parameters we take correspond to the experiments of [4, 7]. In both experiments the radial and axial trapping frequencies differ by several orders of magnitude, suggesting that we use a quasi-2D formalism. This assumption is confirmed by comparing the characteristic oscillator length and the s-wave scattering length [16]. Consequently, we assume that the wavefunction for the condensate can be

factorized into two parts: $\Psi = \psi(x, y) \phi(z)$, where ψ describes the state in the xy -plane, while ϕ is simply the ground state wavefunction of a harmonic oscillator in a trap characterized by ω_z . The wavefunction ψ we normalize to the condensate particle number N_0 and ϕ to unity.

Moreover, we introduce an arbitrary spatial unit x_s , rescaling the spatial variables and the wavefunction ψ to be dimensionless. In the numerical calculation the spatial unit x_s was taken to be equal to the oscillator length $a_{osc,z}$ corresponding to the frequency ω_z .

Substituting this composite wavefunction into the usual Gross-Pitaevskii energy functional and evaluating all integrals with respect to the z direction, we obtain the following expression for the energy E_c of the condensate

$$E_c = N_0 \frac{\omega_z}{2} + \frac{\omega_z^2}{2m x_s^2} \int r^{2D} |\psi|^2 + V_{ext}^{2D}(r) |\psi|^2 + 2 \frac{\mu}{2} \frac{a_s}{a_{osc,z}} \int |\psi|^4 dr; \quad (1)$$

where $r^{2D} = (x, y)$, and the potential term V_{ext}^{2D} is cylindrically symmetric, $(x_s = a_{osc,z}) (x^2 + y^2)$. Interaction between particles is taken into account via the 3D s-wave scattering length, a_s .

The internal energy of the thermal part was approximated by the energy of the trapped but non-interacting Bose-gas [22]

$$E_t = (N - N_0) k_B T_0^{2D} \frac{3}{(3)} \frac{T}{T_0^{2D}}^4; \quad (2)$$

The corresponding entropy contribution S_t can be derived from Eq. (2) using the standard formulae of statistical physics. The validity of this approximation is also justified by experiment [4], where the thermal cloud was not fully within the 2D regime. The total number of particles was close to the theoretical threshold [23], above which the atoms squeeze out to the third dimension even if the condensate dimension in that direction is smaller than the healing length.

Naturally, a more sophisticated treatment is possible and in particular, the presence of the thermal cloud could significantly affect the energy of the vortex core [24]. To go beyond our simple treatment would require a numerically intensive, self-consistent, finite-temperature Hartree-Fock-Bogoliubov treatment [25, 26, 27] which is not in the spirit of this exercise. Here we aim at the simplest possible estimate of the critical temperature for vortex dipole proliferation and we do not expect inaccuracies in the vortex core energy at high temperatures to qualitatively affect our conclusions. This expectation is confirmed by comparison with the (numerical) classical field simulations of Simula and Blakie [20, 28].

In order to evaluate the free energy functional F , we

must also approximate the entropy, which is proportional to the number of physically available microstates. Only the ground state is available for particles in the condensate, therefore $S_c = 0$. On the other hand, if a vortex-dipole is present in the condensate then its entropy can be estimated via a simple geometrical approach (see Fig. 1) such that $S_v = k_B \ln(2x_v/a_s)$, where x_v denotes the distance of the vortex-dipole from the centre. The healing length characterizes the length over which the density depression vanishes and the wave function approaches its bulk value; it is defined by $\xi = \sqrt{4 \frac{\mu}{2} (a_s = a_{osc,z})} \frac{1}{j_{TF}(x_v)}$.

It is now clear that only E_c and S_v have spatial dependence and the remaining energy and entropy expressions simply shift the free-energy by a constant value at a given temperature.




FIG. 1: The dark ring represents the equivalent microstates for the vortex pair displaced at distance x_v from the centre.

Finally, to evaluate E_c in mean-field theory, one needs a wavefunction for the condensate containing a vortex-dipole excitation. The square modulus of the wavefunction ψ without any excitation is chosen to have the Thomas-Fermi density profile on which we superpose the vortex-dipole excitation in the form $\psi(r) = \psi_{TF}(r) \psi'_{-1}(r) \psi'_{+1}(r)$, where the vortex or antivortex wavefunction is

$$\psi'_q(r) = \tanh \frac{\mathbf{r} \cdot \mathbf{r}_q}{r_q} e^{iq \phi(r; r_q)} \quad q = -1; +1:$$

The two values of q correspond to a vortex and antivortex and $\mathbf{r}_q = (x_q; y_q)$ denotes the positions of the localized excitations. The hyperbolic tangent ensures the proper asymptotes both in the vortex core and far from the vortex. The phase $\phi(r; r_q)$ measures the angle of \mathbf{r} with respect to the x -axis. It gives a 2π phase difference if one integrates along a path encircling the vortex core. We note here that the thermal cloud could infiltrate into the cores of vortex-dipoles and may enhance the expulsion of the condensate [24] which, as discussed previously, our model does not take fully into account. The vortex-dipoles appear at the edge of the condensate however, where the disturbance due to the thermal cloud is expected to be small. We therefore expect our simple model to be qualitatively accurate.

Three different vortex-dipole configurations were analyzed, namely a) $(x_{-1}; y_{-1}) = (x_1; y_1)$, b) $(x_{-1}; 0) = (x_1; 0)$ and c) $(x_{-1}; 0) = (x_1; -x_1; 0)$.

In the calculation, the total particle number N and the scaled temperature $T = T_0^{2D}$ were varied. For every pair of N and T we determined the position $x_v = x_1$ of the vortex dipole to be the point where the free-energy reaches its minimum. These minima belonging to configuration a), are represented in Fig. 2 (a).

As expected, the position decreases as the temperature approaches the Bose-Einstein condensation temperature T_0^{2D} , meaning that the minima of free energy are deeper inside the condensate. Simultaneously, however, the increasing particle number opposes the vortex-dipole formation well inside the condensate. Since vortex nucleation is not just a phase-pattern but also involves a density dip, more energy is required to create and maintain a density disturbance in a larger condensate.

In the $T \rightarrow 0$ limit, one would anticipate that x_v would approach R_{TF} . The small deviation from this expectation seen in Fig. 2 (a) is attributable to the numerical procedure, since we require the vortex-dipole remain within the Thomas-Fermi radius.

Theoretically speaking, statistical physics allows for the existence of a vortex-dipole excitation inside the condensate phase with non-zero probability at every finite temperature, since the likelihood of nucleation is proportional to $\exp(-F/k_B T)$. However, there exists a well-defined temperature T_c above which this probability becomes comparable with unity. It is an important question how this temperature varies with respect to the total number of particles. Using the data depicted in Fig. 2 (a)




FIG. 2: (a) Scaled minima of free energy versus scaled temperature. Data are for configuration a). The horizontal dashed line is a reference used in the text. (b) The temperature required for a vortex-dipole to penetrate into the condensate to a distance of $x_v^{(0)}$ as a function of total particle number N . The dashed line is a guide to the eye.

one can infer the behaviour of $T_c(N)$. Let us fix x_v at an arbitrary value $x_v^{(0)}$ (in our case $0.86R_{TF}$). Calculating the intersections of this line with the curves belonging to various particle numbers [see the horizontal dashed line in Fig. 2 (a)], one obtains $(N_i; T_i)$ pairs. This T_i represents the 'critical' temperature for which a system with N_i particles has a minimizing vortex-dipole position at the given distance $x_v^{(0)}$ from the centre. If we increase the temperature the vortex-pair can enter further.

The results are represented in Fig. 2 (b) and show that the temperature naturally approaches the condensate critical temperature as the total number of particles grows. The slope of this graph naturally depends upon the value of $x_v^{(0)}$ but the tendency itself does not and these findings

are comparable with [19].

One can also conclude that it is rather difficult to observe the transition from BEC to BKT state experimentally. At low N the condensate would not be dependably detectable after ballistic expansion, while at high N the vortex-dipole formation in the bulk condensate phase requires a high temperature, which also results in a small condensate and large fluctuations.

In conclusion, our simple model captures the physics of previous more complex analyses. There are two contributions to the free energy which depend on the position of the vortex-dipole; the energy of the condensate and the entropy change due to the excitation. One anticipates that the energy should monotonically decline to the value of Thomas-Fermi energy as x_v approaches R_{TF} , since the vortices deform the Thomas-Fermi density to an ever-decreasing extent. Similarly, the magnitude of the entropy contribution also decreases. This is because since the characteristic size of the vortices $\propto n^{-1/2}$, the healing length rapidly increases. Although the area of the disk $W(x_v)$ of available points in which to place the vortices grows linearly with radius, this does not result in more states available for occupation due to the increasing size of the vortex-dipole. Consequently, these two competing geometrical effects are bound to lead to a minimum in the free energy F .

A similar calculation has been performed in [29] and it was found that for a given angular momentum carried by the vortex dipole, the energy curve has a minimum inside

the condensate. However, a direct numerical comparison with that result is not possible due to the differing approximations used.

Summarizing our result we can say that (i) the configurations of vortex dipoles are not energetically equivalent; configuration a) seems to be able to enter the bulk condensate more easily than the others as the temperature increases. (ii) All types of configuration suggest that vortex-dipoles prefer nucleation at the edge of the condensate, and they can penetrate further in the vicinity of the normal gas phase-transition ($0.8 < T_0^{2D} < 0.9$). This result is consistent with and supports the possibility of vortex-dipole nucleation in the experiment [4] at ENS, Paris and is also consistent with their recent observation of the cross-over from thermal gas to BKT-like phase [30]. (iii) With increasing particle number, the temperature T_c characterising the vortex-dipole nucleation also becomes higher and asymptotically reaches the critical temperature T_0^{2D} of BEC in the ideal gas. Qualitatively, the same tendency was found in [20] using the numerically intensive classical field simulation, in which dynamical effects were taken into account inherently.

The nature of the change from BEC to BKT superfluid, i.e. whether this is a slow continuous change or corresponds to a sharp cross-over, remains an open question and is the subject of further investigations.

We express our thanks for financial support from the Marsden Fund and from the C.N.R.S. of France.

-
- [1] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell, *Science* 269, 198 (1995).
- [2] R. Legere and K. Gibble, *Phys. Rev. Lett.* 81, 5780 (1998).
- [3] N. R. Thomas, N. K. Jrgaard, P. S. Julienne, and A. C. Wilson, *Phys. Rev. Lett.* 93, 173201 (2004).
- [4] S. Stock, Z. Hadzibabic, B. Battelier, M. Cheneau, and J. Dalibard, *Phys. Rev. Lett.* 95, 190403 (2005).
- [5] A. G. Ornitz, J. M. Vogels, A. E. Leanhardt, C. Ramann, T. L. Gustavson, J. R. Abo-Shaeer, A. P. Chikkatur, S. Gupta, S. Inouye, T. Rosenband, and W. Ketterle *Phys. Rev. Lett.* 87, 130402 (2001).
- [6] D. Rychtarik, B. Engeser, H.-C. Nagarl, and R. Grimm, *Phys. Rev. Lett.* 92, 173003 (2004).
- [7] N. L. Smith, W. H. Heathcote, G. H. Echenblakner, E. Nugent, and C. J. Foot, *J. Phys. B: At. Mol. Opt. Phys.* 38, 223 (2005).
- [8] N. D. Mermin and H. W.agner, *Phys. Rev. Lett.* 17, 1133 (1966).
- [9] P. C. Hohenberg, *Phys. Rev.* 158, 383 (1967).
- [10] V. Bagnato and D. Keppner, *Phys. Rev. A* 44, 7439 (1991).
- [11] W. Ketterle and N. J. van Druuten, *Phys. Rev. A* 54, 656 (1996).
- [12] V. L. Berezinskii, *Sov. Phys. JETP* 32, 493 (1971).
- [13] V. L. Berezinskii, *Sov. Phys. JETP* 34, 610 (1972).
- [14] J. M. Kosterlitz and D. J. Thouless, *J. Phys. C: Solid State Phys.* 6, 1181 (1973).
- [15] V. N. Popov, *Functional Integrals in Quantum Field Theory and Statistical Physics*, vol. 8 of *Mathematical Physics and Applied Mathematics* (Springer, 2002).
- [16] D. S. Petrov, M. Holzmann, and G. V. Shlyapnikov, *Phys. Rev. Lett.* 84, 2551 (2000).
- [17] C. Gies, B. P. van Zyl, S. A. Morgan, and D. A. W. Hutchinson, *Phys. Rev. A* 69, 023616 (2004).
- [18] C. Gies and D. A. W. Hutchinson, *Phys. Rev. A* 70, 043606 (2004).
- [19] T. P. Simula, M. D. Lee, and D. A. W. Hutchinson, *Phil. Mag. Lett.* 85, 395 (2005).
- [20] T. P. Simula and P. B. Blakie, *Phys. Rev. Lett.* 96, 020404 (2006).
- [21] N. P. Rokoffev, O. Ruebenacker, and B. Svistunov, *Phys. Rev. Lett.* 87, 270402 (2001).
- [22] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, *Rev. Mod. Phys.* 71, 463 (1999).
- [23] J. P. Fernandez and W. J. Mullin, *J. Low Temp. Phys.* 128, 233 (2002).
- [24] K. Rajagopal, B. Tanatar, P. Vignolo, and M. Tosi, *Phys. Lett. A* 328, 500 (2004).
- [25] A. Gri n, *Phys. Rev. B* 53, 9341 (1996).
- [26] D. A. W. Hutchinson, E. Zaremba, and A. Gri n, *Phys. Rev. Lett.* 78, 1842 (1997).
- [27] T. P. Simula, S. M. M. Vrtanen, and M. M. Salmaa, *Phys. Rev. A* 66, 035601 (2002).
- [28] D. A. W. Hutchinson and P. B. Blakie, *cond-mat/0602614* (2006).

[29] Q. Zhou and H. Zhai, *Phys. Rev. A* **70**, 043619 (2004).

[30] Z. Hadzibabic, P. Krüger, M. Cheneau, B. Battelier, and

J. Dalibard, *Nature* **441**, 1118 (2006).