

Exploring Foundations of Time-Independent Density Functional Theory for Excited-States

Prasanjit Samal* and Manoj K. Harbola†

Department of Physics, Indian Institute of Technology, Kanpur U.P. 208016, India

Abstract

Based on the work of Görling and that of Levy and Nagy, density-functional formalism for many Fermionic excited-states is explored through a careful and rigorous analysis of the excited-state density to external potential mapping. It is shown that the knowledge of the ground-state density is a must to fix the mapping from an excited-state density to the external potential. This is the excited-state counterpart of the Hohenberg-Kohn theorem, where instead of the ground-state density the density of the excited-state gives the true many-body wavefunctions of the system. Further, the excited-state Kohn-Sham system is defined by comparing its non-interacting kinetic energy with the true kinetic energy. The theory is demonstrated by studying a large number of atomic systems.

I. INTRODUCTION

Hohenberg-Kohn-Sham(HKS) density functional theory(DFT)^{1,2,3} has been most widely used to investigate the electronic structures of many-electron systems. It is a theory for dealing with the ground states and their properties^{4,5,6}. Applications of the Hohenberg-Kohn theorem and the Kohn-Sham construction is limited to the ground-state because it is the ground-state density of an electronic system that determines the Hamiltonian, and consequently other physical observables of the electronic system. This suggests both the ground as well as excited-state properties can be determined from the ground-state density through the Hamiltonian operator since it characterizes all the states of a system. On the other hand, to develop an excited-state DFT akin to its ground-state counterpart, it is important to describe an excited-state of a system in terms of the density of that state. Almost for the last two and half decade researchers have investigated^{7,8,9,10,11,12,13,14,15,16,17,18} the possibility of giving a formal foundation to the time-independent excited-state DFT like the HKS DFT does for the ground-states, but it is yet to come to its full fruition.

To make the excited-state calculations feasible within the density functional formalism there are two open questions to be answered: (i) Does there exist a mapping between the excited-state density and the external potential like the ground-states? (ii) Secondly, for the determination of the excited-state energies, is it possible to construct reasonably accurate exchange-correlation energy functionals? We note that although the exact form of the exchange-correlation functional is unknown for the ground-states, there are several accurate and approximate functionals^{19,20,21,22,23} in traditional DFT. The issue is to find such approximate functionals for the excited-states. The second question is partly answered through an attempt made by the present authors²⁴ in the recent past by developing an exchange energy functional within the local-density approximation(LDA) for a particular class of excited-states. This suggests that a correlation functional can also be developed in a similar fashion.

We now address the first question which is the main focus of the present study. Over the past few years, a lot of attention has been paid to the question of mapping^{25,26,27,28,29,30} from an excited-state density $\rho(\vec{r})$ to the external potential $v_{ext}(\vec{r})$, because the entire structure of time-independent excited-state DFT depends on that. A brief account of this is as follows :

The first step in establishing a mapping from an excited-state density to a many electron wavefunction is taken by looking for ρ -stationary states²⁵. These are states Ψ that reproduce a given density ρ_k (density of the k^{th} - excited-state) and simultaneously make the expectation value $\langle \Psi | \hat{T} + \hat{V}_{ee} | \Psi \rangle$ stationary. However, for a given density there are many ρ -stationary states and thus establishing a mapping requires an additional input. Levy and Nagy^{18,26} provide this by requiring that Ψ be orthogonal to $\Psi_j (j < k)$, which are to be determined by the ground-state density ρ_0 for the system under study. Thus the knowledge of the ground-state density is essential to define a map $\rho_k \longrightarrow \Psi_k$. This is reasonable because it is the ground-state density that really fixes the Hamiltonian of a system uniquely. Levy and Nagy then go on to construct a Kohn-Sham system of non-interacting electrons whose m^{th} excited-state produces the given excited-state density. Further they put forth a criterion that the ground-state density of the Kohn-Sham system is closest to the true ground state density of the system in the least square sense.

We have been studying the mapping from an excited-state density to the potential and have been investigating the Levy-Nagy criterion for constructing the Kohn-Sham system for excited-states. Our work in this direction forms the content of this paper. We show:

- (i) an explicit construction of the external potential from an excited-state density using Görling's approach²⁵,
- (ii) that the Levy-Nagy criterion^{18,26} of constructing the Kohn-Sham system is not perfect. We give reasons for it,
- (iii) that for a given state the Kohn-Sham system should be constructed by comparison of the kinetic energies of the true and the non-interacting systems.

II. GÖRLING/LEVY-NAGY FORMALISM

A. Görling's Stationary-State Formulation

We start with Görling's formulation²⁵ of the excited-state problem. The formulation is based on the constrained-search approach³¹ and provides a map from an excited-state density to a many-body wavefunction. For a given excited-state density $\rho(\vec{r})$ a ρ -stationary state $\Psi(\vec{r})$ is given by making the expectation value $\langle \Psi | \hat{T} + \hat{V}_{ee} | \Psi \rangle$ stationary with the constraint that the many-particle wavefunction $\Psi(\vec{r})$ giving the same density $\rho(\vec{r})$. Corresponding to each ρ -stationary state $\Psi(\vec{r})$ there is an external potential $v_{ext}(\vec{r})$. This has been shown by Görling. We give a different proof of it here, with the external potential $v_{ext}(\vec{r})$ arising as the Lagrange multiplier to ensure the constraint of producing the density $\rho(\vec{r})$.

A ρ -stationary wavefunction Ψ gives, by the stationarity principle

$$\langle \delta \Psi | \hat{T} + \hat{V}_{ee} | \Psi \rangle + \langle \Psi | \hat{T} + \hat{V}_{ee} | \delta \Psi \rangle = 0 \quad (1)$$

with the constraint that

$$\delta \rho(\vec{r}) = \int \{ \Psi \delta \Psi^\dagger + \Psi^\dagger \delta \Psi \} d\vec{r}_2 \dots d\vec{r}_N = 0 \quad (2)$$

Writting Eq. 1 in expanded form, we get

$$\int \left\{ \delta \Psi^\dagger \left(\hat{T} + \hat{V}_{ee} \right) \Psi + \Psi^\dagger \left(\hat{T} + \hat{V}_{ee} \right) \delta \Psi \right\} d\vec{r}_1 d\vec{r}_2 \dots d\vec{r}_N = 0 \quad (3)$$

Because of condition (Eq. 2) above this will be satisfied if

$$\int \left(\hat{T} + \hat{V}_{ee} \right) \Psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N) d\vec{r}_2 \dots d\vec{r}_N = \int f_1(\vec{r}_1) \Psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N) d\vec{r}_2 \dots d\vec{r}_N. \quad (4)$$

Thus in general ρ -stationarity of Ψ implies that it satisfies

$$\left(\hat{T} + \hat{V}_{ee} \right) \Psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N) = \left(\sum_i f_i(\vec{r}_i) \right) \Psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N). \quad (5)$$

However, since \hat{T}, \hat{V}_{ee} are symmetric operators and $\Psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N)$ is antisymmetric, it's necessary $(\sum_i f_i(\vec{r}_i))$ must also be symmetric. Thus all f_i 's must be the same function $f(\vec{r})$.

Identifying this function as $f(\vec{r}) = -v(\vec{r}) + E$, where $\lim_{\vec{r} \rightarrow +\infty} v(\vec{r}) = 0$, we get Ψ satisfying

$$\left\{ \hat{T} + \hat{V}_{ee} + \sum_i v(r_i) \right\} \Psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N) = E \Psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N). \quad (6)$$

We note that for different ρ -stationary states $v_{ext}(\vec{r})$ will be different. Thus by applying the constrained search method we get many ρ -stationary states and the corresponding external potentials. The question is which one of these corresponds to a given system. Levy and Nagy identify^{18,26} this system as the one where Ψ is orthogonal to $\Psi_j (j < k)$ for a given ground-state density ρ_0 . Thus in the Levy-Nagy theory^{18,26}, the wavefunction $\Psi[\rho; \rho_0]$ is a bi-functional of ρ and ρ_0 . One subtle point about the Levy-Nagy theory is that if the search for Ψ is restricted to the space orthogonal to $\Psi_j (j < k)$, the variational principle becomes a minimum principle. The prescription above also makes the functional $F[\rho; \rho_0] = \min_{\Psi \rightarrow \rho(\mathbf{r})} \langle \Psi | \hat{T} + \hat{V}_{ee} | \Psi \rangle$ a bi-functional of the excited-state density $\rho(\vec{r})$ as well as the ground-state density $\rho_0(\vec{r})$.




FIG. 1: Shown in the figure are the KS potentials for the excited-state density of $1s^1 2s^0 2p^3$ (5S) state of Be generated for the original and one alternative configuration.

Next we discuss how a Kohn-Sham(KS) system can be realized for an excited-state den-

sity. To get a KS system, the mapping from a given excited-state density to a non-interacting wavefunction is established by making the expectation value $\langle \Phi | \hat{T} | \Phi \rangle$ stationary with respect to variations in Φ with Φ giving the excited-state density. How this is done is described later. As is the case for the interacting systems, there may be several Φ 's and the corresponding KS potentials v_{KS} that give rise to an excited-state density and make $\langle \Phi | \hat{T} | \Phi \rangle$ stationary. Two such potentials for the density of $1s^1 2s^0 2p^3$ (5S) of Be are shown in Fig. 1. The excited-state density used is obtained by solving the excited-state KS equation with the Harbola-Sahni(HS)³² exchange-only potential. The question that arises again is how do we choose one particular KS system to represent a system in its excited-state. An intuitive way would be by comparing the ground-state densities, as was done for the interacting systems. However, the ground-state density of a non-interacting system that reproduces an excited-state density may not be the same as that of the true system (it will be the same only if the electron-electron interaction were absent). Thus Levy and Nagy propose^{18,26} that of the many Fermionic non-interacting systems that give the same excited-state density, the one whose ground-state density resembles the exact one in the least square sense be identified as the Kohn-Sham system. The criterion is obviously exact³⁰ for systems with no electron-electron interaction, as stated above. For interacting-electron systems, the criterion appears to give³⁰ the true Kohn-Sham system, consistent with the orbitals to which the electrons have been excited, as we discuss below.

B. Levy-Nagy Criterion

We have been investigating the Levy-Nagy criterion for many different excited-states of atomic systems. We find that whereas the criterion is satisfied for a large number of excited-states, there are exceptions also. We present and discuss these results below. The results are obtained as follows. Working within the central field approximation, we perform the excited-state calculations using the exchange-only HS method and obtain the excited-state density (KLI-OPM^{33,34,35} gives similar results). We then use the Zhao-Morrison-Parr(ZMP)

method³⁶ to generate the same density with different configurations and test the Levy-Nagy criterion for these configurations. In the ZMP method the Kohn-Sham system for a given density $\rho(\vec{r})$ and a configuration of choice²⁸ is obtained by solving the equation

$$\left\{ -\frac{1}{2}\nabla^2 + \lambda \int \frac{[\sum_j n_j |\phi_j(\vec{r}')|^2 - \rho(\vec{r}')] }{|\vec{r} - \vec{r}'|} d\vec{r}' \right\} \phi_i(\vec{r}) = \varepsilon_i \phi_i(\vec{r}) \quad (7)$$

in the limit of $\lambda \rightarrow \infty$. Here $\{n_j\}$ are the occupation numbers of the orbitals according to the configuration chosen. We choose λ large enough so that not only do the densities match to a high degree of accuracy, the highest occupied eigenvalues ε_{max} of the above equation also matches with the original ε_{max} to within 5%; in fact it is better than within 2% in many of the cases. For example, for the $1s^1 2s^0 2p^3$ excited-state of *Be*, we have generated the same density with three other configurations: $1s^2 2s^0 2p^2$, $1s^2 2s^1 2p^1$ and $1s^1 2s^1 2p^2$. The ε_{max} and the expectation values $\langle \frac{1}{R} \rangle$, $\langle R \rangle$ and $\langle R^2 \rangle$ for different configurations are compared in Table I. To judge the numerical accuracy of our ZMP program, we also generate the excited-state density with the original configuration and compare numbers obtained with the original numbers. We see that with the original configuration, the ε_{max} comes to within 2% of the original value with $\lambda = 5000$ whereas the various expectation values are essentially exact. For the three alternative configurations, the accuracy of $\langle \frac{1}{R} \rangle$, $\langle R \rangle$ and $\langle R^2 \rangle$ is about the same but the ε_{max} values differ slightly more depending on the configuration. The worst case is the $1s^1 2s^1 2p^2$ for which $\varepsilon_{max} = -0.626 \text{ Ry}$ for $\lambda = 30000$. To make sure that the eigenvalue will eventually converge to $\varepsilon_{max} = -0.658 \text{ Ry}$, we performed calculations for different values of λ for this configuration and found that $\varepsilon_{max} = -0.621 \text{ Ry}$, -0.624 Ry and -0.626 Ry for $\lambda = 10000$, 14000 and 30000 , respectively, thereby shifting towards the true eigenvalue albeit very slowly. We also mention that for the configuration $1s^2 2s^1 2p^1$, the uppermost orbital is $2s$ and not $2p$. The local potential in which the electrons are move is then given as:

$$v_{KS}(\vec{r}) = \lambda \int \frac{[\sum_j n_j |\phi_j(\vec{r}')|^2 - \rho(\vec{r}')] }{|\vec{r} - \vec{r}'|} d\vec{r}' . \quad (8)$$

Having found the potential above, we obtain the ground-state density $\tilde{\rho}_0(\vec{r})$ of this potential by occupying the lowest energy orbitals with the given number of electrons. We calculate

its mean square distance from the true ground-state density $\rho_0(\vec{r})$ as³⁰

$$\Delta[\rho_0(\mathbf{r}), \tilde{\rho}_0(\mathbf{r})] = \int_{\infty} \{\rho_0(\mathbf{r}) - \tilde{\rho}_0(\mathbf{r})\}^2 d^3r. \quad (9)$$

By the true ground-state density here, we mean the ground-state density obtained with the Harbola-Sahni exchange potential. The results for Δ for a number of atoms and their excited-states are shown in Tables II & III. The original configuration of the excited-states is shown in the second column. The third column shows the alternative configurations using which we obtain the same density and the fourth column the corresponding $\Delta[\rho_0(\mathbf{r}), \tilde{\rho}_0(\mathbf{r})]$. It is seen from the results that for most of the cases Δ is the smallest for the original configuration but there are cases where Δ is smallest for a different configuration. For example, there is the excited-state $1s^1 2s^0 2p^3$ of *Be* for which the configuration $1s^1 2s^1 2p^2$ gives the smallest Δ . Similarly for the state $1s^2 2s^2 2p^3 3s^2$ of *F*, $1s^2 2s^2 2p^6 3s^2$ of *Ne* and $1s^1 2s^0 2p^3$ of *B*⁺, $1s^1 2s^0 2p^3$ of *Ne*⁶⁺ Δ is the smallest for a configuration other than the original configuration of the system. Thus we find that the Levy-Nagy criterion, as quantified by Eq. 8 above, is not satisfactory in that it leads to a KS system where an excited-state configuration is not consistent with the original system.

C. An Alternative Criterion

Having found that the Levy-Nagy criterion is not fully satisfactory to identify a KS system, we have looked for other ways of doing so, remaining within the Levy-Nagy proposal of comparing the ground-state densities. Thus instead of comparing densities directly, we compare them energetically as follows. After obtaining many different non-interacting systems for an excited-state density, we take their ground-states and calculate the expectation value of the true ground-state Kohn-Sham Hamiltonian (constructed using the HS exchange potential) with respect to these ground-states. Thus the calculation proceeds as follows: we solve the HS equation for the ground-state of a system and obtain the ground-state Kohn-Sham Hamiltonian H_0 . The expectation value $\langle H_0 \rangle$ with respect to the true ground-state orbitals is designated as $\langle H_0 \rangle_{true}$; it is the sum of the eigenvalues of the ground-state orbitals.

Next we take the different non-interacting systems giving an excited-state density, consider their ground-states and calculate the expectation value $\langle H_0 \rangle_{alt.}$ with respect to these states. Because of the variational principle $\langle H_0 \rangle_{alt.}$ should always be above $\langle H_0 \rangle_{true}$. We then identify the true KS system as that for which $\langle H_0 \rangle_{alt.}$ is closest to $\langle H_0 \rangle_{true}$. This comparison is made in Tables IV & V for the same systems as in Tables II and III with HS density. As is clear from the table the alternate criterion is better in that the correspondence between the original system and the Kohn-Sham system is restored for F, Ne and Ne^{6+} . However, new inconsistencies arise in Al, Si and P^+ although in these cases the difference in the numbers for the original and the alternative configuration is very small. On the other hand, the inconsistency in Be and B^+ remains. We note that this criterion is very sensitive to the exchange potential. If calculations are done with KLI-OPM exchange potential, the inconsistency remains only in Be and B^+ systems. It is clear from the discussion in the two sections above that a criterion based on comparison of ground-state densities of excited-state Kohn-Sham systems cannot be satisfactory.

III. PRESENT THEORY

Given the background above, we now present a consistent theory of excited-states within the rubric of density-functional approach. The principal tenets of the theory are:

(i) There is a straightforward way of mapping an excited-state density $\rho(\vec{r})$ to the corresponding many-electron wavefunction $\Psi(\vec{r})$ or the external potential $v_{ext}(\vec{r})$ using the ρ -stationary wavefunctions. The wavefunction depends upon the ground-state density ρ_0 implicitly.

(ii) The Kohn-Sham system is defined through a comparison of the kinetic energy for the excited-states. This avoids any comparison of the ground-state densities which, as seen above, doesn't give a satisfactory way of defining a KS system. We now discuss these two points one by one.

To describe the mapping from an excited-state density $\rho_k(\vec{r})$ to a many-body wavefunc-

tion, we take recourse to the constrained-search approach. This gives, as discussed earlier, many different wavefunctions $\Psi_k(\vec{r})$ and the corresponding external potential $v_{ext}^k(\vec{r})$. If in addition to the excited-state density we also know the ground-state density ρ_0 then $v_{ext}(\vec{r})$ is uniquely determined by the Hohenberg-Kohn¹ theorem. Thus with the knowledge of ρ_0 it is quite straightforward to select a particular Ψ that belongs to a $[\rho, \rho_0]$ combination by comparing $v_{ext}^k(\vec{r})$ with $v_{ext}(\vec{r})$. Alternatively, we can think of it as finding Ψ variationally for a $[\rho, v_{ext}]$ combination because the knowledge of ρ_0 and v_{ext} is equivalent. Through the constrained search above a functional

$$F[\rho; \rho_0] = \langle \Psi[\rho; \rho_0] | \hat{T} + \hat{V}_{ee} | \Psi[\rho; \rho_0] \rangle \quad (10)$$

is also defined. The prescription above is similar to that of Levy and Nagy²⁶ but avoids the orthogonality condition imposed by them.

The densities for different excited state for a given ground-state density ρ_0 or external potential v_{ext} can thus be found as follows: take a density and search for Ψ that makes $\langle \Psi | \hat{T} + v_{ee} | \Psi \rangle$ stationary; check whether the corresponding v_{ext} matches with the given ρ_0 (or v_{ext}); if not, take another density and repeat the procedure until the correct ρ is found. Also because of the proof given in section II above, the Euler equation for the excited-state density is

$$\frac{\delta F[\rho, \rho_0]}{\delta \rho(\vec{r})} + v_{ext}(\vec{r}) = \mu \quad (11)$$

where μ is the Lagrange multiplier to ensure that $\rho_k(\vec{r})$ integrates to the proper number of electrons.

The prescription above for the excited-states in terms of their densities is quite straightforward, particularly because its development is parallel to that for the ground-states. On the other hand, to construct a Kohn-Sham² system for a given density is non-trivial; and to carry out accurate calculations for excited- states it is of prime importance to construct a KS system. Further, a KS system will be meaningful if the orbitals involved in an excitation match with the corresponding excitations in the true system. We have shown above that

the Kohn-Sham system constructed using the Levy-Nagy criterion fails in this regard.

In principle, obtaining a Kohn-Sham system is quite easy. Define the non-interacting kinetic energy $T_s[\rho, \rho_0]$ and use it to further define the exchange-correlation functional as

$$E_{xc}[\rho, \rho_0] = F[\rho, \rho_0] - \frac{1}{2} \int \int \frac{\rho(\vec{r})\rho(\vec{r}')}{|\vec{r} - \vec{r}'|} d\vec{r} d\vec{r}' - T_s[\rho, \rho_0] . \quad (12)$$

Then the Euler equation for the excited-state densities will read

$$\frac{\delta T_s[\rho, \rho_0]}{\delta \rho(\vec{r})} + \int \frac{\rho(\vec{r}')}{|\vec{r} - \vec{r}'|} d\vec{r}' + \frac{\delta E_{xc}[\rho, \rho_0]}{\delta \rho(\vec{r})} + v_{ext} = \mu . \quad (13)$$

which is equivalent to solving

$$\left\{ -\frac{1}{2} \nabla^2 + v_{KS}(\vec{r}) \right\} \phi_i(\vec{r}) = \varepsilon_i \phi_i(\vec{r}) \quad (14)$$

with

$$v_{KS}(\vec{r}) = v_{ext}(\vec{r}) + \int \frac{\rho(\vec{r}')}{|\vec{r} - \vec{r}'|} d\vec{r}' + \frac{\delta E_{xc}[\rho, \rho_0]}{\delta \rho(\vec{r})} . \quad (15)$$

However, it is defining $T_s[\rho, \rho_0]$ that is not easy in the excited-state problem. For the ground-states, $T_s[\rho_0]$ is easily defined as the minimum kinetic energy for a given density obtained by occupying the lowest energy orbitals for a non-interacting system. On the other hand, for the excited-states it is not clear which orbitals to occupy for a given density, particularly because a density can be generated by many different non-interacting systems. Levy-Nagy select one of these systems by comparing the ground-state density of the excited-state non-interacting system with the true ground-state density. However, this criterion is not satisfactory as discussed earlier. Therefore some other criterion has to be evolved to construct the excited-state Kohn-Sham system.

Before searching for other ways of constructing a Kohn-Sham system, we look for reasons that may be responsible for the Levy-Nagy criterion not being fully satisfactory. We argue that we are not being consistent while comparing the ground-state density of an excited-state KS system with the true ground-state density. This is because the ground-state density of the excited-state KS system is not the self-consistent ground-state

density of the $v_{ext}(\vec{r})$ obtained for the excited-state density but of a potential different from $v_{ext}(\vec{r})$. In comparing the ground-state densities, we are thus not comparing the $v_{ext}(\vec{r})$ of the excited-state KS system with the true $v_{ext}(\vec{r})$, and this at times leads to inconsistent results.

In light of the above remarks, it is important that in constructing the Kohn-Sham system only the self-consistently determined quantities corresponding to a given excited-state density be compared. Thus we propose that the KS system be so chosen that it is energetically very close to the original system. To ensure this, we define the KS system as that system for which the non-interacting kinetic energy, obtained through constrained search over the non-interacting wavefunctions, is closest to $T[\rho, \rho_0]$, already obtained through the full constrained search. This then gives the functional $T_s[\rho, \rho_0]$. Thus $T_s[\rho, \rho_0]$ is defined as the kinetic energy that is closest to the true kinetic energy $T[\rho, \rho_0]$ obtained for a given excited-state density ρ . Defining $T_s[\rho, \rho_0]$ in this manner also keeps the DFT exchange-correlation energy close to the conventional quantum mechanical exchange-correlation energy. An added advantage of keeping the difference between the two kinetic energies $T_s[\rho, \rho_0]$ and $T[\rho, \rho_0]$ smallest is that the structure of the Kohn-Sham potential is simple; it is known that contribution of $\underbrace{T - T_s}$ gives more structure to the KS potential.

All the statements in this paragraph are justified on the basis of the differential virial theorem³⁷. Using this theorem the exchange-correlation potential for a given density and the corresponding many body wavefunction Ψ can be written as

$$-\nabla v_{xc}(\vec{r}) = \frac{\left\{ \vec{Z}_{KS}(\vec{r}; [\Gamma_1^{KS}]) - \vec{Z}(\vec{r}; [\Gamma_1]) + \int [\nabla u(\vec{r}, \vec{r}')] [\rho(\vec{r})\rho(\vec{r}') - 2\Gamma_2(\vec{r}, \vec{r}')] d\vec{r}' \right\}}{\rho(\vec{r})} \quad (16)$$

where $\Gamma_1(\Gamma_2)$ is the first(second) order density matrix and the vector fields \vec{Z}, \vec{Z}_{KS} are related to the true and Kohn-Sham kinetic energy density tensors respectively. For a given $[\rho, \rho_0]$, $\vec{Z}, \Gamma_1, \Gamma_2$ are fixed and different configurations chosen give different \vec{Z}_{KS} . These different

configurations can be thought of as arising from a different external potential (as is shown below) or from a different exchange-correlation potential^{27,28}. In any case kinetic energy difference between the true and Kohn-Sham system is³⁸

$$\Delta T = \frac{1}{2} \int \vec{r} \cdot \left\{ \vec{Z}_{KS}(\vec{r}; [\Gamma_1^{KS}]) - \vec{Z}(\vec{r}; [\Gamma_1]) \right\} d\vec{r}. \quad (17)$$

It is this difference that we propose be kept the smallest for the true KS system, and as we show below, it gives the Kohn-Sham system consistent with the original system.

A. Examples

We now demonstrate the ideas presented above with examples. Since we do not know how to perform the general constrained search $\rho \rightarrow \Psi, \delta \langle \Psi | \hat{T} + \hat{v}_{ee} | \Psi \rangle = 0$, we take an indirect path for the purpose of demonstration. In the following we work with atomic excited-state densities generated by HS exchange-only potential and take these as the excited-state density. The densities and the energies obtained by the HS formalism are essentially the same as those of Hartree-Fock(HF)⁴⁰ theory for both the ground as well as the excited-states. Similarly the HS exchange potential is very close to the true local exchange potential of the optimized potential method(OPM). Thus the formalism is well suited to test the ideas presented above. Thus we start with this given $[\rho, v_{ext}]$ combination.

We first demonstrate that an excited-state density is generated by different external potentials depending on the configurations chosen to generate the density. For this we use the ZMP method to get the non-interacting potential giving the density $\rho(\vec{r})$ and subtract from it the Coulomb $v_{coul}(\vec{r})$ and the exchange $v_x^{HS}(\vec{r})$ potential to get the external potential $v_{ext}(\vec{r})$ (*i.e.* $v_{ext}(\vec{r}) = v_{KS}(\vec{r}) - v_{coul}(\vec{r}) - v_x^{HS}(\vec{r})$). The exchange potential for a given set of occupied orbitals is obtained using the Harbola-Sahni formula³⁹ for it. Shown in Fig. 2 are different external potentials thus generated for the $1s^1 2s^0 2p^3$ (5S) of density Be . To check our consistency, we first obtain $v_{ext}(\vec{r})$ for the original configuration ($1s^1 2s^0 2p^3$) and find it correctly to be $-\frac{8}{r} Ry$. The other configurations that we use to obtain the same density




FIG. 2: Shown in the figure are the external potentials for the excited-state density of $1s^1 2s^0 2p^3$ (5S) state of Be corresponding to the original and one alternative configuration.

are $1s^2 2s^0 2p^2$, $1s^2 2s^1 2p^1$, $1s^1 2s^1 2p^2$. We have shown only two potentials corresponding to the configurations $1s^2 2s^1 2p^1$ and $1s^1 2s^0 2p^3$, and compared them with the true external potential $v_{ext} = -\frac{8}{r} Ry$. As discussed earlier, only one of these - that corresponding to the original configuration - matches with the true external potential.

Next we show that the external potentials corresponding to the ground-state density of excited-state KS systems are different from the external potential for the excited-state density. This, as pointed out earlier, sometimes leads to non-satisfaction of the Levy-Nagy criterion. Shown in Fig. 3 are the $\tilde{v}_{ext}(\vec{r})$ corresponding to the ground-state densities of different configurations for the excited-state density of $1s^1 2s^0 2p^3$ (5S) state of Be . These potential are also obtained by subtracting from the Kohn-Sham potential v_{KS} the Coulomb and the Harbola-Sahni exchange potential calculated by occupying the corresponding KS orbitals in the ground-state configuration. We see that whereas the true external potential is $-\frac{8}{r} Ry$, the external potentials corresponding to the ground-states are different. It is this difference that leads to the ground-state densities different from the true one, and also




FIG. 3: Shown in the figure are the external potentials corresponding to the ground-state densities of excited-state KS systems for $1s^1 2s^0 2p^3$ (5S) state of Be . The potentials are compared with the true external potential $v_{ext} = -\frac{8}{r}$.

sometimes to inconsistencies between the KS and the true systems.

Finally we show that a comparison of the kinetic-energies(KE) leads to the appropriate Kohn-Sham system. We demonstrate this with the densities generated using the HS exchange potential and with one example with correlated density. The numbers for the noninteracting kinetic energy for different configurations corresponding to the same excited-state densities as considered earlier are shown in Tables VI and VII and are compared with their original kinetic energy. Since the HS potential is a local potential itself, the correct configuration gives $\Delta T_s = 0$ (slight difference arising due to numerical calculations) and wrong ones a larger value, as is evident from the numbers in the Tables. We see that unlike the Levy-Nagy criterion, a comparison of excited-state KE leads to a proper KS system in all the cases. Of course when we use the correlated densities, the difference ΔT is not going to be *zero* for the proper configuration but should be the smallest. This is clearly seen in the example of $1s2s(^1S)$ state of He atom where the true $T[\rho]$ is 2.146 a.u. . We have used

the ZMP procedure to obtain the KS potentials in this case also and see that KE for the $1s^2$ and the $1s2s$ configurations is 2.044 *a.u.* and 2.153 *a.u.*, respectively. In the latter case it is closer to the true kinetic energy of the system. Thus the configuration $1s2s$ represents the KS system for the $1s2s(^1S)$ *He* density.

IV. CONCLUDING REMARKS

Building on the work of Görling and Levy and Nagy, we have presented in this paper a consistent theory of excited states within the density-functional formalism. The theory is based on constrained-search and defines a bi-density functional $F[\rho, \rho_0]$ without the orthogonality constraint of Levy-Nagy. Further the theory gives a clear definition of the excited-state Kohn-Sham systems as that whose kinetic and exchange-correlation energy components are closest to those of the true system. This avoids the problem of comparing the non-self-consistent ground-state densities, as proposed in the LN theory, so no inconsistency arises in identifying an excited-state Kohn-Sham system.

To conclude, we have analyzed the theoretical foundations of excited-state time-independent density-functional theory and have put Kohn-Sham calculations for excited-states on a firm footing. It is clear from our present work that an excited-state configuration of the KS system corresponds to a similar excited-state of the true system (with the major component of the excited-state wavefunction involving the same orbitals as the KS system). The work should also help in providing guidance in construction of the excited-state exchange-correlation energy functionals to facilitate self-consistent determination of the excited-state energies.

* Electronic address: Fax:+91-512-259 0914 ; E-mail: dilu@iitk.ac.in

† Electronic address: Fax:+91-512-259 0914 ; E-mail: mkh@iitk.ac.in

- ¹ P. Hohenberg and W. Kohn, Phys. Rev. **136**, B864 (1964) .
- ² W. Kohn and L. J. Sham, Phys. Rev. **140**, A1133 (1965) .
- ³ R. O. Jones and O. Gunnarsson , Rev. Mod. Phys. **61**, 689 (1989).
- ⁴ R. G. Parr and W. Yang, *Density-Functional Theory of Atoms and Molecules* (New York: Oxford University Press, 1989).
- ⁵ R. M. Dreizler and E. K. U. Gross, *Density Functional Theory* (Berlin: Springer-Verlag, 1990).
- ⁶ N.H. March, *Electron Density Theory of Atoms and Molecules* (Academic Press, London, 1992).
- ⁷ O. Gunnarsson and B. I. Lundqvist, Phys. Rev. A **13**, 4274 (1976).
- ⁸ T. Ziegler, A. Rauk and E. J. Baerends, Theor. Chim. Acta **43**, 261 (1977).
- ⁹ U. von Barth, Phys. Rev. A **20**, 1963 (1979).
- ¹⁰ A. K. Theophilou, J. Phys. C **12**, 5419 (1979).
- ¹¹ R. K. Pathak, Phys. Rev. A **29**, 978 (1984).
- ¹² J. P. Perdew and M. Levy, Phys. Rev. B **31**, 6264 (1985).
- ¹³ H. Englisch, H. Fieseler and A. Haufe, Phys. Rev. A **37**, 4570 (1988).
- ¹⁴ E. K. U. Gross, L. N. Oliveira and W. Kohn, Phys. Rev. A **37**, 2805 (1988);
E. K. U. Gross, L. N. Oliveira and W. Kohn, Phys. Rev. A **37**, 2809 (1988).
- ¹⁵ K.D. Sen, Chem. Phys. Lett. **188**, 510 (1992).
- ¹⁶ Á. Nagy, Physics Reports **298**, 1 (1998).
- ¹⁷ Ranbir Singh and B. M. Deb, Physics Reports, **311**, 47 (1999) .
- ¹⁸ Á. Nagy, Adv. Quant. Chem. **42**, 363 (2002)
- ¹⁹ A. D. Becke, Phys. Rev. A. **38**, 3098 (1988).
- ²⁰ J. P. Perdew in *P. Ziesche, H. Eschrig, editors, Electronic Structure of Solids '91, Akademie Verlag, Berlin, (1991)*
- ²¹ C. Lee, W. Yang and R. G. Parr, Phys. Rev. B. **37**, 785 (1988).
- ²² J. P. Perdew, K. Burke, and M. Ernzerhof, Phys. Rev. Lett. **77**, 3865 (1996).
- ²³ J. P. Perdew, S. Kurth, A. Zupan, and P. Blaha, Phys. Rev. Lett. **82**, 2544 (1999).
- ²⁴ P. Samal and M. K. Harbola, J. Phys. B **38**, 3765 (2005) .
- ²⁵ A. Görling, Phys. Rev. A **54**, 3912 (1996) ;
A. Görling, Phys. Rev. A **59**, 3359 (1999) .
- ²⁶ M. Levy and Á. Nagy, Phys. Rev. Lett. **83**, 4361 (1999) ;
Á. Nagy and M. Levy, Phys. Rev. A **63**, 052502 (2001) .

- ²⁷ V. Sahni, L. Massa, R. Singh and M. Slamet, Phys. Rev. Lett. **87**, 113002 (2001) .
- ²⁸ M. K. Harbola, Phys. Rev. A **69**, 042512 (2004) .
- ²⁹ R. Gaudoin and K. Burke, Phys. Rev. Lett. **93**, 173001 (2004) , *ibid.* **94**, 029901 (2005) .
- ³⁰ P. Samal, M. K. Harbola and A. Holas, Chem. Phys. Lett. **419**, 217 (2005).
- ³¹ M. Levy, Proc. Natl. Acad. Sci. USA **76**, 6062 (1979);
M. Levy and J. P. Perdew, in *Density Functional Methods in Physics*, edited by R. M. Dreizler and J. da Providencia (Plenum Press, New York, 1985).
- ³² M. K. Harbola and V.Sahni, Phys. Rev. Lett. **62**, 489 (1989) .
- ³³ J. B. Krieger, Y. Li, and G. J. Iafrate, in *Density Functional Theory* , edited by E.K.U. Gross and R.M. Dreizler, (Plenum Press, New York, 1995)
- ³⁴ R. T. Sharp and G. K. Horton, Phys. Rev. **90**, 3876 (1953);
J. D. Talman and W. F. Shadwick, Phys. Rev. A **14**, 36 (1976) .
- ³⁵ Á. Nagy, Int. J. Quantum Chem. **69**, 247 (1998).
- ³⁶ Q. Zhao and R.G. Parr, J. Chem. Phys. **98**, 543 (1993) ;
Q. Zhao, R. C. Morrison and R. G. Parr, Phys. Rev. A **50**, 2138 (1994) .
- ³⁷ A. Holas, N. H. March, Phys. Rev. A **51**, 2040 (1995) .
- ³⁸ V.Sahni, Phys. Rev. A. **55**, 1846 (1997) .
- ³⁹ M.K. Harbola, R.R. Zope and R.K. Pathak, Phys. Rev. A **53**, 3652 (1996).
- ⁴⁰ C. F. Fischer, *The Hartree-Fock method for atoms* (New York: John Wiley, 1977)

TABLE I: Shown in the table ε_{max} and expectation values $\langle \frac{1}{R} \rangle$, $\langle R \rangle$ & $\langle R^2 \rangle$ for various configurations giving the same density as that of $1s^1 2s^0 2p^3$; ⁵ S state of Be . The self-consistently determined values of these physical quantities are : $\varepsilon_{max} = -0.658 \text{ Ry}$, $\langle \frac{1}{R} \rangle = 5.818$, $\langle R \rangle = 6.755$ & $\langle R^2 \rangle = 17.309$.

configurations	λ	$\varepsilon_{max}(\text{Ry.})$	$\langle \frac{1}{R} \rangle$	$\langle R \rangle$	$\langle R^2 \rangle$
$1s^1 2s^0 2p^3$	5000	-0.649	5.818	6.755	17.312
$1s^2 2s^0 2p^2$	5000	-0.649	5.819	6.755	17.312
$1s^2 2s^1 2p^1$	5000	-0.655	5.819	6.755	17.312
$1s^1 2s^1 2p^2$	30,000	-0.626	5.818	6.755	17.310

TABLE II: Value of Δ (See Eq. 8) for different configurations(3^{rd} column) giving the same excited-state density as that for the original configuration(2^{nd} column). Systems where the LN criterion is not satisfactory are indicated with a '*' against them.

atoms/ions	<i>true.config.</i>	<i>alt.config.</i>	$\Delta [\rho_0(\mathbf{r}), \tilde{\rho}_0(\mathbf{r})]$
<i>*Be</i>	$1s^1 2s^0 2p^3$	$1s^1 2s^1 2p^2$	0.0662
		$1s^1 2s^0 2p^3$	0.1627
		$1s^2 2s^1 2p^1$	0.8758
<i>B</i>	$1s^2 2s^0 2p^3$	$1s^2 2s^0 2p^3$	0.0002
		$1s^2 2s^1 2p^2$	0.0065
		$1s^2 2s^2 2p^1$	0.0286
<i>C</i>	$1s^2 2s^1 2p^3$	$1s^2 2s^1 2p^3$	0.0008
		$1s^2 2s^2 2p^2$	0.0156
		$1s^2 2s^0 2p^4$	0.0181
<i>N</i>	$1s^1 2s^0 2p^6$	$1s^1 2s^0 2p^6$	0.2903
		$1s^2 2s^0 2p^5$	6.8461
		$1s^2 2s^1 2p^4$	8.8409
<i>O</i>	$1s^1 2s^1 2p^6$	$1s^1 2s^1 2p^6$	0.3875
		$1s^2 2s^0 2p^6$	8.9609
		$1s^2 2s^1 2p^5$	12.1269
<i>F</i>	$1s^2 2s^1 2p^6$	$1s^2 2s^1 2p^6$	0.0002
		$1s^2 2s^2 2p^5$	0.0621
<i>*F</i>	$1s^2 2s^2 2p^3 3s^2$	$1s^2 2s^2 2p^4 3s^1$	0.0521
		$1s^2 2s^2 2p^3 3s^2$	0.2704
<i>*Ne</i>	$1s^1 2s^1 2p^6 3s^2$	$1s^1 2s^2 2p^6 3s^1$	0.7138
		$1s^1 2s^1 2p^6 3s^2$	1.6176

TABLE III: Caption is the same as in Table II.

atoms/ions	<i>true.config.</i>	<i>alt.config.</i>	$\Delta [\rho_0(\mathbf{r}), \tilde{\rho}_0(\mathbf{r})]$
$*B^+$	$1s^1 2s^0 2p^3$	$1s^1 2s^1 2p^2$	0.0402
		$1s^1 2s^0 2p^3$	0.1931
		$1s^2 2s^1 2p^1$	2.0261
$*Ne^{6+}$	$1s^1 2s^0 2p^3$	$1s^1 2s^1 2p^2$	0.1465
		$1s^1 2s^0 2p^3$	0.3463
Mg	$1s^2 2s^2 2p^6 3s^0 3p^2$	$1s^2 2s^2 2p^6 3s^0 3p^2$	0.0007
		$1s^2 2s^2 2p^6 3s^1 3p^1$	0.0014
		$1s^2 2s^2 2p^6 3s^2 3p^0$	0.0039
Al	$1s^2 2s^2 2p^6 3s^0 3p^3$	$1s^2 2s^2 2p^6 3s^0 3p^3$	0.0008
		$1s^2 2s^2 2p^6 3s^1 3p^2$	0.0022
		$1s^2 2s^2 2p^6 3s^2 3p^1$	0.0073
Si	$1s^2 2s^2 2p^6 3s^1 3p^3$	$1s^2 2s^2 2p^6 3s^1 3p^3$	0.0007
		$1s^2 2s^2 2p^6 3s^2 3p^2$	0.0027
		$1s^2 2s^2 2p^6 3s^0 3p^4$	0.0078
Si^+	$1s^2 2s^2 2p^6 3s^0 3p^3$	$1s^2 2s^2 2p^6 3s^0 3p^3$	0.0004
		$1s^2 2s^2 2p^6 3s^1 3p^2$	0.0042
		$1s^2 2s^2 2p^6 3s^2 3p^1$	0.0148
P^+	$1s^2 2s^2 2p^6 3s^1 3p^3$	$1s^2 2s^2 2p^6 3s^1 3p^3$	0.0005
		$1s^2 2s^2 2p^6 3s^2 3p^2$	0.0053
		$1s^2 2s^2 2p^6 3s^0 3p^4$	0.0099
P	$1s^2 2s^2 2p^6 3s^0 3p^5$	$1s^2 2s^2 2p^6 3s^0 3p^5$	0.0006
		$1s^2 2s^2 2p^6 3s^1 3p^4$	0.0055
		$1s^2 2s^2 2p^6 3s^2 3p^3$	0.0207

TABLE IV: Comparison of the expectation value $-\langle H_0 \rangle_{alt.}$ and $-\langle H_0 \rangle_{exact}$ for various configurations corresponding to a given excited-state density. The first column mentioned are the atoms/ions, the second column the original configuration, fourth column the alternative configurations considered. In the third and last column given the expectations values $-\langle H_0 \rangle_{exact}$ and $-\langle H_0 \rangle_{alt.}$ respectively.

atoms/ions	<i>exact config.</i>	$-\langle H_0 \rangle_{exact}$ a.u.	<i>alt. config.</i>	$-\langle H_0 \rangle_{alt.}$ a.u.
<i>*Be</i>	$1s^1 2s^0 2p^3$	8.9272	$1s^2 2s^1 2p^1$	7.4302
			$1s^1 2s^0 2p^3$	8.8051
			$1s^1 2s^1 2p^2$	8.8601
<i>Be</i>	$1s^2 2s^1 2p^1$	8.9237	$1s^1 2s^1 2p^2$	6.1351
			$1s^2 2s^1 2p^1$	8.9217
<i>B</i>	$1s^2 2s^0 2p^3$	15.1482	$1s^2 2s^2 2p^1$	15.0828
			$1s^2 2s^0 2p^3$	15.1471
			$1s^2 2s^1 2p^2$	15.1365
<i>C</i>	$1s^2 2s^1 2p^3$	22.9759	$1s^2 2s^2 2p^2$	22.9556
			$1s^2 2s^1 2p^3$	22.9723
			$1s^2 2s^0 2p^4$	22.9473
<i>N</i>	$1s^2 2s^1 2p^4$	32.6951	$1s^2 2s^2 2p^3$	32.6682
			$1s^2 2s^1 2p^4$	32.6949
			$1s^2 2s^0 2p^5$	32.6712
<i>O</i>	$1s^2 2s^0 2p^6$	43.3618	$1s^2 2s^2 2p^4$	43.2215
			$1s^2 2s^0 2p^6$	43.3608
			$1s^2 2s^1 2p^5$	43.3252
<i>F</i>	$1s^2 2s^2 2p^3 3s^2$	55.8686	$1s^1 2s^1 2p^6 3s^2$	55.4295
			$1s^2 2s^2 2p^3 3s^2$	55.6436
<i>Ne</i>	$1s^1 2s^1 2p^6 3s^2$	70.1743	$1s^1 2s^2 2p^6 3s^1$	69.2101
			$1s^1 2s^1 2p^6 3s^2$	69.4399

TABLE V: Caption is the same as in Table IV.

atoms/ions	<i>exact config.</i>	$-\langle H_0 \rangle_{exact}$ a.u.	<i>alt. config.</i>	$-\langle H_0 \rangle_{alt}$ a.u.
$*B^+$	$1s^1 2s^0 2p^3$	16.6626	$1s^1 2s^1 2p^2$	16.6106
			$1s^1 2s^0 2p^3$	16.5501
			$1s^2 2s^1 2p^1$	14.2959
Ne^{6+}	$1s^1 2s^0 2p^3$	93.0672	$1s^1 2s^1 2p^2$	92.9276
			$1s^1 2s^0 2p^3$	92.9699
Mg	$1s^2 2s^2 2p^6 3s^0 3p^2$	110.9422	$1s^2 2s^2 2p^6 3s^2 3p^0$	110.9091
			$1s^2 2s^2 2p^6 3s^0 3p^2$	110.9368
			$1s^2 2s^2 2p^6 3s^1 3p^1$	110.9341
$*Al$	$1s^2 2s^2 2p^6 3s^0 3p^3$	137.1905	$1s^2 2s^2 2p^6 3s^1 3p^2$	137.1856
			$1s^2 2s^2 2p^6 3s^0 3p^3$	137.1853
			$1s^2 2s^2 2p^6 3s^2 3p^1$	137.1655
$*Si$	$1s^2 2s^2 2p^6 3s^1 3p^3$	166.2441	$1s^2 2s^2 2p^6 3s^2 3p^2$	166.2411
			$1s^2 2s^2 2p^6 3s^1 3p^3$	166.2393
			$1s^2 2s^2 2p^6 3s^0 3p^4$	166.2201
Si^+	$1s^2 2s^2 2p^6 3s^0 3p^3$	170.5966	$1s^2 2s^2 2p^6 3s^2 3p^1$	170.5714
			$1s^2 2s^2 2p^6 3s^0 3p^3$	170.5941
			$1s^2 2s^2 2p^6 3s^1 3p^2$	170.5923
$*P^+$	$1s^2 2s^2 2p^6 3s^1 3p^3$	203.2722	$1s^2 2s^2 2p^6 3s^2 3p^2$	203.2693
			$1s^2 2s^2 2p^6 3s^1 3p^3$	203.2687
			$1s^2 2s^2 2p^6 3s^0 3p^4$	203.2522
P	$1s^2 2s^2 2p^6 3s^0 3p^5$	198.1074	$1s^2 2s^2 2p^6 3s^2 3p^3$	198.0818
			$1s^2 2s^2 2p^6 3s^0 3p^5$	198.1058
			$1s^2 2s^2 2p^6 3s^1 3p^4$	198.1025
S	$1s^2 2s^2 2p^6 3s^0 3p^6$	232.1008	$1s^2 2s^2 2p^6 3s^2 3p^4$	232.0761
			$1s^2 2s^2 2p^6 3s^0 3p^6$	232.0996
			$1s^2 2s^2 2p^6 3s^1 3p^5$	232.0957

TABLE VI: Shown in the table atoms/ions with the original excited-state configuration in the second column and density of this generated by various alternative configurations in the fourth column. In the third and fifth column given are the values of the kinetic energies corresponding to the original and alternative configurations respectively

atoms/ions	<i>true config.</i>	$T[\rho]$ a.u.	<i>alt. config.</i>	$T_s[\rho]$ a.u.
<i>Be</i>	$1s^1 2s^0 2p^3$	10.1489	$1s^1 2s^1 2p^2$	10.0177
			$1s^1 2s^0 2p^3$	10.1481
			$1s^2 2s^1 2p^1$	8.1357
<i>B</i>	$1s^2 2s^0 2p^3$	24.1249	$1s^2 2s^2 2p^1$	23.7627
			$1s^2 2s^0 2p^3$	24.1211
			$1s^2 2s^1 2p^2$	23.9238
<i>C</i>	$1s^2 2s^1 2p^3$	37.5938	$1s^2 2s^2 2p^2$	37.2985
			$1s^2 2s^1 2p^3$	37.5922
			$1s^2 2s^0 2p^4$	37.9299
<i>N</i>	$1s^1 2s^0 2p^6$	38.5551	$1s^2 2s^0 2p^5$	30.5856
			$1s^1 2s^0 2p^6$	38.5525
			$1s^2 2s^1 2p^4$	30.6238
<i>O</i>	$1s^1 2s^1 2p^6$	54.7136	$1s^2 2s^1 2p^5$	44.6244
			$1s^1 2s^1 2p^6$	54.7095
			$1s^2 2s^0 2p^6$	44.5899
<i>F</i>	$1s^2 2s^1 2p^6$	98.5267	$1s^2 2s^2 2p^5$	97.8733
			$1s^2 2s^1 2p^6$	98.5212
<i>F</i>	$1s^2 2s^2 2p^3 3s^2$	98.2631	$1s^2 2s^2 2p^4 3s^1$	97.8746
			$1s^2 2s^2 2p^3 3s^2$	98.2393
<i>Ne</i>	$1s^1 2s^1 2p^6 3s^2$	94.6521	$1s^1 2s^2 2p^6 3s^1$	93.4337
			$1s^1 2s^1 2p^6 3s^2$	94.6364

TABLE VII: Caption is the same as in Table VI.

atoms/ions	<i>true config.</i>	$T[\rho]$ a.u.	<i>alt. config.</i>	$T_s[\rho]$ a.u.
B^+	$1s^1 2s^0 2p^3$	16.8390	$1s^1 2s^1 2p^2$	16.6581
			$1s^1 2s^0 2p^3$	16.8378
			$1s^2 2s^1 2p^1$	13.2701
Ne^{6+}	$1s^1 2s^0 2p^3$	76.5893	$1s^1 2s^1 2p^2$	76.3318
			$1s^1 2s^0 2p^3$	76.5837
Mg	$1s^2 2s^2 2p^6 3s^0 3p^2$	199.3771	$1s^2 2s^2 2p^6 3s^1 3p^1$	199.2404
			$1s^2 2s^2 2p^6 3s^0 3p^2$	199.3661
			$1s^2 2s^2 2p^6 3s^2 3p^0$	199.1455
Al	$1s^2 2s^2 2p^6 3s^0 3p^3$	241.5112	$1s^2 2s^2 2p^6 3s^1 3p^2$	241.3098
			$1s^2 2s^2 2p^6 3s^0 3p^3$	241.4967
			$1s^2 2s^2 2p^6 3s^2 3p^1$	241.1428
Si	$1s^2 2s^2 2p^6 3s^1 3p^3$	288.7507	$1s^2 2s^2 2p^6 3s^2 3p^2$	288.4802
			$1s^2 2s^2 2p^6 3s^1 3p^3$	288.7335
			$1s^2 2s^2 2p^6 3s^0 3p^4$	288.9977
Si^+	$1s^2 2s^2 2p^6 3s^0 3p^3$	288.0463	$1s^2 2s^2 2p^6 3s^1 3p^2$	287.7594
			$1s^2 2s^2 2p^6 3s^0 3p^3$	288.0259
			$1s^2 2s^2 2p^6 3s^2 3p^1$	287.5146
P^+	$1s^2 2s^2 2p^6 3s^1 3p^3$	340.1993	$1s^2 2s^2 2p^6 3s^2 3p^2$	339.8495
			$1s^2 2s^2 2p^6 3s^1 3p^3$	340.1796
			$1s^2 2s^2 2p^6 3s^0 3p^4$	340.5202
P	$1s^2 2s^2 2p^6 3s^0 3p^5$	339.8574	$1s^2 2s^2 2p^6 3s^1 3p^4$	339.5241
			$1s^2 2s^2 2p^6 3s^0 3p^5$	339.8390
			$1s^2 2s^2 2p^6 3s^2 3p^3$	339.2355