

# Frequency quenching of microwave induced resistance oscillations in a high mobility 2DEG

S. A. Studenikin,<sup>a</sup> M. Byszewski,<sup>b</sup> D. K. Maude,<sup>b</sup> M. Potemski,<sup>b</sup>  
A. S. Sachrajda,<sup>a</sup> M. Hilke,<sup>c</sup> L. N. Pfeiffer,<sup>d</sup> K. W. West<sup>d</sup>

<sup>a</sup>*Institute for Microstructural Sciences, NRC, Ottawa, Ontario K1A-0R6, Canada*

<sup>b</sup>*Grenoble High Magnetic Field Laboratory, MPI/FKF and CNRS, Grenoble 38-042, France*

<sup>c</sup>*Department of Physics, McGill University, Montreal H3A 2T8, Canada*

<sup>d</sup>*Bell Laboratories, Lucent Technologies, Murray Hill, New Jersey 07974-0636, USA*

---

## Abstract

In this work we address experimentally electrodynamics of microwave induced resistance oscillations (MIROs) by studying their frequency dependence. The study has been conducted on a high mobility 2DEG in GaAs/AlGaAs heterostructure. It is found that at frequencies  $\sim 120$  GHz the MIROs start to quench while above  $\sim 250$  GHz they completely disappear. The results are discussed with a model in which plasmons play an intermediate role in the interaction process between the light and single particle electron excitations. The plasmons which are excited by microwaves decay through single particle excitations - electronic transitions between Landau levels involving a spatial displacement, the key component of existing theories for the origin of MIROs. The model also qualitatively explains other unexpected experimental results such as the independence of MIROs on the direction of the circular polarization and the absence of MIROs in absorption/reflection experiments.

*PACS: 73.50.Jt; 73.40.-c; 78.67.-n; 78.20.Ls; 73.43.-f*

---

Keywords: 2DEG; microwaves; zero-resistance states; Landau levels; plasmons.

## 1. Introduction

The recently observed new phenomenon of microwave induced resistance oscillations (MIROs) on very high mobility samples [1] has attracted considerable attention from the scientific community, partly due the fact that under certain conditions MIROs can evolve into zero-resistance states [2,3,4,5]. A number of theoretical models have been suggested to explain this phenomenon based on scattering, due to disorder [6,7], a non-equilibrium distribution function [8], quasi-classical electron orbit dynamics [9], and a plasmon instability [10]. It should be noted that long before the experimental observation of MIROs and zero resistance states, the possibility of such oscillations was discussed by Gladin and Ryzhii in 1970 within a scattering model.[11] All these approaches successfully predict the period and phase of the oscillations. In the first order approximation they arrive to similar final equations which describe quite well the MIROs waveform in the quasi-classical regime of large filling factors. [12] Nonetheless, in spite of substantial theoretical effort, none of the existing theories can explain other essential MIROs characteristics, including: (i) the independence of the MIROs amplitude on the left/right circular polarization [13], (ii) the necessity to assume equal transition probabilities induced by microwaves for electrons between Landau levels (LL) [12], (iii) large nonlinear effects of fractional harmonics [14], (iv) why MIROs are only observed in very high mobility samples while the theories have to introduce disorder (dirtier samples) to explain MIROs, (v) why MIROs are not observed in reflection experiments [15], (vi) why MIROs were never reported in the optical or even sub-millimeter frequency range, (vii) why magneto-plasmons are very much suppressed in MIROs experiments even though electrodynamics clearly plays an important role in these experiments. [15,16]

In this work we examine experimentally the importance of electrodynamics for MIROs experiments on 2DEG samples by studying their evolution over a wide range of frequencies. Below 120 GHz the MIROs waveform is described well by the existing models. At higher frequencies the waveform starts to deviate from the theory and they completely vanish at frequencies beyond 250 GHz. The results may be qualitatively understood within a model where plasmons intermediate the interaction process between the microwave radiation and the conduction electrons. Microwave radiation excites confined plasmon modes in a finite size sample. The plasmons decay through an interaction with the electrons exciting single particle transitions between Landau levels, the key mechanism in existing theoretical models to explain MIROs.

## 2. Experiment and Discussion

The experiment was performed on a GaAs/AlGaAs hetero-structure containing a high mobility two dimensional electron gas (2DEG) confined at the hetero-interface. After a brief illumination with a red LED, the 2DEG attained an electron concentration around  $2.0 \times 10^{11} \text{ cm}^{-2}$  and a mobility of  $8 \times 10^6 \text{ cm}^2/\text{Vs}$  at temperature 2 K. The sample was placed in a  $^4\text{He}$  cryostat equipped with a superconducting solenoid. The

sample was irradiated with microwaves using an oversized stainless steel pipe. A tunable klystron microwave generator (model ΓC-03) in combinations with a frequency doubler was used as the microwave source over 80 to 230 GHz range. The MW power at the sample was maintained at a constant level by simultaneously measuring an Alan Bradley carbon thermo resistor located near the sample.

The experimental traces of MIROs for different frequencies between 80 and 226 GHz are shown in Fig. 1 (a). The corresponding theoretical curves are presented in Fig. 1(b) which were calculated using the following equation: [12]

$$\Delta J_x(B) \propto \int d\varepsilon [n_F(\varepsilon) - n_F(\varepsilon + \hbar\omega)] \nu(\hbar\omega) \frac{\partial}{\partial \varepsilon} [\nu(\varepsilon + \hbar\omega)] \quad (1)$$

where  $\nu(E, B) = \sum_{i=0}^{\infty} \frac{eB}{\pi\hbar} \frac{1}{\pi\Gamma(1 + (\varepsilon - E_i)^2 / \Gamma^2)}$  is the electron density of states in a quantizing magnetic field,  $\varepsilon$  is electron energy in conduction band, and  $\Gamma$  is the width of the Landau levels,  $\omega = 2\pi f$  is the cyclic microwave frequency,  $n_F$  is the Fermi distribution function,  $E_i = (1/2 + i)\hbar\omega_c$  is the Landau energy spectrum, and  $\omega_c = eB/m^*$  is the cyclotron frequency. We note that the waveform described by equation (1) has only a single fitting parameter,  $\Gamma$ . A comparison between Figs. 1 (a) and (b) reveals that the simple theory works relatively well up to 110 GHz while for higher frequencies the MIROs waveforms distort and their amplitudes are monotonically reduced until at frequencies higher than 230 GHz they completely vanish. In ref. [17] it was shown that at even higher frequencies in the THz range the only remaining feature related to cyclotron resonance absorption at  $\omega = \omega_c$  occurs in the photo-response  $\Delta R_{xx}$  due to a bolometric type effect in the quantum Hall regime.

Let us examine the parameter  $\Gamma$  as a function of the microwave frequency. Figure 2 plots the MIROs as a function of  $\omega/\omega_c$  for different microwave frequencies. The experimental traces (points) are fitted by eq. (1) (the solid lines in figure 2). Although in the experiment the first harmonics ( $n=0$  and 1) start to deviate from the theory for frequencies above  $\sim 120$  GHz, it is still possible to fit the higher harmonic oscillations and estimate the parameter  $\Gamma$  from the fit. This is plotted in figure 3 as a function of the MW frequency. Since the fits were performed over approximately the same magnetic field range from  $\sim 0.02$  to 0.4 T, the large changes in  $\Gamma$  cannot be attributed to its magnetic field dependence. From Fig. 3 it is evident that  $\Gamma$  remains constant for frequencies below 120 GHz. We have performed similar frequency dependence measurements on a second sample and confirmed that  $\Gamma$  remains constant down to our lowest frequencies of 20 GHz. [17] In Fig. 3 the parameter  $\Gamma$  increases sharply from 25 to 60  $\mu\text{eV}$  over the frequency range from 120 to 156 GHz while at 226 GHz it increases even more rapidly until the MIROs completely vanish at higher frequencies. Such behaviour needs to be understood and, it clearly cannot be explained by the existing theoretical models [6,7,8,9] for which the microwave radiation interacts directly with the conduction electrons, since for these models the Landau level width,  $\Gamma$  cannot depend on any external parameter including, in our case, the microwave frequency. In fact, the theory predicts that the MIROs amplitude and the number of observable oscillations should increase with increasing frequency (Fig. 1(b))

in clear contradiction with the experiment, Fig. 1(a). Let us discuss the origin of this discrepancy and other unresolved experimental observations mentioned above within the context of 2D plasmons.

A key aspect of existing theoretical approaches is an assumption regarding the 2DEG microwave absorption process. The relevant theoretical models [6,7,8] assume that MIROs result from single particle excitations with a spatial displacement between occupied and empty LL electron states in the conduction band. To explain experimental waveforms it is necessary to assume that the absorption strength of the single particle excitations is independent of  $\Delta n = n_f - n_i$ , where  $n_f$  and  $n_i$  are the final and initial LL numbers. It is remarkable how well the MIROs waveform can be described with this counterintuitive assumption.[12] It should be noted that the direct absorption of light by harmonics of the cyclotron resonance is prohibited to first order and is usually very weak due to higher order corrections. A common conjecture is that this absorption is activated by disorder, which contradicts the experimental observation that MIROs are observed clearly only in samples with the highest possible mobilities meaning the least disorder. It is clear, therefore, that the physics of MIROs is more complex than that described by these existing models. The absorption process is mostly likely mediated by a collective phenomenon. This was underlined by Mikhailov [10,16,18]] very soon after the first MIROs reports, who pointed out the importance of plasmon excitations in the MIROs regime, for frequencies below the cyclotron frequency. Indeed, a number of experiments have demonstrated that microwave absorption by a 2DEG in the magnetic field range below the cyclotron resonance is due to excitations of confined, bulk as well as edge, plasmon modes. [12,16,19,20] On the other hand, there exists experimental evidence that magnetoplasmons do interact with higher harmonics of the cyclotron resonance [21,22], e.g. due to nonlocal effects [23]

Therefore, it is reasonable to assume that microwave radiation primarily induces plasmon excitations in finite size samples, which then decay into single-particle inter Landau level excitations, which in turn (consistent with the theoretical models) lead to the observation of MIROs. Although, a fully consistent picture which integrates the physics of plasmon excitations with the appearance of MIROs still remains to be theoretically developed, it is clear qualitatively that such an approach could indeed explain all of the unexplained experimental observations mentioned in the introduction.

We now turn to the principal effects reported in this paper. The increased damping of MIROs with frequency is to be expected within this picture as a result of the frequency dependent broadening of the plasmon excitations. Broadening of the intermediate plasmon excitations may be considered as an effective broadening of the excitation frequency,  $\omega$  in eq. (1). Numerical simulations using eq. (1) confirmed that a broadening of this excitation frequency is equivalent to a Landau level broadening. Therefore, if the width of the plasmon excitations,  $\Gamma_p$ , increases with frequency [25], the MIROs will experience an additional damping when  $\Gamma_p$  becomes comparable to or larger than  $\Gamma$ . Within this model it can be explained why MIROs are not observed in optics or in the far infrared regime as due to an essential plasmon lifetime decreases at higher frequencies,  $f$ . The number of excited plasmon modes increases quadratically with frequency as  $N_p \approx (f/f_p)^2$  where  $f_p = \sqrt{n_s e^2 / 2am^* \epsilon}$  is the fundamental plasmon mode,  $m^*$  is the electron effective mass,  $\epsilon$  is the effective dielectric constant of surrounding medium, and  $a$  is the

width of a rectangular sample. [18,24] In our sample ( $a=1.5$  mm)  $f_p \approx 18$  GHz and therefore at  $f=226$  GHz we have  $N_p \approx 158$ . Such large numbers of  $N_p$  may lead to a decrease in the plasmon oscillator strength and, significantly that new decay mechanisms may occur for such high modes leading to the line broadening observed in Fig. 3.

It is known that 2D plasmons have a very much reduced damping, in particular, the classical Landau damping is suppressed simply due to the energy and momentum conservation laws. [25] In very high quality, high mobility samples such as those used for MIROs experiments at low temperatures the lifetime of the plasmon excitation,  $\tau_p$ , becomes very long (with  $\Gamma_p = \hbar/\tau_p$ ), since the standard decay mechanisms are very much suppressed e.g. due to phonons and disorder. Under these circumstances, other decay channels through inter-Landau level excitations involving a spatial displacement, which does not depend on disorder, may become dominant. This explains why MIROs are observed only on the highest mobility samples at low temperatures. In addition, highly degenerate system like 2DEGs ( $E_F \gg k_B T$ ) is not sensitive to electron heating, therefore 2DEG helps further to enhance the MIROs by playing a selective role masking all absorption processes but the inter-Landau transitions with spatial displacements. In the same way, it is clear why the electron transition probability does not depend on the LL index change  $\Delta n$  [12] and why MIROs do not depend on the circular polarization [13]. The conduction electrons do not interact with microwaves directly but via plasmons so that the selection rules relevant for the interaction between light and free electrons do not apply.

### 3. Conclusion

We have experimentally studied the evolution of MIROs over a wide range of frequencies. At MW frequencies below  $\sim 120$  GHz the MIROs waveform is well described by existing theoretical models, for which the width of the Landau levels is the only fitting parameter. The LL width extracted by this procedure remains constant for  $f < 120$  GHz but rapidly increases for higher frequencies. The results can be qualitatively understood within a model for which the interaction between the microwave radiation and electrons is intermediated by plasmon excitations. The microwave radiation excites plasmons, and plasmons decay through the electron transitions between Landau levels giving rise to MIROs.

## Figures




Fig. 1. MIROs ( $\Delta R_{xx}=R_{xx}(\text{MWs on})-R_{xx}(\text{no MWs})$ ) for different MW frequencies from 80 to 226 GHz (a) experiment, (b) theory eq. (1) with the same Landau level width  $\Gamma=30 \mu\text{eV}$ . Traces are shifted vertically for clarity.




Fig. 2. Experimental traces of MIROs for different frequencies plotted as a function of  $\omega/\omega_c$  (green points) fitted with eq. (1) – thin solid lines.




Fig. 3. The Landau level width determined from MIROs.

- 
- [<sup>1</sup>] M. A. Zudov, R.R. Du, J. A. Simmons, and J. L. Reno, Phys. Rev. B **64**, 201311 (2001).  
[<sup>2</sup>] R. G. Mani, J. H. Smet, K. von Klitzing, V. Narayanamurti, W. B. Johnson, and V. Umansky, Nature **420**, 646 (2002).  
[<sup>3</sup>] M. A. Zudov, R. R. Du, L. N. Pfeiffer, and K. W. West, Phys. Rev. Lett. **90**, 046807 (2003).  
[<sup>4</sup>] S. A. Studenikin, M. Potemski, P. T. Coleridge, A. S. Sachrajda, and Z. R. Wasilewski, Solid State Commun. **129**, 341 (2004).  
[<sup>5</sup>] S. I. Dorozhkin, JETP Letters **77**, 577 (2003).  
[<sup>6</sup>] A. C. Durst, S. Sachdev, N. Read, and S. M. Girvin, Phys. Rev. Lett. **91**, 86803 (2003).  
[<sup>7</sup>] V. Ryzhii, Phys. Rev. B **68** (2003) 193402.  
[<sup>8</sup>] A. Dmitriev, M. G. Vavilov, I. L. Aleiner, A. D. Mirlin, and D. G. Polyakov et al. Phys. Rev. B **71**, 115316 (2005).  
[<sup>9</sup>] J. Inarrea, and G. Platero, Phys. Rev. Lett. **94** (2005) 016806.  
[<sup>10</sup>] S. A. Mikhailov, cond-mat/0303130 (2003).  
[<sup>11</sup>] A. D. Gladin and V.I. Ryzhii, Soviet Physics JETP **30**, 534 (1970).  
[<sup>12</sup>] S. A. Studenikin, M. Potemski, A. Sachrajda, M. Hilke, L. N. Pfeiffer, and K. W. West, Phys. Rev. B **71**, 245313 (2005).  
[<sup>13</sup>] J. H. Smet, B. Gorshunov, C. Jiang, L. Pfeiffer, K. West, V. Umansky, M. Dressel, R. Meisels, F. Kuchar, and K. von Klitzing, PRL **95**, 116804 (2005).  
[<sup>14</sup>] M. A. Zudov, R. R. Du, L. N. Pfeiffer, and K. W. West, Phys. Rev. B **73**, 041303 (2006).  
[<sup>15</sup>] S. A. Studenikin, M. Potemski, A. Sachrajda, M. Hilke, L. N. Pfeiffer, and K. W. West, IEEE Transactions on Nanotechnology **4**, 124 (2005).  
[<sup>16</sup>] S. A. Mikhailov, Phys. Rev. B **70** (2004) 165311  
[<sup>17</sup>] S. A. Studenikin et al. to be published in Physica E (2006).  
[<sup>18</sup>] S. A. Mikhailov, and N. A. Savostianova, Phys. Rev. B **71**, 035320 (2005).  
[<sup>19</sup>] E. Vasiliadou, G. Muller, D. Heitmann, D. Weiss, K. v. Klitzing, H. Nickel, W. Schlapp, and R. Losch, Phys. Rev. B **48**, 17145 (1993).  
[<sup>20</sup>] I. V. Kukushkin, J. H. Smet, V. A. Kovalskii, S. I. Gubarev, K. v. Klitzing, and W. Wegscheider, Phys. Rev. B **72**, 161317 (2005).  
[<sup>21</sup>] R. Krahne, M. Hochgrafe, Ch. Heyn, D. Heitmann, M. Hauser, and K. Eberl, Phys. Rev. B **62**, 15345 (2000).  
[<sup>22</sup>] E. Batke, D. Heitmann, and C. W. Tu, Phys. Rev. B **34**, 6951 (1986).  
[<sup>23</sup>] A. V. Chaplik, and D. Heitmann, J. Phys. C: Solid State Phys. **18**, 3357 (1985).  
[<sup>24</sup>] V. A. Volkov, and S. A. Mikhailov, "Electrodynamics of two-dimensional electron systems in high magnetic fields" in Landau Level Spectroscopy, Eds. G. Landwehr and E.I. Rashba, Elsevier Science Publishers, v.27.2, p.855 (1991)  
[<sup>25</sup>] C. A. Ullrich, and G. Vignale, Phys. Rev. B **58**, 7141 (1998).