

Conductance and density of states anomalies as resonances of energy bands in disordered coupled chains

L. Aloatti¹ and G. Grosso²

¹Max-Planck Institut für Metallforschung, Heisenbergstr. 3, D-70569 Stuttgart, Germany and Institut für Theoretische und Angewandte Physik, Universität Stuttgart, Pfaffenwaldring 57, D-70569 Stuttgart, Germany

²NEST-INFM and Dipartimento di Fisica "E. Fermi", Università di Pisa, Via F. Buonarroti 2, I-56127 Pisa, Italy

We show that off-diagonal nearest neighbor disorder in quasi-one-dimensional single particle tight-binding coupled chains leads to anomalies in the density of states and in the mean conductance, that can be interpreted as due to specific resonances of the band structure of the perfect system underlying the disordered one. We demonstrate that this phenomenology may appear not necessarily at the energy $E = 0$ as reported so far in the literature and we show that also the even-odd chain number effect on the mean conductance is ruled by the same resonances. For different cases we provide a series expansion for the resonant contribution to the mean conductance. These expansions allow to make predictions well verified numerically.

The existence of a divergent density of states at the center of the spectrum ($E = 0$) of a one-dimensional phonon model with off-diagonal disorder goes back to the pioneer paper of Dyson in 1953 [1]. The effect of this kind of disorder on the divergence of the localization length at the energy $E = 0$ was then evidenced [2, 3], in contrast with the general wisdom that in strictly one dimensional systems with uncorrelated diagonal or off-diagonal disorder all the states are localized [4, 5].

Recently further interest has grown on off-diagonal disorder in quasi-one-dimensional systems composed of N coupled chains. In this case it has been shown that for odd N the localization length and the density of states diverge at the center of the band, at zero energy, while they remain finite for even N [6].

It has been claimed that the origin of delocalization of the states at $E = 0$, is the existence of an additional (chiral) sublattice symmetry present in system with off-diagonal disorder only [7].

To our knowledge only the point $E = 0$ has been investigated in the literature. In this paper we look for the possibility of occurrence of the anomalies described above at other energy points. We show that such anomalies can arise in every point of the energy spectrum, provided a well defined condition on the energy bands of the perfect underlying chain system is fulfilled.

We start considering an infinitely extended periodic quantum wire composed by N interacting chains; diagonal and off-diagonal disorder is then introduced in a strand of length L of the wire (the lattice constant is taken as unit of length).

The Schrödinger equation for the multichain system considered is:

$$U_{n,n} + T_{n-1,n-1} + T_{n,n+1}^Y = E_n; \quad 8n; \quad (1)$$

where U_n is the matrix connecting the sites belonging to the n -th column and T_n is the matrix specifying the hopping amplitudes from the column n to the next one.

The order of these matrices is equal to the number of coupled chains, N . The index $n \in \mathbb{Z}$ labels the columns along the wire and the \mathbf{t}_n are N -dimensional vectors.

In order to introduce the disorder we write them matrices U_n and T_n in the following form:

$$U_n = U + W_n^U; \quad T_n = T + W_n^T; \quad (2)$$

where U and T are the matrices, independent of n , defining the periodic chain system and W_n^U and W_n^T are randomly distributed matrices. We take $W_n^U \neq 0$ only for $n = 1; \dots; L$ and $W_n^T \neq 0$ for $n = 1; \dots; L-1$. The distinction (2) defines the ordered part and the perturbation part (disorder) of the Hamiltonian, namely $H = H_0 + W$. All the elements of the random matrices have the form

$$(W_n^{U(T)})_{ij} = w_{ij}^{U(T)} e^{i \phi_{ij}^{U(T)}} \quad (3)$$

where all the $w_{ij}^{U(T)}$ and $\phi_{ij}^{U(T)}$ are independent and uniformly distributed real random variables, with zero mean and second moment equal to one. The term $w_{ij}^{U(T)}$ and $\phi_{ij}^{U(T)}$ do not depend on the column index n and regulate the modulus and the phase of the disorder.

Let us consider first the periodic chain system described by the Hamiltonian H_0 . Let $Z(\mathbf{B})$ be the matrix diagonalizing the hermitian matrix:

$$Q = T e^{i\mathbf{B}} + U + T^Y e^{i\mathbf{B}}; \quad (4)$$

where $\mathbf{B} \in [\pi; \pi]$ is the "Bloch angle", and the eigenvalues $E_i(\mathbf{B})$ form the band structure of the unperturbed system. The states

$$|z_n^{(j)}\rangle = Z^{(j)} e^{i\mathbf{B}}; \quad (5)$$

where $Z^{(j)}$ denotes the j -th column of the matrix $Z(\mathbf{B})$, form a complete set of eigenstates of H_0 .

The Landauer conductance $g(E)$ of the disordered wire can be calculated by means of the Møller operator [9]

$$g(E) = 1 + \frac{1}{(H - E + i)^2} W = 1 + G^R(E) W; \quad (6)$$

name the occurrence of this condition as " σ -coupling of the energy bands".

It is possible to see that, as far as the condition $(\epsilon_+^+)^2 = 1$ is satisfied, not only the fourth order of the average conductance acquires an additional term, but also every even-order term of the expansion in the disorder strength, so determining the magnification of the average conductance. We define the sum of all these diagrams as the "resonant contribution to the average conductance" $hg_{i_{\text{res}}}$.

For one-dimensional systems the resonance occurs at $E = 0$. With $T_{1,1} = \frac{1}{2}$ (which implies, at $E = 0$, $G_{1,1} = i$ and $\epsilon_+^+ = \frac{1}{2}$) and for diagonal disorder of intensity $w_{1,1}^U = w_d$ the sum of all the diagrams up to the twelfth order gives:

$$hg(E=0)_{i_{\text{res}}} = \frac{2}{2!}w_d^4L^2 + \frac{42}{3!}w_d^6L^3 + \frac{888}{4!}w_d^8L^4 + \frac{24336}{5!}w_d^{10}L^5 + \frac{1521631}{6!}w_d^{12}L^6 + \dots \quad (11)$$

With similar manipulations for real off-diagonal disorder of intensity $w_{1,1}^T = w_f$ we obtain:

$$hg(E=0)_{i_{\text{res}}} = \frac{32}{2!}w_f^4L^2 + \frac{1664}{3!}w_f^6L^3 + \frac{112640}{4!}w_f^8L^4 + \frac{10698752}{5!}w_f^{10}L^5 + \frac{1392902144}{6!}w_f^{12}L^6 + \dots \quad (12)$$

Interestingly, we have noticed numerically that for any given value of w_d there exists a value of w_f such that the resulting conductances $g(E)$ have almost the same off-resonance shape, while the height of the peak for off-diagonal disorder is much bigger than the other. In addition, we note that for off-diagonal disorder all the diagrams contributing to a fixed order in the expansion have the same sign, while for diagonal disorder there is a mutual cancellation of diagrams.

We have extended the previous approach to quantum wires composed by N different chains all with equal nearest-neighbor interactions and containing a disordered region always of the same kind and length. We have observed a strong dependence of the average conductance at $E = 0$ from the number N of chains. In particular, for N odd and a wide range of real off-diagonal disorder [8], $hg(E=0; N)$ is considerably greater than for the even case $N+1$. By comparing this result with the conductance evaluated at slightly different energies (where the resonance is destroyed), we can conclude that this even-odd effect of the conductance is just caused by the resonant terms discussed above.

In the case of more than one coupled quantum chains, the rule for calculating the diagrams which are relevant is very similar to the one depicted for the one dimensional case. The only difference consists in the fact that now there are more angles which can propagate and additional care has to be taken in selecting the paths contributing to the resonance. We give in Eq. (13) below the fourth

order expansion of the resonant contribution to the conductance of the two coupled chains with two resonating bands:

$$hg_{i_{\text{res}}} = \frac{L^2}{2!} \text{Tr} (P_1 M_m P_1 M_n P_2 M_m P_2 M_n P_1 + P_1 M_m P_2 M_n P_1 M_m P_2 M_n P_1 + P_1 M_m P_2 M_n P_2 M_m P_1 M_n P_1 + P_2 M_m P_2 M_n P_1 M_m P_1 M_n P_2 + P_2 M_m P_1 M_n P_2 M_m P_1 M_n P_2 + P_2 M_m P_1 M_n P_1 M_m P_2 M_n P_2) i + \text{cc} \quad (13)$$

where n and m are two arbitrarily chosen different column indexes and:

$$M_i = Z^Y G W_{U,i} Z + F Z^Y G W_{T,i} Z + Z^Y G W_{T,i} Z F^+ ;$$

$$P_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} ; P_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} ; F = \begin{pmatrix} e^{i\epsilon_1} & 0 \\ 0 & e^{i\epsilon_2} \end{pmatrix} :$$

The result of the previous trace is in general quite cumbersome, so we report it only for the case where the Bloch angles are $\epsilon_1 = \epsilon_2 = \frac{\pi}{3}$, and $\epsilon_+^+ = \epsilon_-^- = \frac{2}{3}$, and the matrices Z and G are given by:

$$Z = \begin{pmatrix} \cos & \sin \\ \sin & \cos \end{pmatrix} ; G = \begin{pmatrix} \frac{i}{3} & 0 \\ 0 & -\frac{i}{3} \end{pmatrix} : \quad (14)$$

It is easy to verify that for $\epsilon_+^+ = \frac{\pi}{4}$ we recover the conductance at $E = 0$ of two coupled chains with first neighbor hoppings all equal to 1.

Introducing the disorder, of equal intensity, only on the hoppings corresponding to the matrix elements $(W_n^T)_{1,1}$ and $(W_n^T)_{2,2}$, we obtain:

$$hg(\epsilon)_{i_{\text{res}}} = \frac{11}{6} \langle h(W_{1,1}^T)^2 i^2 L^2 \sin^4 + O(w^6 L^3) \rangle : \quad (15)$$

This \sin^4 behavior is in good qualitative agreement with the numerical data reported in Fig. 2. The most interesting cases are those for which the σ -coupling between the bands may occur for energies different from the center of the energy spectrum or only between a subgroup of all the bands. In this cases, in fact, the (chiral) sublattice symmetry claimed in the references [7] is broken. An example of this occurs for

$$U = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} ; T = \begin{pmatrix} 1 & 0.3 \\ 0.3 & 1 \end{pmatrix} : \quad (16)$$

In Fig. 3 (a) we report the density of states and in Fig. 3 (b) the average conductance for the two-chains system with off-diagonal disorder. The energy band structure of the perfect underlying system is reported in Fig. 3 (c). As a further example let us focus on the peak at $E = 1$ where all the bands are present but only one resonates. In this case, an equation similar to Eq. 13 gives:

$$hg_{i_{\text{res}}} / \langle h(W_{1,1}^T)^2 i + h(W_{2,2}^T)^2 i - h(W_{1,2}^U)^2 i^2 L^2 + \dots \rangle$$




FIG. 2: Average conductance for two coupled chains as function of the Bloch angle $\theta_2^+ = \theta_2$ and of the angle defining the matrix Z in (14). The angle $\theta_1^+ = \theta_1$ are fixed to be $\pi/3$. The peaks appearing for $\theta_2^+ = \pi/3$ are due to the resonance contribution and, as function of α , it follows qualitatively the \sin^4 predicted by (15). A gray-scale has been used to emphasize the peaks.




FIG. 3: Average DOS in arbitrary units (a), average conductance (b), and band structure (c) of the two coupled chains defined by (16). The density of states corresponds to the disordered region (device) isolated from the leads. There are three energies at which π -coupling occurs. In correspondence of these energies, the average conductance and the density of states are peaked.

The latter expression shows clearly that these two kinds of off-diagonal disorder (on the T and on the U matrices) compete with each other, and this is consistent with our

numerical simulations [8].

From the evaluation of the spectrum of eigenvalues for several configurations we have obtained the average DOS depicted in Fig. 3 (a). This figure evidences a direct connection between the peaks in the conductance and in the density of states. As far as we know, such a connection has been demonstrated for strictly one dimensional systems only [10].

Finally we justify why we kept in the development of the mean conductance only terms of the form $w^{2p}L^q$ with $p = q$ and no smaller values of q . The reason is that the average conductance $\langle g(E; w; L) \rangle$ converges at a fixed non-trivial value for every energy when $L \rightarrow \infty$ with w^2L constant (w represents the global strength of disorder). In practice we have verified that this convergence is very fast [8].

We would like to thank A. Cresti for having provided a very fast code, based on the Keldysh formalism [11], used as a check of our code and as a source of large statistics of the conductance.

-
- [1] F. J. Dyson, Phys. Rev. 92, 1331 (1953).
 - [2] G. Theodorou and M. H. Cohen, Phys. Rev. B 1, 4597 (1976).
 - [3] T. P. Eggarter and R. Riedinger, Phys. Rev. B 18, 569 (1978).
 - [4] R. E. Borland, Proc. R. Soc. Lond. A 274, 529 (1963).
 - [5] C. M. Soukoulis and E. N. Economou, Phys. Rev. B 24, 5698 (1981).
 - [6] P. W. Brouwer, C. Mudry, B. D. Simons, and A. Altland, Phys. Rev. Lett. 81, 862 (1998); P. W. Brouwer, C. Mudry, and A. Furusaki, Phys. Rev. Lett. 84, 2913 (2000); C. Mudry, P. W. Brouwer, and A. Furusaki, Phys. Rev. B 62, 8249 (2000); G. Chiappe and M. J. Sanchez, Phys. Rev. B 68, 155408 (2003).
 - [7] R. Oppermann and F. Wegner, Z. Phys. B 4, 327 (1979); Timothy A. L. Ziman, Phys. Rev. B 26, R7066 (1982); M. Inui, S. A. Trugman, and E. Abraham, Phys. Rev. B 49, 3190 (1994).
 - [8] L. Albatti and G. Grosso, to appear (2006).
 - [9] See e.g. M. L. Goldberger, and K. M. Watson, Collision theory (J. Wiley, New York, 1964), Chap. 5
 - [10] D. J. Thouless, J. Phys. C 5, 77 (1972).
 - [11] A. Cresti, R. Farchioni, G. Grosso and G. Pastori Parravicini, Phys. Rev. B 68, 75306 (2003).
 - [12] Necessary and sufficient condition is that the matrices U and T are both diagonalizable and that they have a common basis of eigenvectors. A sufficient condition is $T = T^\dagger$ and $[U; T] = 0$. In these cases the bands are cosine-like, namely $E(\theta) = \langle U \rangle + 2j \langle T \rangle \cos(\theta)$, where $\langle U; T \rangle$ are the eigenvalues of the matrices U and T and $\theta = \arctan[\langle T \rangle / \langle U \rangle]$.