

Mass and heat fluxes for a binary granular mixture at low-density

Vicente Garzo

Departamento de Física, Universidad de Extremadura, E-06071 Badajoz, Spain

Jose María Montanero

Departamento de Electrónica e Ingeniería Electromecánica,
Universidad de Extremadura, E-06071 Badajoz, Spain

James W. Duffy^z

Department of Physics, University of Florida, Gainesville, Florida 32611
(Dated: March 23, 2024)

The Navier-Stokes order hydrodynamic equations for a low density granular mixture obtained previously from the Chapman-Enskog solution to the Boltzmann equation are considered further. The six transport coefficients associated with mass and heat flux in a binary mixture are given as functions of the mass ratio, size ratio, composition, and coefficients of restitution. Their quantitative variation across this parameter set is demonstrated using low order Sonine polynomial approximations to solve the exact integral equations. The results are also used to quantify the violation of the Onsager reciprocal relations for a granular mixture. Finally, the stability of the homogeneous cooling state is discussed.

I. INTRODUCTION

The relevance and context of a hydrodynamic description for granular gases remains a controversial topic. At sufficiently low density the origin of hydrodynamics can be studied from Boltzmann kinetic theory.^{1,2} To obtain hydrodynamics from the Boltzmann kinetic equation the essential assumption is the existence of a "normal" solution, defined to be one for which all space and time dependence occurs through the macroscopic hydrodynamic fields.³ This solution, together with the macroscopic balance equations, leads to a closed set of hydrodynamic equations for these fields. The Chapman-Enskog method provides a constructive means to obtain an approximation to such a solution for states whose spatial gradients are not too large. In this general context, the study of hydrodynamics for granular gases is the same as that for normal gases.

The details of the Chapman-Enskog method have been carried out for a one component gas to obtain the Navier-Stokes order hydrodynamic equations, together with exact integral equations determining the transport coefficients occurring in these equations.⁴ These integral equations have been solved approximately using Sonine polynomial expansions and compared to both Direct Monte Carlo simulation of the Boltzmann equation and molecular dynamics simulation of the gas.⁵ Good agreement is obtained even for relatively strong degrees of dissipation. The results support the formal theoretical analysis and the claim that hydrodynamics is not limited to the quasi-elastic limit.

The analysis for multicomponent granular gases is much more complicated than for a one component gas. Most of the previous attempts⁶ were made for nearly elastic spheres where the equipartition of energy can be considered as an acceptable assumption. In addition, according to this level of approximation, the inelasticity is only accounted for by the presence of a sink term in the energy balance equation, so that the expressions for the transport coefficients are the same as those obtained for normal fluids.⁷ However, the failure of energy equipartition in multicomponent granular gases⁸ has been also confirmed by computer simulations⁹ and even observed in real experiments.¹⁰ Although the possibility of nonequipartition was already pointed out many years ago,¹¹ it has not been until recently that a systematic study of the effect of nonequipartition on transport has been carefully analyzed. In this context, Garzo and Duffy¹² have carried out a derivation of the Navier-Stokes hydrodynamic equations for a binary mixture at low-density that accounts for nonequipartition of granular energy. These equations and associated transport coefficients provide a somewhat more stringent test of the analysis since the parameter space is much larger. There are now many more transport coefficients, given as functions of the three independent coefficients of restitution, size ratio, mass ratio, and composition. As in the one component case, explicit expressions for the transport coefficients requires also to consider Sonine polynomial expansions. The accuracy of this approach has been confirmed by comparison with Monte Carlo simulations of the Boltzmann equation in the cases of the shear viscosity¹³ and the tracer diffusion¹⁴ coefficients. Exceptions to this agreement are extreme mass or size ratios and strong dissipation, although these discrepancies between theory and simulation diminish as one considers more terms in the Sonine polynomial approximation.¹⁴

Since the dependence of the shear viscosity coefficient on the parameters of the mixture (masses, sizes, concentration, coefficients of restitution) has been widely studied in a previous work,¹³ a primary objective here is to demonstrate

the variation of the six transport coefficients associated with the mass and heat flux in this parameter space, using the same Sonine polynomial approximation as was found applicable for the one component gas. To set the context for these quantitative results two qualitative and potentially confusing issues are briefly noted at the outset.

There is some ambiguity regarding the hydrodynamic temperature in a mixture since temperatures for each species can be defined in addition to the global temperature. What should be the hydrodynamic fields? For normal gases the answer is clear. These are set by the slow variables (at large space and time scales) associated with conserved quantities. In addition to species number and momentum, only the total kinetic energy is conserved, so only one global temperature occurs as a hydrodynamic field. The energy is no longer conserved for granular gases, but it remains a slow variable if the cooling rate is not too large. Thus, in this case as well, only the global temperature should appear among the hydrodynamic fields.

Nevertheless, the species temperatures play a new and interesting secondary role. For a normal gas, there is a rapid velocity relaxation in each fluid cell to a local equilibrium state on the time scale of a few collisions. Subsequently, the equilibration among cells occurs via the hydrodynamic equations. In each cell the species velocity distributions are characterized by the species temperatures. These are approximately the same due to equipartition, and the hydrodynamic relaxation occurs for the single common temperature.³ A similar rapid velocity relaxation occurs for granular gases in each small cell, but to a universal state different from local equilibrium and one for which equipartition no longer occurs. Hence, the species temperatures T_i are different from each other and from the overall temperature T of the cell.⁸ Nevertheless, the time dependence of all temperatures is the same in this and subsequent states, $T_i(t) = T(t)$. This implies that the species temperatures do not provide any new dynamical degree of freedom. They still characterize the shape of the partial velocity distributions and affect the quantitative averages calculated with these distributions. The transport coefficients for granular mixtures therefore have new quantitative effects arising from the time independent temperature ratio α_i for each species.¹² This dependence is illustrated here as well.

In some earlier works,^{11,15} additional equations for each species temperature have been included among the hydrodynamic set. This is possible since the overall temperature is determined from these by $T(t) = \sum_i x_i T_i(t)$, where x_i is the concentration of species i . However, this is an unnecessary complication, describing additional kinetics beyond hydrodynamics that is relevant only on the time scale of a few collisions. As described above (and supported by molecular dynamics simulations¹⁶) the dynamics of all temperatures quickly reduces to that of $T(t)$. The remaining time independent determination of the α_i follows directly from the condition that the cooling rates of all temperatures be the same.

A second confusing issue is the context of the Navier-Stokes equations considered here. The derivation of the Navier-Stokes order transport coefficients does not limit their application to weak inelasticity. For this reason the results reported below include a domain of both weak and strong inelasticity, $0.5 \leq \alpha \leq 1$. The Navier-Stokes hydrodynamic equations themselves may or may not be limited with respect to inelasticity, depending on the particular states considered. The derivation of these equations by the Chapman-Enskog method assumes that relative changes in the hydrodynamic fields over distances of the order of a mean free path are small. For normal gases this can be controlled by the initial or boundary conditions. It is more complicated for granular gases. In some cases (e.g., steady states such as the simple shear flow problem¹⁷) the boundary conditions imply a relationship between the coefficient of restitution and some hydrodynamic gradient so the two cannot be chosen independently. Consequently, there are examples for which the Navier-Stokes approximation is never valid or is restricted to the quasi-elastic limit.¹⁷ However, the transport coefficients characterizing the Navier-Stokes order hydrodynamic equations are well-defined functions of α , regardless of the applicability of those equations (e.g., the coefficients of a Taylor series for some function $f(\mathbf{x})$ in powers of x may be defined for all x at each order, even when the series cannot be truncated at that order).

The plan of the paper is as follows. First, in Sec. II the hydrodynamic equations and associated fluxes to Navier-Stokes order are recalled, and the expressions for the transport coefficients for heat and mass transport are given to leading Sonine polynomial approximation. The elastic limit is discussed to aid in the interpretation of these expressions. Next, in Sec. III the results for these six coefficients are illustrated for a common coefficient of restitution and same size ratio as functions of α at a composition $x_1 = 0.2$ for several values of the mass ratio. With the exception of thermal conductivity, the deviations from normal gas values are largest at small α and large mass ratio. The usual Onsager relations among these coefficients for normal gases is then noted and tested for the granular gas in Sec. IV. Since the underlying basis for these relations (time reversal symmetry) no longer holds for granular systems, the expected violation is demonstrated as a function of α for the same conditions. The stability of a spatially homogeneous solution to the mixture hydrodynamic equations is then studied, and some comments on the implications of the instability found are offered. Finally, the results are summarized and discussed in the last section.

II. BOLTZMANN KINETIC THEORY FOR THE MASS AND HEAT FLUXES

We consider a binary mixture of inelastic, smooth, hard spheres of masses m_1 and m_2 , and diameters σ_1 and σ_2 . The inelasticity of collisions among all pairs is characterized by three independent constant coefficients of normal restitution α_{11} , α_{22} , and $\alpha_{12} = \alpha_{21}$, where α_{ij} is the coefficient of restitution for collisions between particles of species i and j . It is assumed that the density of each species is sufficiently low that their velocity distribution functions are accurately described by the coupled set of inelastic Boltzmann kinetic equations. The precise form of these equations is given in Ref. 12 and will not be required here. These equations imply the exact macroscopic balance equations for the particle number density of each species, $n_i(\mathbf{r};t)$, flow velocity $\mathbf{u}(\mathbf{r};t)$, and temperature, $T(\mathbf{r};t)$,¹²

$$D_t n_i + n_i \nabla \cdot \mathbf{u} + \frac{\mathbf{r} \cdot \nabla n_i}{m_i} = 0; \quad i = 1, 2 \quad (1)$$

$$D_t \mathbf{u} + \nabla \cdot \mathbf{P} = 0; \quad (2)$$

$$D_t T + \frac{T}{n} \nabla \cdot \mathbf{u} + \frac{\mathbf{r} \cdot \nabla T}{m_i} + \frac{2}{3n} (\nabla \cdot \mathbf{q} + \mathbf{P} : \nabla \mathbf{u}) = -\dot{T}; \quad (3)$$

In the above equations, $D_t = \partial_t + \mathbf{u} \cdot \nabla$ is the material derivative, $\rho = m_1 n_1 + m_2 n_2$ is the total mass density, n_i is the particle number flux for species i , \mathbf{q} is the heat flux, \mathbf{P} is the pressure tensor, and \dot{T} is the cooling rate. For the two component mixture considered here there are six independent fields, n_1, n_2, T, \mathbf{u} . To obtain a closed set of hydrodynamic equations, expressions for $n_i, \mathbf{q}, \mathbf{P}$, and \dot{T} must be given in terms of these fields. Such expressions are called "constitutive equations". It is convenient to give these constitutive equations in terms of a different set of experimentally more accessible fields, x_1, p, T, \mathbf{u} , where $x_1 = n_1/(n_1 + n_2)$ is the composition of species 1, and $p = (n_1 + n_2)T$ is the hydrostatic pressure. This is simply a change of variables, so that Eqs. (1)-(3) become

$$D_t x_1 + \frac{\mathbf{r} \cdot \nabla x_1}{n^2 m_1 m_2} = 0; \quad (4)$$

$$D_t p + p \nabla \cdot \mathbf{u} + \frac{2}{3} (\nabla \cdot \mathbf{q} + \mathbf{P} : \nabla \mathbf{u}) = -\dot{p}; \quad (5)$$

$$D_t \mathbf{u} + \nabla \cdot \mathbf{P} = 0; \quad (6)$$

$$D_t T + \frac{T}{n} \nabla \cdot \mathbf{u} + \frac{\mathbf{r} \cdot \nabla T}{m_i} + \frac{2}{3n} (\nabla \cdot \mathbf{q} + \mathbf{P} : \nabla \mathbf{u}) = -\dot{T}; \quad (7)$$

The constitutive equations up to the Navier-Stokes order have been obtained from the Boltzmann equation in Ref. 12 with the results

$$\mathbf{j}_i = \frac{m_1 m_2 n}{\rho} \left(D \mathbf{r} x_1 - \frac{1}{p} D p \mathbf{r} \cdot \mathbf{r} - \frac{1}{T} D T \mathbf{r} \cdot \mathbf{r} \right); \quad \mathbf{j}_2 = -\mathbf{j}_1; \quad (8)$$

$$\mathbf{q} = -T^2 D^0 \mathbf{r} x_1 - L \mathbf{r} \cdot \mathbf{r} \cdot \nabla T; \quad (9)$$

$$P_{k\ell} = p \delta_{k\ell} + \frac{2}{3} \left(\mathbf{r} \cdot \mathbf{u}_k + \mathbf{r} \cdot \mathbf{u}_\ell \right) \delta_{k\ell} + \frac{2}{3} \mathbf{r} \cdot \mathbf{r} \cdot \mathbf{u}; \quad (10)$$

$$= \rho_0 + O(r^2) \quad (11)$$

The transport coefficients $\rho_0, D_p, D^0, L, \dots$ verify a set of coupled linear integral equations which can be solved approximately by using the leading terms in a Sonine polynomial expansion. This solution provides explicit expressions for the transport coefficients in terms of the coefficients of restitution and the parameters of the mixture (masses, sizes, and composition). The above expressions for mass and heat fluxes can be defined in a variety of equivalent

ways depending on the choice of driving forces used. For systems with elastic collisions, the specific set of gradients contributing to each flux is restricted by fluid symmetry, Onsager's relations (time reversal invariance), and the form of entropy production.¹⁸ In this case, one usual representation leads to the mass and heat fluxes proportional to $(r_i)_T$ and r_T , where i (defined below) is the chemical potential per unit mass. However, for inelastic systems only fluid symmetry holds and so there is more flexibility in representing the fluxes and identifying the corresponding transport coefficients. In particular, a third contribution proportional to r_p appears in both fluxes. Some care is required in comparing transport coefficients in different representations using different independent gradients for the driving forces.

The cooling rate to lowest order in the gradients is $\dot{Q} = \dot{Q}_0(x_1; p; T)$. There are no contributions to first order in the gradient for the low density Boltzmann equation. The general form including second order gradient contributions is displayed in Appendix A. These second order terms have been calculated for a one component gas⁴ and found to be very small. Here, these second order contributions to the cooling rate will be neglected.

Substitution of the Navier-Stokes constitutive equations, (8)-(10), into the exact balance equations, (4)-(7), gives the Navier-Stokes hydrodynamic equations for a binary mixture

$$D_t x_1 = \frac{r}{n^2 m_1 m_2} - \frac{m_1 m_2 n}{n^2 m_1 m_2} D r x_1 + \frac{D_p}{p} r p + \frac{D^0}{T} r T; \quad (12)$$

$$\begin{aligned} (D_t + \frac{5}{3} p r) u = \frac{2}{3} r T^2 D^0 r x_1 + L r p + r T \\ + \frac{2}{3} r \cdot u_k + r_k u, \quad \frac{2}{3} k \cdot r - u r \cdot u_k; \end{aligned} \quad (13)$$

$$\begin{aligned} (D_t + \frac{2}{3} p r) T + u = \frac{T m_2}{n m_1 m_2} \frac{m_1}{m_2} r - \frac{m_1 m_2 n}{n^2 m_1 m_2} D r x_1 + \frac{D_p}{p} r p + \frac{D^0}{T} r T \\ + \frac{2}{3 n} r T^2 D^0 r x_1 + L r p + r T \\ + \frac{2}{3 n} r \cdot u_k + r_k u, \quad \frac{2}{3} k \cdot r - u r \cdot u_k; \end{aligned} \quad (14)$$

$$D_t u + \frac{1}{3} r \cdot p = \frac{1}{3} r_k \cdot r \cdot u_k + r_k u, \quad \frac{2}{3} k \cdot r - u : \quad (15)$$

For the chosen set of fields $n = p = T$ and $\dot{Q} = p[(m_1 - m_2)x_1 + m_2] = T$. These equations are exact to second order in the spatial gradients for a low density Boltzmann gas.

A. Mass flux

The mass flux contains three transport coefficients, the diffusion coefficient D , the pressure diffusion coefficient D_p , and the thermal diffusion coefficient D^0 . Explicit expressions for these were obtained in Ref. 12 using a first Sonine approximation. Dimensionless forms are defined by

$$D = \frac{T}{m_1 m_2 \lambda_0} \bar{D}; \quad D_p = \frac{n T}{\lambda_0} \bar{D}_p; \quad D^0 = \frac{n T}{\lambda_0} \bar{D}^0; \quad (16)$$

Here, $\lambda_0 = \frac{p}{n} \frac{1}{v_0}$, $\lambda_{12} = (\lambda_1 + \lambda_2) = 2$, and $v_0 = \frac{p}{2T(m_1 + m_2)} = \frac{p}{m_1 m_2}$ is a thermal velocity defined in terms of the temperature T of the mixture. The explicit forms are then

$$\bar{D} = \frac{\partial}{\partial x_1} x_{1-1} \bigg|_{p,T} + \frac{\partial}{\partial x_1} \frac{1}{2} \bar{D}_p \frac{1}{2} \bar{D}^0; \quad (17)$$

$$\bar{D}_p = x_{1-1} \frac{(1 + \lambda_{12})}{1 + \lambda_{12}} \frac{3}{2} + \frac{2}{2} \bar{D}^0; \quad (18)$$

$$D^0 = \frac{1}{2} D_p; \quad (19)$$

In these equations,

$$\alpha_1 = \frac{T_1}{T} = \frac{1}{1 + x_1 \left(\frac{m_1}{m_2} - 1 \right)}; \quad \alpha_2 = \frac{T_2}{T} = \frac{1}{1 + x_1 \left(\frac{m_1}{m_2} - 1 \right)}; \quad (20)$$

where $\alpha = m_1/m_2$ is the mass ratio, $\alpha_{ij} = m_i/(m_i + m_j)$; $\alpha = x_1/x_2$, and $\alpha = T_1/T_2$. The detailed forms for the temperature ratio; dimensionless collision rate, and dimensionless cooling rate are given in Appendix A. Since $\alpha_{11} = \alpha_{22}$ and $\alpha x_1 = \alpha x_2$, it is expected that D should be symmetric with respect to interchange of particles 1 and 2 while D_p and D^0 should be antisymmetric. This can be easily verified by noting that $x_1 x_2 + x_2 x_1 = 1$.

In the case of elastic collisions $\alpha_{ij} = 1$, $\alpha = 0$, $\alpha = 1$, and Eqs. (17)-(19) become

$$D = \frac{3}{8} \frac{1 + \frac{1}{12} \left(\frac{m_1}{m_2} - 1 \right)}{\left(\frac{m_1}{m_2} + 1 \right)}; \quad D_p = x_1 \frac{\left(\frac{m_1}{m_2} - 1 \right)}{1 + \frac{1}{12} \left(\frac{m_1}{m_2} - 1 \right)} D; \quad D^0 = 0; \quad (21)$$

These coincide with known results obtained for elastic collisions in the first Sonine approximation.³ Recently, it has been shown that the estimate given by the first Sonine approximation for the diffusion coefficient D (in the very dilute concentration limit $x_1 \rightarrow 0$) compares quite well with Monte Carlo simulations,¹⁴ except for the cases in which the gas particles are much heavier and/or much larger than impurities. For these extreme cases, the second Sonine approximation to D improves the accuracy of the kinetic theory results.¹⁴

B. Heat flux

The heat flux requires going up to the second Sonine approximation. The transport coefficients D^0 , L , and appearing in the heat flux (9) are given by¹²

$$D^0 = \frac{5}{2} \frac{n}{(m_1 + m_2)^0} \left[\frac{x_1}{12} d_1^0 + \frac{x_2}{21} d_2^0 \right] - \frac{1}{12} - \frac{2}{21} D; \quad (22)$$

$$L = \frac{5}{2} \frac{T}{(m_1 + m_2)^0} \left[\frac{x_1}{12} \lambda_1 + \frac{x_2}{21} \lambda_2 \right] - \frac{1}{12} - \frac{2}{21} D_p; \quad (23)$$

$$= \frac{5}{2} \frac{nT}{(m_1 + m_2)^0} \left[\frac{x_1}{12} \lambda_1 + \frac{x_2}{21} \lambda_2 \right] - \frac{1}{12} - \frac{2}{21} D^0; \quad (24)$$

where the expressions for the (dimensionless) Sonine coefficients d_i^0 ; λ_i ; α_{ij} are displayed in Appendix A. In Eqs. (22)-(24) it is understood that the coefficients D , D_p , and D^0 are given by Eqs. (17)-(19), respectively (first Sonine approximation). As expected, our results show that D^0 is antisymmetric with respect to the change 1 \leftrightarrow 2 while L and D are symmetric. Consequently, in the case of mechanically equivalent particles ($m_1 = m_2 = m$, $\alpha = 1$, $\alpha_{ij} = 1$), the coefficient D^0 vanishes.

An equivalent representation is given in terms of the heat flow J_q defined as

$$J_q = q - \frac{5}{2} T \sum_i \frac{j_i}{m_i} = q - \frac{5}{2} T \frac{m_2 - m_1}{m_1 m_2} j; \quad (25)$$

where in the second equality use has been made of the requirement $j_1 = -j_2$. The difference between q and J_q is a heat flow due to diffusion. In addition, for elastic collisions, J_q is the flux conjugate to the temperature gradient in the form of the entropy production where the contribution coming from the mass flux couples only to the gradient of the chemical potentials. The thermal conductivity in a mixture is generally measured in the absence of diffusion, i.e., when $j = 0$. To identify this coefficient, we have to express J_q in terms of j , rT , and rp . The corresponding coefficient of rT defines the thermal conductivity.¹⁸ According to Eq. (8), the gradient of mole fraction rx_1 is

$$rx_1 = \frac{j}{m_1 m_2 n D} - \frac{2}{m_1 m_2 p D} \frac{D^0}{rT} - \frac{2}{m_1 m_2 n p D} \frac{D_p}{rp}; \quad (26)$$

The expression of J_q is obtained by substituting Eq. (9) into Eq. (25) and eliminating $r x_1$ by using the identity (26). Thus, the heat flow is given by

$$J_q = -r T + \frac{p}{m_1 m_2 n^2} T \frac{d}{dx} L_p r p; \quad (27)$$

where

$$= \frac{2 T^2}{m_1 m_2 p} \frac{D^0}{D}; \quad (28)$$

$$T = T \frac{D^0}{D} - \frac{5 n}{2} (m_2 - m_1); \quad (29)$$

$$L_p = L - \frac{2 T}{n^2 m_1 m_2} \frac{D^0 D_p}{D}; \quad (30)$$

As in the elastic case, the coefficient is the thermal conductivity while T is called the thermal-diffusion factor or Dufour coefficient. There is a new contribution proportional to $r p$ not present in the elastic case that defines the transport coefficient L_p .

For elastic collisions, $L_p = 0$ ¹⁹ and the expressions derived here for T and L_p coincide with those obtained for a gas-mixture of elastic hard spheres.³ Furthermore, in the case of mechanically equivalent particles, the Dufour coefficient T vanishes as expected and the heat flow (27) can be written as

$$J_q = -r T - r n; \quad (31)$$

where

$$-r = -r + n L_p = \frac{25}{32} \frac{m T}{n} \frac{1}{2} \frac{1+c}{2}; \quad (32)$$

$$-r = T L_p = \frac{75 T}{32 n} \frac{m T}{n} \frac{1}{2} \frac{2}{3} \frac{1+c}{2} + \frac{1}{3} \frac{c}{3} (2-3)^{-1}; \quad (33)$$

$$= \frac{1}{3} (1+c)^{-1} + \frac{33}{16} (1-c)^{-1} + \frac{19}{1024} c; \quad (34)$$

$$c = \frac{32 (1-c)^{-1} (1-2^2)}{81-17+30^2 (1-c)^{-1}}; \quad = \frac{5}{12} (1-c^2)^{-1} + \frac{3}{32} c; \quad (35)$$

Note that in writing Eq. (31) use has been made of the relation $r p = n r T + T r n$. Equations (31)-(35) are the same as those obtained for the one component granular gas.⁴ This confirms the relevant known limiting cases for the granular mixture results described here.

III. TRANSPORT COEFFICIENTS

The transport coefficients depend on many parameters: $f x_1; T; m_1 = m_2; \alpha = 2; \eta_1; \eta_2; \eta_3$. This complexity exists in the elastic limit as well, so the primary new feature is the dependence on the coefficients of restitution α_{ij} being different from unity. To illustrate the differences between granular and normal gases the transport coefficients are normalized to their values in the elastic limit. Then, the dependence on the overall temperature scales out. Also, only the simplest case of a common coefficient of restitution ($\alpha_{11} = \alpha_{22} = \alpha_{12}$) and common size ($\eta_1 = \eta_2 = \eta_3$) is considered. This reduces the parameter set to three quantities: $f m_1 = m_2; x_1; g$.

In Figs. 1-6, we plot the above transport coefficients as functions of the coefficient of restitution for $x_1 = 0.2$, and several values of the mass ratio. It is understood that all coefficients have been reduced with respect to their elastic values, except in the cases of D^0 and L_p since both coefficients vanish for elastic collisions. In these latter two




FIG. 1: Plot of the reduced mutual diffusion coefficient $D(\alpha)/D(1)$ as a function of the coefficient of restitution α for $x_1 = 0.2$, $\beta = 1$ and $\gamma = 0.5$ (a), $\gamma = 1$ (b), and $\gamma = 4$ (c). The dashed lines correspond to the approximation $c_1 = c_2 = 0$.




FIG. 2: Plot of the reduced pressure diffusion coefficient $D_p(\alpha)/D_p(1)$ as a function of the coefficient of restitution α for $x_1 = 0.2$, $\beta = 1$ and $\gamma = 0.5$ (a), $\gamma = 2$ (b), and $\gamma = 4$ (c).

cases, we have considered the reduced coefficients D^0 defined by Eq. (17) and $L_p = [(5/2)T_0/(m_1 + m_2)]^{1/2} L_p$. Figure 1 shows the mutual diffusion coefficient as a function of α for three mass ratios $\gamma = 0.5, 1$; and 4 . There is a monotonic increase of the coefficient with decreasing α in all cases. Moreover, that effect increases as the mass of the dilute species increases. This is consistent with an observed singular behavior in the extreme case of tracer diffusion for a massive particle.²⁰ The velocity distributions in a granular gas are no longer Maxwellian,⁸ and the difference is measured by the coefficients c_i that appear in the expressions for the transport coefficients.¹² The dashed curves in Fig. 1 correspond to $c_1 = c_2 = 0$, the Maxwell limit. In this case it is seen that the distortion of the Maxwellian is not very important for this transport coefficient (see however, discussion of Fig. 4 below). Figure 2 shows that the pressure diffusion coefficient has a very similar behavior. The thermal diffusion coefficient vanishes in the elastic limit and remains small and slightly negative when the dilute species has small mass ratio, as illustrated in Fig. 3. However as the mass ratio becomes large it becomes large and positive for strong dissipation. The effect of different species temperatures is also shown on this graph. The dashed curves correspond to setting $\beta = T_1/T_2 = 1$. This is seen to yield large errors, particularly in the case of large mass ratio (temperature differences are greater for mechanically different particles), indicating the real quantitative effect of two different species temperatures in granular gases.

The thermal conductivity is shown in Fig. 4, with the same monotonic increase with increasing dissipation. The dashed lines ($c_i = 0$) indicate a significant effect of the distortion of the reference distribution function from its Maxwellian form. Presumably, this is due to the fact that the thermal conductivity depends on a higher velocity moment than the mutual diffusion coefficient D and is more sensitive to the larger distortions at higher velocities. We also observe that there is little mass dependence when the dilute species is lighter. However, when the dilute




FIG. 3: Plot of the reduced thermaldiffusion coefficient $D^{T*}(\alpha)$ as a function of the coefficient of restitution α for $x_1 = 0.2$, $\beta = 1$ and $\gamma = 0.5$ (a), $\gamma = 2$ (b), and $\gamma = 4$ (c). The dashed lines correspond to $c_1 = c_2 = 0$.




FIG. 4: Plot of the reduced thermal conductivity coefficient $\kappa(\alpha)/\kappa(1)$ as a function of the coefficient of restitution α for $x_1 = 0.2$, $\beta = 1$ and $\gamma = 0.5$ (a), $\gamma = 1$ (b), and $\gamma = 4$ (c). The dashed lines correspond to the approximation $c_1 = c_2 = 0$.

species is more massive there is a significant decrease in the thermal conductivity, opposite to the case of diffusion. In contrast, the Dufour coefficient does have a dependence on the mass ratio more like diffusion (Fig. 5). Finally, the coefficient L_p is illustrated in Figure 6. A gain there is weak dependence on the mass ratio until it becomes larger for the dilute species.

In summary, the mass and heat flux transport coefficients for a granular mixture differ significantly from those for a normal gas mixture even at moderate dissipation. In most cases (thermal conductivity is an exception) the differences increase with decreasing α , depend weakly on the mass ratio when the dilute species ($x_1 = x_2 < 1$) is lighter than the excess species ($m_1 = m_2 < 1$) but increase significantly in the opposite case ($m_1 = m_2 > 1$).

IV. ONSAGER'S RECIPROCAL RELATIONS

In the usual language of the linear irreversible thermodynamics for ordinary fluids,¹⁸ the constitutive equations for the mass flux (8) and heat flow (27) is written

$$\mathbf{j}_i = \sum_j L_{ij} \left(-\frac{\nabla \mu_j}{T} \right) - L_{iq} \frac{\nabla T}{T^2} - C_p \nabla p; \quad (36)$$




FIG. 5: Plot of the reduced Dufour coefficient $\kappa_T(\alpha) = \kappa_T(1)$ as a function of the coefficient of restitution α for $x_1 = 0.2$, $\eta = 1$ and $\epsilon = 0.5$ (a), $\epsilon = 2$ (b), and $\epsilon = 4$ (c).




FIG. 6: Plot of the reduced coefficient $L_p^*(\alpha)$ as a function of the coefficient of restitution α for $x_1 = 0.2$, $\eta = 1$ and $\epsilon = 0.5$ (a), $\epsilon = 1$ (b), and $\epsilon = 4$ (c).

$$J_q = -L_{qq} \nabla T + \sum_i L_{qi} \frac{\nabla \mu_i}{T} + C_p^0 \nabla p; \quad (37)$$

where

$$\frac{\nabla \mu_i}{T} = \frac{1}{m_i} \nabla \ln(x_i p); \quad (38)$$

μ_i being the chemical potential per unit mass. Here, the coefficients L_{ij} are the so-called Onsager phenomenological coefficients. For normal fluids, Onsager showed¹⁸ that time reversal invariance of the underlying microscopic equations of motion implies important restrictions on the above set of transport coefficients

$$L_{ij} = L_{ji}; \quad L_{iq} = L_{qi}; \quad C_p = C_p^0 = 0; \quad (39)$$

The first two symmetries are called reciprocal relations as they relate transport coefficients for different processes. The last two are statements that the pressure gradient does not appear in any of the fluxes even though it is admitted by symmetry. Even for a one component fluid, Onsager's theorem is significant as it leads to Fourier's law for the heat flow rather than (31), i.e. $\nabla p = 0$. Since there is no time reversal symmetry for granular fluids, Eqs. (39) cannot be expected to apply. However, since explicit expressions for all transport coefficients are at hand, the quantitative extent of the violation can be explored.

To make connection with the previous sections it is first necessary to transform Eqs. (36)-(38) to the variables x_1 ; p ; T : Since $r x_1 = r x_2$, Eq. (38) implies

$$\frac{(r_1)_T}{T} = \frac{(r_2)_T}{T} = \frac{n}{1/2} r x_1 + \frac{n_1 n_2}{n} (m_2 - m_1) r \ln p : \quad (40)$$

The coefficients L_{ij} then can be easily obtained in terms of those of the previous sections. The result is

$$L_{11} = L_{12} = L_{21} = \frac{m_1 m_2}{2} D ; \quad L_{1q} = T D^0 ; \quad (41)$$

$$L_{q1} = L_{q2} = \frac{T^2}{n} D^0 + \frac{5T}{2} \frac{1/2}{2} (m_2 - m_1) D ; \quad L_{qq} = \frac{5}{2} \frac{m_2}{m_1 m_2} D^0 ; \quad (42)$$

$$C_p = -D_p = \frac{1/2}{p} (m_2 - m_1) D ; \quad (43)$$

$$C_p^0 = L = \frac{5T}{2p} \frac{m_2}{m_1 m_2} C_p = \frac{n_1 n_2}{np} T^2 (m_2 - m_1) D^0 : \quad (44)$$

The Onsager's relation $L_{12} = L_{21}$ is already evident since the diffusion coefficient D is symmetric under the change $1 \leftrightarrow 2$, as discussed following Eq. (20)

Imposing Onsager's relation $L_{1q} = L_{q1}$ yields

$$D^0 = \frac{5}{2} \frac{n}{T} (m_2 - m_1) D + \frac{n^2}{T} D^0 ; \quad (45)$$

while the condition $C_p = C_p^0 = 0$ leads to the following additional requirements

$$D_p = \frac{1/2}{3} (m_2 - m_1) D ; \quad (46)$$

$$\frac{5T}{2p} (m_1 - m_2) \frac{n_1 n_2}{2} (m_2 - m_1) D - \frac{D_p}{m_1 m_2} = pL = \frac{n_1 n_2}{n} T^2 (m_2 - m_1) D^0 : \quad (47)$$

Since the relations (45)-(47) involve transport coefficients that have been determined in the first Sonine approximation, we restrict here our discussion to this level of approximation. In this case, $d_1^{(0)} = \delta_1 = \delta_2 = 0$ so that Onsager's theorem, Eqs. (45)-(47), gives the conditions

$$P(ij) = [1 - 1 + (1 - 2)] \frac{D}{5} + \frac{1}{5} \frac{(1 +)}{(1 +)} D_p = 0 ; \quad (48)$$

$$Q(ij) = D_p - x_i \frac{(1)}{(1 +)} D = 0 ; \quad (49)$$

$$R(ij) = \frac{1 + }{(1 - 2)} Q(ij) = 0 : \quad (50)$$

In the elastic limit, the reduced coefficients D_p and D are given by Eq. (19) and these conditions are verified. Also, for mechanically equivalent particles with arbitrary γ , $\delta_1 = 1$ and $D_p = 0$ so that $P() = Q() = R() = 0$. Nevertheless, beyond these limit cases, Onsager's relations do not apply (as expected). At this macroscopic level the origin of this failure is due to the cooling of the reference state as well as the occurrence of different kinetic temperatures for both species. Figures 7, 8, and 9 show the dependence of the quantities P , Q , and R , respectively, on the (common) coefficient of restitution ϵ_{ij} for mass ratios $\gamma = 0.5$; 2; and 4.




FIG .7: Plot of $P(\alpha)$ as a function of α for $x_1 = 0.2$, $! = 1$ and $= 0.5$ (a), $= 2$ (b), and $= 4$ (c).




FIG .8: Plot of $Q(\alpha)$ as a function of α for $x_1 = 0.2$, $! = 1$ and $= 0.5$ (a), $= 2$ (b), and $= 4$ (c).




FIG .9: Plot of $R(\alpha)$ as a function of α for $x_1 = 0.2$, $! = 1$ and $= 0.5$ (a), $= 2$ (b), and $= 4$ (c).

V. LINEARIZED HYDRODYNAMIC EQUATIONS AND STABILITY

In contrast to normal fluids, the Navier-Stokes hydrodynamic equations (12)-(15) have non-trivial solutions even for spatially homogeneous states,

$$\partial_t x_{1H} = 0 = \partial_t u_H; \quad (51)$$

$$[\partial_t + (x_{1H}; T_H; p_H)] T_H = 0; \quad [\partial_t + (x_{1H}; T_H; p_H)] p_H = 0; \quad (52)$$

where the subscript H denotes the homogeneous state. Since the dependence of the cooling rate $(x_{1H}; T_H; p_H)$ on $x_{1H}; T_H; p_H$ is known (see Appendix A), these first order nonlinear equations can be solved for the time dependence of the homogeneous state. The result is the familiar Haff's law cooling law for $T(t)$ at constant density.² As discussed above, each species temperature also has the same time dependence but each with a different value⁸. In this section, the hydrodynamics for small initial spatial perturbations of this homogeneous cooling state (HCS) is discussed. For normal fluids such perturbations decay in time according to the hydrodynamic modes of diffusion (shear, thermal, mass) and damped sound propagation. The analysis is for fixed coefficients of restitution different from unity in the long wavelength limit. It will be seen here that the corresponding modes for a granular fluid are then quite different from those for a normal fluid. An alternative analysis with fixed long wavelength and coefficient of restitution approaching unity leads to usual normal fluid modes. Thus, the nature of hydrodynamic modes is non uniform with respect to the inelasticity and the wavelength of the perturbation.

Let $y(r; t) = y(r; t) - y_H(t)$ denote the deviation of $x_1; u; T; p$ from their values in the HCS. If the initial spatial perturbation is sufficiently small, then for some initial time interval these deviations will remain small and the hydrodynamic equations (12)-(15) can be linearized with respect to $y(r; t)$. This leads to partial differential equations with coefficients that are independent of space but which depend on time since the HCS is cooling. As in the one component case,^{4,21} this time dependence can be eliminated through a change in the time and space variables, and a scaling of the hydrodynamic fields. We introduce the following dimensionless space and time variables:

$$\begin{aligned} Z &= \int_0^t dt' v_H(t') \\ r^0 &= r - v_H \int_0^t dt' \end{aligned} \quad (53)$$

where $v_{0H} = \sqrt{\frac{2T_H(m_1 + m_2)}{m_1 m_2}}$ is the thermal velocity introduced above, $v_H = \sqrt{\frac{2}{m_H}} \frac{1}{\lambda_H}$ is an effective mean free path, and $\lambda_H(t) = \frac{v_H(t)}{v_{0H}(t)}$ is the effective collision frequency. The dimensionless time scale is therefore an average number of collisions up to the time t . A set of Fourier transformed dimensionless variables are then defined by

$$y_k(\tau) = \int dr^0 e^{ik \cdot r^0} y(r^0; \tau); \quad (54)$$

$$x_k(\tau) = \frac{x_{1k}(\tau)}{x_{1H}}; \quad u_k(\tau) = \frac{u_k(\tau)}{v_{0H}(\tau)}; \quad (55)$$

$$T_k(\tau) = \frac{T_k(\tau)}{T_H(\tau)}; \quad p_k(\tau) = \frac{p_k(\tau)}{p_H(\tau)}; \quad (56)$$

In terms of these variables the linearized hydrodynamic equations for the set $\{x_k; u_k; T_k; p_k\}$ separate into a degenerate pair of equations for the two transverse velocity components $w_{k\perp}$ (orthogonal to k)

$$\frac{\partial}{\partial \tau} \left(\frac{1}{2} + k^2 \right) w_{k\perp} = 0; \quad (57)$$

and a coupled set of equations for $x_k; u_k; T_k$, and the longitudinal velocity component $w_{k\parallel}$ (parallel to k)

$$\frac{\partial}{\partial \tau} \begin{pmatrix} x_k \\ u_k \\ T_k \end{pmatrix} = \begin{pmatrix} M^{(0)} + ikM^{(1)} + k^2 M^{(2)} \end{pmatrix} \begin{pmatrix} x_k \\ u_k \\ T_k \end{pmatrix}; \quad (58)$$

where now $z_k(\tau)$ denotes the four variables $x_k; u_k; T_k; w_{k\parallel}$. The matrices in this equation are

$$M^{(0)} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ x_1 \frac{\partial}{\partial x_1} & \frac{\partial}{\partial \tau} & 0 & 0 \\ x_1 \frac{\partial}{\partial x_1} & \frac{\partial}{\partial \tau} & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{2} \end{pmatrix}; \quad (59)$$

$$M^{(1)} = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} \frac{x_1 + x_2}{x_1 + x_2} & 0 & 0 \end{pmatrix}; \quad (60)$$

$$M^{(2)} = \begin{pmatrix} 0 & \frac{1}{2} \frac{x_1 + x_2}{1+} D & \frac{1}{2x_1} \frac{x_1 + x_2}{1+} D^0 & \frac{1}{2x_1} \frac{x_1 + x_2}{1+} D_p & 0 & 1 \\ \frac{2}{3} D^0 & \frac{1}{2(1+)} D & \frac{1}{2(1+)} D^0 & \frac{2}{3} & \frac{2}{3} L + \frac{1}{2(1+)} D_p & 0 \\ \frac{2}{3} x_1 D^0 & 0 & \frac{2}{3} & 0 & \frac{2}{3} L & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{4}{3} \end{pmatrix}; \quad (61)$$

In these equations, $x_1 = n_{1H}/n_H$, $x_2 = n_{2H}/n_H$ and the reduced transport coefficients D , D_p , and D^0 are given by Eqs. (17)–(19), respectively. Moreover, the reduced Navier–Stokes transport coefficients are

$$D^0 = \frac{\eta_H}{n_H v_{0H}^2}; \quad (62)$$

$$D^0 = \frac{\eta_H T_H D^0}{n_H v_{0H}^2}; \quad L = \frac{\eta_H L}{v_{0H}^2}; \quad = \frac{\eta_H}{n_H v_{0H}^2}; \quad (63)$$

where $\eta_H = m_1 n_{1H} + m_2 n_{2H}$.

It is instructive to consider first the solutions to these equations in the extreme long wavelength limit, $k = 0$. In this case, the eigenvalues or hydrodynamic modes are given by

$$s_2 = \frac{1}{2}; \quad s_n = 0; 0; \frac{1}{2}; \frac{1}{2}; \quad (64)$$

where s_n refers to the longitudinal modes. Two of the eigenvalues are positive, corresponding to growth of the initial perturbation in time. Thus, some of the solutions are unstable. The two zero eigenvalues represent marginal stability solutions, while the negative eigenvalue gives stable solutions. For general initial perturbations all modes are excited. These modes correspond to evolution of the fluid due to uniform perturbations of the HCS, i.e. a global change in the HCS parameters. The unstable modes are seen to arise from the initial perturbations $w_{kz}(0)$ or $w_{kjj}(0)$. The marginal modes correspond to changes in the composition at fixed pressure, density, and velocity, and to changes in n_{kz} at constant composition and velocity. The decaying mode corresponds to changes in the temperature or pressure for $n_k = n_k$. The unstable modes may appear trivial since they are due entirely to the normalization of the fluid velocity by the time dependent thermal velocity. However, this normalization is required by the scaling of the entire set of equations to obtain time independent coefficients.

At finite wave vectors, these instabilities give rise to real growth of spatial perturbations. The linear growth of the transverse modes is simply given by

$$w_{kz}(t) = w_{kz}(0) \exp\left(\frac{1}{2} k^2 t\right); \quad (65)$$

The instability for the two shear modes is removed at sufficiently large $k > k_z^c$, where

$$k_z^c = \frac{1}{2}; \quad (66)$$

The wave vector dependence of the remaining four modes is more complex. This is illustrated in Fig. 10 showing the real parts of the modes $s(k)$ for $x_{ij} = 0.9$; $x_1 = x_2 = 1$, $x_1 = 0.2$, and $m_1 = m_2 = 4$. The $k = 0$ values are those of (64), corresponding to six hydrodynamic modes with two different degeneracies. The shear mode degeneracy remains at finite k but the other is removed at any finite k . At sufficiently large k a pair of real modes become equal and become a complex conjugate pair at all larger wave vectors, like a sound mode. The smaller of the unstable modes is that associated with the longitudinal velocity, which couples to the scalar hydrodynamic fields. It becomes negative at a wave vector smaller than that of Eq. (66) and gives the threshold for development of spatial instabilities.

The results obtained here for the mixture show no new surprises relative to the earlier work for a one component gas,^{4,21} with only the addition of the stable mass diffusion mode. Of course, the quantitative features can be quite different since there are additional degrees of freedom with the parameter set $\{x_1, T; m_1 = m_2; x_1 = x_2; x_{ij}g\}$. Also, the manner in which these linear instabilities are enhanced by the nonlinearities may be different from that for the one component case.




FIG. 10: Dispersion relations for $\alpha = 0.9$, $x_1 = 0.2$, $\beta = 1$ and $\gamma = 4$.

VI. SUMMARY AND DISCUSSION

The Navier-Stokes order hydrodynamic equations have been discussed for a low density granular binary mixture. The form of the momentum flux is the same as for a one component gas, with only the value of the viscosity changed (see Appendix A). Since the dependence of the viscosity on the parameters of the mixture has been widely explored in a previous paper,¹³ attention has been focused here on the mass and heat fluxes and their associated transport coefficients. There is no phenomenology involved as the equations and the transport coefficients have been derived systematically from the inelastic Boltzmann equation by the Chapman-Enskog procedure. Consequently, there is no a priori limitation on the degree of inelasticity, size and mass ratios, or composition. For practical purposes, the integral equations determining the transport coefficients have been solved by truncated expansions in Sonine polynomials. This is expected to fail at extreme values of size or mass ratio,^{13,14} but the results are quite accurate otherwise.

The hydrodynamic equations are the same as for a normal gas, except for a sink in the energy equation due to granular cooling, and additional transport coefficients in the mass and heat flux constitutive equations. The latter arise because the usual restrictions of irreversible thermodynamics no longer apply. These restrictions include Onsager reciprocal relations among various transport coefficients, and the extent to which these are violated has been demonstrated in Section 4. It has been verified that the results described here reduce to those for a normal mixture in the elastic limit,³ and to those for a one component granular gas⁴ when the species are mechanically identical.

As is the case for a normal gas, the hydrodynamic fields include only the global temperature even though two species temperatures can be defined. For a normal gas the species temperatures rapidly approach the global temperature due to equipartition. For the granular gas the species temperatures approach different values, but with the same time dependence as the global temperature. The transport coefficients have an additional dependence on the composition due to this time independent ratio of species temperatures. This has been illustrated in Fig. 3 for the thermal diffusion coefficient, where the effect is seen to be large for large mass ratio (implying very different species temperatures). Generally, it was seen that the deviation due to inelasticity is enhanced for greater mechanical differences between the species.

The linear equations for small perturbations of the special homogeneous cooling solution to the hydrodynamic equations were obtained and discussed. In order to characterize the solutions in terms of modes, it is necessary to introduce dimensionless fields so that the time dependence of the reference HCS is eliminated. The resulting equations exhibit a long wavelength instability for three of the modes. This is quite similar to the case of a one component granular gas,^{4,21} and in fact the same modes are unstable here. The additional diffusion mode for two species behaves as for a normal fluid. The consequences of this instability for a binary mixture were not studied here. This entails an analysis of the dominant nonlinearities which has not been performed as yet. Since there are additional degrees of freedom would be interesting to see if the density clustering that occurs for a one component system is more complex here (e.g., species segregation).

The cooling rate and frequency are also functions of the hydrodynamic field x_1 . However, all the parameters above are independent of the temperature and density.

2. Heat and momentum flux parameters

As in the last section effects due to the distortion of the reference Maxwellian are neglected. In the case of the heat flux q , Eq. (9), the transport coefficients $D^{(0)}$, L , and A are given by Eqs. (22)-(24), respectively. By using matrix notation, the (dimensionless) Sonine coefficients $d_i^{(0)}$, γ_i , and α_i verify the coupled set of six equations¹²

$$X^{(0)} = Y \quad (A 6)$$

where $X^{(0)}$ is the column matrix

$$X = \begin{pmatrix} d_1^{(0)} \\ d_2^{(0)} \\ \gamma_1 \\ \gamma_2 \\ \alpha_1 \\ \alpha_2 \end{pmatrix}; \quad (A 7)$$

and Y is the matrix

$$Y = \begin{pmatrix} 0 & 11 & \frac{3}{2} & 12 & \frac{\partial}{\partial x_1} p;T & 0 & \frac{\partial}{\partial x_1} p;T & 0 \\ \frac{\partial}{\partial x_1} p;T & 0 & 0 & 0 & 11 & \frac{5}{2} & 12 & \frac{5}{2} \\ 0 & 0 & 0 & 0 & 21 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 22 & 0 & 11 & 12 \\ 0 & 0 & 0 & 0 & 0 & =2 & 21 & 22 \end{pmatrix} : \quad (A 8)$$

The column matrix Y has the elements

$$Y_1 = D_{12} \frac{1}{x_1^2} \frac{\partial}{\partial x_1} p;T; \quad Y_2 = D_{21} \frac{1}{x_2^2} \frac{\partial}{\partial x_1} p;T; \quad (A 9)$$

$$Y_3 = D_p \frac{1}{x_1^2}; \quad Y_4 = D_p \frac{1}{x_2^2}; \quad (A 10)$$

$$Y_5 = \frac{1}{1} + D^{(0)}_{12} \frac{1}{x_1^2}; \quad Y_6 = \frac{1}{2} D^{(0)}_{21} \frac{1}{x_2^2}; \quad (A 11)$$

The dimensionless collision integrals γ_{12} , γ_{11} , and γ_{22} are given, respectively, by

$$\gamma_{12} = \frac{4}{3} \frac{r}{2} \frac{1}{12} \frac{1}{1} \frac{1}{2} (1 + \frac{2}{11}) + \frac{4}{15} \frac{1}{21} \frac{1}{1} (1 + \frac{1}{12}) (1 + \frac{1}{2}) \frac{1}{2} (1 + \frac{2}{12}) \frac{x_2}{x_1} A + B; \quad (A 12)$$

$$\gamma_{11} = \frac{16}{15} x_1 \frac{1}{12} (2 + \frac{1}{12}) \frac{1}{2} (1 + \frac{1}{11}) + \frac{33}{16} (1 + \frac{1}{11}) + \frac{2}{15} x_2 \frac{1}{21} (1 + \frac{1}{12}) \frac{1}{2 (1 + \frac{1}{2})} E + 5 \frac{1 + \frac{2}{1}}{1} A; \quad (A 13)$$

$$\gamma_{22} = \frac{2}{15} x_2 \frac{1}{12} (1 + \frac{1}{12}) \frac{1}{2 (1 + \frac{1}{2})} F + 5 \frac{1 + \frac{2}{2}}{2} B; \quad (A 14)$$

In the above equations we have introduced the quantities²³

$$A = 5(2_{12} + 2_1) + 2_{21}(1 + 2_2) 5(1_{12}) 2(7_{12} 11)_{12} 1^1 + 18 \frac{2}{12} 1^1 + 2 \frac{2}{21} 2 \frac{2}{12} 3_{12} + 4 \frac{1}{1} (1 + 2_2)^2 5 \frac{2}{1} 1^1 (1 + 2_2) \quad (A15)$$

$$B = 5(2_{12} 1) + 2_{21}(1 + 2_2) 5(1_{12}) + 2(7_{12} 11)_{12} 2^1 + 18 \frac{2}{12} 2^1 2 \frac{2}{21} 2 \frac{2}{12} 3_{12} + 4 \frac{1}{2} (1 + 2_2)^2 + 5(1 + 2_2) \quad (A16)$$

$$E = 2 \frac{2}{21} 1^2 (1 + 2_2)^2 2 \frac{2}{12} 3_{12} + 4 (5_1 + 8_2) 2_1 (1 + 2_2) 2_{12} 1^2 (5_1 + 8_2) (7_{12} 11) + 2 \frac{2}{1} 1^1 (29_{12} 37) 25(1_{12}) + 18 \frac{2}{12} 1^2 (5_1 + 8_2) + 2_{12} 1^1 (25_1 + 66_2) + 5 \frac{2}{1} 1^1 (11_1 + 6_2) 5(1 + 2_2) 1^2 2 (5_1 + 6_2) \quad (A17)$$

$$F = 2 \frac{2}{21} 2^2 (1 + 2_2)^2 2 \frac{2}{12} 3_{12} + 4 (8_1 + 5_2) 2_1 (1 + 2_2) 2_{12} 2^2 (8_1 + 5_2) (7_{12} 11) 2_1 2^1 (29_{12} 37) + 25(1_{12}) + 18 \frac{2}{12} 2^2 (8_1 + 5_2) 2_{12} 2^1 (66_1 + 25_2) + 5 \frac{1}{1} 1^1 (6_1 + 11_2) 5(1 + 2_2) 2^1 (6_1 + 5_2) \quad (A18)$$

Here, $_{12} = \frac{1}{2} (2_{12} + 2_{21})$. The corresponding expressions for $_{21}$, $_{22}$, and $_{21}$ can be inferred from Eqs. (A12)-(A18) by interchanging 1 & 2. For elastic collisions, the expressions (A12)-(A18) reduce to those obtained for hard sphere mixtures.²⁴

The solution to Eq. (A6) is

$$X = \frac{1}{\rho} Y_0 \quad (A19)$$

This relation provides an explicit expression for the coefficients $d_1^{(0)}$, λ_1 , and μ_1 in terms of the coefficients of restitution and the parameters of the mixture. Their explicit forms are

$$d_1^{(0)} = \frac{1}{2} [2_{12} Y_2 - Y_1 (2_{22} - 3_1)]_{12} 2_{21} 11_{22} + 2(11_1 + 22_2) 4^2 + 2 \frac{\partial}{\partial x_1} \left(\frac{Y_3 + Y_5}{p;T} \right) (Y_3 + Y_5) 2_{12} 2_{21} + 2 \frac{2}{22} (7_{22} - 6_1) 2_{12} \frac{\partial}{\partial x_1} \left(\frac{Y_4 + Y_6}{p;T} \right) (Y_4 + Y_6) (2_{11} + 2_{22} - 7_1) ; \quad (A20)$$

$$\lambda_1 = \frac{1}{2} [2Y_3 2(12_{21} 11_{22})_{22} + (7_{11} 22 5_{12} 2_{21} + 2 \frac{2}{22} 6_{11} 7_{22} + 6^2) + 2Y_4 12_{22} 2_{12} 2_{21} 2_{11} 22 + 2(11_1 + 22_2) 2^2 + 2Y_5 2_{12} 2_{21} + 22(2_{22} - 7_1) + 6^2 2_{12} Y_6 [2(11_1 + 22_2) - 7_1] ; \quad (A21)$$

$$\mu_1 = \frac{1}{2} [Y_3 2_{12} 2_{21} + 22(2_{22} - 7_1) + 6^2 + 12_{22} Y_4 [2(11_1 + 22_2) - 7_1] + Y_5 4_{12} 2_{21} (2_{22} - 7_1) + 2 \frac{2}{22} (5_{12} 2_{11} + 2_{11} (7_{22} - 6^2) + 5^2 (6_{12} - 7_{22})) + 12 Y_6 [4_{12} 2_{21} + 2_{11} (5_{12} 2_{22}) + (10_{22} - 23_1)] g ; \quad (A22)$$

where

$$= 4(12_{21} 11_{22}) + 6(11_1 + 22_2) 9^2 12_{21} 11_{22} + 2(11_1 + 22_2) 4^2 ; \quad (A23)$$

The expressions for $d_2^{(0)}$, λ_2 , and μ_2 can be obtained from Eqs. (A20)-(A22) by setting 1 & 2. From the above expressions one can easily get the transport coefficients D^0 , L and η from Eqs. (22)-(24), respectively. They are functions of x_1 but independent of temperature and pressure.

The pressure tensor $P_{k\ell}$ is given by

$$P_{k\ell} = p \delta_{k\ell} - \eta (\partial_k u_\ell + \partial_\ell u_k) + \frac{2}{3} \delta_{k\ell} \partial_r u_r; \quad (\text{A } 24)$$

where η is the shear viscosity coefficient. Its expression can be written as¹²

$$\eta = \frac{nT}{0} \left(x_1^2 \frac{\partial^2}{\partial x_1^2} + x_2^2 \frac{\partial^2}{\partial x_2^2} \right); \quad (\text{A } 25)$$

with

$$\eta_1 = \frac{2 \frac{\partial^2}{\partial x_1^2} (2 \frac{\partial^2}{\partial x_2^2}) + 4 \frac{\partial^2}{\partial x_1 \partial x_2} (1 \frac{\partial^2}{\partial x_1^2} + 1 \frac{\partial^2}{\partial x_2^2})}{1 \frac{\partial^2}{\partial x_1^2} + 1 \frac{\partial^2}{\partial x_2^2} + 2 \frac{\partial^2}{\partial x_1 \partial x_2}}; \quad (\text{A } 26)$$

$$\eta_2 = \frac{2 \frac{\partial^2}{\partial x_1 \partial x_2} (2 \frac{\partial^2}{\partial x_1^2}) + 4 \frac{\partial^2}{\partial x_1 \partial x_2} (1 \frac{\partial^2}{\partial x_1^2} + 1 \frac{\partial^2}{\partial x_2^2})}{1 \frac{\partial^2}{\partial x_1^2} + 1 \frac{\partial^2}{\partial x_2^2} + 2 \frac{\partial^2}{\partial x_1 \partial x_2}}; \quad (\text{A } 27)$$

The dimensionless quantities η_{ij} are given by^{12,13}

$$\begin{aligned} \eta_{11} = & \frac{16}{5} x_1^2 \frac{1}{12} \frac{\partial^2}{\partial x_1^2} + \frac{1}{4} (1 \frac{\partial^2}{\partial x_1^2} + 1 \frac{\partial^2}{\partial x_2^2})^2 \\ & + \frac{8}{15} x_2^2 \frac{1}{12} (1 \frac{\partial^2}{\partial x_1^2} + 1 \frac{\partial^2}{\partial x_2^2})^2 + \frac{1}{6} \frac{\partial^2}{\partial x_1^2} (1 \frac{\partial^2}{\partial x_2^2} + 1 \frac{\partial^2}{\partial x_1 \partial x_2}) (1 \frac{\partial^2}{\partial x_1^2} + 1 \frac{\partial^2}{\partial x_2^2}) \\ & + \frac{3}{2} \frac{\partial^2}{\partial x_1^2} (1 \frac{\partial^2}{\partial x_2^2} + 1 \frac{\partial^2}{\partial x_1 \partial x_2}) (3 \frac{\partial^2}{\partial x_1^2} + 5 \frac{\partial^2}{\partial x_1 \partial x_2})^{1=2}; \end{aligned} \quad (\text{A } 28)$$

$$\begin{aligned} \eta_{22} = & \frac{8}{15} x_2^2 \frac{1}{12} (1 \frac{\partial^2}{\partial x_1^2} + 1 \frac{\partial^2}{\partial x_2^2})^2 + \frac{1}{6} \frac{\partial^2}{\partial x_2^2} (1 \frac{\partial^2}{\partial x_1^2} + 1 \frac{\partial^2}{\partial x_1 \partial x_2}) (1 \frac{\partial^2}{\partial x_1^2} + 1 \frac{\partial^2}{\partial x_2^2}) \\ & + \frac{3}{2} \frac{\partial^2}{\partial x_2^2} (1 \frac{\partial^2}{\partial x_1^2} + 1 \frac{\partial^2}{\partial x_1 \partial x_2}) (3 \frac{\partial^2}{\partial x_1^2} + 5 \frac{\partial^2}{\partial x_1 \partial x_2})^{1=2}; \end{aligned} \quad (\text{A } 29)$$

The corresponding expressions for η_{22} and η_{21} can be inferred from Eqs. (A 26) and (A 27) by interchanging 1 & 2.

The program for calculating the cooling rates, the temperature ratio and the transport coefficients of the binary mixture can be obtained on request from the authors.

E-mail: vicenteg@unex.es

^y E-mail: jnm@unex.es

^z E-mail: duffy@physu.edu

¹ A. G. Goldshtein and M. Shapiro, "Mechanics of collisional motion of granular materials. Part 1. General hydrodynamic equations," *J. Fluid Mech.* 282, 41 (1995); J. J. Brey, J. W. Duffy, and A. Santos, "Dissipative dynamics for hard spheres," *J. Stat. Phys.* 87, 1051 (1997).

² N. V. Brilliantov and T. Poschel, *Kinetic Theory of Granular Gases* (Oxford University Press, Oxford, 2004).

³ S. Chapman and T. G. Cowling, *The Mathematical Theory of Nonuniform Gases* (Cambridge University Press, Cambridge, 1970).

⁴ J. J. Brey, J. W. Duffy, C. S. Kim, and A. Santos, "Hydrodynamics for granular flow at low density," *Phys. Rev. E* 58, 4638 (1998); J. J. Brey and D. Cubero, "Hydrodynamic transport coefficients of granular gases," in *Granular Gases in Lectures Notes in Physics*, edited by T. Poschel and S. Luding (Springer, Berlin, 2001), p. 59.

⁵ See for instance, J. J. Brey, M. J. Ruiz-Montero, and D. Cubero, "On the validity of linear hydrodynamics for low-density granular flows described by the Boltzmann equation," *Europhys. Lett.* 48, 359 (1999); J. J. Brey, M. J. Ruiz-Montero, D. Cubero, and R. Garcia-Roja, "Self-diffusion in freely evolving granular gases," *Phys. Fluids* 12, 876 (2000); V. Garzo and J. M. Montanero, "Transport coefficients of a heated granular gas," *Physica A* 313, 336 (2002); J. M. Montanero, A. Santos, and V. Garzo, "DSMC evaluation of the Navier-Stokes shear viscosity of a granular fluid," in *Rarefied Gas Dynamics 24*, edited by M. Capitelli (AIP Conference Proceedings, 72, 2005), p. 803.

- ⁶ J. T. Jenkins and F. M. Ancini, \Kinetic theory for binary mixtures of smooth, nearly elastic spheres," *Phys. Fluids A* 1, 2050 (1989); P. Zamankhan, \Kinetic theory for multicomponent dense mixtures of slightly inelastic spherical particles," *Phys. Rev. E* 52, 4877 (1995); B. A. Marson and J. T. W. Illits, \Thermal diffusion in binary mixtures of smooth, nearly elastic spheres with and without gravity," *Phys. Fluids* 10, 1324 (1998); J. T. W. Illits and B. A. Marson, \Kinetic theory of a binary mixture of nearly elastic disks," *Phys. Fluids* 11, 3116 (1999); M. Alam, J. T. W. Illits, B. A. Marson, and S. Luding, \Kinetic theory of a binary mixture of nearly elastic disks with size and mass disparity," *Phys. Fluids* 14, 4085 (2002); B. A. Marson and J. T. Jenkins, \Binary mixtures of inelastic hard spheres: Simplified constitutive theory," *Phys. Fluids* 16, 4543 (2004.)
- ⁷ M. Lopez de Haro, E. G. D. Cohen, and J. M. K. Incaid, \The Enskog theory for multicomponent mixtures. I. Linear response theory," *J. Chem. Phys.* 78, 2746 (1983).
- ⁸ V. Garzo and J. W. Duffy, \Homogeneous cooling state for a granular mixture," *Phys. Rev. E* 60, 5706 (1999).
- ⁹ See for instance, J. M. Montanero and V. Garzo, \Monte Carlo simulation of the homogeneous cooling state for a granular mixture," *Granular Matter* 4, 17 (2002); A. Barrat and E. Trizac, \Lack of energy equipartition in homogeneous heated binary granular mixtures," *Granular Matter* 4, 57 (2002); R. C. Lelland and C. M. Hrenya, \Simulations of a binary-sized mixture of inelastic grains in rapid shear flow," *Phys. Rev. E* 65, 031301 (2002); S. R. Dahl, C. M. Hrenya, V. Garzo, and J. W. Duffy, \Kinetic temperatures for a granular mixture," *Phys. Rev. E* 66, 041301 (2002); R. Pagnani, U. M. B. Marconi, and A. Puglisi, \Driven low density granular mixtures," *Phys. Rev. E* 66, 051304 (2002); D. Paolotti, C. Cattuto, U. M. B. Marconi, and A. Puglisi, \Dynamical properties of vibrated granular mixtures," *Granular Matter* 5, 75 (2003); P. K. Rouskop and J. Talbot, \Mass and size effects in three-dimensional vibrated granular mixtures," *Phys. Rev. E* 68, 021304 (2003); H. Wang, G. Jin, and Y. Ma, \Simulation study on kinetic temperatures of vibrated binary granular mixtures," *Phys. Rev. E* 68, 031301 (2003); J. J. Brey, M. J. Ruiz-Montero, and F. Moreno, \Energy partition and segregation for an intruder in a vibrated granular system under gravity," *Phys. Rev. Lett.* 95, 098001 (2005).
- ¹⁰ R. D. Wildman and D. J. Parker, \Coexistence of two granular temperatures in binary vibrated beds," *Phys. Rev. Lett.* 88, 064301 (2002); K. Feitosa and N. Menon, \Breakdown of energy equipartition in a 2D binary vibrated granular gas," *Phys. Rev. Lett.* 88, 198301 (2002).
- ¹¹ J. Jenkins and F. M. Ancini, \Balance laws and constitutive relations for plane flows of a dense binary mixture of smooth, nearly elastic, circular disks," *J. Appl. Mech.* 54, 27 (1987).
- ¹² V. Garzo and J. W. Duffy, \Hydrodynamics for a granular binary mixture at low density," *Phys. Fluids* 14, 1476 (2002).
- ¹³ J. M. Montanero and V. Garzo, \Shear viscosity for a heated granular binary mixture at low-density," *Phys. Rev. E* 67, 021308 (2003).
- ¹⁴ V. Garzo and J. M. Montanero, \Diffusion of impurities in a granular gas," *Phys. Rev. E* 69, 021301 (2004).
- ¹⁵ L. Huilin, D. Gidaspow, and E. Manger, \Kinetic theory of vibrated binary granular mixtures," *Phys. Rev. E* 64, 061301 (2001); M. F. Ramahan, J. Naser, and P. J. Witt, \An unequal temperature kinetic theory: description of granular flow with multiple particle classes," *Powder Tech.* 138, 82 (2003); J. E. Galvin, S. R. Dahl, and C. M. Hrenya, \On the influence of the equipartition-of-energy assumption in kinetic theories for rapidly-flowing, granular mixtures," *J. Fluid Mech.* 528, 207 (2005).
- ¹⁶ S. R. Dahl, C. M. Hrenya, V. Garzo, and J. W. Duffy, \Kinetic temperatures for a granular mixture," *Phys. Rev. E* 66, 041301 (2002).
- ¹⁷ A. Santos, V. Garzo, and J. Duffy, \Inherent rheology of a granular fluid in uniform shear flow," *Phys. Rev. E* 69, 061303 (2004).
- ¹⁸ S. R. de Groot and P. Mazur, *Nonequilibrium Thermodynamics* (Dover, New York, 1984).
- ¹⁹ Note that this coefficient is not strictly zero in the approximation used here due to the fact that while the set $\{f_i^{(0)}; i; i_g\}$ has been evaluated in the second Sonine approximation, the coefficients D_{\parallel} , D_{\perp} , and D^0 have been estimated in the first Sonine approximation. However, the magnitude of L_p is negligible.
- ²⁰ A. Santos and J. W. Duffy, \Critical behavior of a heavy particle in a granular fluid," *Phys. Rev. Lett.* 86, 4823 (2001); \Nonequilibrium phase transition for a heavy particle in a granular fluid," *Phys. Rev. E* 64, 051305 (2001).
- ²¹ V. Garzo, \Instabilities in a free granular fluid described by the Enskog equation," *Phys. Rev. E* 72, 021106 (2005).
- ²² V. Garzo and J. W. Duffy, \Dense fluid transport for inelastic hard spheres," *Phys. Rev. E* 59, 5895 (1999).
- ²³ Some misprints occur in the expressions given in Ref. 12 for these quantities. The expressions given here correct such results.
- ²⁴ See Ref. 3, Chap. 9, Eqs. (9.6,1)-(9.6,10).