

Rabi oscillations in a qubit coupled to a quantum two-level system

S. Ashhab,¹ J. R. Johansson,¹ and Franco Nori^{1,2}

¹Frontier Research System, The Institute of Physical and Chemical Research (RIKEN), Wako-shi, Saitama 351-0198, Japan

²Center for Theoretical Physics, C S C S, Department of Physics,
University of Michigan, Ann Arbor, Michigan 48109-1040, USA

(dated: December 31, 2021)

We consider the problem of a qubit driven by a harmonically oscillating external field while it is coupled to a quantum two-level system. We perform a systematic numerical analysis of the problem by varying the relevant parameters. The numerical calculations agree with the predictions of a simple intuitive picture, namely one that takes into consideration the four-level energy spectrum, the simple principles of Rabi oscillations and the basic effects of decoherence. Furthermore, they reveal a number of other interesting phenomena. We provide explanations for the various features that we observe in the numerical calculations and discuss how they can be used in experiment. In particular, we suggest an experimental procedure to characterize an environment of two-level systems.

I. INTRODUCTION

There have been remarkable advances in the field of superconductor-based quantum information processing in recent years [1]. Coherent oscillations and basic gate operations have been observed in systems of single qubits and two interacting qubits [2, 3, 4, 5, 6, 7, 8, 9]. One of the most important operations that are used in manipulating qubits is the application of an oscillating external field on resonance with the qubit to drive Rabi oscillations [3, 4, 5, 6, 10]. A closely related problem with great promise of possible applications is that of a qubit coupled to a quantum harmonic oscillator mode [11, 12, 13, 14].

Qubits are always coupled to uncontrollable degrees of freedom that cause decoherence in its dynamics. One generally thinks of the environment as slowly reducing the coherence of the qubit, typically as a monotonically decreasing decay function. In some recent experiments, however, oscillations in the qubit have been observed that imply it is strongly coupled to quantum degrees of freedom with long decoherence times [10, 15]. The effects of those degrees of freedom have been successfully described by modelling them as quantum two-level systems (TLSs) [16, 17, 18, 19, 20]. Since, as mentioned above, Rabi oscillations are a simple and powerful method to manipulate the quantum state of a qubit, it is important to understand the behaviour of a qubit that is driven on or close to resonance in the presence of such a TLS. Furthermore, we shall show below that driving the qubit close to resonance can be used to extract more parameters about an environment of TLSs than has been done in experiment so far. The results of this study are also relevant to the problem of Rabi oscillations in a qubit that is interacting with other surrounding qubits.

Some theoretical treatments and analysis of special cases of the problem at hand were given in Refs. [15, 19]. In this paper we perform a more systematic analysis in order to reach a more complete understanding of this phenomenon. We shall present a few simple physical principles that can be used to understand several aspects of the

behaviour of this system with different possible choices of the relevant parameters. Those principles are (1) the four-level energy spectrum of the qubit+TLS system, (2) the basic properties of the Rabi-oscillation dynamics and (3) the basic effects of decoherence. We shall then perform numerical calculations that will agree with that intuitive picture and also will reveal other results that are more difficult to definitively predict otherwise. Finally, we suggest an experimental procedure where the driven qubit dynamics can be used to characterize the environment of TLSs.

The paper is organized as follows: in Sec. II we introduce the model system and the Hamiltonian that describes it. In Sec. III we present a few simple arguments that will be used as a foundation for our numerical analysis of Sec. IV, which will confirm that intuitive picture and reveal other less intuitively predictable results (note that a reader who is sufficiently familiar with the subject matter can skip Sec. III). In Sec. V we discuss how our results can be used in experiment. We finally conclude our discussion in Sec. VI.

II. MODEL SYSTEM

The model system that we shall study in this paper is composed of a harmonically driven qubit, a quantum TLS and their weakly coupled environment [21]. We assume that the qubit and the TLS interact with their own (uncorrelated) environments that would cause decoherence even in the absence of qubit-TLS coupling. The Hamiltonian of the system is given by:

$$\hat{H}(t) = \hat{H}_q(t) + \hat{H}_{\text{TLS}} + \hat{H}_I + \hat{H}_{\text{ENV}}; \quad (1)$$

where \hat{H}_q and \hat{H}_{TLS} are the qubit and TLS Hamiltonians, respectively; \hat{H}_I describes the coupling between the qubit and the TLS, and \hat{H}_{ENV} describes all the degrees of freedom in the environment and their coupling to the

qubit and TLS. The (time-dependent) qubit Hamiltonian is given by:

$$\hat{H}_q(t) = \frac{q}{2} \hat{\sigma}_x^{(q)} - \frac{q}{2} \hat{\sigma}_z^{(q)} + F \cos(\omega t) \sin \theta \hat{\sigma}_x^{(q)} + \cos \theta \hat{\sigma}_z^{(q)}; \quad (2)$$

where q and q are the adjustable static control parameters of the qubit, $\hat{\sigma}^{(q)}$ are the Pauli spin matrices of the qubit, F and ω are the amplitude (in energy units) and frequency, respectively, of the driving field, and θ is an angle that describes the orientation of the external field relative to the qubit $\hat{\sigma}_z$ axis. Although we have used a rather general form to describe the coupling between the qubit and the driving field, we shall see in Sec. III that one only needs a single, easily extractable parameter to characterize the amplitude of the driving field. We assume that the TLS is not coupled to the external driving field, and its Hamiltonian is given by:

$$\hat{H}_{\text{TLS}} = \frac{J_{\text{TLS}}}{2} \hat{\sigma}_x^{(\text{TLS})} - \frac{J_{\text{TLS}}}{2} \hat{\sigma}_z^{(\text{TLS})}; \quad (3)$$

where the definition of the parameters and operators is similar to those of the qubit, except that the TLS parameters are uncontrollable. Note that our assumption that the TLS is not coupled to the driving field can be valid even in cases where the physical nature of the TLS and the driving field leads to such coupling, since we generally consider a microscopic TLS, rendering any coupling to the external field negligible.

The energy splitting between the two quantum states of each subsystem, in the absence of coupling between them, is given by:

$$E = \frac{q}{2} \sqrt{1 + \frac{q^2}{J^2}}; \quad (4)$$

where the index refers to either the qubit or the TLS. The corresponding ground and excited states are, respectively, given by:

$$\begin{aligned} |g\rangle_i &= \cos \frac{\theta}{2} |0\rangle_i + \sin \frac{\theta}{2} |1\rangle_i \\ |e\rangle_i &= \sin \frac{\theta}{2} |0\rangle_i - \cos \frac{\theta}{2} |1\rangle_i; \end{aligned} \quad (5)$$

where the angle θ is given by the criterion $\tan \theta = \frac{q}{J}$. We take the interaction Hamiltonian between the qubit and the TLS to be of the form:

$$\hat{H}_I = \frac{g}{2} \hat{\sigma}_z^{(q)} \hat{\sigma}_z^{(\text{TLS})}; \quad (6)$$

where g is the (uncontrollable) coupling strength between the qubit and the TLS. Note that any interaction Hamiltonian that is a product of a qubit observable times a

TLS observable can be recast into the above form using a simple basis transformation, keeping in mind that such a basis transformation also changes the values of q , J_{TLS} and θ .

We assume that all the coupling terms in \hat{H}_{ENV} are small enough that its effect on the dynamics of the qubit+TLS system can be treated within the framework of the Markovian Bloch-Redfield master equation approach. We shall use a noise power spectrum that can describe both dephasing and relaxation with independently adjustable rates, and shall present our results in terms of those decoherence rates. For definiteness in the numerical calculations, we take the coupling of the qubit and the TLS to their respective environments to be described by the operators $\hat{\sigma}_z^{(i)}$, where i refers to the qubit and the TLS. Note, however, that since we use the relaxation and dephasing rates to quantify decoherence, our results are independent of the choice of system-environment coupling operators.

III. INTUITIVE PICTURE

We start our analysis of the problem by presenting a few physical principles that prove very helpful in intuitively predicting the behaviour of the above-described system. Note that the arguments given in this section are well known [22, 23, 24]. For the sake of clarity, however, we present them explicitly and discuss their roles in the problem at hand.

A. Energy levels and eigenstates

The first element that one needs to consider is the energy levels of the combined qubit+TLS system. In order for a given experimental sample to function as a qubit, the qubit-TLS coupling strength must be much smaller than the energy splitting of the qubit E_q . We therefore take that limit, as well as the limit E_{TLS} , and straightforwardly find the energy levels to be given by:

$$\begin{aligned} E_1 &= \frac{E_{\text{TLS}} + E_q}{2} - \frac{cc}{2} \\ E_2 &= \frac{1}{2} \sqrt{q^2 + (E_{\text{TLS}} - E_q)^2} + \frac{2}{2} \frac{ss}{2} + \frac{cc}{2} \\ E_3 &= + \frac{1}{2} \sqrt{q^2 + (E_{\text{TLS}} - E_q)^2} + \frac{2}{2} \frac{ss}{2} + \frac{cc}{2} \\ E_4 &= + \frac{E_{\text{TLS}} + E_q}{2} - \frac{cc}{2}; \end{aligned} \quad (7)$$

where $cc = \cos \theta \cos \theta_{\text{TLS}}$, $ss = \sin \theta \sin \theta_{\text{TLS}}$. The corresponding eigenstates are given by:

$$\begin{aligned} |g\rangle_i &= |g\rangle_i \\ |e\rangle_i &= \cos \frac{\theta}{2} |e\rangle_i + \sin \frac{\theta}{2} |g\rangle_i \end{aligned}$$

$$\begin{aligned}\beta_i &= \sin \frac{\theta'}{2} |e_i\rangle - \cos \frac{\theta'}{2} |g_i\rangle \\ \alpha_i &= |g_i\rangle;\end{aligned}\quad (8)$$

where the first symbol refers to the qubit state and the second one refers to the TLS state in their respective uncoupled bases, the angle θ' is given by the criterion $\tan \theta' = \frac{V}{\Delta E_{\text{TLS}} - E_q}$, and for definiteness in the form of the states $|\alpha_i\rangle$ and $|\beta_i\rangle$ we have assumed that $E_{\text{TLS}} > E_q$.

Note that the mean-field shift of the qubit resonance frequency, ω_{cc} , is present regardless of the values of the qubit and TLS energy splittings. The avoided-crossing structure involving states $|\alpha_i\rangle$ and $|\beta_i\rangle$, however, is only relevant when the qubit and TLS energies are almost equal. One can therefore use spectroscopy of the four-level structure to experimentally measure the TLS energy splitting E_{TLS} and angle θ_{TLS} , as will be discussed in more detail in Sec. V.

B. Rabi oscillations

If a two-level system (e.g. a qubit) with energy splitting Δ_0 , initially in its ground state, is driven by a harmonically-oscillating weak field with a frequency ω close to its energy splitting (up to a factor of \hbar) as described by Eq. (3), its probability to be found in the excited state at a later time t is given by:

$$P_e = \frac{\Delta_0^2}{\Delta_0^2 + (\hbar\omega - \Delta_0)^2} \frac{1 - \cos(\Omega t)}{2} \quad (9)$$

where $\Omega = \sqrt{\Delta_0^2 + (\hbar\omega - \Delta_0)^2}$, and the on-resonance Rabi frequency $\Omega_0 = F \sqrt{\sin^2(\theta_f - \theta_q)}$ (we take $\hbar = 1$) [24]. We therefore see that maximum oscillations with full gate conversion probability are obtained when the driving is resonant with the qubit energy splitting. We also see that the width of the Rabi peak in the frequency domain is given by Δ_0 . Simple Rabi oscillations can also be observed in a multi-level system if the driving frequency is on resonance with one of the relevant energy splittings but off resonance with all others.

C. The effect of decoherence

In an undriven system, the effect of decoherence is to push the density matrix describing the system towards its thermal-equilibrium value with time scales given by the characteristic dephasing and relaxation times. The effects of decoherence, especially dephasing, can be thought of in terms of a broadening of the energy levels. In particular, if the energy separation between the states $|\alpha_i\rangle$ and $|\beta_i\rangle$ of Sec. III-A is smaller than the typical decoherence rates in the problem, any effect related to that energy separation becomes unobservable. Alternatively, one could say that only processes that occur on

a time scale faster than the decoherence times can be observed.

It is worth taking a moment to look in some more detail at the problem of a resonantly-driven qubit coupled to a dissipative environment, which is usually studied under the name of Bloch-equations [24, 25]. If the Rabi frequency is much smaller than the decoherence rates, the qubit will remain in its thermal equilibrium state, since any deviations from that state caused by the driving field will be dissipated immediately. If, on the other hand, the Rabi frequency is much larger than the decoherence rates, the system will perform damped Rabi oscillations, and it will end up close to the maximally mixed state in which both states $|\alpha_i\rangle$ and $|\beta_i\rangle$ have equal occupation probability. In that case, one could say that decoherence succeeds in making us lose track of the quantum state of the qubit but fails to dissipate the energy of the qubit, since more energy will always be available from the driving field.

D. Combined picture

We now take the three elements presented above and combine them to obtain a simple intuitive picture of the problem at hand.

Let us for a moment neglect the effects of decoherence and only consider the case $\omega = \Delta_0$. The driving field tries to flip the state of the qubit alone. However, two of the relevant eigenstates are entangled states, namely $|\alpha_i\rangle$ and $|\beta_i\rangle$. One can therefore expect that if the width of the Rabi peak, or in other words the on-resonance Rabi frequency, is much larger than the energy separation between the states $|\alpha_i\rangle$ and $|\beta_i\rangle$, the qubit will start oscillating much faster than the TLS can respond, and the initial dynamics will look similar to that of the uncoupled system. Only after many oscillations and a time of the order of $(E_3 - E_2)^{-1}$ will one start to see the effects of the qubit-TLS interaction. If, on the other hand, the Rabi frequency is much smaller than the energy separation between the states $|\alpha_i\rangle$ and $|\beta_i\rangle$, the driving field can excite at most one of those two states, depending on the driving frequency. In that case the qubit-TLS interactions are strong enough that the TLS can follow adiabatically the time evolution of the qubit. In the intermediate region, one expects that if the driving frequency is closer to one of the two transition frequencies $E_2 - E_1$ and $E_3 - E_1$, beating behaviour will be seen right from the beginning. If one looks at the Rabi peak in the frequency domain, e.g. by plotting the maximum gate qubit-state conversion probability as a function of frequency, the single peak of the weak-coupling limit separates into two peaks as the qubit-TLS coupling strength becomes comparable to and exceeds the on-resonance Rabi frequency.

We do not expect weak to moderate levels of decoherence to cause any qualitative changes in the qubit dynamics other than, for example, imposing a decaying envelope on the qubit excitation probability. As mentioned above, features that are narrower (in frequency) than the




FIG. 2: Maximum qubit excitation probability $P_{f,max}^{(q)}$ between $t=0$ and $t=20 =_0$ for $\delta =_0 = 0.5$ (a), 2 (b), 2.5 (c) and 5 (d). $\varphi = \pi/4$, and $\tau_{TLS} = \pi/6$.




FIG. 3: Maximum qubit excitation probability $P_{f,max}^{(q)}$ between $t=0$ and $t=20 =_0$ for the four-level system (solid line) and the reduced three-level system (dashed line). $\delta =_0 = 2$, $\varphi = \pi/4$, and $\tau_{TLS} = \pi/5$.

By looking at Fig. 1, one might say that the $\delta =_0 = 0$ peak clearly corresponds to a two-photon process coupling states $|j\rangle_i$ and $|i\rangle_i$. In fact, for further demonstration that this is the case, we have included in Fig. 4 the probability of the combined qubit+TLS system to be in state $|i\rangle_i$. This peak is easiest to observe in the intermediate coupling regime. In the weak-coupling limit, the qubit and TLS are essentially decoupled, especially on the time scale of qubit dynamics. In the strong coupling limit, one can argue that a Raman transition will give rise to that peak. However, noting that the width of that peak is of the order of the smaller of the values $\delta_{0=ss}^2$ and $\delta_{0=cc}^2$, we can see that it becomes increas-

ingly narrow in that limit. In other words, the virtual intermediate state after the absorption of one photon is far enough in energy from the states $|j\rangle_i$ and $|i\rangle_i$ to make the peak invisibly narrow. It is rather surprising, however, that in the intermediate-coupling regime the peak reaches unit qubit conversion probability, even though the transitions to states $|j\rangle_i$ and $|i\rangle_i$ are real, rather than being virtual transitions whose role is merely to mediate the coupling between states $|j\rangle_i$ and $|i\rangle_i$. We have verified that the (almost) unit height of the peak is quite robust against changes in the angles φ and τ_{TLS} for a wide range in δ , even when that peak coincides with the top of one of the two main peaks. In fact, the Hamiltonian \hat{H}_e can be diagonalized rather straightforwardly in the case $\delta =_0 = 0$, and one can see that there is no symmetry that requires full conversion between the states $|j\rangle_i; N$ and $|i\rangle_i; N$. The lack of any special relations between the energy differences in the eigenvalues of \hat{H}_e , however, suggests that almost full conversion should be achieved in a reasonable amount of time.




FIG. 4: Qubit excitation probability $P_{f,t}^{(q)}$ as a function of time (solid line) for $\delta =_0 = 0$ (a), 0.92 (b), 1.61 (c) and 1.77 (d). The dashed line is the occupation probability of state $|i\rangle_i; N$. $\delta =_0 = 2$, $\varphi = \pi/4$, and $\tau_{TLS} = \pi/5$.

The asymmetry between the two main peaks in Figs. 2 and 3 can also be explained by the fact that in one of those peaks state $|i\rangle_i$ is also involved in the dynamics and it increases the quantity $P_{f,max}^{(q)}$. As above, we have included in Figs. 4(b) and 4(c) the probability of the combined qubit+TLS system to be in state $|i\rangle_i$.

In order to explain the dips in Figs. 2 and 3, we note that the plotted quantity, $P_{f,max}^{(q)}$, is the sum of four terms (in the reduced three-level system): a constant and three oscillating terms. The frequencies of those terms correspond to the energy differences in the diagonalized 3×3 Hamiltonian. The dips occur at frequencies where the two largest frequencies are integer multiples of the smallest

est one. Away from any such point, $P_{\uparrow, \text{max}}^{(q)}$ will reach a value equal to the sum of the amplitudes of the four terms. Exactly at those points, however, such a constructive buildup of amplitudes is not always possible, and a dip is generally obtained. The width of that dip decreases and vanishes asymptotically as we increase the simulation time, although the depth remains unaffected.

We also studied the case where the qubit and TLS energy splittings were different. As can be expected, the effects of the TLS decrease as it moves away from resonance with the qubit. That is most clearly reflected in the two-peak structure, where one of the two main peaks becomes substantially smaller than the other. The two-photon peak was still clearly observable in plots corresponding to the same quantity plotted in Fig. 2, i.e. plots of $P_{\uparrow, \text{max}}^{(q)}$ vs. $\delta\omega \neq 0$, even when the detuning between the qubit and the TLS was a few times larger than the coupling strength and the on-resonance Rabi frequency.




FIG. 5: Maximum qubit excitation probability $P_{\uparrow, \text{max}}^{(q)}$ between $t = 0$ and $t = 20 = 0$. The solid line corresponds to the case of no decoherence. The dotted ($T_{1/2}^{(q)} = 0.1$, $T_{1/2}^{(TLS)} = 0.2$), dashed ($T_{1/2}^{(q)} = 0$ and $T_{1/2}^{(TLS)} = 2.0$) and dash-dotted ($T_{1/2}^{(q)} = 0$ and $T_{1/2}^{(TLS)} = 0$) lines correspond to different decoherence regimes. $T_{1/2}^{(q)} = 2$, $T_{1/2}^{(TLS)} = 4$, and $T_{1/2}^{(TLS)} = 5$. Γ_1 and Γ_2 are the relaxation and dephasing rates of the subsystem, respectively.

The truncated dressed-state picture with four energy levels is insufficient to study the effects of decoherence. For example, relaxation from state $|2\rangle$ to $|1\rangle$ does not necessarily have to involve emission of a photon into the driving field mode. We therefore study the effects of decoherence by treating the driving field classically. We then solve a Bloch-Redfield master equation with a time-dependent Hamiltonian and externally-imposed dephasing and relaxation times, as was done in Ref. [20]. In Fig. 5 we reproduce the four-level results of Fig. 3, i.e. $P_{\uparrow, \text{max}}^{(q)}$ vs. $\delta\omega \neq 0$ with no decoherence, along with the same quantity obtained when we take into account the ef-

fects of decoherence. For a moderate level of decoherence, we see that the qubit excitation probability is somewhat reduced and all the features that are narrower than the decoherence rates are suppressed partially or completely by the effects of decoherence. For large qubit decoherence rates, the qubit excitation probability is greatly reduced close to resonance, where the Rabi frequency takes its lowest values. The shallow dip in the dash-dotted line in Fig. (5) occurs because for those frequencies and in the absence of decoherence the maximum amplitude is only reached after several oscillations, whereas it is reached during the first few oscillations outside that region. For large TLS decoherence rates, the TLS becomes weakly coupled to the qubit, and a single peak is recovered in the qubit dynamics (with a height larger than either the two split peaks). All of these effects are in agreement with the simple picture presented in Sec. III.

V. EXPERIMENTAL CONSIDERATIONS

In the early experiments on phase qubits coupled to TLSs [10, 15], the qubit relaxation rate $\Gamma_1^{(q)}$ (40M Hz) was comparable to the splitting between the two Rabi peaks Ω_{ss} (20-70M Hz), whereas the on-resonance Rabi frequency Ω_0 was tunable from 30M Hz to 400M Hz (note that, as discussed in Sec. III, the Rabi frequency cannot be reduced to values much lower than the decoherence rates, or Rabi oscillations would disappear altogether). The large relaxation rates in those experiments would make several effects discussed in this paper unobservable. The constraint that Ω_0 could not be reduced below 30M Hz made the strong-coupling regime, where $\Omega_0 \gg \Omega_{ss}$, inaccessible. The weak-coupling regime, where $\Omega_0 \ll \Omega_{ss}$, was easily accessible in those experiments. However, as can be seen from Fig. 2, it shows only a minor signature of the TLS. Although the intermediate-coupling regime was also accessible, as evidenced by the observation of the splitting of the Rabi peak into two peaks, observation of the two-photon process and the additional dips of Fig. 2 discussed above would have required a time at least comparable to the qubit relaxation time. That would have made them difficult to distinguish from experimental fluctuations.

With the new qubit design of Ref. [26], the qubit relaxation time has been increased by a factor of 20. The constraint that Ω_0 must be at least comparable to $\Omega_{1/2}^{(q)}$ no longer prevents accessibility of the strong-coupling regime. Furthermore, since our simulations were run for a period of time corresponding to approximately ten Rabi oscillation cycles, i.e. shorter than the relaxation time observed in that experiment, all the effects that were discussed above should be observable, including the observation of the two-photon peak and the transition from the weak- to the strong-coupling regimes by varying the driving amplitude.

We finally consider one possible application of our results to experiments on phase qubits, namely the problem

of characterizing the environment composed of TLSs. As we shall show shortly, characterizing the TLS parameters and the nature of the qubit-TLS coupling are not independent questions. The energy splitting of a given TLS, E_{TLS} , can be obtained easily from the location of the qubit-TLS resonance as the qubit energy splitting is varied. One can then obtain the distribution of values of E_{TLS} for a large number of TLSs, as was in fact done in Ref. [26]. The splitting of the Rabi resonance peak into two peaks by itself, however, is insufficient to determine the values of Γ_{TLS} and Γ_{TLS} separately. By observing the location of the two-photon peak, in addition to the locations of the two main peaks, one would be able to determine both Γ_{cc} and Γ_{ss} for a given TLS, as can be seen from Fig. 1. Those values can then be used to calculate both E_{TLS} and $\Gamma_{\text{TLS}} = \arctan(\Gamma_{\text{ss}}/\Gamma_{\text{cc}})$ of that TLS. The distribution of values of Γ_{TLS} can then be used to test models of the environment, such as the one given in Ref. [17] to describe the results of Ref. [27].

In order to reach the above conclusion, we have made the assumption that the distribution of values of Γ_{TLS} for those TLSs with sufficiently strong coupling to the qubit is representative of all TLSs. Since it is generally believed that strong coupling is a result of proximity to the junction, the above assumption is quite plausible, as long as the other TLSs share the same nature. Although it is possible that there might be two different types of TLSs of different nature in a qubit's environment, identifying that possibility would also be helpful in understanding the nature of the environment. We have also assumed that q does not take the special value $q=2$ (note that, based on the arguments of Refs. [17, 27], we are also assuming that generally $\Gamma_{\text{TLS}} \neq 2$). That assumption would not raise any concern when dealing with charge or flux qubits, where both q_c and q_f can be adjusted in a single experiment, provided an appropriate design is used. However, the situation is trickier with phase qubits. The results in that case depend on the nature of the qubit-TLS coupling, which we discuss next.

The two mechanisms that are currently considered the most likely candidates to describe the qubit-TLS coupling are through either (1) a dependence of the Josephson junction's critical current on the TLS state or (2) Coulomb interactions between a charged TLS and the charge across the junction. In the former case, one has an effective value of q that is different from $q=2$ [28], and the assumption of an intermediate value of q is justified. In the case of coupling through Coulomb interactions, on the other hand, one effectively has $q=2$, and therefore Γ_{cc} vanishes for all the TLSs. In that case the two-

photon peak would always appear at the midpoint (to a good approximation) between the two main Rabi peaks. Although that would prevent the determination of the distribution of values of Γ_{TLS} , it would be a strong indication that Coulomb interactions with the charge across the junction are responsible for the qubit-TLS coupling rather than the critical current dependence on the TLS state. Note also that if it turns out that this is in fact the case, and the distribution of values of Γ_{TLS} cannot be extracted from the experimental results, that distribution might be irrelevant to the question of decoherence in phase qubits.

VI. CONCLUSION

We have studied the problem of a harmonically-driven qubit that is interacting with an uncontrollable two-level system and a background environment. We have presented a simple picture to understand the majority of the phenomena that are observed in this system. That picture is composed of three elements: (1) the four-level energy spectrum of the qubit+TLS system, (2) the basic properties of the Rabi-oscillation dynamics and (3) the basic effects of decoherence. We have confirmed the predictions of that picture using a systematic numerical analysis where we have varied a number of relevant parameters. We have also found unexpected features in the resonance-peak structure. We have analyzed the behaviour of the system and provided simple explanations in those cases as well. Our results can be tested with available experimental systems. Furthermore, they can be used in experimental attempts to characterize the TLSs surrounding a qubit, which can then be used as part of possible techniques to eliminate the TLSs' detrimental effects on the qubit operation.

Acknowledgments

This work was supported in part by the Army Research Office (ARO), Laboratory of Physical Sciences (LPS), National Security Agency (NSA) and Advanced Research and Development Activity (ARDA) under Air Force Office of Research (AFOSR) contract number F49620-02-1-0334; and also supported by the National Science Foundation grant No. EIA-0130383. One of us (S.A.) was supported by a fellowship from the Japan Society for the Promotion of Science (JSPS).

[1] There are now several reviews on the subject. See e.g. J. Q. You and F. N. Nori, *Phys. Today* 58 (11), 42 (2005); Y. Makhlin, G. Schon, and A. Shnirman, *Rev. Mod. Phys.* 73, 357 (2001).

[2] Y. Nakamura, Yu. A. Pashkin, and J. S. Tsai, *Nature*

398, 786 (1999).

[3] Y. Nakamura, Yu. A. Pashkin, and J. S. Tsai, *Phys. Rev. Lett.* 87, 246601 (2001).

[4] D. Vion, A. Aassime, A. Cottet, P. Joyez, H. Pothier, C. Urbina, D. Esteve, and M. H. Devoret, *Science* 296, 886

- (2002).
- [5] Y. Yu, S. Han, X. Chu, S.-I. Chu, and Z. Wang, *Science* 296, 889 (2002).
- [6] J. M. Martinis, S. Nam, J. Aumentado, and C. Urbina, *Phys. Rev. Lett.* 89, 117901 (2002).
- [7] I. Chiorescu, Y. Nakamura, C. J. P. Hammond, and J. E. Mooij, *Science* 299, 1869 (2003).
- [8] Yu. A. Pashkin, T. Yamamoto, O. Astaev, Y. Nakamura, D. V. Averin, J. S. Tsai, *Nature* 421, 823 (2003).
- [9] T. Yamamoto, Yu. A. Pashkin, O. Astaev, Y. Nakamura, J. S. Tsai, *Nature* 425, 941 (2003).
- [10] R. W. Simmonds, K. M. Lang, D. A. Hite, D. P. Pappas, and J. M. Martinis, *Phys. Rev. Lett.* 93, 077003 (2004).
- [11] J. Q. You and F. Nori, *Phys. Rev. B* 68, 064509 (2003).
- [12] I. Chiorescu, P. Bertet, K. Semba, Y. Nakamura, C. J. P. M. Hammond, and J. E. Mooij, *Nature* 431, 159 (2004).
- [13] A. Wallraff, D. I. Schuster, A. Blais, L. Frunzio, R. S. Huang, J. Majer, S. Kumar, S. M. Girvin, and R. J. Schoelkopf, *Nature* 431, 162 (2004).
- [14] J. Johansson, S. Saito, T. Mieno, H. Nakano, M. Ueda, K. Semba, and H. Takayanagi, *Phys. Rev. Lett.* 96, 127006 (2006).
- [15] K. B. Cooper, M. Steffen, R. McDermott, R. W. Simmonds, S. Oh, D. A. Hite, D. P. Pappas, and J. M. Martinis, *Phys. Rev. Lett.* 93, 180401 (2004).
- [16] L.-C. Ku and C. C. Yu, *Phys. Rev. B* 72, 024526 (2005).
- [17] A. Shnirman, G. Schon, I. Martin, and Y. Makhlin, *Phys. Rev. Lett.* 94, 127002 (2005).
- [18] L. Faoro, J. Bergli, B. L. Altshuler, and Y. M. Galperin, *Phys. Rev. Lett.* 95, 046805 (2005).
- [19] Y. M. Galperin, D. V. Shantsev, J. Bergli, and B. L. Altshuler, *Europhys. Lett.* 71, 21 (2005).
- [20] S. Ashhab, J. R. Johansson, and F. Nori, *cond-mat/0512677*.
- [21] We shall make a number of assumptions that might seem too specific at certain points. We have been careful, however, not to make a choice of parameters that causes any of the interesting physical phenomena to disappear. For a more detailed discussion of our assumptions, see [20].
- [22] G. Baym, *Lectures on Quantum Mechanics*, (Addison Wesley, New York, 1990).
- [23] See, e.g., C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, *Atom-Photon Interactions*, (John Wiley & Sons, New York, 1992).
- [24] For a detailed discussion of the problem, see, e.g., C. P. Slichter, *Principles of magnetic resonance*, (Springer, New York, 1996).
- [25] For studies of certain aspects of this problem in superconducting qubits, see: A. Yu. Smirnov, *Phys. Rev. B* 67, 155104 (2003); N. Kosugi, S. Matsuo, K. Konno, and N. Hatkenaka, *Phys. Rev. B* 72, 172509 (2005).
- [26] J. M. Martinis, K. B. Cooper, R. McDermott, M. Steffen, M. Ansmann, K. D. Osborn, K. Cicak, S. Oh, D. P. Pappas, R. W. Simmonds, and C. C. Yu, *Phys. Rev. Lett.* 95, 210503 (2005).
- [27] O. Astaev, Yu. A. Pashkin, Y. Nakamura, T. Yamamoto, and J. S. Tsai, *Phys. Rev. Lett.* 93, 267007 (2004).
- [28] Further arguments regarding the value of q are given in A. M. Zagoskin, S. Ashhab, J. R. Johansson, and F. Nori, *cond-mat/0603753*.