Interference as a Probe of Spin Incoherence in Strongly Interacting Quantum Wires

M. Kindermann,¹ P. W. Brouwer,¹ and A. J. Millis²

¹ Laboratory of Atomic and Solid State Physics, Cornell University, Ithaca, New York 14853

² Department of Physics, Columbia University, 538 West 120th Street, New York, New York 10027

(Dated: March 2006)

We show that interference experiments can be used to identify the spin-incoherent regime of strongly interacting one-dimensional conductors. Two qualitative signatures of spin-incoherence are found: a strong magnetic field dependence of the interference contrast and an anomalous scaling of the interference contrast with the applied voltage, with a temperature and magnetic field dependent scaling exponent. The experiments distinguish the spin-incoherent from the spin-polarized regime, and so may be useful in deciding between alternative explanations proposed for the anomalous conductance quantization observed in quantum point contacts and quantum wires at low density.

PACS numbers: 73.63.Nm,71.10.Pm,71.27.+a

One dimensional conductors are of intense current interest because they generically display strong "Luttinger liquid" interaction effects, such as scaling with nontrivial powerlaws [1]. Over the last decade high quality quantum wires have been fabricated and shown to display characteristic Luttinger liquid properties [2, 3]. A particularly interesting limit is the "spin-incoherent" regime [4, 5, 6, 7, 8, 9, 10, 11] in which the energy scale J of spin excitations is much less than the temperature T. Spinincoherent behavior is a generic feature of the low density limit of an interacting one-dimensional electron gas, but it may also be realized in ultra-thin conductors at high electron density [12].

An important motivation for study of the spinincoherent regime is provided by the discovery of anomalous conductance quantization in quantum wires and quantum point contacts [13, 14, 15, 16, 17, 18]. As the electron density is reduced in these systems, the conductance exhibits a series of plateaus at integer multiples of $2e^2/h$ [19, 20]. At the lowest densities, however, an additional plateau is observed, at a reduced conductance value between $0.5 \times 2e^2/h$ and $0.7 \times 2e^2/h$. While conductance quantization at multiples of $2e^2/h$ can be well understood within the framework of non-interacting electrons [19, 20] electron-electron interactions are believed to be responsible for the reduced conductance at the lowest plateau.

Two explanations of the reduced conductance have been proposed: a spin-polarization of conduction electrons [21, 22, 23] and the formation of a Wigner crystal with small exchange energy J at low electron density [24, 25]. Matveev showed that the spin-incoherent regime at $kT \gg J$ was characterized by a conductance plateau with a reduced conductance, consistent with experiments. Despite their different physical content spin-polarization versus spin-incoherence — both models are consistent with a wide range of experimental observations, both of conductance and of current fluctuations [11, 26]. While a spontaneous spin-polarization can be experimentally detected in magnetic focusing ex-



FIG. 1: First interference geometry. In spin-polarized conductors amplitudes for an electron to tunnel from the left to the right wire at the points x and x' interfere. Their phase difference $\varphi = e\Phi/\hbar c$ depends on the flux Φ through the interference loop. In the spin-incoherent regime they generically involve different spin configurations that prevent interference.

periments [27], the technique cannot easily be applied to longer wires such as carbon nanotubes [15, 16, 18], where the Lieb-Schulz-Mattis theorem suggests that a ferromagnetic state is unlikely. A direct probe of the existence of the spin-incoherent regime is therefore needed.

In this Letter we show that quantum interference experiments provide a powerful probe of the presence and properties of spin-incoherence in one dimensional conductors. We consider the standard interference geometry sketched in Fig 1: a loop enclosing a magnetic flux Φ . We shall be interested in the interference contrast C: the relative change of current I with the phase $\varphi = e\Phi/\hbar c$

$$C = \frac{\sqrt{\langle (I(\varphi) - \langle I \rangle_{\varphi})^2 \rangle_{\varphi}}}{\langle I \rangle_{\varphi}}, \quad \langle \dots \rangle_{\varphi} = \int_0^{2\pi} \frac{d\varphi}{2\pi} \dots \quad (1)$$

Typically the interference contrast decays exponentially with the linear dimension l of the interference region provided that l is greater than a phase coherence length l_{φ} . In strongly interacting one dimensional systems electrons decompose into charge and spin excitations with generically different velocities v and v_{σ} . The coherence length $l_{\varphi}^{\sigma} \sim v_{\sigma}/T$ for spin excitations may become much smaller than that of charge excitations, $l_{\varphi}^{\rho} \sim v/T$. In the

 $\mathbf{2}$

limit $l_{\varphi}^{\rho} \gg l \gg l_{\varphi}^{\sigma}$, where charge excitations can move coherently through the interference region but spin excitations cannot, the interference contrast is thus strongly suppressed by interactions in the system. This is particularly evident in the spin-incoherent regime of impenetrable electrons $(l_{\varphi}^{\sigma} \lesssim a)$, where a is of the order of the inter-electron spacing). The spin state of the conductor is then static and tunneling events at different points will in general result in different spin orderings, as depicted in Fig. 1. Interference will only be possible if all N electrons in the interference loop have the same spin state. This occurs with the probability 2^{-N} and it becomes very unlikely at large N. A magnetic field with a Zeeman energy $E_Z \gg kT$ large enough to polarize the electron spins restores the interference. This loss of coherence at arbitrarily low temperatures contrasts with the decay of persistent currents in isolated rings in the same regime that occurs only at finite temperatures $kT \gtrsim v/lN$ [28].

More formally, an interference experiment in the geometry of Fig. 1 probes the amplitude for an electron to propagate from point x to x' via different wires, and thus probes the exponential decay of the single-electron Green function with distance predicted for the spin-incoherent regime [4, 6]. This decay can alternatively be observed in momentum resolved tunneling [3, 7]. It is not a feature of the spin-incoherent regime $l_{\varphi}^{\sigma} \lesssim a$ alone, but occurs whenever temperatures are large enough that $l_{\omega}^{\sigma} < l$ [29]. We show that in an alternative geometry (shown in Fig. 2) the scaling of interference effects with voltage may be used to distinguish the true spin-incoherent regime $l_{\varphi}^{\sigma} \lesssim a$ from the regime $l_{\varphi}^{\sigma} \lesssim l$. Because of space limitations we present explicit calculations only in the spin-incoherent limit $l_{\varphi}^{\sigma} \lesssim a$, but we will compare with the other limit $a \lesssim l_{\varphi}^{\sigma} \lesssim l$ at appropriate points.

We consider two interference geometries: one geometry where electrons tunnel between semi-infinite wires and one where they tunnel between effectively infinite wires, as shown in Figs. 1 and 2. Following Refs. [6, 7] we model a spin-incoherent wire by a spinless Luttinger liquid and a static spin background,

$$H = H_1 + H_2 + H_T,$$

$$H_{\alpha} = v \int_0^{\infty} \frac{dx}{2\pi} \left[g^{-1} (\partial_x \theta_{\alpha})^2 + g (\partial_x \phi_{\alpha})^2 \right],$$

$$H_T = e^{ieV\tau} \sum_{\sigma} \left[t \, \psi_{1\sigma}^{\dagger}(x_1) \psi_{2\sigma}(x_2) + e^{i\varphi} t' \psi_{1\sigma}^{\dagger}(x'_1) \psi_{2\sigma}(x'_2) + \text{h.c.} \right]. \quad (2)$$

Here, $\alpha = 1, 2, V$ is the bias voltage, τ the time argument and t and t' are the tunneling amplitudes. We have chosen units such that $\hbar = 1$. The boson fields obey the standard commutation relation $[\theta_{\alpha}(x), \phi_{\alpha}(x')] = -i\pi \Theta(x - x')$. The velocity v is a property of the spinless charge carriers c_{α} . The electron fields $\psi_{\alpha\sigma}$ are expressed in terms of these fermions c_{α} and operators $S_{\alpha\sigma}(x)$ that add a spin σ to the spin background of wire α at position x as $\psi_{\alpha\sigma}(x) = c^{\dagger}_{\alpha}(x)S_{\alpha\sigma}(x)$ with

$$c_{\alpha}(x) = \frac{\eta_{\alpha}}{\sqrt{2\pi a}} \sum_{n=\pm 1} e^{in[\theta_{\alpha}(x)+k_F x]} e^{i\phi_{\alpha}(x)}$$
(3)

in bosonized form. Here, η_{α} are Majorana fermions and k_F is the Fermi wavevector of the fermions c_{α} . The Hamiltonian (2) can be derived microscopically as the low-energy theory of a Hubbard model with infinite onsite repulsion U and an additional long-range interaction described by a purely forward scattering density-density coupling. An infinite-U Hubbard model is described in terms of spinless holes that become the fermions c_{α} introduced above and a static spin background [30].

The total tunnel current $I = I_{\rm dir} + I_{\rm int}$ is the sum of a φ -independent contribution $I_{\rm dir}$ and a φ -dependent interference contribution $I_{\rm int}$. The two contributions are expressed in terms of the electron Green functions as

$$I_{\rm dir} = 2|t|^2 \sum_{\sigma} \int d\tau \, e^{-ieV\tau} \left[G_{1\sigma}^{<}(x,x,\tau) G_{2\sigma}^{>}(x,x,-\tau) - G_{1\sigma}^{>}(x,x,\tau) G_{2\sigma}^{<}(x,x,-\tau) \right] + (t \to t', \, x \to x'), \tag{4}$$

$$f_{\rm int} = 2t^* t' e^{i\varphi} \sum_{\sigma} \int d\tau \, e^{-ieV\tau} \left[G_{1\sigma}^<(x, x', \tau) G_{2\sigma}^>(x', x, -\tau) - G_{1\sigma}^>(x, x', \tau) G_{2\sigma}^<(x', x, -\tau) \right] + \text{c.c.}, \tag{5}$$

$$G^{>}_{\alpha\sigma}(x,x',\tau) = -i\langle\psi_{\alpha\sigma}(x,\tau)\psi^{\dagger}_{\alpha\sigma}(x',0)\rangle$$
(6)
$$= -i\langle S_{\alpha\sigma}(x,\tau)c^{\dagger}_{\alpha}(x,\tau)c_{\alpha}(x',0)S^{\dagger}_{\alpha\sigma}(x',0)\rangle,$$

1

and $G^{<}$ is defined correspondingly. The spin expectation value in Eq. (6) is non-vanishing only if adding a spin σ at position x' and time 0 and removing a spin of the same orientation at position x and time τ does not alter the spin background. Since the spin background in our model of impenetrable electrons follows moving charges, this occurs with probability $p_{\sigma}^{|N_{x\alpha}(\tau)-N_{x'\alpha}(0)|}$, where $N_{x\alpha}$ is the number of electrons to the left of point x and

$$p_{\uparrow} = 1 - p_{\downarrow} = \frac{1}{1 + \exp(-E_Z/kT)}$$
 (7)

is the probability for a spin to point along the direction of the applied magnetic field. Consequently the spin and the charge expectation values in Eq. (6) do not factorize as in the case of spin-coherent Luttinger liquids. This underlies most of the phenomena that we discuss in this Letter. We thus obtain [6, 7]

$$G^{>}_{\alpha\sigma}(x,x',\tau) = -i\sum_{k} p^{|k|}_{\sigma} \frac{d\xi}{2\pi} e^{i\xi k}$$

$$\times \langle e^{-i\xi N_{x\alpha}(\tau)} c^{\dagger}_{\alpha}(x,\tau) c_{\alpha}(x',0) e^{i\xi N_{x'\alpha}(0)} \rangle.$$
(8)

 $G^{<}$ is calculated similarly. After bosonization one has $N_{x\alpha}(\tau) = [k_F x + \theta_{\alpha}(x,\tau)]/\pi$. In this representation, the discrete nature of $N_{x\alpha}$ is lost, so that our results will only be approximate. In keeping with the continuous nature of the bosonized theory, we replace the summation over k in Eq. (8) by an integral. With this replacement, our calculation using bosonization approaches the exact results in the spin-polarized limit $p_{\sigma} \to 1$, when the integration over k enforces $\xi = 0$ in Eq. (8).

In order to prevent the interference current I_{int} to be smeared out by the applied bias voltage, interference experiments need to be done with a bias voltage $eV \ll v/l$. Therefore, all results presented in this Letter are for the limit $v\tau \gg l$. For definiteness, we also assume $p_{\uparrow} \ge p_{\downarrow}$.

We first analyze the geometry of Fig. 1, where tunneling occurs between the bulk of one wire and the end of the other wire. We evaluate the Green functions Eq. (8) in semi-infinite wires following Ref. [31]. We give the Green functions of the source wire, $\alpha = 1$; Green functions for $\alpha = 2$ are obtained by interchanging the coordinates xand x'. For temperatures $kT \ll \tau^{-1}$ we find

$$G_{1\sigma}^{>}(x, x', \tau) \sim \int dk \, \frac{l^{1/4g} \sin(k_F \delta x) \sin(\pi k + k_F \delta x)}{w (iv\tau + a)^{1/g}} \\ \times p_{\sigma}^{|k-1/4|} e^{-(k - \langle N_l \rangle)^2 / w^2}, \qquad (9)$$

where $\delta x \ll a$ is the distance of x' to the end of the wire, $l \gg a$ is the distance between the tunneling points x and x' measured along the wire, $\langle N_l \rangle = k_F l/\pi$, and we abbreviated $w = [g \ln(2l/a)]^{1/2}/\pi$. Equation (9) and all other expressions for Green functions below are up to a numerical proportionality factor that depends on the high-energy cutoff a. For $\langle N_l \rangle \gg \max\{w, w^2 | \ln p_\sigma | \}$, a condition that is fulfilled for repulsive interactions if $\langle N_l \rangle \gg 1$, |k| in Eq. (9) can be replaced by k. We then perform the integration over k and find

$$G^{>}_{1\sigma}(x,x',\tau) \sim p_{\sigma}^{\langle N_l \rangle - 1/4} \frac{\langle N_l \rangle^{1/4g - g/4 + g \ln^2 p_{\sigma}/4\pi^2}}{(iv\tau + a)^{1/g}} (10)$$
$$\times \sin(k_F \delta x) \sin(k_F l + \pi w^2 \ln p_{\sigma}/2).$$

To evaluate $I_{\rm dir}$ we need two more Green functions,

$$G_{1\sigma}^{>}(x,x,\tau) \sim \int dk \, \frac{l^{1/2g} p_{\sigma}^{|k|} \cos \pi k \, e^{-k^2/2w^2}}{w(iv\tau+a)^{1/g}}, (11)$$
$$G_{1\sigma}^{>}(x',x',\tau) \sim \frac{\sin^2(k_F \delta x)}{(iv\tau+a)^{1/g}}.$$
(12)

Upon substitution into Eqs. (4) and (5), we conclude that at $kT \ll eV I_{\rm dir}$ and $I_{\rm int}$ have the same voltage dependence, $I \propto (eV)^{2/g-1}$. The interference current $I_{\rm int}$ is, however, reduced relative to $I_{\rm dir}$ through the incoherence of the electron spins. The interference contrast Cdepends on the average number $N = 2\langle N_l \rangle$ of electrons inside the interference loop,

$$C \sim p_{\uparrow}^N N^{-g/2+g\ln^2 p_{\uparrow}/2\pi^2},\tag{13}$$

where we assumed $(p_{\downarrow}/p_{\uparrow})^N \ll 1$. For temperatures $kT \gg eV$, one has $I_{\rm dir} \sim I_{\rm int} \sim eV(kT)^{2/g-2}$ while the interference contrast is still given by Eq. (13).

The first factor in Eq. (13) describes the exponential dependence of the interference contrast on N that has been anticipated by the argument in the introduction $(p_{\uparrow} = p_{\downarrow} = 1/2)$ in the absence of a magnetic field). It leads to an exponential suppression of C at $E_Z \leq kT \ln N$ (for large N). This remarkable magnetic field dependence that is absent in the spin-polarized case can be distinguished from the effect of magnetic impurities by its strong dependence on the electron density in the wire via N (tunable by a gate voltage). In the regime of $a \leq l_{\varphi}^{\sigma} \leq l$ an exponential suppression of C similar to that of Eq. (13) occurs. Experimentally it can be distinguished from the spin-incoherent regime $(kT \gg J)$ by the field scale required to polarize the electron gas and thus to restore interference which is raised to $E_Z \approx J$.

The second factor in Eq. (13) adds a power law dependence of C on N that is due to quantum fluctuations of N. A similar scaling with the distance from a boundary has been found in a different context in Ref. [32], but the scaling exponent found here depends on the applied magnetic field and the temperature. While this anomalous scaling is masked in the geometry of Fig. 1 by the much stronger exponential dependence p^N_{\uparrow} , it becomes observable in the geometry of Fig. 2, which we now discuss.

In the geometry shown in the inset of Fig. 2, tunneling takes place between the bulk regions of both wires, with the distance between the tunneling points x, x' and the ends of the wires being much larger than v/eV. Calculating the Green functions as before, we find

$$G_{1\sigma}^{>}(x,x,\tau) = G_{1\sigma}^{>}(x',x',\tau)$$
(14)
 $\sim \int dk \frac{p_{\sigma}^{|k|} \cos \pi k \, e^{-k^2/2w'^2}}{w'(iv\tau+a)^{1/2g}},$

where $w' = [g \ln(iv\tau/a)]^{1/2}/\pi$, and

$$G_{1\sigma}^{>}(x,x',\tau) \sim p_{\sigma}^{\langle N_l \rangle} \frac{\cos(k_F l + \pi w'^2 \ln p_{\sigma})}{(iv\tau + a)^{g/2 + 1/2g - g \ln^2 p_{\sigma}/2\pi^2}}.$$
 (15)

In the derivation of Eqs. (14) and (15) we again assumed $\langle N_l \rangle \gg \max[|w'|, |w'|^2| \ln p_{\sigma}|]$. Equations (14) and (15) reduce to the Green functions obtained in Ref. [6] in the limit of zero magnetic field, $p_{\sigma} = 1/2$. For



FIG. 2: In our second geometry (inset) the interference contrast C obeys a powerlaw with a temperature and magnetic field dependent exponent $g - g \ln^2 p_{\uparrow}/\pi^2$. We show the normalized interference contrast $c = C(V)/C(V_0)$ compared to its value without magnetic field $c_{B=0}$, for various p_{\uparrow} (g = 1).

g = 1, when describing an infinite-U Hubbard model, they moreover coincide with the Green functions found in Ref. [4] without bosonization and the resulting loss of the discreteness of charge. For not too large magnetic fields, $E_Z \ll gkT \ln(v\tau/2a)$, the k-integral in Eq. (14) is to a good approximation τ -independent. The resulting voltage dependence of the tunneling current is different than for the geometry of Fig. 1: we find $I_{\rm dir} \propto (eV)^{1/g-1}$, while the bias dependence of $I_{\rm int}$ exhibits the anomalous scaling observed in the length dependence of Eq. (13),

$$I_{\rm int} \sim p_{\uparrow}^N (eV)^{g+1/g-g\ln^2 p_{\uparrow}/\pi^2 - 1},$$
 (16)

where, again, $p_{\uparrow} \geq 1/2$. The scaling of I_{int} with eV is temperature and magnetic field dependent through p_{\uparrow} , Eq. (7). This is a defining signature of spin-incoherence in quantum wires and in particular serves to distinguish the truly spin-incoherent limit $kT \gg J$ from the regime $a \leq l_{\varphi}^{\sigma} \leq l$. The interference contrast in this geometry acquires a voltage dependence

$$C \sim p_{\uparrow}^{N} (eV)^{g-g \ln^2 p_{\uparrow}/\pi^2}.$$
 (17)

Also such scaling of the interference contrast with the applied voltage is absent in spin-polarized conductors. The interference contrast C crosses over to the spin-polarized behavior at large magnetic fields $E_Z \gtrsim gkT \ln(v\tau/2a)$ through the τ -dependence of the k-integral in Eq. (14).

Our calculation has been done using equilibrium Green functions in (semi-)infinite wires. It applies to wires of finite length L if $eV \gg v/L$. The equilibrium assumption is justified for spin relaxation times $\tau_{\rm s}$ in the wire that are shorter than the transit time, $\tau_{\rm s} \ll k_F Le/I$.

In conclusion, we have proposed interference experiments for distinguishing the spin-polarized from the spinincoherent regime of strongly interacting wires. We identified two unique signatures of spin-incoherence: A strong dependence of the interference contrast on an applied magnetic field and a power law dependence of the interference contrast on the applied voltage, if the tunneling takes place between bulk regions of the quantum wires. The scaling exponent of the interference current is surprisingly temperature and magnetic field dependent, another feature that is unknown from spinpolarized Luttinger liquids. These clear and qualitative signatures of spin-incoherence make interference experiments promising tools in the search for the mechanism of the observed conductance anomalies in interacting onedimensional conductors.

This work was supported by the NSF under grants no. DMR 0431350 (AM) and DMR 0334499 and by the Packard Foundation (PB and MK).

- [1] T. Giamarchi, ed., *Quantum Physics in One Dimension* (Oxford University Press, 2002).
- [2] Z. Yao, H. Postma, L. Balents, and C. Dekker, Nature 402, 273 (1999).
- [3] O. Auslaender, et al., Science **295**, 825 (2002).
- [4] V. V. Cheianov and M. B. Zvonarev, Phys. Rev. Lett. 92, 176401 (2004).
- [5] V. V. Cheianov and M. B. Zvonarev, J. Phys. A: Math. Gen. 37, 2261 (2004).
- [6] G. A. Fiete and L. Balents, Phys. Rev. Lett. 93, 226401 (2004).
- [7] G. A. Fiete, K. Le Hur, and L. Balents, Phys. Rev. B 72, 125416 (2005).
- [8] G. A. Fiete, J. Qian, Y. Tserkovnyak, and B. I. Halperin, Phys. Rev. B 72, 045315 (2005).
- [9] G. A. Fiete, K. Le Hur, and L. Balents, condmat/0511715 (2005).
- [10] V. V. Cheianov, H. Smith, and M. B. Zvonarev, Phys. Rev. A 71, 033610 (2005).
- [11] M. Kindermann and P. W. Brouwer, cond-mat/0506455 (2005).
- [12] M. M. Fogler, Phys. Rev. B 71, 161304(R) (2005).
- [13] K. J. Thomas, et al., Phys. Rev. Lett. 77, 135 (1996).
- [14] S. M. Cronenwett, et al., Phys. Rev. Lett. 88, 226805 (1988).
- [15] B. E. Kane, et al., Appl. Phys. Lett. 72, 3506 (1998).
- [16] D. J. Reilly, et al., Phys. Rev. B 63, 121311(R) (2001).
- [17] K. J. Thomas, et al., Phys. Rev. B 61, 13365(R) (2000).
- [18] M. J. Biercuk, et al., Phys. Rev. Lett. 94, 026801 (2005).
- [19] B. J. van Wees, et al., Phys. Rev. Lett. 60, 848 (1988).
- [20] D. A. Wharam, et al., J. Phys. C **21** (1988).
- [21] D. J. Reilly, et al., Phys. Rev. Lett. 89, 246801 (2002).
- [22] Y. Meir, K. Hirose, and N. S. Wingreen, Phys. Rev. Lett. 89, 196802 (2002).
- [23] A. D. Klironomos, J. S. Meyer, and K. A. Matveev, condmat/0507387 (2005).
- [24] K. A. Matveev, Phys. Rev. Lett. 92, 106801 (2004).
- [25] K. A. Matveev, Phys. Rev. B **70**, 245319 (2004).
- [26] P. Roche, et al., Phys. Rev. Lett. 93, 116602 (2004).
- [27] L. Rokhinson, L. Pfeiffer, and K. West, condmat/0509448 (2005).
- [28] S. Viefers, P. Koskinen, P. S. Deo, and M. Manninen, Physica E 21, 1 (2004).
- [29] K. Le Hur, Phys. Rev. Lett. **95**, 076801 (2005).
- [30] J. Bernasconi, M. J. Rice, W. R. Schneider, and S. Strässler, Phys. Rev. B 12, 1090 (1975).
- [31] M. Fabrizio and A. O. Gogolin, Phys. Rev. B 51, 17827 (1995).
- [32] S. Eggert and I. Affleck, Phys. Rev. Lett. 75, 934 (1995).