

# Comment on "Connection between the Burgers equation with an elastic forcing term and a stochastic process"

Piotr Garbaczewski

Institute of Physics, University of Zielona Gora, 65-516 Zielona Gora, Poland

In the above mentioned paper by E. Moreau and O. Vallee [Phys. Rev. E 73, 016112, (2006)], the one-dimensional Burgers equation with an elastic (attractive) forcing term has been claimed to be connected with the Ornstein-Uhlenbeck process. We point out that this connection is valid only in case of the repulsive forcing.

PACS numbers: 02.50.Ey, 05.90.+m, 05.45.-a

Let us consider the Langevin equation for the one-dimensional stochastic process in the external conservative force field  $F(x) = -dV(x)/dx$ :

$$\frac{dx}{dt} = F(x) + \sqrt{2} b(t) \quad (1)$$

where  $b(t)$  stands for the normalized white noise:  $b(t)i = 0$ ,  $b(t^0)b(t^1)i = \delta(t - t^0)$ . The corresponding Fokker-Planck equation for the probability density  $p(x;t)$  reads:

$$\partial_t p = \partial_{xx} p + \partial_x (F p) \quad (2)$$

and by means of a standard substitution  $u(x;t) = p(x;t) \exp[2V(x)]$ , [1], can be transformed into the generalized diffusion equation for an auxiliary function  $u(x;t)$ :

$$\partial_t u = \partial_{xx} u + V(x) u \quad (3)$$

where

$$V(x) = \frac{1}{2} \frac{F^2}{F} + \partial_x F \quad (4)$$

As discussed in detail in Refs. [3, 4, 5], given the so-called forward drift  $b(x;t)$  of the Markovian diffusion process, in the above identified with  $b(x;t) = F(x)$ , one readily infers the so-called backward drift of this process

$$b_-(x;t) = b(x;t) - 2 \partial_x (\ln p)(x) \quad (5)$$

which is known to solve the forced Burgers equation:

$$\partial_t b_- + b_- \partial_x b_- - \partial_{xx} b_- = F \quad (6)$$

where  $F = -2 \partial_x V$ .

For the Ornstein-Uhlenbeck process  $b(x) = F(x) = -x$  and accordingly

$$V(x) = \frac{x^2}{4} - \frac{x}{2} \quad (7)$$

Substituting the inferred  $V(x)$  into Eq. (3), we get Eq. (32) of Ref. [2]. We observe that the velocity field  $u(x;t)$ , defined by Eq. (35) of Ref. [2], does coincide with our  $b_-(x;t)$ , provided we set  $b(x;t) = -x$  in Eq. (5).

The related  $V(x)$  gives rise to

$$F(x) = 2 \partial_x V(x) = -x \quad (8)$$

while an original elastic forcing problem addressed by [2], Eq. (2) therein, has the form:

$$\partial_t u + u \partial_x u - \partial_{xx} u = -x^2 \quad (9)$$

and clearly differs from Eq. (6), with the necessarily arising  $-x^2$  on its right-hand-side, by an innocent-looking but crucial in the present context sign of the forcing term.

Let us add that this particular sign issue has received due attention in Ref. [6]. A specific class of diffusion-type processes has been considered that would account for standard Newtonian accelerations (of the form  $\partial_x W$  with  $W(x)$  bounded from below) on the right-hand-side of Eq. (6). It is in principle possible at the price of introducing an additional pressure-type forcing term.

Acknowledgement: This note has been supported by the Polish Ministry of Scientific Research and Information Technology under the (solicited) grant No PBZ-M IN-008/P 03/2003.

- [1] H. Risken, The Fokker-Planck equation, (Springer-Verlag, Berlin, 1989)  
[2] E. Moreau and O. Vallee, Phys. Rev. E 73, 016112, (2006)  
[3] P. Garbaczewski and G. Kondrat, Phys. Rev. Lett. 77, 2608, (1996)

- [4] P. Garbaczewski, G. Kondrat and R. Olikiewicz, Phys. Rev. E 55, 1401, (1997)  
[5] P. Muratore Ginanneschi, Physica D 115, 341, (1998)  
[6] P. Garbaczewski, Phys. Rev. E 57, 569, (1998)