

Orbital Functionals in Current-Spin-Density-Functional Theory

N. Helbig,^{1,2,3} S. Kurth,¹ S. Pittalis,¹ E. Rasanen,¹ and E. K. U. Gross^{1,2}

¹Institut für Theoretische Physik, Freie Universität Berlin, Am Müllendamm 14, D-14195 Berlin, Germany

²Fritz-Haber-Institut der Max-Planck-Gesellschaft, Faradayweg 4-6, D-14195 Berlin, Germany

³Unité de Physico-Chimie et de Physique des Matériaux, Université Catholique de Louvain, B-1348 Louvain-la-Neuve, Belgium

The description of interacting many-electron systems in external magnetic fields is presented in the framework of non-collinear current-spin-density-functional theory for exchange-correlation energy functionals which are explicit functionals of two-component Kohn-Sham spinors. The integral equations for the exchange-correlation components of the scalar and vector potentials and the magnetic field are derived in the formalism of the optimized effective potential method and a simplifying approximation to these equations is suggested. Using the exact-exchange energy functional, the formalism is applied to study two-dimensional quantum dots in magnetic fields. An excellent agreement with quantum Monte Carlo results is obtained, and the role of the exchange vector potential is found to be minor.

PACS numbers: 71.15.Mb, 71.15.-m

Since its introduction in 1964, density functional theory (DFT) [1, 2] has become a standard tool to calculate the electronic structure of atoms, molecules, and solids from first principles. Early on, the original DFT formulation has been extended to the case of spin-polarized systems [3] which also provides a description of many-electron systems in an external magnetic field. However, in this spin-DFT (SDFT) framework the magnetic field only couples to the spin but not to the orbital degrees of freedom, i.e., the coupling of the electronic momenta to the vector potential associated with the external magnetic field is not taken into account. A proper treatment of this coupling requires extension to current-spin-density-functional theory (CSDFT) [4, 5] in terms of three basic variables: the electron density $n(\mathbf{r})$, the spin magnetization density $\mathbf{m}(\mathbf{r})$, and the paramagnetic current density $\mathbf{j}_p(\mathbf{r})$. These densities are conjugate variables to the electrostatic potential, the magnetic field, and the vector potential, respectively.

In order to be applicable in practice, DFT of any flavor requires an approximation to the exchange-correlation (xc) energy functional. The use of the local-vorticity approximation [4, 5], which is an extension of the local-spin-density approximation (LSDA), in CSDFT is problematic: the xc energy per particle of a uniform electron gas exhibits derivative discontinuities whenever a new Landau level is occupied in an increasing external magnetic field. This leads to discontinuities in the corresponding xc potential [6]. These discontinuities then incorrectly appear when the local values of the inhomogeneous density and vorticity coincide with the corresponding values of the homogeneous electron gas. A popular way to circumvent this problem is to use functionals which interpolate between the limits of weak and high magnetic fields [7, 8].

In this work, we show that explicitly orbital-dependent functionals, which are successfully used in DFT and

collinear SDFT [9], are natural candidates to approximate the xc energy in CSDFT for two reasons: first, they are constructed without recourse to the model of the uniform electron gas and second, they are ideally suited to describe orbital effects such as the filling of Landau levels. In this way, the problem inherent in any uniform-gas-derived functional for CSDFT is avoided in a natural way.

The use of orbital functionals requires the so-called optimized effective potential (OEP) method [10] to calculate the effective potentials. Here, we extend the OEP formalism which recently has been generalized to non-collinear SDFT [11] to the CSDFT framework and obtain three coupled integral equations for the xc contribution to the electrostatic and vector potentials as well as to the magnetic field. Generalizing the approach of Krieger, Li and Iafrate [12, 13], we introduce a simplifying approximation which transforms the OEP integral equations into a set of algebraic equations. An extension of DFT, similar to the one given here, is given in [14], where a larger set of basic densities [15] is considered to include spin-orbit coupling. In another work [16] our formalism has been used to study open-shell atoms at zero external magnetic field. Here, we demonstrate the applicability of our formalism by calculating the electronic structure of a quantum dot (QD) [17] in an external magnetic field. We find excellent agreement with quantum Monte Carlo (QMC) results.

The Kohn-Sham (KS) equation in CSDFT reads (atomic units are used throughout)

$$\left[-\frac{1}{2} \nabla^2 + v_s(\mathbf{r}) + v_H(\mathbf{r}) + v_{xc}(\mathbf{r}) + \mathbf{B}_s(\mathbf{r}) \cdot \nabla \right] \psi_{\mathbf{k}} = \epsilon_{\mathbf{k}} \psi_{\mathbf{k}} \quad (1)$$

The KS potentials are $v_s(\mathbf{r}) = v_0(\mathbf{r}) + v_H(\mathbf{r}) + v_{xc}(\mathbf{r}) + 1/2c^2 [A_0^2(\mathbf{r}) - A_s^2(\mathbf{r})]$, $B_s(\mathbf{r}) = B_0(\mathbf{r}) + B_{xc}(\mathbf{r})$, and $A_s(\mathbf{r}) = A_0(\mathbf{r}) + A_{xc}(\mathbf{r})$, where the xc potentials are

functional derivatives of the xc energy E_{xc} with respect to the corresponding densities

$$v_{xc}(\mathbf{r}) = \frac{E_{xc}[\mathbf{n}; \mathbf{m}; \mathbf{j}_p]}{n(\mathbf{r})}; \quad (2)$$

$$B_{xc}(\mathbf{r}) = \frac{E_{xc}[\mathbf{n}; \mathbf{m}; \mathbf{j}_p]}{m(\mathbf{r})}; \quad (3)$$

$$\frac{1}{c} A_{xc}(\mathbf{r}) = \frac{E_{xc}[\mathbf{n}; \mathbf{m}; \mathbf{j}_p]}{\mathbf{j}_p(\mathbf{r})}; \quad (4)$$

The self-consistency cycle is closed by calculating the density

$$n(\mathbf{r}) = \sum_{k=1}^{\mathcal{X}^{cc}} \psi_k^y(\mathbf{r}) \psi_k^y(\mathbf{r}); \quad (5)$$

the magnetization density

$$m(\mathbf{r}) = \sum_{k=1}^{\mathcal{X}^{cc}} \psi_k^y(\mathbf{r}) \psi_k^y(\mathbf{r}); \quad (6)$$

and the paramagnetic current density

$$\mathbf{j}_p(\mathbf{r}) = \frac{1}{2i} \sum_{k=1}^{\mathcal{X}^{cc}} \psi_k^y(\mathbf{r}) \nabla \psi_k^y(\mathbf{r}) - (\nabla \psi_k^y(\mathbf{r})) \psi_k^y(\mathbf{r}); \quad (7)$$

The ground state total energy of the interacting system can then be computed from

$$\begin{aligned} E[\mathbf{n}; \mathbf{m}; \mathbf{j}_p] &= T_s[\mathbf{n}; \mathbf{m}; \mathbf{j}_p] + U[\mathbf{n}] + E_{xc}[\mathbf{n}; \mathbf{m}; \mathbf{j}_p] \\ &+ \int d^3r n(\mathbf{r}) v_0(\mathbf{r}) + \int d^3r m(\mathbf{r}) B_0(\mathbf{r}) \\ &+ \frac{1}{c} \int d^3r \mathbf{j}_p(\mathbf{r}) A_0(\mathbf{r}) + \frac{1}{2c^2} \int d^3r n(\mathbf{r}) A_0^2(\mathbf{r}); \quad (8) \end{aligned}$$

where T_s and U are the kinetic energy of the KS system and the Hartree energy, respectively.

Gauge invariance of the energy functional implies that E_{xc} depends on the current only through the vorticity $\mathbf{j}_p(\mathbf{r}) = \nabla \psi_p(\mathbf{r}) \times \nabla \psi_p(\mathbf{r})$, i.e., $E_{xc}[\mathbf{n}; \mathbf{j}_p; \mathbf{m}] = E_{xc}[\mathbf{n}; \mathbf{j}_p; \mathbf{m}]$ [5]. This immediately leads to the following relation for the xc vector potential

$$\nabla \cdot (\mathbf{n}(\mathbf{r}) A_{xc}(\mathbf{r})) = 0; \quad (9)$$

If one has an approximate E_{xc} which is given explicitly in terms of the densities, the calculation of the corresponding xc potentials via Eqs. (2)–(4) is straightforward. Here, however, we deal with approximations to the xc energy which are explicit functionals of the KS spinor orbitals ψ_k . These functionals are, via the Hohenberg-Kohn theorem, implicit functionals of the densities. In the spirit of the original OEP formalism, we derive the

corresponding OEP integral equations for the xc potentials by requiring that the effective fields optimize the value of the ground-state total energy (8). Therefore, the functional derivatives of the total energy with respect to the three KS potentials vanish. This procedure leads to three OEP equations which are most conveniently written as [16]

$$\sum_{k=1}^{\mathcal{X}^{cc}} \psi_k^y(\mathbf{r}) \psi_k^y(\mathbf{r}) + h.c. = 0; \quad (10)$$

$$\sum_{k=1}^{\mathcal{X}^{cc}} \psi_k^y(\mathbf{r}) \psi_k^y(\mathbf{r}) + h.c. = 0; \quad (11)$$

and

$$\frac{1}{2i} \sum_{k=1}^{\mathcal{X}^{cc}} \psi_k^y(\mathbf{r}) \nabla \psi_k^y(\mathbf{r}) - (\nabla \psi_k^y(\mathbf{r})) \psi_k^y(\mathbf{r}) + h.c. = 0; \quad (12)$$

where we have defined the so-called "orbital shifts" [9, 18]

$$\psi_k^y(\mathbf{r}) = \sum_{\substack{j=1 \\ j \neq k}}^{\mathcal{X}} \frac{D_{kj}^y}{\psi_j^y(\mathbf{r})}; \quad (13)$$

and

$$\begin{aligned} D_{kj}^y &= \int d^3r^0 v_{xc}(\mathbf{r}^0) \psi_j^y(\mathbf{r}^0) \psi_k^y(\mathbf{r}^0) \\ &+ \frac{1}{2ic} A_{xc}(\mathbf{r}^0) \psi_j^y(\mathbf{r}^0) \psi_k^y(\mathbf{r}^0) - \psi_j^y(\mathbf{r}^0) \nabla \psi_k^y(\mathbf{r}^0) \\ &+ \sum_{B} B_{xc}(\mathbf{r}^0) \psi_j^y(\mathbf{r}^0) \psi_k^y(\mathbf{r}^0) - \psi_j^y(\mathbf{r}^0) \frac{E_{xc}}{\psi_k^y(\mathbf{r}^0)}; \quad (14) \end{aligned}$$

The orbital shifts ψ_k^y have the structure of a first-order shift from the unperturbed orbital ψ_k under a perturbation whose matrix elements are given by D_{kj}^y . Physically, the OEP equations (10)–(12) then imply that the densities do not change under this perturbation. If A_{xc} is set to zero, Eqs. (10) and (11) reduce exactly to the OEP equations for non-collinear SDFT [11].

Eqs. (10)–(12) are a set of coupled integral equations for the three unknown xc potentials. In standard (S)DFT, the OEP equations can be solved by a direct computation of the orbital shifts [18]. A similar technique has been used to solve the OEP equations in non-collinear SDFT [11]. Instead of taking this route, we employ the Krieger-Li-Iafrate (KLI) approach as a simplifying approximation [12, 13]. In SDFT it often yields potentials which are very close to the full OEP ones. The KLI approximation can be derived by replacing the energy denominator $\psi_k^y \psi_j^y$ in the definition (13) of ψ_k^y by a constant $\psi_k^y \psi_j^y$. This leads to the KLI equations for the xc potentials which are written in matrix form as

$$D(\mathbf{r}) V_{xc}(\mathbf{r}) = R(\mathbf{r}); \quad (15)$$

Here, we have defined the seven-component vector $V_{xc}(r)$ as

$$V_{xc}^T(r) = v_{xc}(r); B_{xc}^T(r); \frac{1}{c} A_{xc}^T(r) \quad (16)$$

and the 7×7 matrix $D(r)$ has the structure

$$D(r) = \begin{pmatrix} 0 & & & & & & \\ & n(r) & m^T(r) & \frac{1}{c} j_p^T(r) & & & \\ & m(r) & \frac{2}{c} n(r) \mathbb{I} & J^T(r) A & & & \\ & \frac{1}{c} j_p(r) & J^T(r) & N(r) & & & \end{pmatrix};$$

where \mathbb{I} is the 3×3 unit matrix. The matrix elements of the 3×3 matrices J and N are defined by

$$J_k(r) = \frac{B_{xc}}{2i} \sum_{k=1}^{X^{cc}} \frac{y_k(r)}{r} \left(\frac{\partial}{\partial r} \frac{y_k(r)}{r} - \frac{\partial}{\partial r} \frac{y_k(r)}{r} \right) \quad (17)$$

and

$$N_k(r) = \frac{1}{2} \sum_{k=1}^{X^{cc}} \left(\frac{\partial}{\partial r} \frac{y_k(r)}{r} \frac{\partial}{\partial r} \frac{y_k(r)}{r} + \frac{\partial}{\partial r} \frac{y_k(r)}{r} \frac{\partial}{\partial r} \frac{y_k(r)}{r} \right) + \frac{1}{4n(r)} \left(\frac{\partial n(r)}{\partial r} \right)^2; \quad (18)$$

where $k = 1; 2; 3$ corresponds to the Cartesian coordinates x, y , and z , respectively. The seven components of the vector $R(r)$ on the rhs of Eq. (15) are given by

$$R_1(r) = \frac{1}{2} \sum_{k=1}^{X^{cc}} \frac{y_k(r)}{r} \left(\frac{E_{xc}}{y_k(r)} + n_k(r) D_{kk}^y + h.c. \right); \quad (19)$$

$$R_{1+}(r) = \frac{1}{2} \sum_{k=1}^{X^{cc}} \frac{y_k(r)}{r} \left(\frac{E_{xc}}{y_k(r)} + m_k(r) D_{kk}^y + h.c. \right); \quad (20)$$

$$R_{4+}(r) = \frac{1}{2} \sum_{k=1}^{X^{cc}} \frac{1}{2i} \frac{y_k(r)}{r} \left(\frac{\partial}{\partial r} \frac{E_{xc}}{y_k(r)} + \frac{1}{2i} \frac{\partial}{\partial r} \frac{E_{xc}}{y_k(r)} + \frac{1}{c} \frac{\partial}{\partial r} \frac{E_{xc}}{y_k(r)} + j_{pk}(r) D_{kk}^y + h.c. \right); \quad (21)$$

with the density $n_k(r)$, magnetization density $m_k(r)$ and paramagnetic current density $j_{pk}(r)$ of the single orbital k . In order to arrive at this form of the KLI equations we made use of Eq. (9).

To demonstrate the applicability of our formalism, we use the KLI equations to solve the electronic structure of a quasi-two-dimensional semiconductor QD [17] in an external magnetic field. The QD is described as a many-electron system restricted to the xy plane and confined in that plane by an external parabolic potential $v_0 =$

$\frac{1}{2} m \omega_0^2 r^2$ with $r^2 = x^2 + y^2$. The external magnetic field is defined to be uniform and perpendicular to the xy plane, i.e., $B_0(r) = r A_0(r) = B_0 e_z$ with $A_0(r) = B_0 e = 2$. We apply the effective mass approximation with the material parameters for GaAs, i.e., the effective mass $m = 0.067$, the dielectric constant, $\epsilon = 12.4$, and the effective gyromagnetic ratio $g = 0.44$.

In order to simplify the implementation we assume collinearity, i.e., both the KS magnetic field B_s and the magnetization m only have non-vanishing z -components. Then the Pauli-type KS equation becomes diagonal in spin space and can be decoupled into two separate equations for the spin-up and spin-down orbitals $\psi_k(r)$. We further assume that the xc potentials preserve the cylindrical symmetry of the problem, i.e., $v_{xc}(r) = v_{xc}(r) = v_{xc}(r) + B g B_{xc}(r)$, where upper signs are for spin-up and lower signs for spin-down electrons, and $A_{xc}(r) = A_{xc}(r) e$. Due to the cylindrical symmetry we can separate the wave function into radial and angular parts as $\psi_{jl}(r) = \exp(i l \phi) R_{jl}(r)$, where the radial wave functions $R_{jl}(r)$ are eigenfunctions of the Hamiltonian

$$\hat{H}_{s1} = \frac{1}{2m} \left(\frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} - \frac{l^2}{r^2} \right) + \frac{1}{2} \omega_0^2 r^2 + m \frac{r^2}{2} + \frac{1}{m c} \frac{A_{xc}(r)}{r} B m g B_0 + v_H(r) + v_{xc}(r) \quad (22)$$

with the total confinement $\omega_0^2 = \frac{P}{2} \left(\frac{1}{a_0^2} + \frac{1}{c^2} \right)$, and the cyclotron frequency $\omega_c = B_0 / m c$. The radial wave functions are expanded in the basis of eigenfunctions of the corresponding non-interacting problem, i.e., the eigenfunctions of the Hamiltonian (22) with the Hartree and all xc potentials set to zero. In the following we have employed the exact-exchange (EXX) approximation to E_{xc} , i.e.,

$$E_x^{EXX} = \frac{1}{2} \sum_{f, g} \sum_{k, l} \int d^3 r \int d^3 r' \frac{r_{k1}(r) r_{l1}(r')}{|r - r'|} \quad (23)$$

For the given symmetry, the 7×7 KLI matrix equation (15) reduces to a 3×3 matrix equation for the two scalar exchange potentials v_x and the angular component of the exchange vector potential A_x . It can be shown that the KLI equations automatically lead to a vanishing exchange vector potential (and therefore CSDFT reduces to the well-known SDFT result) if the system is in a state with vanishing current density [16].

Figure 1 shows the total energy of a six-electron QD ($\omega_0 = 5$ meV) as a function of B_0 . The kinks correspond to changes in the ground-state configuration ($L_z; S_z$). Apart from the fully-polarized ($S_z = 3$) states, the EXX energies are considerably too large when compared with the accurate QMC results [8]. EXX also




FIG. 1: Magnetic field dependence of the total ground-state energy ($E_{\text{tot}} - 6Q = 6\sqrt{l_0^2 + m_0^2} = 4$) in the six-electron quantum dot calculated with the exact-exchange (EXX), EXX plus LSDA correlation (EXX+cLSDA), and with EXX plus the corrected LSDA correlation (EXX+c[#]LSDA). The LSDA and quantum Monte Carlo (QMC) results [8] are shown for comparison.

leads to an erroneous occurrence of the (5;2) ground-state at $B_0 = 1.5 \dots 2.0$ T. However, adding the LSDA correlation (cLSDA) [19] post-hoc to the EXX energies yields the correct sequence of states as a function of B_0 . This is a major improvement over the cLSDA-corrected Hartree-Fock calculation which does not give the correct ground states for a similar system [20]. As seen in Fig. 1, the energies of EXX+cLSDA are consistently too low by $1.0 \dots 1.5$ meV. Hence, considering the good agreement between QMC and the conventional LSDA, our results demonstrate the inherent tendency of the LSDA to cancel out its respective errors in exchange and correlation. However, we are able to remove most of the error in cLSDA by applying a corrected functional where the like-spin-correlation has been removed (EXX+c[#]LSDA) as suggested by Becke [21].

Figure 2 shows the paramagnetic current j_p and the diamagnetic current $j_d(r) = n(r)A_0(r) - m$ at $B_0 = 11$ T for the (21;3) state. The total current $j = j_p + j_d$ changes sign at $r \approx 350$ a.u. due to the existence of a single vortex at the center of the QD that we find in agreement with the LSDA and numerically exact calculations [22]. In Fig. 2 we also show the exchange vector potential A_x . Its maximum, located near the edge of the QD, reaches a value which is 10% of that of the external vector potential A_0 . Despite the considerable relative magnitude of A_x , its effect on the physical quantities like the total energy, density, and current density is negligible. A similar result has been observed for open-shell atoms [16] and also for QDs on the level of LSDA [8].

In summary, we have extended the optimized effective potential method to current-spin-density functional theory allowing the use of orbital functionals in a current-density functional treatment. A major advantage of




FIG. 2: Angular components of the paramagnetic and diamagnetic currents, j_p and j_d , and their sum for the (21;3) state at $B_0 = 11$ T. The dashed line shows the exchange vector potential for the same configuration.

these functionals, as we have demonstrated for the exact-exchange functional, is that the filling of Landau levels is taken into account in a natural way and causes no numerical difficulties. We emphasize that the same approximation for E_{xc} is used over the whole range of external magnetic fields without any interpolation between high- and low-field approximations. Application of our approach to quantum dots yields excellent agreement of the ground-state energies, densities, and currents with the previously published data. In particular, adding a corrected LSDA correlation to the EXX energies in a post-hoc way leads to more accurate energies than those of conventional LSDA. Further development of the correlation functionals and their self-consistent implementation are left for future research.

The authors like to thank H. Saarikoski and A. Harju for providing us with their data published in Fig. 3 of Ref. [8]. We gratefully acknowledge financial support through the Deutsche Forschungsgemeinschaft and by the EU's Sixth Framework Program through the Nanoquanta Network of Excellence (NMP4-CT-2004-500198).

-
- [1] P. Hohenberg and W. Kohn, Phys. Rev. 136, B864 (1964).
 - [2] W. Kohn and L. Sham, Phys. Rev. 140, A1133 (1965).
 - [3] U. von Barth and L. Hedin, J. Phys. C 5, 1629 (1972).
 - [4] G. Vignale and M. Rasolt, Phys. Rev. Lett. 59, 2360 (1987).
 - [5] G. Vignale and M. Rasolt, Phys. Rev. B 37, 10685 (1988).
 - [6] P. Skudlarski and G. Vignale, Phys. Rev. B 48, 8547 (1993).
 - [7] A. Wensauer and U. Rossler, Phys. Rev. B 69, 155302 (2004).
 - [8] H. Saarikoski et al, Phys. Rev. B 67, 205327 (2003).
 - [9] T. Grabo, T. Kreibich, S. Kurth, and E. K. J. Gross, in Strong Coulomb Correlations in Electronic Structure:

- Beyond the Local Density Approximation, edited by V. Anisimov (Gordon & Breach, Tokyo, 2000), pp. 203-311.
- [10] J.D. Talman and W.F. Shadwick, *Phys. Rev. A* **14**, 36 (1976).
- [11] S. Sharma et al., cond-mat/0510800 (2005).
- [12] R.T. Sharp and G.K. Horton, *Phys. Rev.* **90**, 317 (1953).
- [13] J.B. Krieger, Y. Li, and G.J. Iafrate, *Phys. Rev. A* **46**, 5453 (1992).
- [14] S. Rohra and A. Gorling, *Phys. Rev. Lett.* **97**, 013005 (2006).
- [15] K. Bencheikh, *J. Phys. A: Math. Gen.* **36**, 11929 (2003).
- [16] S. Pittalis, S. Kurth, N. Helbig, and E.K.J. Gross, *Phys. Rev. A* **74**, 062511 (2006).
- [17] S.M. Reimann, M. Manninen, *Rev. Mod. Phys.* **74**, 1283 (2002).
- [18] S. Kummel and J.P. Perdew, *Phys. Rev. Lett.* **90**, 043004 (2003); *Phys. Rev. B* **68**, 035103 (2003).
- [19] C. Attaccalite, S. Moroni, P. Gori-giorgi, and G. Bachelet, *Phys. Rev. Lett.* **88**, 256601 (2002).
- [20] W. Xin-Qiang, *J. Phys.: Condens. Matter* **12**, 4207 (2000).
- [21] A. Becke, *Int. J. Quant. Chem.* **27**, 585 (1985).
- [22] H. Saarikoski, A. Harju, M.J. Puska, and R.M. Nieminen, *Phys. Rev. Lett.* **93**, 116802 (2004).