

## Bosonic representation of spin operators in the field-induced critical phase of spin-1 Haldane chains

Masahiro Sato

Synchrotron Radiation Research Center, Japan Atomic Energy Agency, Sayo,  
Hyogo 679-5148, Japan and CREST JST, Japan

**Abstract.** A bosonized-field-theory representation of spin operators in the uniform-field-induced Tomonaga-Luttinger-liquid (TLL) phase in spin-1 Haldane chains is formulated by means of the non-Abelian (Tsvetlik's Majorana fermion theory) and the Abelian bosonizations and Furusaki-Zhang technique [Phys. Rev. B **60**, 1175 (1999)]. It contains massive magnon fields as well as massless boson fields. From it, asymptotic forms of spin correlation functions in the TLL phase are completely determined. Applying the formula, we further discuss effects of perturbations (bond alternation, single-ion anisotropy terms, staggered fields, etc) for the TLL state, and the string order parameter. Throughout the paper, we often consider in some detail how the symmetry operations of the Haldane chains are translated in the low-energy effective theory.

PACS numbers: 75.10.Jm, 75.10.Pq, 75.50.Ee

## 1. Introduction

For (quasi) one-dimensional (1D) magnets, sophisticated theoretical tools have been developed in the last decades; for example, bosonizations [1, 2, 3], conformal field theory (CFT) [4], Bethe ansatz [5, 6], and several numerical methods (exact diagonalization, density-matrix renormalization group (DMRG), quantum Monte Carlo method (QMC), etc). Thanks to them, we have obtained a deep understanding of 1D spin systems. Of course, they have succeeded in explaining a number of experimental results of quasi 1D magnets. In this paper, we try to further develop a low-energy field theory description for spin-1 antiferromagnetic (AF) chains in a uniform magnetic field, which are a fundamental model in 1D spin systems.

Before discussing spin-1 AF chains, for comparison, let us briefly review the well-established low-energy effective theory (so-called Abelian bosonization) for the spin-1/2 XXZ chain [7, 8], which is one of the well-investigated and realistic models in spin systems. The Hamiltonian is defined by

$$\hat{H}_{\text{xxz}} = J \sum_j \frac{1}{2} (S_j^+ S_{j+1} + \text{h.c.}) + \sum_j S_j^z S_{j+1}^z - H \sum_j S_j^z; \quad (1)$$

where  $S_j$  is spin-1/2 operator on site  $j$ ,  $J > 0$  is the exchange interaction,  $\sum_j S_j^z S_{j+1}^z$  is the anisotropy parameter ( $\sum_j S_j^z S_{j+1}^z = 1$  case is the SU(2) Heisenberg chain), and  $H = 0$  is a uniform field. In a wide parameter regime including the zero-field line ( $1 < \sum_j S_j^z S_{j+1}^z < 2$  and  $H = 0$ ) [3, 8], the low-energy physics is governed by a one-component Tomonaga-Luttinger liquid (TLL) phase, where the spin correlation functions are of an algebraic decay type. For this critical phase, a beautiful field theory description has been established as follows. The low-energy effective Hamiltonian is a massless Gaussian model,

$$\hat{H}_{\text{xxz}}^e = \int dx \frac{v}{2} K (\partial_x \phi)^2 + \frac{1}{K} (\partial_x \theta)^2; \quad (2)$$

where  $x = j a_0$  ( $a_0$  is the lattice constant),  $\phi(x)$  is the scalar bosonic field,  $\theta(x)$  is the dual of  $\phi$ ,  $v$  is equal to the massless spinon velocity of the chain (1), and  $K$  is the TLL parameter ( $K = 1$  and  $K = 1/2$  respectively correspond to the XY (free fermion) point  $\sum_j S_j^z S_{j+1}^z = 0$  and the SU(2) Heisenberg one  $\sum_j S_j^z S_{j+1}^z = 1$ ) [9]. In this effective theory framework, the spin operator  $S_j$  is approximated [3, 7, 8, 10] as

$$\begin{aligned} S_j^z = a_0 & \left[ \frac{M}{a_0} + \frac{1}{2} \partial_x \phi + (-1)^j A_1^z \sin \frac{P}{4} + 2M \frac{x}{a_0} + \dots \right]; \\ S_j^+ = a_0 & \left[ e^{i\phi} - (-1)^j A_0 + A_1 \sin \frac{P}{4} + 2M \frac{x}{a_0} + \dots \right]; \end{aligned} \quad (3)$$

where  $M$  is the magnetization per site  $\langle S_j^z \rangle$ , and  $A_n$  and  $A_n^z$  are nonuniversal real constants. The magnetization  $M$ , the velocity  $v$ , and the TLL parameter  $K$  can be determined as a function of the parameters ( $J; \sum_j S_j^z S_{j+1}^z; H$ ), via the Bethe ansatz integral equations for the XXZ chain. If these three parameters are given, one can correctly know the asymptotic behavior of any spin correlation functions through the formula (3). Since the accurate values of  $A_n$  and  $A_n^z$  have recently been estimated [11, 10], it is also possible to obtain the amplitudes of the spin correlation functions. This effective theory, especially the bosonized spin formula (3), is very useful in analyzing not only the XXZ chain itself, but also several perturbation effects

(bond alternations, next-nearest-neighboring interactions, anisotropy term  $s$ , etc) for the XXZ chain and coupled spin-1/2 chain systems.

Now, we turn to spin-1 AF chains and their effective theories. In this paper, we mainly consider the following simple spin-1 AF Heisenberg chain,

$$\hat{H}_{S=1} = J \sum_j S_j \cdot S_{j+1} - H \sum_j S_j^z; \quad (4)$$

where  $S_j$  is "spin-1" operator on site  $j$ . In contrast to the spin-1/2 case, the spin-1 AF chain (4) without a uniform field ( $H = 0$ ) possesses a finite excitation gap (call Haldane gap) on its disordered ground state, which is described by the famous valence-bond-solid picture [12]. The low-lying excitation consists of a massive spin-1 magnon triplet around wave number  $p = \pm a_0$ . The spin correlation functions exhibit an exponential decay. This drastic difference between the spin-1/2 and spin-1 cases was first predicted by Haldane in 1983 [13]: he developed the nonlinear sigma model (NLSM) approach [14, 15, 7] for spin systems in order to explain the above low-energy physics of the spin-1 chain. After Haldane's conjecture, based on the non-Abelian bosonization approach to 1D spin systems [16, 17, 18], Tsvetlik proposed a Majorana (real) fermion theory [19, 1, 2, 20], which also has an ability to demonstrate the low-energy properties of the spin-1 AF chains. Both theories are now a standard field theory method of studying 1D spin-1 AF systems [21], like the Abelian bosonization for spin-1/2 chains.

When a uniform field  $H$  is applied in the spin-1 SU(2) AF chain (4), the low-lying magnon bands are split due to the Zeeman coupling. Provided that the bottom of one band crosses the energy level of the disordered ground state, a magnon condensed state and a finite magnetization occur [22, 23, 24, 25]. It is well known that this quantum phase transition is of a commensurate-incommensurate (C-IC) type [26, 3], and the resultant phase can be regarded as a TLL. Therefore, it is expected that, just like the formula (3), one can obtain a bosonized-spin expression in this TLL phase. However, as far as we know, any clear bosonization formulas of spin-1 operators in this magnon-condensed phase have not been established yet. Namely, such a formula has not been argued enough so far [27].

The main purpose of this paper is to derive a satisfactory bosonization formula for spin-1 operators in the above TLL phase. To this end, we start with Tsvetlik's Majorana fermion theory. This theory makes it possible to treat the Zeeman term in a nonperturbative way. Moreover, it can carefully deal with the uniform component of spins (the part around wave number  $p = 0$ ) as well as the staggered ones, in contrast to the NLSM approach. (We explain these properties of the fermion theory in later sections). To make the new bosonization formula, the bosonized-spin expression for the uniform-field-induced TLL phase in a two-leg spin-1/2 AF ladder [28] and its derivation, which was performed by Furusaki and Zhang [29], is very instructive and helpful. We will rely on these results. We will often discuss the symmetry correspondences between the original spin-1 AF chain (4) and the effective field theory. Like the formula (3), our resulting formula would be powerful in studying various (quasi) 1D spin-1 AF systems with magnons condensed. Recently, magnon-condensed states in quasi 1D gapped magnets, including spin-1 compounds such as NDMAP [30, 31, 32, 33, 34, 35], NDMAZ [36], TMNIN [37] and NTENP [38, 39], have been intensively investigated by some experiment groups. Our new formula hence is expected to be effective for understanding results of such experiments.

The organization of the rest of the paper is as follows. In Sec. 2, we review

Tsvetlik's Majorana fermion theory for the spin-1 AF chain without external fields. Section 3 is devoted to discussing the effective field theory for the uniform-field-induced TLL in spin-1 AF chains, which is based on the Majorana fermion theory. These two sections are needed as the preparation to construct a field-theory formula for spin operators in the TLL phase, although several parts of them have been already discussed in literature [1, 2, 19, 20, 40]. In Sec. 4, which is the main content of this paper, employing the contents of Secs. 2 and 3, we derive the formula of spin operators. Moreover, using it, we estimate the asymptotic forms of spin correlation functions in the TLL phase. Section 5 provides easy applications of our field-theory formula of spin operators. We consider the nonlocal string order parameter [41, 40], effects of some perturbation terms (bond alternation, a few anisotropy terms, etc) for the TLL state, and the magnon-decay process caused by anisotropic perturbations. We summarize the results of this paper in Sec. 6. Appendices may prove useful when reading the main text.

## 2. Spin-1 AF chain without external fields

In this section, we review the non-Abelian bosonization (Tsvetlik's Majorana fermion theory) method for the spin-1 Heisenberg AF chain with  $H = 0$ , especially focusing on the symmetries of the chain.

### 2.1. WZNW model and Majorana fermion theory

First, we define the spin-1 bilinear-biquadratic chain,

$$\hat{H} = J \sum_j S_j \cdot S_{j+1} + (\sum_j S_j \cdot S_{j+1})^2 : \quad (5)$$

Several studies [42] reveal that the system belongs to the "Haldane phase" in the regime  $J_2 < J_1$ , and the points  $J_2 = J_1$  correspond to a quantum criticality. In the Haldane phase, the ground state is disordered (namely, all symmetries are conserved) and the lowest excitations have a finite Haldane gap on it. It is shown [16, 43, 44] that the low-energy physics of the critical point  $J_2 = J_1$  is captured by an  $SU(2)$  level-2 Wess-Zumino-Novikov-Witten (WZNW) model [1, 2], a model of CFT with central charge  $c = 3=2$ , plus irrelevant perturbations preserving the spin  $SU(2)$  symmetry. This WZNW model possesses a Majorana (real) fermion description, i.e., the WZNW model with  $c = 3=2$  is equivalent to three copies of massless real fermions, each of which is regarded as a critical Ising system (a minimal model of CFT) with  $c = 1=2$  [17, 28, 1, 2]. Using the fermion (or Ising) picture, one can represent the Hamiltonian of the WZNW<sub>2</sub> model as follows:

$$\hat{H}_{\text{wznw}} = \int dx \frac{i}{2} v_0 \sum_{q=1,2,3} \left( \psi_{L,q}^\dagger \partial_x \psi_{L,q} - \psi_{R,q}^\dagger \partial_x \psi_{R,q} \right) ; \quad (6)$$

where  $\psi_{L(R),q}^\dagger(x) = \psi_{L(R),q}^\dagger(y)$  is the left (right) moving field of the  $q$ th real fermion with scaling dimension  $\Delta_s = 1=2$ , and  $v_0$  is the "light" velocity of the fermions. We here normalize the fermions  $\psi_{L(R),q}^\dagger$  through the anticommutation relation  $\{\psi_{L(R),q}^\dagger(x), \psi_{L(R),q}^\dagger(y)\} = \delta(x-y)$ , where the delta function stands for  $\int dx \delta(x-y) = 1$  [is a constant of  $O(a_0)$ ]. At the WZNW point  $J_2 = J_1$ , the spin operator is approximated as the following sum of the uniform and the staggered components [16, 19, 1, 2],

$$S_j^q = a_0 \left( J^q(x) + (-1)^j C_0 N^q(x); \quad (q = 1; 2; 3 \text{ or } x; y; z); \quad (7) \right)$$



## 2.2. Symmetries of the spin-1 AF Heisenberg chain

The effective Hamiltonian (9) and the field-theory expression of the spin (7) tell us how the symmetries of the spin-1 Heisenberg chain are translated in the field theory language [16, 18, 20]. Here, we consider four kinds of symmetry: the spin rotation  $S_j \rightarrow R S_j$  ( $R$  is an  $SO(3)$  matrix), the one-site translation  $S_j \rightarrow S_{j+1}$ , the time reversal  $S_j \rightarrow S_j$ , and the site-parity transformation  $S_j \rightarrow S_{-j}$ . Note that the link-parity symmetry is the combination of the one-site translational and site-parity symmetries.

(i) The spin rotation may be realized by

$$\tilde{\psi}_{L,R} \rightarrow R \tilde{\psi}_{L,R}; \quad N \rightarrow RN; \quad (12)$$

where  $\tilde{\psi} = \psi^t(1; 2; 3)$  and  $N = \psi^t(N^1; N^2; N^3)$ . The "rotation" of fermions induces  $\mathcal{J} \rightarrow R \mathcal{J}$  and  $\mathcal{J} \rightarrow R \mathcal{J}$ , where  $\mathcal{J} = \psi^t(J^1; J^2; J^3)$  and  $\mathcal{J} = \psi^t(J^1; J^2; J^3)$ . The latter transformation in Eq. (12) could be regarded as

$$\tilde{\psi} \rightarrow R \tilde{\psi}; \quad (13)$$

where  $\tilde{\psi} = \psi^t(2; 1; 3)$ . If both the mass and the interaction terms are absent (i.e., the system is completely critical), the chiral  $SO(3) \times SO(3)$  symmetry (independent rotations of the left and right elds) appears.

(ii) The one-site translation may correspond to

$$\tilde{\psi}_{L,R}(x) \rightarrow \tilde{\psi}_{L,R}(x + a_0); \quad N(x) \rightarrow N(x + a_0); \quad (14)$$

Since the fermions  $\psi_{L,R}^q$  describe the magnon excitations around wave number  $p = \pi a_0$  in the original chain (4), we should insert  $\gamma$  in the first transformation of Eq. (14). The second transformation can be realized by

$$\begin{aligned} \tilde{\psi}(x) &\rightarrow \tilde{\psi}(x + a) \\ \tilde{\psi}(x) &\rightarrow \tilde{\psi}(x + a_0) \end{aligned}; \quad \text{or} \quad \begin{aligned} \tilde{\psi}(x) &\rightarrow \tilde{\psi}(x + a) \\ \tilde{\psi}(x) &\rightarrow \tilde{\psi}(x + a) \end{aligned}; \quad (15)$$

where  $\tilde{\psi} = \psi^t(1; 2; 3)$ . Due to this transformation, the factor  $(-1)^j$  in front of the staggered component of the spin (7) changes to  $(-1)^{j+1}$ .

(iii) The time reversal may correspond to

$$\tilde{\psi}_{L,R} \rightarrow \tilde{\psi}_{R,L}; \quad N \rightarrow N; \quad i \rightarrow i; \quad (16)$$

The second mapping  $N \rightarrow N$  could be interpreted as

$$\begin{aligned} \tilde{\psi} &\rightarrow \tilde{\psi} \\ \tilde{\psi} &\rightarrow \tilde{\psi} \end{aligned}; \quad \text{or} \quad \begin{aligned} \tilde{\psi} &\rightarrow \tilde{\psi} \\ \tilde{\psi} &\rightarrow \tilde{\psi} \end{aligned}; \quad (17)$$

(iv) As a proper mapping of the continuous elds towards the site-parity transformation, we can propose

$$\begin{aligned} \tilde{\psi}_L(x) &\rightarrow \tilde{\psi}_R(-x); \\ \tilde{\psi}_R(x) &\rightarrow \tilde{\psi}_L(-x); \end{aligned}; \quad N(x) \rightarrow N(-x); \quad (18)$$

All the transformations (i)–(iv) leaves the effective Hamiltonian (9) invariant. In other words, the symmetries (i)–(iv) allow the existence of the mass term and the current-current interaction one.

## 2.3. Dirac fermion and Abelian bosonization

Besides the relationship between the WZNW model and three copies of massless real fermions, it is well known that two copies of the real fermions are equivalent to a massless Gaussian theory with TLL parameter 1 or a massless Dirac (complex) fermion, except for a few subtle aspects [4]. Namely, one can bosonize two copies of the real fermions. In this subsection, we apply such a bosonization procedure for the effective theory (9). The results are useful in the later sections.

Let us define the following Dirac fermion,

$$\psi_L = \frac{1}{\sqrt{2}} (\psi_L^1 + i \psi_L^2); \quad \psi_R = \frac{1}{\sqrt{2}} (\psi_R^1 + i \psi_R^2); \quad (19)$$

which obey  $\psi_L^\dagger(x); \psi_L^\dagger(y)g = 0$  and  $\psi_L^\dagger(x); \psi_L^\dagger(y)g = \delta(x-y)$ . Using this, we can rewrite the bilinear (free) part of the Hamiltonian (9) as follows:

$$\begin{aligned} \hat{H}_{\text{free}} &= \int dx \text{iv}_0 \left( \psi_L^\dagger \partial_x \psi_L + \psi_R^\dagger \partial_x \psi_R + m i (\psi_L^\dagger \psi_R - \psi_R^\dagger \psi_L) \right) \\ &\quad + \frac{i}{2} v_0 \left( \psi_L^3 \partial_x \psi_L^3 + \psi_R^3 \partial_x \psi_R^3 + m i \psi_L^3 \psi_R^3 \right) \\ &= \hat{H}[\psi] + \hat{H}[\psi^3]; \end{aligned} \quad (20)$$

Applying the standard Abelian bosonization method (see Appendix A) to the Dirac-fermion part  $\hat{H}[\psi]$ , we obtain the following sine-Gordon Hamiltonian,

$$\hat{H}[\psi] \rightarrow \hat{H}[\psi] = \int dx \frac{v_0}{2} \left( (\partial_x \phi)^2 + (\partial_x \theta)^2 + \frac{m}{4} \cos\left(\frac{p}{4}\phi\right) \right); \quad (21)$$

where  $\phi$  is the scalar field,  $\theta$  is the dual of  $\phi$ ,  $a_0$  is the nonuniversal short-distance cutoff, the relevant cos term bears the Haldane gap, and we simply dropped the Klein factors. The fields  $(\psi_L^1; \psi_L^2)$  and  $(\psi_L^{1;2}; \psi_L^{2;1})$  in the spin operator (7) can also be rewritten in the Dirac-fermion or the boson languages. The uniform component in Eq. (7) is

$$\begin{aligned} J_L^1 + J_R^1 &= \frac{1}{\sqrt{2}} (\psi_L^\dagger \psi_L) \psi_L^3 + \frac{1}{\sqrt{2}} (\psi_R^\dagger \psi_R) \psi_R^3 \\ &= \frac{i}{\sqrt{2}} \psi_L^3 \sin\left(\frac{p}{4}\phi\right) + \psi_R^3 \sin\left(\frac{p}{4}\phi\right); \\ J_L^2 + J_R^2 &= \frac{i}{\sqrt{2}} (\psi_L^\dagger + \psi_L) \psi_L^3 + \frac{i}{\sqrt{2}} (\psi_R^\dagger + \psi_R) \psi_R^3 \\ &= \frac{i}{\sqrt{2}} \psi_L^3 \cos\left(\frac{p}{4}\phi\right) + \psi_R^3 \cos\left(\frac{p}{4}\phi\right); \\ J_L^3 + J_R^3 &= \psi_L^\dagger \psi_L : \psi_R^\dagger \psi_R = \frac{1}{\sqrt{2}} \partial_x \phi; \end{aligned} \quad (22)$$

where  $K$  is the Klein factor for the fermion  $\psi$ , and we assumed  $\psi_L^\dagger; \psi_R^\dagger g = 0$ . The symbol  $::$  denotes a normal-ordered product;  $\psi_L^\dagger \psi_L = \psi_L^\dagger \psi_L - \langle \psi_L^\dagger \psi_L \rangle$ . From the formula (A.6), the staggered component of the spin [50] is bosonized as

$$\begin{aligned} N^1 &= \psi_L^3 \sin\left(\frac{p}{4}\phi\right); \\ N^2 &= \psi_L^3 \cos\left(\frac{p}{4}\phi\right); \\ N^3 &= \psi_L^3 \cos\left(\frac{p}{4}\phi\right); \end{aligned} \quad (23)$$

Employing Eqs. (19)–(23) and Abelian bosonization formulas (see Appendix A), let us consider how symmetry operations of the original spin-1 chain (4) are represented within the above Dirac-fermion and boson framework, as in the last subsection.

(i) Let us define the following U(1) rotation around the spin z axis,

$$\begin{pmatrix} 0 & 1 & 0 \\ S_j^x & \cos & \sin & 0 \\ S_j^y & \sin & \cos & 0 \\ S_j^z & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ A \\ A \\ S_j^z \end{pmatrix} = \begin{pmatrix} 1 \\ S_j^x \\ S_j^y \\ S_j^z \end{pmatrix} \quad (24)$$

where  $\theta$  is a real number. One can easily find that this corresponds to

$$L_{j,R} \rightarrow e^{i\theta} L_{j,R}; \quad (x) \rightarrow (x) - \theta \quad (25)$$

This transformation is nothing but the U(1) symmetry that is explicitly preserved via the Abelian bosonization procedure.

(ii) The one-site translation  $S_j \rightarrow S_{j+1}$  could be realized as

$$\begin{pmatrix} L_{j,R} \\ S_j^x \\ L_{j,R} \end{pmatrix} (x) \rightarrow \begin{pmatrix} L_{j,R} \\ S_j^x \\ L_{j,R} \end{pmatrix} (x + a_0); \quad \begin{pmatrix} (x) \\ (x) \\ (x) \end{pmatrix} \rightarrow \begin{pmatrix} (x + a_0) \\ (x + a_0) \\ (x + a_0) \end{pmatrix} \quad (26)$$

The second and third transformations change the staggered factor  $(-1)^j$  in Eq. (7) into  $(-1)^{j+1}$ . These two correspond to the first proposal in Eq. (15).

(iii) The time reversal  $S_j \rightarrow S_j$  could be interpreted as

$$\begin{pmatrix} L_{j,R} \\ S_j^y \\ L_{j,R} \end{pmatrix} \rightarrow \begin{pmatrix} R_{j,L} \\ S_j^y \\ R_{j,L} \end{pmatrix}; \quad \begin{pmatrix} L_{j,R} \\ (x) \\ L_{j,R} \end{pmatrix} \rightarrow \begin{pmatrix} R_{j,L} \\ (x) \\ R_{j,L} \end{pmatrix}; \quad \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} \rightarrow \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix}; \quad i \rightarrow -i \quad (27)$$

The second and third mappings are equivalent to the second proposal in Eq. (17).

(iv) The site-parity transformation  $S_j \rightarrow S_j$  might correspond to [51]

$$\begin{pmatrix} L_{j,R} \\ S_j^z \\ L_{j,R} \end{pmatrix} (x) \rightarrow \begin{pmatrix} R_{j,L} \\ S_j^z \\ R_{j,L} \end{pmatrix} (x); \quad \begin{pmatrix} L_{j,R} \\ (x) \\ L_{j,R} \end{pmatrix} \rightarrow \begin{pmatrix} R_{j,L} \\ (x) \\ R_{j,L} \end{pmatrix}; \quad \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} \rightarrow \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix}; \quad \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} \rightarrow \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} \quad (28)$$

### 3. Effective field theory for uniform - eld-induced TLL in spin-1 AF chain

Based on the Majumara fermion theory explained in the last section, we consider the uniform - eld effects in the spin-1 AF chain (4), especially the uniform - eld-induced critical phase.

From the formula (7), the Zeeman term in the chain (4) is approximated as the following fermion bilinear form,

$$H = \sum_j S_j^z \int dx iH \left( \frac{1}{L} \frac{\partial}{\partial L} + \frac{1}{R} \frac{\partial}{\partial R} \right); \quad (29)$$

Therefore, it is possible to nonperturbatively treat the Zeeman term, together with the free part (20), within the field theory scheme. This is a significant advantage of the Majumara fermion theory. Take notice here that the Zeeman coupling reduces the spin SU(2) symmetry to the U(1) one around the spin z axis, and destroys the time-reversal symmetry; the remaining symmetries are the U(1), the one-site translation, and the site-parity ones. The right-hand side in Eq. (29) is indeed not invariant under the time reversal and any spin rotations except for the U(1) one.

Supposing the inter-magnon interaction may be negligible at the starting point, the effective Hamiltonian for the spin-1 AF chain (4) with a finite  $H_z$  is

$$\begin{aligned} \hat{H}_H^e = & \int dx \left[ i v_0 \left( \frac{y}{L} \partial_x \psi_L - \frac{y}{R} \partial_x \psi_R \right) \right. \\ & + m i \left( \frac{y}{L} \psi_R - \frac{y}{R} \psi_L \right) + H \left( \frac{y}{L} \psi_L + \frac{y}{R} \psi_R \right) \\ & \left. + \frac{i}{2} v_0 \left( \psi_L^3 \partial_x \psi_L - \psi_R^3 \partial_x \psi_R \right) + m i \psi_L^3 \psi_R^3 \right]; \end{aligned} \quad (30)$$

In order to diagonalize this Hamiltonian, and see its band structure, we introduce the Fourier transformations of fermion fields as follows:

$$\begin{aligned} \psi_L^3(x) &= \frac{1}{\sqrt{L}} \int_{-X}^X e^{ikx} \tilde{\psi}^3(k); \quad \tilde{\psi}^3(k) = \frac{1}{\sqrt{L}} \int dx e^{-ikx} \psi_L^3(x); \\ \psi_R^3(x) &= \frac{1}{\sqrt{L}} \int_{-X}^X e^{ikx} \tilde{\psi}^3(k); \quad \tilde{\psi}^3(k) = \frac{1}{\sqrt{L}} \int dx e^{-ikx} \psi_R^3(x); \end{aligned} \quad (31)$$

where  $L = N a_0$  is the system size ( $N$ : total number of site), and  $k$  ( $|k| < \frac{\pi}{a_0}$  is the ultraviolet cut-off) is the wave number; a related quantity  $p = k + \frac{\pi}{a_0}$  is interpreted as the wave number of the original lattice system (4). Since  $\tilde{\psi}^3$  is real,  $\tilde{\psi}^3(k) = \tilde{\psi}^3(-k)$  holds. New fermions  $\tilde{\psi}^3$  and  $\tilde{\psi}^3$  obey the anticommutation relations  $f \tilde{\psi}^3(k); \tilde{\psi}^3_0(k^0)g = \delta_{k,k^0}; f \tilde{\psi}^3(k); \tilde{\psi}^3_0(k^0)g = \delta_{k,k^0}$ , and  $f \tilde{\psi}^3(k); \tilde{\psi}^3_0(k^0)g = 0$ . Substituting Eq. (31) into the Hamiltonian (30), we obtain

$$\hat{H}_H^e = \sum_{k < \frac{\pi}{2}} \tilde{\psi}_k^y M_k \tilde{\psi}_k + \sum_{0 < k < \frac{\pi}{2}} \tilde{\psi}_k^y M_k \tilde{\psi}_k; \quad (32)$$

where  $\tilde{\psi}_k = \begin{pmatrix} \tilde{\psi}_R \\ \tilde{\psi}_L \end{pmatrix}$ ,  $\tilde{\psi}_k = \begin{pmatrix} \tilde{\psi}_R^3 \\ \tilde{\psi}_L^3 \end{pmatrix}$ , and

$$M_k = \begin{pmatrix} kv_0 + H & im \\ im & kv_0 + H \end{pmatrix}; \quad M_k = \begin{pmatrix} kv_0 & im \\ im & kv_0 \end{pmatrix}; \quad (33)$$

Subsequently, performing the following Bogoliubov transformation,

$$\tilde{\psi}_k = U_k \begin{pmatrix} \tilde{\psi}_-(k) \\ \tilde{\psi}_+(k) \end{pmatrix}; \quad \tilde{\psi}_k = U_k \begin{pmatrix} \tilde{\psi}_0(k) \\ \tilde{\psi}_0(-k) \end{pmatrix}; \quad (34)$$

where

$$\begin{aligned} U_k &= \frac{m}{2 \epsilon_0(k)} \begin{pmatrix} \epsilon_0(k) & kv_0 \\ i \epsilon_0(k) & kv_0 \end{pmatrix}^{1=2} \begin{pmatrix} \epsilon_0(k) + kv_0 \\ i \epsilon_0(k) + kv_0 \end{pmatrix}^{1=2} \\ & \quad \frac{1}{Q} \begin{pmatrix} \epsilon_0(k) & kv_0 \\ i \epsilon_0(k) & kv_0 \end{pmatrix}^{1=2} \begin{pmatrix} \epsilon_0(k) + kv_0 \\ i \epsilon_0(k) + kv_0 \end{pmatrix}^{1=2} m; \\ \epsilon_0(k) &= \sqrt{k^2 v_0^2 + m^2}; \end{aligned} \quad (35)$$

one can finally obtain the diagonalized Hamiltonian,

$$\begin{aligned} \hat{H}_H^e = & \sum_{k < \frac{\pi}{2}} \tilde{\psi}_k^y \tilde{\psi}_k \tilde{\psi}_k + \sum_{0 < k < \frac{\pi}{2}} \tilde{\psi}_k^y \tilde{\psi}_0(k) \tilde{\psi}_0(k) \\ & + \sum_{k < \frac{\pi}{2}} \tilde{\psi}_k^y \tilde{\psi}_+(k) \tilde{\psi}_+(k); \end{aligned} \quad (36)$$

where  $\epsilon_{\pm}(\mathbf{k}) = \epsilon_{\pm 0}(\mathbf{k}) - H$ . While three bands  $\epsilon_{\pm 0}(\mathbf{k})$  are degenerate at  $H = 0$ , a finite field  $H$  mixes the first and third fermions  $\psi^1$  and  $\psi^3$ , and leaves these bands split. The dispersions  $\epsilon_{\pm 0}(\mathbf{k})$  indicate that the Majumara fermion theory correctly reproduces the Zeeman splitting. Fields  $\tilde{\psi}_+$ ,  $\tilde{\psi}_0^y$  and  $\tilde{\psi}_-$  may respectively be regarded as  $S^z = 1; 0$  and  $S^z = -1$  magnon creation operators. Here, let us consider how the symmetry operations are represented in the picture of the magnon fields  $\tilde{\psi}_{\pm 0}$ . Using the transformations (25) and (26) and the relationship between  $(\psi^1, \psi^3)$  and  $\tilde{\psi}_{\pm 0}$  [Eqs. (31) and (34)], one finds that the  $U(1)$  rotation along the spin  $z$  axis and the one-site translation respectively correspond to

$$\tilde{\psi}_{\pm} \rightarrow e^{i\alpha} \tilde{\psi}_{\pm}; \quad (37)$$

$$\tilde{\psi}_{\pm}(\mathbf{k}) \rightarrow e^{i\mathbf{k}a_0} \tilde{\psi}_{\pm}(\mathbf{k}); \quad \tilde{\psi}_0(\mathbf{k}) \rightarrow \begin{cases} e^{i\mathbf{k}a_0} \tilde{\psi}_0(\mathbf{k}) & (\mathbf{k} > 0) \\ e^{-i\mathbf{k}a_0} \tilde{\psi}_0(\mathbf{k}) & (\mathbf{k} < 0) \end{cases}; \quad (38)$$

If we neglect the  $\mathbf{k}$ -dependence of  $U_{\mathbf{k}}$  or consider only fermion fields around  $\mathbf{k} = 0$ , we could interpret that the site-parity transformation in Eq. (28) is realized by

$$\begin{aligned} \tilde{\psi}_+(\mathbf{k}) &\rightarrow i\tilde{\psi}_-(\mathbf{k}); & \tilde{\psi}_0(\mathbf{k}) &\rightarrow i\tilde{\psi}_0(\mathbf{k}); \\ \tilde{\psi}_0(\mathbf{k}) &\rightarrow \begin{cases} i\tilde{\psi}_0(\mathbf{k}) & (\mathbf{k} > 0) \\ i\tilde{\psi}_0(\mathbf{k}) & (\mathbf{k} < 0) \end{cases}; \end{aligned} \quad (39)$$

The Hamiltonian (36) is clearly invariant under these transformations. One has to keep in mind that the symmetry operation (39) is valid only around  $\mathbf{k} = 0$ .

As  $H$  exceeds the Haldane gap  $m$ , the band  $\epsilon_+(\mathbf{k})$  crosses the zero energy line. This just corresponds to the C-IC transition of the spin-1 AF chain (4) [22, 23, 24, 25]. Then the system enters in a critical TLL phase with the magnon  $\tilde{\psi}_+$  condensed. The two remaining bands  $\epsilon_{\pm 0}(\mathbf{k})$  are still massive and completely empty of magnons:  $\langle \psi^3 \rangle \neq 0$  and  $\langle \psi^1 \rangle = 0$  still hold. Focusing on the low-energy physics in this case of  $H > m$ , we can approximate the  $S^z = 1$  magnon field  $\tilde{\psi}_+$  as a new Dirac fermion with a linear dispersion as in Fig. 1. The left and right movers  $L(x)$  and  $R(x)$  of the Dirac fermion are defined as

$$\begin{aligned} L(x) &= \frac{1}{\sqrt{L}} \sum_{\mathbf{k} < k_F} e^{i\mathbf{k}x} \tilde{\Gamma}(\mathbf{k}); & R(x) &= \frac{1}{\sqrt{L}} \sum_{\mathbf{k} > k_F} e^{i\mathbf{k}x} \tilde{R}(\mathbf{k}); \\ \tilde{\Gamma}(\mathbf{k}) &= \tilde{\psi}_+(\mathbf{k} + k_F); & \tilde{R}(\mathbf{k}) &= \tilde{\psi}_+(\mathbf{k} - k_F); \end{aligned} \quad (40)$$

where the Fermi wave number  $k_F = \frac{p}{H^2 - m^2} = v_0$  is determined from  $\epsilon_+(k_F) = 0$ , and the cut-off should be much smaller than  $\Lambda$ . Using the Dirac fermion, one can rewrite the effective Hamiltonian (36) in the case of  $H > m$ , in the real space, as follows:

$$\begin{aligned} \hat{H}_{H > m}^e &= \int dx \left[ i v_F L^y \partial_x L - R^y \partial_x R \right. \\ &\quad + \frac{i}{2} v_0 \left( L^3 \partial_x^3 L - R^3 \partial_x^3 R \right) + m i L^3 R^3 \\ &\quad \left. + \gamma \frac{v_0^2}{2m} \partial_x^2 + m + H + O(\partial_x^4) \right] \\ &= \hat{H}[L; R] + \hat{H}[\psi^3] + \hat{H}[\psi^1]; \end{aligned} \quad (41)$$

where  $v_F = \partial_{\mathbf{k}} \epsilon_+|_{\mathbf{k}=k_F} = \frac{p}{H^2 - m^2} = v_0$  is the Fermi velocity, and  $\psi^i(x) = \frac{1}{\sqrt{L}} \sum_{\mathbf{k}} e^{i\mathbf{k}x} \tilde{\psi}^i(\mathbf{k})$ . Take notice that the linearized dispersion and the definition of  $L$  and  $R$  become less reliable as the field  $H$  becomes closer to the lower critical value  $m$  ( $v_F \rightarrow 0$ ).




Figure 1. Spin-1 magnon band structure at  $H > m$ .

So far we have omitted the inter-magnon interaction terms in this section. They would yield interactions among  $L, R, S$  and  $\sigma$  (see the next section). A better method of dealing with such terms is the Abelian bosonization. Through it, the Dirac-fermion part in the effective theory (41) is mapped to the following Gaussian model,

$$\hat{H}[L;R] = \int dx \frac{v}{2} K (\partial_x \phi)^2 + \frac{1}{K} (\partial_x \chi)^2; \quad (42)$$

where  $\phi$  is a scalar field,  $\chi$  is the dual of  $\phi$ ,  $v$  is the renormalized Fermi velocity, and  $K$  is the TLL parameter: although  $K = 1$  and  $v = v_F$  at the stage of Eq. (41), both parameters are subject to the renormalization due to irrelevant interaction effects neglected here. The Hamiltonian (42) just describes the low-energy physics of the TLL phase in the chain (4).

#### 4. Field-theory representation of spin operators in $H > m$

Making use of the contents in the preceding sections, we construct a field-theory representation of spin operators in the uniform-field-induced TLL phase in the spin-1 AF chain (4).

##### 4.1. Spin uniform component and Magnetization

First, we consider the uniform component of the spin,  $S = S_L + S_R$  in Eq. (7). In the low-energy limit, Eqs. (34) and (40) allow us to approximate the "old" Dirac fermion as follows:

$$\begin{aligned} \psi_R(x) &= \frac{1}{\sqrt{2}} (\psi(x) + e^{ik_F x} U_+ \psi(x) + e^{-ik_F x} U_- \psi(x)); \\ \psi_L(x) &= \frac{i}{\sqrt{2}} (\psi(x) - ie^{ik_F x} U_- \psi(x) - ie^{-ik_F x} U_+ \psi(x)); \end{aligned} \quad (43)$$

where the prefactor  $1/\sqrt{2}$  of the first term is the  $(1;1)$   $[(2;1)]$  component of the matrix  $U_{k=0}$ , and  $U_{\pm} = m / (2H \pm \sqrt{H^2 - m^2})$  is the  $(1;1)$  component of  $U_{k_F}$ . Note that  $U_{\pm}$  have properties (i)  $U_{\pm} = 1/\sqrt{2}$  at  $H = m + 0$ , (ii)  $U_+^2 + U_-^2 = 1$ , and (iii)  $U_+ U_- = \frac{m}{2H}$ . From this relationship between  $U_{\pm}$  and  $(L;R; \sigma)$  [see also Eqs. (37)-(39)], we know how the latter three fields are transformed by symmetry operations. The  $U(1)$  rotation in Eq. (25) corresponds to

$$\begin{aligned} \psi &\rightarrow e^{i\theta} \psi; & L &\rightarrow e^{-i\theta} L \\ \psi &\rightarrow e^{-i\theta} \psi; & R &\rightarrow e^{i\theta} R \end{aligned} \quad (44)$$

The one-site translation in Eq. (26) is mapped to

$$\begin{aligned} L(x) & \rightarrow L(x + a_0); & R(x) & \rightarrow R(x + a_0) \end{aligned} \quad (45)$$

The site-parity transformation in Eq. (28),  $L(x) \rightarrow R(x)$  and  $R(x) \rightarrow L(x)$ , can be reproduced by

$$\begin{aligned} L(x) & \rightarrow iR(x); & R(x) & \rightarrow iL(x) \end{aligned} \quad (46)$$

One can confirm that the Hamiltonian (41) is invariant under these transformations.

Using the relation (43), we can rewrite fields  $\psi_{L,R}^{\pm}$  in terms of  $L, R$ , and  $\chi$ . As a result, the currents  $J_{L,R}$  are reexpressed as

$$\begin{aligned} J_L^1(x) &= \frac{\hbar}{2} (\psi_L^\dagger \psi_L + \psi_R^\dagger \psi_R) + \frac{i}{2} U_- (e^{ik_F x} \psi_L^\dagger \psi_L + e^{ik_F x} \psi_R^\dagger \psi_R) \\ &+ \frac{i}{2} U_+ (e^{ik_F x} \psi_R^\dagger \psi_L + e^{ik_F x} \psi_L^\dagger \psi_R) \quad (47a) \end{aligned}$$

$$\begin{aligned} J_L^2(x) &= \frac{\hbar}{2} (\psi_L^\dagger \psi_L - \psi_R^\dagger \psi_R) + \frac{1}{2} U_- (e^{ik_F x} \psi_L^\dagger \psi_L - e^{ik_F x} \psi_R^\dagger \psi_R) \\ &+ \frac{1}{2} U_+ (e^{ik_F x} \psi_R^\dagger \psi_L - e^{ik_F x} \psi_L^\dagger \psi_R) \quad (47b) \end{aligned}$$

$$\begin{aligned} J_L^3(x) &= \frac{1}{4} (\psi_L^\dagger \psi_L + \psi_R^\dagger \psi_R) + \frac{U_-^2}{2} (\psi_L^\dagger \psi_L + \psi_R^\dagger \psi_R) + \frac{U_+^2}{2} (\psi_R^\dagger \psi_R + \psi_L^\dagger \psi_L) \\ &+ \frac{U_- \hbar}{2} e^{ik_F x} \psi_L^\dagger \psi_L + \frac{U_+ \hbar}{2} e^{ik_F x} \psi_R^\dagger \psi_R + \hbar \chi \\ &+ U_+ U_- e^{2ik_F x} \psi_L^\dagger \psi_R + \hbar \chi \quad (47c) \end{aligned}$$

$$\begin{aligned} J_R^1(x) &= \frac{\hbar}{2} (\psi_L^\dagger \psi_L + \psi_R^\dagger \psi_R) + \frac{1}{2} U_+ (e^{ik_F x} \psi_L^\dagger \psi_L + e^{ik_F x} \psi_R^\dagger \psi_R) \\ &+ \frac{1}{2} U_- (e^{ik_F x} \psi_R^\dagger \psi_L + e^{ik_F x} \psi_L^\dagger \psi_R) \quad (47d) \end{aligned}$$

$$\begin{aligned} J_R^2(x) &= \frac{\hbar}{2} (\psi_L^\dagger \psi_L - \psi_R^\dagger \psi_R) + \frac{1}{2} U_+ (e^{ik_F x} \psi_L^\dagger \psi_L - e^{ik_F x} \psi_R^\dagger \psi_R) \\ &+ \frac{1}{2} U_- (e^{ik_F x} \psi_R^\dagger \psi_L - e^{ik_F x} \psi_L^\dagger \psi_R) \quad (47e) \end{aligned}$$

$$\begin{aligned} J_R^3(x) &= \frac{1}{4} (\psi_L^\dagger \psi_L + \psi_R^\dagger \psi_R) + \frac{U_+^2}{2} (\psi_L^\dagger \psi_L + \psi_R^\dagger \psi_R) + \frac{U_-^2}{2} (\psi_R^\dagger \psi_R + \psi_L^\dagger \psi_L) \\ &+ \frac{U_+ \hbar}{2} e^{ik_F x} \psi_L^\dagger \psi_L + \frac{U_- \hbar}{2} e^{ik_F x} \psi_R^\dagger \psi_R + \hbar \chi \\ &+ U_+ U_- e^{2ik_F x} \psi_L^\dagger \psi_R + \hbar \chi \quad (47f) \end{aligned}$$

It is easily found that the above  $J_{L,R}$  are appropriately transformed for the symmetry operations (44), (25), (45), (26), (46) and (28). In addition, the Hermitian nature of all current operators is preserved in Eq. (47). Even if, instead of Eq. (43), we use precise representations of continuous fields in the Fourier space such as Eqs. (31) and (40), we can finally arrive at the same expression as Eq. (47).

The Fermi wave number  $k_F$  is related with the uniform magnetization  $M = \hbar S_j^z$ . It is given by the formula (47) or the Fourier-space representation of  $J^3 = J_L^3 + J_R^3$ .

The result is

$$\begin{aligned}
 M &= \frac{1}{N} \sum_j S_j^z = \frac{1}{N} \int dx h J^3(x) \\
 &= \sum_{\mathbf{k} < \mathbf{k}_F} h \tilde{v}_+(\mathbf{k}) \tilde{v}_+(\mathbf{k}) = \frac{k_F a_0}{2} :
 \end{aligned} \tag{48}$$

From the derivation process of Eq. (48), we also find that the first three terms in Eqs. (47c) and (47f) should respectively be regarded as

$$\begin{aligned}
 &\frac{1}{4} \sum_y \sum_y + \frac{U^2}{2} L L^y L^y L + \frac{U_+^2}{2} R R^y R^y R \\
 &: U^2 : L^y L : U_+^2 : R^y R : + \frac{M}{2a_0} \frac{1}{2} \sum_y ; \\
 &\frac{1}{4} \sum_y \sum_y + \frac{U^2}{2} L L^y L^y L + \frac{U^2}{2} R R^y R^y R \\
 &: U_+^2 : L^y L : U^2 : R^y R : + \frac{M}{2a_0} \frac{1}{2} \sum_y :
 \end{aligned} \tag{49}$$

Utilizing the results (47) –(49), and applying the Abelian bosonization to the Dirac fermion  $(L;R)$ , we can straightforwardly lead to the following partially bosonized currents  $J_{L;R}$ :

$$\begin{aligned}
 J_L^1(x) &= \frac{h}{2} (\sum_y + ) + \frac{iU}{\mathcal{P}=\mathcal{E}_0} \cos(\frac{p}{4} L) \cos(\frac{p}{4} L) \quad (M x=a_0) \\
 &+ \frac{iU_+}{\mathcal{P}=\mathcal{E}_0} \cos(\frac{p}{4} R) \cos(\frac{p}{4} R) \quad (M x=a_0) \quad \frac{i}{L} ; \\
 J_L^2(x) &= \frac{h}{2} (\sum_y) + \frac{iU}{\mathcal{P}=\mathcal{E}_0} \sin(\frac{p}{4} L) \sin(\frac{p}{4} L) \quad (M x=a_0) \\
 &+ \frac{iU_+}{\mathcal{P}=\mathcal{E}_0} \sin(\frac{p}{4} R) \sin(\frac{p}{4} R) \quad (M x=a_0) \quad \frac{i}{L} ; \\
 J_L^3(x) &= \frac{U^2}{\mathcal{P}=\mathcal{E}_x} L + \frac{U_+^2}{\mathcal{P}=\mathcal{E}_x} R + \frac{M}{2a_0} \frac{1}{2} \sum_y \\
 &+ \frac{U}{\mathcal{P}=\mathcal{E}_0} \sum_y h e^{i \frac{p}{4} L + i M x=a_0} + h x \quad \frac{i}{L} \\
 &+ \frac{U_+}{\mathcal{P}=\mathcal{E}_0} \sum_y h e^{i \frac{p}{4} R - i M x=a_0} + h x \quad \frac{i}{R} \\
 &\frac{U_+ U}{\mathcal{P}=\mathcal{E}_0} L R \sin(\frac{p}{4} L R) \quad (2 M x=a_0); \\
 J_R^1(x) &= \frac{h}{2} (\sum_y) + \frac{iU_+}{\mathcal{P}=\mathcal{E}_0} \sin(\frac{p}{4} L) \sin(\frac{p}{4} L) \quad (M x=a_0) \\
 &\frac{iU}{\mathcal{P}=\mathcal{E}_0} \sin(\frac{p}{4} R) \sin(\frac{p}{4} R) \quad (M x=a_0) \quad \frac{i}{R} ; \\
 J_R^2(x) &= \frac{h}{2} (\sum_y + ) + \frac{iU_+}{\mathcal{P}=\mathcal{E}_0} \cos(\frac{p}{4} L) \cos(\frac{p}{4} L) \quad (M x=a_0)
 \end{aligned}$$

$$\begin{aligned}
& + \frac{iU}{\mathcal{P}^2} \cos\left(\frac{p}{4} M x=a_0\right) \frac{i}{R}{}^3; \\
J_R^3(x) &= \frac{U_+^2}{\mathcal{P}^2} \theta_{x-L} \frac{U_-^2}{\mathcal{P}^2} \theta_{x-R} + \frac{M}{2a_0} \frac{1}{2} y \\
& \frac{U_+}{\mathcal{P}^2} \frac{h}{4} y_L e^{i\frac{p}{4} L + i M x=a_0} + h x \frac{i}{R} \\
& \frac{U_-}{\mathcal{P}^2} \frac{h}{4} y_R e^{i\frac{p}{4} R - i M x=a_0} + h x \frac{i}{L} \\
& \frac{U_+ U_-}{i} \frac{1}{\mathcal{P}^2} \sin\left(\frac{p}{4} 2 M x=a_0\right); \quad (50)
\end{aligned}$$

where  $\theta = 1 = 0$  is the short-distance cut off, and  $\theta_{L(R)}$  is the Klein factor for the field  $L(R)$ . Therefore, the bosonized uniform components of the spin are written as

$$\begin{aligned}
J^3(x) &= \frac{M}{a_0} \frac{1}{\mathcal{P}^2} \theta_{x-L} y \frac{iC_0^z}{L} \sin\left(\frac{p}{4} 2 M x=a_0\right) \\
& + C_1^z y_L e^{i\frac{p}{4} L + i M x=a_0} + h x \\
& C_1^z y_R e^{i\frac{p}{4} R - i M x=a_0} + h x + \quad ; \quad (51a)
\end{aligned}$$

$$\begin{aligned}
J^+(x) &= J^1(x) + iJ^2(x) \\
& i + e^{i\frac{p}{4} L} iC_1 e^{i\frac{p}{4} L + M x=a_0} \\
& + iC_2 e^{i\frac{p}{4} R - M x=a_0} + \frac{i}{L} + \frac{3}{L} \\
& + e^{i\frac{p}{4} L} C_2 e^{i\frac{p}{4} L + M x=a_0} \\
& + C_1 e^{i\frac{p}{4} R - M x=a_0} + \frac{i}{R} + \frac{3}{R}; \quad (51b)
\end{aligned}$$

where  $C_n$  and  $C_n^z$  are nonuniversal constants, which contain the effects of irrelevant terms neglected at the stage (41). When the irrelevant terms are not taken into account, constants  $C_0^z$ ,  $C_1^z$ ,  $C_1$ , and  $C_2$  are respectively proportional to  $U_+ U_-$ ,  $U_- U_+$ ,  $U_-$ , and  $U_+$ . Therefore, it is inferred that  $C_1^z \neq 0$  at  $H \neq m + 0$ . In Eqs. (50) and (51), we used the formula (A.3). If, instead of it, we use another formula (A.4), we can add more irrelevant terms to Eqs. (50) and (51). Namely, we may perform the replacement,

$$\begin{aligned}
e^{i\frac{p}{4} L + i M x=a_0} & \sum_{n=0}^{\infty} C_n e^{i(2n+1)\frac{p}{4} L - M x=a_0} e^{i\frac{p}{4} L}; \\
e^{i\frac{p}{4} R - i M x=a_0} & \sum_{n=0}^{\infty} C_n e^{i(2n+1)\frac{p}{4} R - M x=a_0} e^{i\frac{p}{4} R}; \quad (52)
\end{aligned}$$

where  $C_n$  is a nonuniversal constant. Equation (51) is a main result in this paper. We expect that replacing the Klein factors in Eq. (51) with a constant is admitted in most situations. (A typical situation is when one investigates asymptotic features of spin correlation functions. See Sec. 4.4.)

Finally, using the results (47)–(50), let us investigate how the current-current interaction term modifies the parameter  $K$  of the effective theory (42). Since the

SU(2) spin-rotation symmetry is reduced to the U(1) one in the case with a finite  $H$ , the term  $J_L J_R$  might be modified to  $\gamma (J_L^1 J_R^1 + J_L^2 J_R^2) + \kappa J_L^3 J_R^3$ , in which we assume that both  $\gamma$  and  $\kappa$  are positive. Equations (47) and (49) lead to the following representation of the current-current interaction [52]:

$$\begin{aligned}
J_L^3(x_+) J_R^3(x) &= U^2 :R^Y R^X : + U_+^2 :L^Y L^X : + U_+^2 :R^Y R^X : + U^2 :L^Y L^X : \\
&+ U_+^2 U^2 L^Y R(x_+) R^Y L(x) + R^Y L(x_+) L^Y R(x) \\
&\frac{M}{2a_0} :L^Y L^X : + :R^Y R^X : \\
&+ \frac{U_+^2 + U^2}{2} \gamma :L^Y L^X : + :R^Y R^X : \\
&\frac{U_+ U}{2} \hbar \gamma (L(x_+) L^Y(x) + R(x_+) R^Y(x)) + \hbar \kappa : \\
&+ \frac{U^2}{2} U_+^2 \gamma (x_+) \gamma(x) L R + \hbar \kappa : \\
&+ \text{oscillating terms} + \text{others}; \tag{53a}
\end{aligned}$$

$$\begin{aligned}
J_L^1(x_+) J_R^1(x) + J_L^2(x_+) J_R^2(x) &= iU_+ U L^Y L^X - L L^Y + R^Y R^X - R R^Y \frac{3}{L} \frac{3}{R} \\
&+ \text{oscillating terms} + \text{others}; \tag{53b}
\end{aligned}$$

where we used  $x_+ = x + \frac{1}{2} \frac{v}{H}$  ( $\frac{1}{2}$  is a small parameter of  $O(\frac{1}{H})$ ), "oscillating terms" have a factor  $e^{inMx/a_0}$  ( $n$ : integer), which are irrelevant except for the case that  $M$  is close to a special commensurate value, and "others" are constructed by only massive fields ( $\psi^3$ ;  $\bar{\psi}$ ). The Hamiltonian (41) plus the above current-current interaction is regarded as a more accurate low-energy effective theory for the uniform- $\hbar$ -driven TLL phase. Integrating out the part of massive fields ( $\psi^3$ ;  $\bar{\psi}$ ) in the effective theory [53], we obtain a Hamiltonian for the interacting Dirac fermion ( $L$ ;  $R$ ) that just corresponds to the Gaussian theory (42). In the Gaussian theory framework, the first and second terms of Eq. (53a) directly contribute to the renormalization of the TLL parameter  $K$  and the velocity  $v$  as follows. The bosonized forms of these two terms are written as

$$\begin{aligned}
U^2 :R^Y R^X : + U_+^2 :L^Y L^X : + U_+^2 :R^Y R^X : + U^2 :L^Y L^X : \\
! \frac{1}{4} (\partial_x)^2 \frac{(U^2 - U_+^2)^2}{4} (\partial_x)^2 \\
+ \frac{U^4 - U_+^4}{4} [\partial_x \partial_x - \partial_x \partial_x]; \tag{54a}
\end{aligned}$$

$$\begin{aligned}
U_+^2 U^2 L^Y R(x_+) R^Y L(x) + R^Y L(x_+) L^Y R(x) \\
! \frac{U_+^2 U^2}{(2\hbar)^2} e^{i\frac{p}{4}(x_+) - (x)} + e^{i\frac{p}{4}(x_+) - (x)} \\
\frac{m^2}{4H^2} (\partial_x)^2 + \text{const} + \dots; \tag{54b}
\end{aligned}$$

where we used  $U_+^2 + U^2 = 1$  and  $U_+ U = \frac{m}{2H}$ , and assumed  $\frac{1}{2} = 0$ . The  $\frac{1}{2}$  term in Eq. (54a) could be negligible or be replaced with a constant: such a procedure is supported by the equal-time commutation relation  $[\psi(x); \psi(y)] = \frac{1}{2} \text{sgn}(x - y)$ . Consequently, two terms in Eq. (54) lead to an additional boson kinetic term,

$$\frac{1}{4} \left( \kappa - \frac{m^2}{H^2} \gamma \right) (\partial_x)^2 + \frac{\kappa}{4} (U^2 - U_+^2)^2 (\partial_x)^2; \tag{55}$$



where  $D$  is an arbitrary constant [55]. Then, employing approximated results (58), (60), (43), and the bosonization formula  $L;R e^{i\frac{p}{4}L;R}$  in the right-hand sides in Eq. (62), we can arrive at a desirable representation of  $e^{i\frac{p}{4}L;R}$ . This method was proposed in Ref. [29].

Following the above idea, we obtain

$$e^{i\frac{p}{4}L;R} = \frac{r}{4} e^{i\frac{p}{4}L;R} U (i_L L + R R) + U_+ (i_L R e^{i\frac{p}{4}L;R} + R L e^{i\frac{p}{4}L;R}) + \frac{p}{2} (i_L e^{i\frac{p}{4}L;R} + R e^{i\frac{p}{4}L;R}); \tag{63a}$$

$$e^{i\frac{p}{4}L;R} = \frac{r}{4} U (i_L L + R R) e^{i\frac{p}{4}L;R} + U_+ (i_L R + R L) e^{i\frac{p}{4}L;R} + \frac{p}{2} (i_L e^{i\frac{p}{4}L;R} + R e^{i\frac{p}{4}L;R}); \tag{63b}$$

where we set  $D = 1/2$ , and  $K_{L(R)}$  is the Klein factor of the field  $L(R)$  [56]. It is found that the naive expectations (58) and (60) are qualitatively consistent with the result (63). From this formula, our target, the staggered component of the spin may be represented as

$$N^3(x) = \frac{3}{4} \cos(\frac{p}{4}L;R) + D_1^z \cos(\frac{p}{4}L;R) + D_1^{z0} (i_L + R) e^{i\frac{p}{4}L;R} + h.c.; \tag{64a}$$

$$N^+(x) = N^1(x) + iN^2(x) = i \frac{3}{4} e^{i\frac{p}{4}L;R} + D_1^z e^{i\frac{p}{4}L;R} + D_2 \cos(\frac{p}{4}L;R) + D_1^0 (i_L e^{i\frac{p}{4}L;R} + R e^{i\frac{p}{4}L;R}); \tag{64b}$$

where  $D_n^{(z,0)}$  are nonuniversal constants, which include effects of irrelevant terms, and we naively omitted some of Klein factors. We used the formula  $L;R e^{i\frac{p}{4}L;R}$  in Eqs. (63) and (64). Instead of that, applying the formula (A.4), we can introduce more irrelevant terms in Eqs. (63) and (64) [see the replacement (52)]. For instance, we might replace  $\cos(\frac{p}{4}L;R)$  in Eq. (64a) with

$$\sum_{n=0}^{\infty} \tilde{D}_n \cos((2n+1)\frac{p}{4}L;R); \tag{65}$$

where  $\tilde{D}_n$  is a nonuniversal constant.

4.3. Symmetries

Through several considerations, we obtained a field-theory formula for spin operators, Eqs. (51) and (64). Using the formula, transformations (44)–(46) and the effective Hamiltonians (41) and (42), we can consider how the symmetries of the spin-1 AF chain (4) are represented in the partially-bosonized effective theory framework.

(i) As we discussed already, a U(1) rotation around the spin z axis,  $S_j^+ \rightarrow e^{i\phi} S_j^+$ , could be realized by

$$S_j^+ \rightarrow e^{i\phi} S_j^+; \quad S_j^z \rightarrow S_j^z + \frac{\phi}{2\pi} \quad (66)$$

(ii) The one-site translation  $S_j \rightarrow S_{j+1}$  may correspond to

$$\begin{aligned} \psi_L(x) &\rightarrow \psi_L(x+a_0); & \psi_R(x) &\rightarrow \psi_R(x+a_0); \\ \psi_L(x) &\rightarrow \psi_L(x+a_0) + \frac{p}{M}; & \psi_R(x) &\rightarrow \psi_R(x+a_0) - \frac{p}{M}; \end{aligned} \quad (67)$$

These transformations cooperatively change the staggered factor  $(-1)^j$  in front of  $N^z(x)$  into  $(-1)^{j+1}$ .

(iii) The site-parity transformation  $S_j \rightarrow S_{-j}$  might be regarded as [51]

$$\begin{aligned} \psi_L(x) &\rightarrow i \psi_L(-x); & \psi_R(x) &\rightarrow \psi_R(-x); \\ \psi_R(x) &\rightarrow i \psi_R(-x); & \psi_L(x) &\rightarrow \psi_L(-x); \end{aligned} \quad (68)$$

These symmetry operations might tell us which additional terms are allowed to be present in formulas (51) and (64): for example, the staggered component  $N^z(x)$  may include  $(\psi_L + \psi_R) e^{in\frac{p}{M}} + \text{h.c.}$ ,  $(\psi_L - \psi_R) e^{iy}$ , etc ( $n$ : an integer). These terms are properly transformed for all the symmetry operations.

If we integrate out massive fields in the partition function and replace all massive-field parts in spin operators with their expectation values, the effective Hamiltonian becomes the Gaussian model (42) and the reduced spin operators containing only massless boson fields are given by

$$\begin{aligned} J^z(x) &\rightarrow \frac{M}{a_0} \frac{1}{p} \cos \frac{p}{4} + C_0^z \cos \frac{p}{4} - 2M \psi_L \psi_R + \dots; \\ N^+(x) &\rightarrow e^{i\frac{p}{M}} h^3 i D_1 + D_2 \cos \frac{p}{4} - 2M \psi_L \psi_R + \dots; \\ (J^+(x))^2 &\rightarrow e^{2i\frac{p}{M}} C_1 + C_2 \cos \frac{p}{4} - 2M \psi_L \psi_R + \dots; \\ (N^z(x))^2 &\rightarrow D_1 + D_2 \cos \frac{p}{4} - 2M \psi_L \psi_R + \dots; \end{aligned} \quad (69)$$

where  $C_n$  and  $D_n$  are nonuniversal constants, all the Klein factors are naively eliminated, and we used  $(\psi_L)^2 = \text{const}$ ,  $h^3 i \psi_L \psi_R = O(a_0^{-1})$ , and  $h^3 \psi_L \psi_R = O(a_0^{-1})$  (see Appendix B). This result indicates that the compactification radius of  $\psi_L$  and that of  $\psi_R$  are respectively  $l = \frac{p}{4}$  and  $l = \frac{p}{2}$ . In this bosonized-theory framework, we could interpret that the U(1) rotation  $S_j^+ \rightarrow e^{i\phi} S_j^+$ , the one-site translation  $S_j \rightarrow S_{j+1}$ , and

the site-parity operation  $S_j \rightarrow S_{-j}$ , respectively, corresponds to

$$S_j \rightarrow S_{-j}; \quad (70a)$$

$$S_j \rightarrow S_{-j} + \frac{p}{2}; \quad S_j \rightarrow S_{-j} - \frac{p}{2} M; \quad (70b)$$

$$S_j \rightarrow S_{-j}; \quad S_j \rightarrow S_{-j} + \frac{p}{2}; \quad (70c)$$

The symmetries of Eq. (70) strongly restrict the emergence of the relevant vertex operators in the effective field theory (42): the  $U(1)$  symmetry (70a) [the translational symmetry (70b)] forbids  $e^{in\frac{p}{2}}$  [ $e^{in\frac{p}{4}}$ ] to exist in the effective Hamiltonian [57]. Consequently, the TLL state stably remains.

One can see that the above transformations (70) are very similar to those of the effective bosonization theory for spin-1/2 AF chains (see Appendix C).

#### 4.4. Asymptotic behavior of spin correlation functions

Applying the formulas (51) and (64), let us investigate equal-time spin correlation functions in the field-induced TLL phase. The asymptotic forms of equal-time two-point functions of  $J$  and  $N$  are evaluated as

$$\begin{aligned} \langle J^3(x)J^3(0) \rangle &= \frac{M^2}{a_0} + \frac{1}{h} \langle \xi_x \xi_x \rangle(0) \\ &+ A^2 \cos \left( 2M \frac{x}{a_0} \right) \langle e^{i\frac{p}{4}x} e^{i\frac{p}{4}x} \rangle(0) \\ &= \frac{M^2}{a_0} + \frac{K}{2^2} \frac{1}{x^2} \\ &+ A^2 \cos \left( 2M \frac{x}{a_0} \right) \frac{0}{x^{2K}} + \dots; \end{aligned} \quad (71a)$$

$$\begin{aligned} \langle J^+(x)J^-(0) \rangle &= A \cos \left( M \frac{x}{a_0} \right) \langle e^{i\frac{p}{2}x} e^{-i\frac{p}{2}x} \rangle(0) \\ &+ h_{L=R}^3 \langle \xi_{L=R}^3 \rangle(0) \\ &+ \cos \left( M \frac{x}{a_0} \right) \frac{0}{x^{\frac{K+K}{2}-1}} K_1 \frac{x}{c} + \dots; \end{aligned} \quad (71b)$$

$$\begin{aligned} \langle N^3(x)N^3(0) \rangle &= B^2 \cos \left( M \frac{x}{a_0} \right) \langle h^3(x) \rangle(0) \\ &+ \langle e^{i\frac{p}{2}x} e^{i\frac{p}{2}x} \rangle(0) \\ &+ \cos \left( M \frac{x}{a_0} \right) \frac{0}{x^{K-2}} K_0 \frac{x}{c} + \dots; \end{aligned} \quad (71c)$$

$$\begin{aligned} \langle N^+(x)N^-(0) \rangle &= B_1 \langle h^3(x) \rangle(0) \langle e^{i\frac{p}{2}x} e^{-i\frac{p}{2}x} \rangle(0) \\ &+ B_2 \cos \left( 2M \frac{x}{a_0} \right) \langle e^{i\frac{p}{4}x} e^{i\frac{p}{4}x} \rangle(0) \\ &+ \frac{0}{x^{1-(2K)}} + B_2^0 \cos \left( 2M \frac{x}{a_0} \right) \\ &+ \frac{0}{x^{\frac{1}{2K}+2K}} + \dots; \end{aligned} \quad (71d)$$

where  $A_{(n)}^{(z)}$  and  $B_{(n)}^{(z;0)}$  are nonuniversal constants,  $K$  is the modified Bessel function, and  $x = v_0/m$ . In the calculation of Eq. (71), we used Eqs. (10) [see the comment [48]], (A.3), (B.3)–(B.5). From this result, longitudinal and transverse spin-spin correlators are determined as

$$\begin{aligned} \langle hS_j^z S_0^z \rangle &= M^2 \frac{K}{2^z} \frac{a_0^2}{x} + C_0^z \cos(2Mj) \frac{a_0^{2K}}{x} \\ &+ D_1^z \cos((1+M)j) \frac{a_0^{(K+1)2}}{x} e^{-x} + \dots \end{aligned} \quad ; \quad (72a)$$

$$\begin{aligned} \langle hS_j^+ S_0^- \rangle &= C_1 \cos(Mj) \frac{a_0^{(K+K-1+1)2}}{x} e^{-x} \\ &+ D_1 (1)^j \frac{a_0^{1=(2K)}}{x} \\ &+ D_2 \cos((1+2M)j) \frac{a_0^{\frac{1}{2K}+2K}}{x} + \dots \end{aligned} \quad ; \quad (72b)$$

where  $x = ja_0/m$ , and  $C_q^{(z)}$  and  $D_q^{(z)}$  are nonuniversal constants which are related to  $C_n^{(z)}$  and  $D_n^{(z;0)}$  in Eqs. (51) and (64). The first two terms in  $\langle hS_j^z S_0^z \rangle$  and the second term in  $\langle hS_j^+ S_0^- \rangle$  can also be derived by the NLSM plus Ginzburg-Landau approach [23, 24], but it is difficult to obtain all the other terms within the same approach. One can verify that the critical exponent of the incommensurate part around wave number  $p = 2k$  ( $k_F = M/a_0$ ) in  $\langle hS_j^z S_0^z \rangle$ ,  $z = 2K$ , and that of the staggered part in  $\langle hS_j^+ S_0^- \rangle$ ,  $z = 1=(2K)$ , satisfy the famous relation  $\nu_z = 1$  [58, 59, 23, 29, 10]. In addition, it is found that the contribution around  $p = a_0/k_F$  in  $\langle hS_j^z S_0^z \rangle$  and that around  $p = k$  in  $\langle hS_j^+ S_0^- \rangle$  exhibit an exponential decay. Comparing Eq. (72) with the spin correlators of a two-leg spin-1/2 ladder in a uniform field in Ref. [29] would be instructive (although our calculations in Eqs. (71) and (72) are rougher than those in Ref. [29]). Besides Eq. (72), using the formulas (51) and (64), one can calculate various physical quantities (susceptibilities, dynamical structure factors, NMR relaxation rates, etc) in the TLL phase [24].

In all the calculations of this subsection, we assumed that the three systems  $\hat{H}[\mu]$ ,  $\hat{H}[\mu^3]$ , and  $\hat{H}[\mu^2]$  are independent of each other. Although small interactions among these systems would actually be present, it is expected that their effects in the low-energy, long-distance physics are almost negligible and could be absorbed into some parameters such as  $K$ ,  $v_0$ ,  $m$ , etc.

## 5. Some applications of the field-theory representation of spin operators

In this section, utilizing the derived formulas (51) and (64), we discuss some topics for the low-energy physics in/around the uniform-field-induced TLL phase.

### 5.1. String order parameter

As a quantity characterizing the Haldane phase ( $H < m$ ), there is the nonlocal string order parameter which can detect the so-called "hidden AF long-range order" in the phase [60, 61, 8]. It is defined by

$$O(i; j) = \lim_{j \rightarrow \infty} \frac{1}{j!} \sum_{j_1, \dots, j_{j-1}} S_i \exp \left( i \sum_{n=i+1}^{j-1} S_n \right) S_j \quad ; \quad (73)$$

In the continuous-eld-theory framework, we can predict [41, 40] that the string parameter is approximated as follows:

$$O_z(i; j) = h^{-1} (x_i)^{-2} (x_i)^{-1} (x_j)^{-2} (x_j)^{-1} i; \quad (74)$$

where  $x_i = ia_0$  and  $x_j = ja_0$ . Actually, the right-hand side is a finite, non-zero value in the Haldane phase where  $h \neq 0$ .

From the formula (A.6), the z component of the string parameter is rewritten as

$$O_z(i; j) = h \cos\left(\frac{p}{2} (x_i)\right) \cos\left(\frac{p}{2} (x_j)\right) i; \quad (75)$$

In the eld-induced TLL phase ( $H > m$ ), the vertex operator  $\exp\left(\frac{p}{2} (x)\right)$  is represented by using elds  $(; ; )$  [see Eqs. (63b) and (64a)]. Hence, it is predicted that in the TLL phase,  $O_z(i; j)$  behaves as

$$O_z(i; j) = h \cos\left(\frac{p}{2} (x_i) - M i\right) \cos\left(\frac{p}{2} (x_j) - M j\right) i + \cos\left[M (i - j)\right] \frac{a_0}{x_i - x_j} + \dots : \quad (76)$$

namely,  $O_z(i; j)$  is shown to exhibit power decay in the TLL phase.

## 5.2. SU(2)-invariant perturbations

In Secs. 5.2-5.4, we investigate typical perturbations for the critical TLL state. In particular, we focus on whether or not the perturbation terms yield a first excitation gap, and the symmetries of them.

In this subsection, we discuss two terms: the bond alternation  $\sum_j (1)^j J S_j S_{j+1}$  ( $j, j+1$ ) and the next-nearest-neighbor (NNN) exchange  $\sum_j J_2 S_j S_{j+2}$  ( $j, j+2$ ), which are invariant under the global SU(2) transformation. Because a global U(1) symmetry, a part of the SU(2) one, is usually necessary for the realization of the TLL i.e., a  $c = 1$  CFT [1, 2, 4, 8], and (as we mentioned in Sec. 4.3) it prohibits all vertex operators with the dual eld  $(x)$  from emerging in the effective Hamiltonian, we expect, without any calculations, that the TLL phase survives even when these perturbations are applied. Let us consider the two terms below in more detail.

From the continuous-eld formula (7), we can expect that the bond alternation term is approximated as

$$\begin{aligned} \sum_j (1)^j J S_j S_{j+1} &= \sum_j J (1)^j \frac{1}{2} S_j^+ S_{j+1} + h \mathcal{X} + S_j^z S_{j+1}^z \\ &= \sum_j \int dx \left( (1)^j \frac{h}{2} J^+(x) J^-(x+a_0) + h \mathcal{X} + J^z(x) J^z(x+a_0) \right. \\ &\quad \left. + \frac{1}{2} N^+(x) N^-(x+a_0) + h \mathcal{X} + N^z(x) N^z(x+a_0) \right. \\ &\quad \left. + \frac{1}{2} J^+(x) N^-(x+a_0) + N^+(x) J^-(x+a_0) + h \mathcal{X} \right. \\ &\quad \left. + J^z(x) N^z(x+a_0) + N^z(x) J^z(x+a_0) \right) : \end{aligned} \quad (77)$$

Let us substitute the formulas (51) and (64) into the above result, although such a procedure sometimes causes mistakes (see the comment [55]). As a result, we obtain

$$\sum_j (1)^j J S_j S_{j+1} = \sum_j \int dx \sum_{n=1}^{\infty} \frac{h}{n} \cos\left(\frac{p}{4} (x) - 2 M j\right) + \dots$$

$$+ (1)^j \tilde{n} \cos(n \frac{p}{4} - 2Mj) + \tilde{n}^i + \dots; \tag{78}$$

where  $\tilde{n}^{(0)}$  and  $\tilde{n}^{(i)}$  are nonuniversal constants. Here, we used  $(\delta(x))^2 = \text{const}$ , and the operator product expansion (OPE)  $\delta_L \delta_R = \delta + \dots$  [4, 62], and then integrated out all the massive fields in the partition function (see the comment [53]). In order to more restrict the form of Eq. (78), we utilize the symmetry argument. For the one-site translation and the site-parity operation, the bond alternation term changes its sign. Its bosonized form should also have the same property; namely, we require the right-hand side in Eq. (78) to change the sign for  $\delta(x) \rightarrow \delta(x + a_0) e^{-i\pi M}$  and  $\delta(x) \rightarrow \delta(-x)$  [see Eq. (70)]. Consequently, we set  $\tilde{n}^0 = 0$  and  $\tilde{n} = -2$ . The resultant form of Eq. (78) is

$$\sum_j (1)^j J S_j S_{j+1} \dots J \int_{-\infty}^{\infty} dx \sum_{n=1}^{\infty} (1)^j \tilde{n} \sin(n \frac{p}{4} - 2Mj) + \dots; \tag{79}$$

This bosonized form indicates that in general, the bond alternation is irrelevant in the TLL phase due to the staggered factor  $(1)^j$  and the phase  $2\pi \frac{M}{4} j$ . However, when  $M = 1/2$  (half of the saturation), a relevant interaction  $\sin(\frac{p}{4})$  originates from the  $n = 1$  term in Eq. (79) because of the cancellation of two factors  $(1)^j$  and  $2Mj$ . The scaling dimension  $\nu$  of  $\sin(\frac{p}{4})$  is  $K$  ( $1 < K < 1.5$  [25]). At this case of  $M = 1/2$ , the low-energy physics may be described by a sine-Gordon theory, and an infinitesimal bond alternation induces a finite excitation gap and a finite dimerization parameter  $h S_{2n} S_{2n+1} S_{2n+1} S_{2n+2} i$ . The existence of the gap further means that the bond alternation brings an  $M = 1/2$  plateau in the uniform magnetization process. The scaling argument near a criticality [63] shows that the bond-alternation-induced gap and dimerization parameter respectively behave as

$$j^{-1/(2-K)}; \tag{80}$$

$$h S_{2n} S_{2n+1} S_{2n+1} S_{2n+2} i \sim \text{sgn}(\tilde{J}) j^{K/(2-K)};$$

for a small  $j$ . Because of the inequality  $1/(2-K) > 1$ , the gap gradually grows with increasing  $j$ . These predictions for the (small) bond alternation are consistent with previous numerical [64] and analytical [65] works.

Next, let us turn to the NNN coupling ( $J_2$ ) term. Through an argument similar to that above, we arrive in the following result:

$$\sum_j J_2 S_j S_{j+2} \dots J_2 \int_{-\infty}^{\infty} dx \frac{a_0}{p} (\partial_x)^2 \frac{2M}{p} \partial_x + \sum_{n=1}^{\infty} \tilde{n} \cos(n \frac{p}{4} - 2Mj) + \dots; \tag{81}$$

where  $\tilde{n}$  is a nonuniversal constant. We used the symmetry argument: the right-hand side in Eq. (81) is invariant under  $\delta(x) \rightarrow \delta(x + a_0) e^{-i\pi M}$  and  $\delta(x) \rightarrow \delta(-x)$ . Since the bosonized NNN coupling does not contain any relevant operators for arbitrary magnetization values, we conclude that the TLL phase remains even when a sufficiently small NNN exchange perturbation is introduced. The derivative term  $\partial_x$  is absorbed into the Gaussian part via  $\int_{-\infty}^{\infty} dx \text{const} \int dx$  and provides a small correction of the magnetization  $M$ . While the boson quadratic term  $(\partial_x)^2$  makes the velocity  $v$  and

the TLL parameter  $K$  modify. After easy calculations, we can see that when  $J_z > 0$  ( $J_z < 0$ ),  $M$  and  $K$  decrease (increase), but  $v$  increases (decreases).

### 5.3. Axially symmetric terms

In this subsection, we consider three kinds of  $U(1)$ -symmetric perturbation terms: the single-ion anisotropy  $D_z \sum_j (S_j^z)^2$  (so-called  $D$  term), the XXZ type anisotropy  $J_z \sum_j S_j^z S_{j+1}^z$ , and the staggered-eld Zeeman term along the spin  $z$  axis  $h_z \sum_j (-1)^j S_j^z$  ( $\mathcal{D}_z \sum_j \mathfrak{h}_z j - J$  and  $j_z j - 1$ ). As in the cases of the bond alternation and the NNN coupling, there is a high possibility that the TLL phase survives as these perturbations are added.

Following the similar argument to that in the last subsection, we can bosonize the three terms as follows:

$$D_z \sum_j (S_j^z)^2 \rightarrow D_z \int dx \frac{n}{a_0} (\partial_x \phi)^2 - \frac{2M}{\pi} \partial_x \phi + \sum_{n=1}^{\infty} d_n \cos(n \frac{\pi}{4} (2M - j)) + \dots; \quad (82a)$$

$$J_z \sum_j S_j^z S_{j+1}^z \rightarrow J_z \int dx \frac{n}{a_0} (\partial_x \phi)^2 - \frac{2M}{\pi} \partial_x \phi + \sum_{n=1}^{\infty} \tilde{d}_n \cos(n \frac{\pi}{4} (2M - j)) + \dots; \quad (82b)$$

$$h_z \sum_j (-1)^j S_j^z \rightarrow h_z \int dx \sum_{n=1}^{\infty} (-1)^j z_n \cos(n \frac{\pi}{4} (2M - j)) + \dots; \quad (82c)$$

where  $d_n$ ,  $\tilde{d}_n$  and  $z_n$  are nonuniversal constants. For instance, we required the bosonization form (82c) of the staggered-eld term to change its sign for the one-site translation  $\phi(x) \rightarrow \phi(x + a_0) - \frac{\pi}{2M}$  and to be invariant under the site-parity transformation  $\phi(x) \rightarrow \phi(-x)$ . The results (82a) and (82b) suggest that the  $D$  term and the XXZ exchange play almost the same roles in the low-energy, long-distance physics of the TLL phase. If  $D_z > 0$  ( $D_z < 0$ ),  $M$  and  $K$  decrease (increase), while  $v$  increases (decreases). As expected, these three terms generally do not destroy the TLL state. However, as in the case of the bond alternation term, when the uniform magnetization  $M$  becomes close to  $1/2$ , the staggered-eld term involves a relevant term  $\cos(\frac{\pi}{4})$  with  $s = K$ . Therefore, a staggered-eld-induced gap  $h_z$  opens and  $h_z S_j^z$  obtains a staggered component at  $M = 1/2$ . From the standard scaling argument, the gap and the magnetization are shown to behave as

$$h_z \sim \mathfrak{h}_z j^{-(2-K)}; \quad h_z S_j^z \sim \frac{1}{2} + (-1)^j f \frac{H}{J} \text{sgn}(h_z) \mathfrak{h}_z j^{K-(2-K)}; \quad (83)$$

where  $f$  is a function.

5.4. Axial-symmetry-breaking terms

Here, we discuss three axial-symmetry-breaking terms: the  $D_x$  term with the  $x$  component of spin  $D_x = \sum_j (S_j^x)^2$ , another kind of the single-ion anisotropy (so-called E term)  $E = \sum_j [(S_j^x)^2 - (S_j^y)^2] = \frac{E}{2} \sum_j [(S_j^+)^2 + (S_j^-)^2]$ , and the staggered-field term along the spin  $x$  axis  $h_x = \sum_j (-1)^j S_j^x$  ( $\mathcal{D}_x = \sum_j \mathcal{E}_j \mathcal{J}_x$ ,  $\mathcal{J} = J$ ). Since the three terms destroy the axial  $U(1)$  symmetry (i.e., vertex operators with  $(x)$  are allowed to be present in the effective Hamiltonian. It is inferred that such vertex operators cause an instability of the TLL state, and then a finite excitation gap occurs. Here, note that the  $D_x$  and E terms are invariant under the rotation  $S_j^+ \rightarrow S_j^+ e^{i\phi}$  (i.e.,  $P^-$ ) [20], whereas the  $h_x$  term obtains a minus sign via the same rotation. Furthermore, the  $\pi$ -rotation leaves the E term change the sign.

Through some calculations, the three perturbation terms are bosonized as

$$D_x = \sum_j (S_j^x)^2 = D_x \int dx \cos\left(\frac{p}{4}\right) \sum_{n=0}^{\infty} d_n^{(1)} \cos\left(n\left(\frac{p}{4} - 2Mj\right)\right) + \cos\left(4\frac{p}{4}\right) \sum_{n=0}^{\infty} d_n^{(2)} \cos\left(n\left(\frac{p}{4} - 2Mj\right)\right) + \sum_{n=0}^{\infty} d_n^{(3)} \cos\left(n\left(\frac{p}{4} - 2Mj\right)\right) + \dots; \quad (84a)$$

$$E = \sum_j [(S_j^x)^2 - (S_j^y)^2] = E \int dx \cos\left(\frac{p}{4}\right) \sum_{n=0}^{\infty} e_n \cos\left(n\left(\frac{p}{4} - 2Mj\right)\right) + \dots; \quad (84b)$$

$$h_x = \sum_j (-1)^j S_j^x = h_x \int dx \cos\left(\frac{p}{4}\right) \sum_{n=0}^{\infty} x_n \cos\left(n\left(\frac{p}{4} - 2Mj\right)\right) + \dots; \quad (84c)$$

where  $d_n^{(1)}$ ,  $e_n$  and  $x_n$  are nonuniversal constants. (Since the bosonized form of the anisotropic exchange  $\mathcal{J} = \sum_j S_j^x S_{j+1}^x$  is the same type as Eq. (84a), we do not discuss it here.) One should note the following properties: (i)  $\cos\left(\frac{p}{4}\right) = \cos\left(\frac{p}{4}\right)$ ,  $\cos\left(\frac{p}{4}\right) = \cos\left(\frac{p}{4}\right)$  and  $\cos\left(4\frac{p}{4}\right) = \cos\left(4\frac{p}{4}\right)$  for the  $\pi$ -rotation, and (ii)  $\cos\left(\frac{p}{4}\right) = \cos\left(\frac{p}{4}\right)$  for the  $\pi$ -rotation.

The most relevant operator in both  $D_x$  and E terms is always  $\cos\left(\frac{p}{4}\right)$  with  $s = 1 - K < 2$ . Thus, the low-energy properties can be explained by a sine-Gordon model, and a gap emerges. Supposing  $d_0^{(1)}$  and  $e_0$  are positive, the potential  $\cos\left(\frac{p}{4}\right)$  pins the phase field to  $\frac{p}{4} = 2\pi(0 \text{ or } \pi)$  modulo  $2\pi$  for  $D_x; E > 0$  ( $D_x; E < 0$ ). In such a case of  $\frac{p}{4} = 2\pi(0 \text{ or } \pi)$ , it is expected that  $h S_j^x i = (-1)^j \cos\left(\frac{p}{4}\right) i = 0$  ( $\neq 0$ ) and  $h S_j^y i = (-1)^j \sin\left(\frac{p}{4}\right) i = 0$  ( $\neq 0$ ). From this prediction and the scaling argument [ $s$  of  $e^{i\frac{p}{4}}$  is  $1 - (4K)$ ], we conclude that for a small  $D_x$  term, the gap  $\Delta_{D_x}$  and the transverse component of the spin moment increase as follows:

$$\Delta_{D_x} \propto \mathcal{D}_x J^{K/(2K-1)};$$

$$\begin{aligned}
 \begin{aligned}
 hS_j^x i &= 0 \\
 hS_j^y i &= (1) D_x^{1=(8K-4)} ; \quad (\text{for } D_x > 0); \\
 hS_j^x i &= (1) D_x^{1=(8K-4)} \\
 hS_j^y i &= 0 ; \quad (\text{for } D_x < 0):
 \end{aligned}
 \end{aligned}
 \tag{85}$$

Of course, for a small E term, the similar results hold: we may replace  $D_x$  to E in Eq. (85). The staggered moment (i.e., a Neel order) along the spin x or y axis shows that the  $D_x$  (or E) term causes the spontaneous breakdown of the one-site translational symmetry. Because of  $K = (2K - 1) < 1$  and  $1 = (8K - 4) < 1$ , the  $D_x$ - or E-term-induced gap and the transverse moment  $hS_j^{xy} i$  rapidly increases with the growth of  $D_x$  or E.

The staggered-eld  $h_x$  term contains the relevant term  $\cos(\frac{p}{s})$  with  $s = 1 = (4K) < 2$ . Therefore, the  $h_x$ -induced gap  $h_x$  and the staggered magnetization are shown to behave as

$$h_x \sim J_x^{4K = (8K - 1)}; \quad hS_j^x i \sim (1) \text{sgn}(h_x) J_x^{1=(8K - 1)}; \tag{86}$$

For the spin-1/2 AF Heisenberg chain, which low-energy sector is also described by a TLL theory (see Introduction), a staggered-eld also yields a gap and a staggered moment. Oshikawa and Aleck [66] show that in the spin-1/2 case,  $h_x \sim J_x^{2=3}$  and  $hS_j^x i \sim (1) \text{sgn}(h_x) J_x^{1=3}$ . The result (86) thus indicates that the growth of both the gap and the staggered moment in the spin-1 case is much sharper than that in the spin-1/2 case.

### 5.5. Magnon decay induced by axial-symmetry-breaking terms

In this subsection, we briefly mention the magnon-decay processes, which have already in some detail discussed in Ref. [20]. In order to consider such processes, let us go back to the effective Hamiltonian (36), where  $\sim_+$ ,  $\sim_0^y$  and  $\sim^y$  respectively denote the  $S^z = +1, 0$  and  $-1$  magnon creation operators. The marginally irrelevant term, omitted in Eq. (36), just contribute to the magnon decay.

First, we focus on the U(1)-symmetric AF chain (4) without any perturbations. Since the one-magnon excitations are present only around  $p = \pi - a_0$ , only the decay from a magnon to an odd number of magnons is possible. If H is increased so that  $3_+(0) < 0_-(0)$  ( $H > m=2$ ) [ $3_+(0) < 0_-(0)$  ( $H > 2m=3$ )] are satisfied, a magnon  $\sim^y$  [ $\sim_0^y$ ] is energetically permitted to decay into three magnons  $\sim_+ \sim_+ \sim_+$  via the term. However, since  $\sim_+$ ,  $\sim_0^y$  and  $\sim^y$  possess different eigenvalues of  $S^z$  (namely, three kinds of one-magnon states are in different sectors of the Hilbert space), this type of the decay is forbidden. Indeed, for a U(1) rotation  $S_j^+ \rightarrow e^{i\theta} S_j^+$ ,  $\sim_+ [\sim_0]$  is transformed as  $\sim_+ \rightarrow e^{i\theta} \sim_+ [\sim_0 \rightarrow \sim_0]$  (invariant), while the product  $\sim_+ \sim_+ \sim_+$  obey a different rotation  $\sim_+ \sim_+ \sim_+ \rightarrow e^{3i\theta} \sim_+ \sim_+ \sim_+$ . Three kinds of magnons ( $\sim_+$ ,  $\sim_0^y$ ,  $\sim^y$ ) therefore would be well-defined quasiparticles in the U(1)-symmetric system (4).

On the other hand, when a U(1)-symmetry-breaking term is introduced, there is a possibility that the above decay processes are allowed. In the case with the  $D_x$  or E terms, the  $\pi$ -rotation symmetry ( $S_j^+ \rightarrow S_j^+$ ;  $\theta = \pi$ ), a part of the U(1) rotation, survives. For this rotation,  $\sim_+$  and  $\sim_+ \sim_+ \sim_+$  are odd, whereas  $\sim_0$  is even. Furthermore, from Eqs. (38) and (39), we find that for example a sum of four terms  $\sim(k_1) \sim(k_2) \sim(k_1) \sim(k_2) \sim(k_{1,2}) \sim(k_{1,2}) + h.c.$  is invariant under both the one-site translation and the site-parity operation. (Because the site-parity operation (39) is the result of the approximation, the requirement of the invariance

under this transformation might be too strong.) As a result, the process  $\sim^y \rightarrow \sim_+ \sim_+ \sim_+$  is permissible. It is hence inferred that as sufficiently strong  $D_x$  or  $E$  terms are present in the system,  $\sim^y$  magnons become ill-defined particles, and we should eliminate the magnon fields from the Hamiltonian (41) and the field-theory formulas of the spin, Eqs. (51) and (64). In the case with the  $h_x$  staggered-field term, symmetries of the rotation and the one-site translation are also broken (the two-site translational symmetry remains). Therefore, the magnon decay would be more promoted. When  $H$  is further increased and the  $\sim_+$ -magnon condensation occurs ( $H > m$ ), other types of the magnon decay are energetically admitted: for instance,  $\sim^y \rightarrow \sim_0^y \sim_0^y \sim_+ \sim_+$ .

From the simple discussion above, one sees that as  $H$  is sufficiently strong, a large axial-symmetry-breaking perturbation tends to make the lifetime of massive magnons shorten. However, if such a perturbation is small enough, our effective theory framework would still be reliable and have the ability to explain various low-energy properties of spin-1 AF chains.

## 6. Summary

In this paper, based on the Majorana fermion theory, we have reconsidered the field theory description of the spin-1 AF chain (4), and derived an explicit field-theory form of spin operators in the uniform-field-driven TLL phase in the chain (4), i.e., Eqs. (51) and (64) [the corresponding effective Hamiltonian is Eqs. (41) and (42)]. From the formula, we have completely determined the asymptotic forms of spin correlation functions (Sec. 4.4). Furthermore, applying the formula, we have investigated the string order parameter and effects of some perturbation terms (the bond alternation, the next-nearest interaction, anisotropy terms) in Sec. 5. We have estimated the excitation gaps and some physical quantities (staggered moments and the dimerization parameter) generated from the perturbations. From Sec. 2 to Sec. 5, we have often argued how symmetries of the spin-1 AF chain are represented in the effective field theory world.

Our results, especially Eqs. (51) and (64), must be useful in analyzing and understanding various spin-1 AF chains with magnons condensed (i.e., with a finite magnetization) and the extended models of them (e.g., spin-1 AF ladders, spatially anisotropic 2D or 3D spin-1 AF systems). The results of Sec. 5 guarantee this expectation. We will apply the contents of this paper to other spin-1 AF systems in the near future.

Determining nonuniversal coefficients in Eqs. (51), (64) and (69), especially those in front of terms including only massless bosons, is important for more quantitative predictions of (quasi) 1D spin-1 systems. A powerful way of the determination is to accurately evaluate the long-distance behavior of spin correlation functions by means of a numerical method such as DMRG and QMC [10].

## Acknowledgments

This work is supported by a Grant-in-Aid for Scientific Research (B) (No. 17340100) from the Ministry of Education, Culture, Sports Science and Technology of Japan.

Appendix A. Abelian bosonization for fermion systems

Here, we briefly summarize the Abelian bosonization. As mentioned in Sec. 2, in (1+1)D case, a massless Dirac (complex) fermion or two species of massless Majorana (real) fermions (critical Ising models) are equivalent to a massless bosonic Gaussian theory. The former Hamiltonian is written as

$$\begin{aligned} \hat{H}[\psi] &= \int dx \, i c \left( \psi_L^\dagger \partial_x \psi_L - \psi_R^\dagger \partial_x \psi_R \right) \\ &= \int dx \sum_{q=1,2} \frac{i}{2} c \left( \psi_L^q \partial_x \psi_L^q - \psi_R^q \partial_x \psi_R^q \right) = \hat{H}[\phi^1; \phi^2]; \end{aligned} \quad (A.1)$$

where  $\psi_L^q$  and  $\psi_R^q$  are respectively the chiral components of the Dirac fermion and the real one, and  $c$  is the Fermi velocity. These fermions obey anticommutation relations:  $\psi_L^q(x); \psi_L^q(y) = 0$ ;  $\psi_L^q(x); \psi_L^q(y) = 0$  and  $\psi_L^q(x); \psi_L^q(y) = 0$ ;  $\psi_L^q(x); \psi_L^q(y) = 0$ . The corresponding Hamiltonian of the Gaussian theory is

$$\hat{H}[\phi] = \int dx \, \frac{c}{2} \left( (\partial_x \phi)^2 + (\partial_x \phi)^2 \right) = \int dx \, c \left( (\partial_x \phi_L)^2 + (\partial_x \phi_R)^2 \right); \quad (A.2)$$

where  $\phi = \phi_L + \phi_R$  is the real scalar field,  $\phi_L = \phi_L - \phi_R$  is the dual field of  $\phi$ , and  $\phi_{L(R)}$  is the left (right) moving part of  $\phi$ . The fields  $\phi$  and  $\phi_L, \phi_R$  satisfy the canonical commutation relation  $[\phi(x); \phi(y)] = \frac{i}{2} \text{sgn}(x-y)$ . Chiral fields  $\phi_{L,R}$  obey  $[\phi_L(x); \phi_R(y)] = 0$  and  $[\phi_{L=R}(x); \phi_{L=R}(y)] = \frac{i}{4} \text{sgn}(x-y)$ . In condensed-matter physics, the Hamiltonian (A.1) usually originates from a microscopic system in solids (e.g., a lattice system such as the Hubbard chain and the Heisenberg one) after a coarse-graining or a renormalization procedure.

Among these fermion and boson fields, operator identities hold. The fermion annihilation (or creation) operators are bosonized as

$$\psi_L = \frac{1}{\sqrt{2}} \exp\left(i \frac{p}{4} \phi_L\right); \quad \psi_R = \frac{1}{\sqrt{2}} \exp\left(i \frac{p}{4} \phi_R\right); \quad (A.3)$$

where  $\psi_{L,R}$  are Klein factors which satisfy  $\psi_L^2 = 2$ ,  $\psi_R^2 = 2$ , and are necessary for the boson vertex (exponential) operators to reproduce the correct anticommutation relation between  $\psi_L$  and  $\psi_R$ . The parameter  $p$  is a short-distance cutoff, which depends on details of the microscopic model considered. (Note that it is possible to construct another formula without Klein factors, although it requires a modification of commutation relations among bosons. See Refs. [67, 68].) Following Haldane's harmonic-uid approach [69, 3, 10], one can obtain an alternative bosonized form of  $\psi_L$  and  $\psi_R$ : when a real-space fermion field  $\psi(x)$  in the considering microscopic system is approximated as  $e^{ik_F x} \psi_R + e^{ik_F x} \psi_L$ , one may bosonize  $\psi_{L,R}$  as

$$\begin{aligned} e^{ik_F x} \psi_L &= \frac{1}{\sqrt{2}} \sum_{n=0}^{\infty} e^{i(2n+1)(\phi^- + k_F x)} e^{i \frac{p}{4} \phi^-}; \\ e^{ik_F x} \psi_R &= \frac{1}{\sqrt{2}} \sum_{n=0}^{\infty} e^{i(2n+1)(\phi^- + k_F x)} e^{i \frac{p}{4} \phi^-}; \end{aligned} \quad (A.4)$$

The quantity  $k_F$  is the Fermi wave number in the microscopic system. The  $n=0$  most relevant terms correspond to Eq. (A.3). The chiral U(1) currents  $J_L = \frac{1}{2} \psi_L^\dagger \psi_L$ :

and  $J_R = : \psi_R^\dagger \psi_R :$  are written as

$$J_L = \frac{1}{2} \psi_L^\dagger \psi_L; \quad J_R = \frac{1}{2} \psi_R^\dagger \psi_R; \quad (A.5)$$

In addition to the correspondences between the fermion and the boson, it is known [1, 4, 28, 29, 62, 70] that the Ising order and disorder fields can be bosonized as [71]

$$\begin{aligned} \sigma &\sim \cos\left(\frac{\phi}{2}\right); & \psi &\sim \cos\left(\frac{\phi}{2}\right); \\ \psi &\sim \sin\left(\frac{\phi}{2}\right); & \sigma &\sim \sin\left(\frac{\phi}{2}\right); \end{aligned} \quad (A.6)$$

Following these Abelian bosonization rules, one can bosonize 1D interacting Dirac fermion systems as well as the free massless fermion (A.1). If the interaction terms are all irrelevant in the sense of the renormalization group, the effective Hamiltonian at the low-energy limit is still a Gaussian type with the velocity  $c$  and coefficients of  $(\partial_x \phi)^2$  and  $(\partial_x \psi)^2$  renormalized. Conventionally [3], the resultant Hamiltonian is written as

$$\hat{H}^e[\phi] = \int dx \frac{c^0}{2} g (\partial_x \phi)^2 + \frac{1}{g} (\partial_x \psi)^2; \quad (A.7)$$

where  $c^0$  is the renormalized velocity, and  $g$  is called the TLL parameter (the Hamiltonian (A.2) corresponds to a  $g = 1$  theory). When a system is reduced to this type at the low-energy limit, we say that the system belongs to the TLL universality [69, 1, 2, 3]. The Gaussian theory [1, 2, 3, 4, 8] yields

$$\begin{aligned} \langle \psi(x) \psi(0) \rangle &= \frac{K}{2} \frac{1}{x^2}; \\ \langle e^{iq\phi(x)} e^{-iq\phi(0)} \rangle &= \frac{q^2 g = 2}{q^2 q^0} \frac{1}{x}; \\ \langle e^{iq\psi(x)} e^{-iq\psi(0)} \rangle &= \frac{q^2 = (2g)}{q^2 q^0} \frac{1}{x}; \end{aligned} \quad (A.8)$$

where  $q$  and  $q^0$  are an integer (note the comment [50]). This result shows that the scaling dimensions of  $\psi$ ,  $e^{iq\phi}$  and  $e^{iq\psi}$  are respectively 1,  $q^2 g = 4$  and  $q^2 = (4g)$ .

### Appendix B. Correlation functions of massive fermion theories

In this Appendix, we evaluate correlation functions of two massive fermion systems,

$$\begin{aligned} \hat{H}[\psi] &= \int dx \frac{1}{2} v_0 \left[ \psi_L^\dagger \partial_x \psi_L + \psi_R^\dagger \partial_x \psi_R + m \psi_L^\dagger \psi_R \right]; \\ &= \int dx \psi_0^\dagger(k) \tilde{\psi}_0(k) \tilde{\psi}_0(k); \end{aligned} \quad (B.1)$$

$$\begin{aligned} \hat{H}[\phi] &= \int dx \psi^\dagger \left[ \frac{v_0^2}{2m} \partial_x^2 + m + H + O(\partial_x^4) \right] \psi; \\ &= \int dx \psi^\dagger(k) \tilde{\psi}(k) \tilde{\psi}(k); \end{aligned} \quad (B.2)$$

where  $\psi_{L,R}$  is the chiral real fermion field, and other fields  $\psi$  and  $\tilde{\psi}_0$  are defined in Sec. 3.

First, we consider the Majorana fermion system (B.1). At  $T = 0$ ,  $\langle h_{0}^{-y}(k) \sim_{0}(k^0) \rangle = 0$  and  $\langle h_{0}^{-y}(k) \sim_{0}(k^0) \rangle = \langle k, k^0 \rangle$ . Therefore, the two-point function of  $\psi_L^3$  is calculated as

$$\begin{aligned} \langle h_L^3(x) \psi_L^3(0) \rangle &= \frac{1}{L} \sum_{k>0} \frac{1}{2v_0(k)} \left( v_0(k) - kv_0 \right) e^{ikx} + \left( v_0(k) + kv_0 \right) e^{-ikx} \\ &= \frac{1}{2} \langle x \rangle + \frac{1}{2} \int_0^x \frac{dk}{2} \frac{kv_0}{v_0(k)} e^{-ikx} \\ &= \frac{2m}{v_0} K_1(x=c); \quad (\text{at } x < c); \end{aligned} \tag{B.3}$$

where we used Eqs. (31), (34) and (35). Here,  $c$  is the ultraviolet cut off,  $c = v_0 m$ , and  $K$  is the modified Bessel function ( $K(z) \sim \frac{1}{2} e^{-z}$  at  $z \gg 1$ ). Similarly, one can obtain

$$\begin{aligned} \langle h_R^3(x) \psi_R^3(0) \rangle &= \langle h_L^3(x) \psi_L^3(0) \rangle; \\ \langle h_L^3(x) \psi_R^3(0) \rangle &= \frac{m}{2v_0} K_0(x=c); \quad (\text{at } x < c); \end{aligned} \tag{B.4}$$

In another system (B.2), the similar relations  $\langle h^{-y}(k) \sim(k^0) \rangle = 0$  and  $\langle h^{-y}(k) \sim(k^0) \rangle = \langle k, k^0 \rangle$  hold at  $T = 0$ . One hence easily finds

$$\begin{aligned} \langle h^y(x) \psi(0) \rangle &= 0; \\ \langle h(x) \psi^y(0) \rangle &= \langle x \rangle - \frac{x^2}{2}; \end{aligned} \tag{B.5}$$

where  $c$  is the short-distance cut off.

In addition to these results, one can of course compute any correlation functions of the systems (B.1) and (B.2), using Wick's theorem, etc.

### Appendix C. Symmetries of the spin-1/2 XXZ chain

The spin-1/2 XXZ chain (1) has the same global symmetries as those of the spin-1 AF chain (4): the  $U(1)$  rotation around the spin  $z$  axis, the one-site translation, and the site-parity transformation. In the Abelian bosonization framework (see Eqs. (2) and (3)), these three symmetries could be realized by the following transformations of boson fields and [7, 72].

(i) The  $U(1)$  rotation  $S_j^+ \rightarrow e^i S_j^+$  corresponds to

$$\phi \rightarrow \phi + \frac{2\pi}{L}; \tag{C.1}$$

(ii) The one-site translation  $S_j \rightarrow S_{j+1}$  corresponds to

$$\begin{aligned} \phi(x) &\rightarrow \phi(x+a_0) + \frac{2\pi}{L} \left( M + \frac{1}{2} \right); \\ \psi(x) &\rightarrow \psi(x+a_0) + \frac{2\pi}{L}; \end{aligned} \tag{C.2}$$

(iii) The site-parity transformation  $S_j \rightarrow S_{-j}$  corresponds to

$$\begin{aligned} \phi(x) &\rightarrow \phi(-x) + \frac{2\pi}{L}; \\ \psi(x) &\rightarrow \psi(-x); \end{aligned} \tag{C.3}$$

## References

- [1] Gogolin A O, Nersisyan A A and Tsvetlik A M, 1998 *Bosonization and Strongly Correlated Systems* (Cambridge University Press, Cambridge, England).
- [2] Tsvetlik A M, 2003 *Quantum Field Theory in Condensed Matter Physics*, 2nd ed, (Cambridge University Press, Cambridge, England).
- [3] Giamarchi T, 2004 *Quantum Physics in One Dimension* (Oxford University Press, New York).
- [4] Francesco P D, Mathieu P and Senechal D, 1997 *Conformal Field Theory* (Springer-Verlag, New York).
- [5] Korepin V E, Bogoliubov N M, and Izergin A G, 1993 *Quantum Inverse Scattering Method and Correlation Functions* (Cambridge University Press, Cambridge, England).
- [6] Takahashi M, 1999 *Thermodynamics of One-Dimensional Solvable Models* (Cambridge University Press, Cambridge, England).
- [7] A eck I, 1990 in *Fields, Strings and Critical Phenomena*, edited by E. Brezin and J. Zinn-Justin, p. 564 (North-Holland, Amsterdam).
- [8] *Quantum Magnetism* edited by Schollwöck U, Richter J, Farnell D J J and Bishop R F, 2004 (Springer-Verlag, Berlin).
- [9] The relationship between our notation using  $K$  and another using a compactification radius  $R$  for the field is given by  $K = 1/(4R^2)$ . In addition to these two, there are other normalization forms in the Abelian bosonization. See for instance Ref. [3].
- [10] Hikihara T and Furusaki A, 2004, *Phys. Rev. B* 69, 064427.
- [11] Lukyanov S and Zamolodchikov A, 1997, *Nucl. Phys. B* 493, 571.
- [12] A eck I, Kennedy T, Lieb E H and Tasaki H, 1987 *Phys. Rev. Lett.* 59, 799; 1988 *Commun. Math. Phys.* 115, 477.
- [13] Haldane F D M, 1983 *Phys. Rev. Lett.* 50, 1153; 1983 *Phys. Lett.* 93A, 464.
- [14] Fradkin E, 1991 *Field Theories of Condensed Matter Systems* (Addison-Wesley).
- [15] Auerbach A, 1994 *Interacting Electrons and Quantum Magnetism* (Springer-Verlag, New York).
- [16] A eck I, 1986, *Nucl. Phys. B* 265, [FS15], 409.
- [17] Zamolodchikov A B and Fateev V A, 1986, *Yad. Fiz.* 43, 1031 [1986, *Sov. J. Nucl. Phys.* 43, 657].
- [18] A eck I and Haldane F D M, 1987, *Phys. Rev. B* 36, 5291.
- [19] Tsvetlik A M, 1990 *Phys. Rev. B* 42, 10499.
- [20] Essler F H L and A eck I, 2004 *J. Stat. Mech.* P12006.
- [21] Besides the NLSM and the Majumdar-Femion theory, there is Schulz's method which try to explain the low-energy properties of spin-S AF chains by using coupled  $2S$  spin-1/2 chains (Schulz H J, 1986, *Phys. Rev. B* 34, 6372). However, since it contains extra degree of freedom and has more subtle points than the former two, generally the predictions of Schulz's method would be less reliable than those of the NLSM and the femion theory.
- [22] A eck I, 1990, *Phys. Rev. B* 41, 6697.
- [23] A eck I, 1991, *Phys. Rev. B* 43, 3215.
- [24] Konik R M and Fendley P, 2002, *Phys. Rev. B* 66, 144416.
- [25] Fath G, 2003 *Phys. Rev. B* 68, 134445.
- [26] Schulz H J, 1980 *Phys. Rev. B* 22, 5274.
- [27] A part of the bosonized-spin expression has been already completed. It however contains only a few relevant terms. See Refs. [22, 23, 24, 25, 20].
- [28] Shelton D G, Nersisyan A A and Tsvetlik A M, 1996 *Phys. Rev. B* 53, 8521.
- [29] Furusaki A and Zhang S C, 1999 *Phys. Rev. B* 60, 1175.
- [30] Chen Y, Honda Z, Zheludev A, Broholm C, Katsumata K and Shapiro S M, 2001 *Phys. Rev. Lett.* 86, 1618.
- [31] Zheludev A, Honda Z, Chen Y, Broholm C L and Katsumata K, 2002 *Phys. Rev. Lett.* 88, 077206.
- [32] Zheludev A, Honda Z, Broholm C L, Katsumata K, Shapiro S M, Kolezhuk K, Park S and Qui Y, 2003 *Phys. Rev. B* 68, 134438.
- [33] Hagiwara M, Honda Z, Katsumata K, Kolezhuk A K and Mikeska H -J, 2003 *Phys. Rev. Lett.* 91, 177601.
- [34] Zheludev A, Shapiro S M, Honda Z, Katsumata K, Grenier B, Ressouche E, Regnault L -P, Chen Y, Vondervisch P, Mikeska H -J and Kolezhuk A K, 2004 *Phys. Rev. B* 69, 054414.
- [35] Tsujii H, Honda Z, Andratska B, Katsumata K and Takano Y, 2005 *Phys. Rev. B* 71, 014426.
- [36] Zheludev A, Honda Z, Katsumata K, Feyereherm R and Prokes K, 2001 *Europhys. Lett.* 55, 868.
- [37] T. Goto, T. Ishikawa, Y. Shimooka, and Y. Fujii, 2006 *Phys. Rev. B* 73, 214406.
- [38] M. Hagiwara, L. P. Regnault, A. Zheludev, A. Stunault, N. Metoki, T. Suzuki, S. Suga, K.

- Kakurai, Y., Koike, P., Vorderwiesch, and J.-H. Chung, 2005 Phys. Rev. Lett. 94, 177202.
- [39] M. Hagiwara, H. Tsujii, C. R. Rotundu, B. Andraka, Y. Takano, N. Tatewaka, T. C. Kobayashi, T. Suzuki, and S. Suga, 2006 Phys. Rev. Lett. 96, 147203.
- [40] M. Sato, 2005 Phys. Rev. B 71, 024402.
- [41] M. Nakamura, 2003 Physica B 329-333, 1000.
- [42] For more detail of the bilinear-biquadratic chain, refer to Schollwöck U., Jolicoeur Th and Garel T., 1996 Phys. Rev. B 53, 3304; Fath G and Soto A., 2000 Phys. Rev. B 62, 3778; Allen D and Senechal D., 2000 Phys. Rev. B 61, 12134; Ref. [40].
- [43] Avdeev L V., 1990 J. Phys. A 23, L485.
- [44] Takhtajan L., 1982 Phys. Lett. 87A, 479; Babujian J., 1982 Phys. Lett. 90A, 479.
- [45] Qin S, Wang X and Yu L., 1997 Phys. Rev. B 56, R14251.
- [46] Todo S and Kato K., 2001 Phys. Rev. Lett. 87, 047203.
- [47] Wu T T, McCoy B, Tracy C A and Barouch E., 1976 Phys. Rev. B 13, 316.
- [48] We used Eqs. (2.31a) and (2.31b) in Ref. [47], and the asymptotic expansion formula of modified Bessel function,  $K_0(z) \sim \frac{1}{\sqrt{2z}} e^{-z} [1 + (4^{-2} - 1) = (8z) + (4^{-2} - 1)(4^{-2} - 3) = (2!(8z)^2) + \dots]$ .
- [49] Sørensen E S and Aelck I., 1994 Phys. Rev. B 49, 15771.
- [50] In the standard compact boson theory (see Refs. [1, 2, 4, 7, 8]) as an effective theory for a lattice system, the coexistence of  $e^{i\phi}$  and  $e^{i\psi}$ , which are present in Eqs. (22) and (23), is prohibited, because the former vertex operator requires the compactification radius  $R = 1 = \frac{1}{4}$  for the scalar field, but the latter does  $R = 1 = \frac{1}{4}$ . However, since the present theory consists of the boson part and the remaining Ising theory ( $^3; ^3; ^3$ ), we think that the "entanglement" between these two parts (i.e., three copies of Ising systems) allows such a coexistence. In other words, the field is expected not to be simply compactified, as long as we concurrently consider both the boson and the Ising parts.
- [51] Imposing a nontrivial transformation on Klein factors might be strange. Therefore, there is the possibility that we can find an alternating, more suitable transformation.
- [52] There are a few mistakes in the calculation of  $J_L J_R$  in Section IV C of Ref. [40], although they do not change the main results in Ref. [40].
- [53] A systematic method of the integration over the massive fields is the cumulant expansion for the partition function. For example, refer to Wang Y.-J., Essler F H L, Fabrizio M and Nersisyan A A., 2002 Phys. Rev. B 66, 024412. For the present effective theory, the cumulant expansion would be considered as a  $k; \nu$  expansion. Thus, it would be valid as far as the Fermi velocity  $v (v_F)$  is sufficiently large, i.e.,  $H$  is sufficiently larger than the Haldane gap  $m$ .
- [54] The emergence of  $(\partial_x)^2$  might be a failure in the field theory strategy. See Sections 6.1.2 and 7.2.1 in Ref. [1].
- [55] We should carefully treat a product between two continuous fields that coordinates are very close each other [e.g.,  $e^{i\phi(x)} e^{i\psi(x)}$  in Eq. (62a)], because such fields possess no information on the scale of the lattice constant  $a_0$ . Therefore, prefactors in all terms of Eqs. (62) and (63) would not be reliable at least quantitatively.
- [56] Note that  $L=R$  and  $L=R$  commute with each other in Eq. (62). Thus, we should interpret that  $L=R$  and  $L=R$  (or  $L=R$ ) also commute with each other.
- [57] Oshikawa M., Yamakawa M and Aelck I., 1997 Phys. Rev. Lett. 78, 1984.
- [58] F. D. M. Haldane, 1980 Phys. Rev. Lett. 45, 1358.
- [59] T. Sakai and M. Takahashi, 1991 J. Phys. Soc. Jpn. 60, 3615.
- [60] den Nijs M and Rommelse K., 1989 Phys. Rev. B 40, 4709.
- [61] Kennedy T and Tasaki H., 1992 Phys. Rev. B 45, 304; 1992 Commun. Math. Phys. 147, 431.
- [62] Francesco P D, Saleur H and Zuber J B., 1987 Nucl. Phys. B 290, 527.
- [63] For example see Cardy J L., 1996, Scaling and Renormalization in Statistical Physics (Cambridge Univ. Press, Cambridge, England).
- [64] Tonegawa T, Nakao T and Kaburagi M., 1996, J. Phys. Soc. Jpn. 65, 3317.
- [65] Totsuka K., 1998 Phys. Lett. A 228, 103.
- [66] Oshikawa M and Aelck I., 1997 Phys. Rev. Lett. 79, 2883; 1999 Phys. Rev. B 60, 1038.
- [67] Shankar R., 1995 Acta Phys. Pol. B 12, 1835
- [68] Delft J v and Schoeller H., 1998 Annalen der Physik (Leipzig) 4, 225; cond-mat/9805275.
- [69] Haldane F D M., 1981 Phys. Rev. Lett. 47, 1840.
- [70] Fabrizio M, Gogolin A O and Nersisyan A A., 2000 Nucl. Phys. B 580, 647.
- [71] In order to more accurately treat the commutation relations among vertex operators, fermions  $^q$  and Ising fields ( $^q, ^q$ ), one had better insert a certain quantity such as Klein factors in Eq. (A.6). For example see Ref. [70].
- [72] Eggert S and Aelck I., 1992 Phys. Rev. B 46, 10866.