

The Fractional Quantum Hall States of Dirac Electrons in Graphene

Vadim M. Apalkov

Department of Physics and Astronomy, Georgia State University, Atlanta, Georgia 30303, USA

Tapash Chakraborty^z

Department of Physics and Astronomy, University of Manitoba, Winnipeg, Canada R3T 2N2

(Dated: March 23, 2022)

We have investigated the fractional quantum Hall states of Dirac electrons in a graphene layer in different Landau levels. The relativistic nature of the energy dispersion relation of electrons in graphene significantly modifies the inter-electron interactions. This results in a specific dependence of the ground state energy and the energy gaps for electrons on the Landau-level index. For the valley-polarized states, i.e. at $\nu = 1/m$, m being an odd integer, the energy gaps have the largest values in the $n = 1$ Landau level. For the valley-unpolarized states, e.g., for the $2=3$ state, the energy gaps are suppressed for $n = 1$ as compared to those at $n = 0$. For both $n = 1$ and $n = 0$ the ground state of the $2=3$ system is fully valley-unpolarized.

PACS numbers: 73.43.f, 73.43.Lp, 73.21.b

A two-dimensional electron system in a single layer of graphite (graphene) is known to exhibit many remarkable properties. From the band structure studies [1] it was established early on that, to a good approximation, the energy dispersion of electrons in graphene is linear near the points at the corners of the Brillouin zone where the valence band and the conduction band meet. As a consequence, the low-energy excitations follow the Dirac-Weyl equations for massless relativistic particles [2]. In an external magnetic field, the electron system also shows unique properties that are different from those of the standard non-relativistic electron systems [2, 3, 4]. Recent experimental demonstration of some of those properties, in particular, the discovery of the integer quantum Hall effect [5, 6] that was predicted in earlier theoretical works [7] has caused intense interest in the electronic properties of the Dirac electrons in graphene [8]. However, effects of electron correlations in this system have not been reported yet. In this paper, we report on the nature of the fractional quantum Hall states of Dirac electrons in graphene.

A unit cell of the two-dimensional (2D) graphene honeycomb lattice contains two carbon atoms, say A and B. The dynamics of electrons in graphene is described by a tight-binding Hamiltonian with the nearest-neighbor hopping. In the continuum limit this Hamiltonian generates the band structure with two bands and the Fermi levels are located at two inequivalent points, $K = (2\pi/a)(1/3; 1/3)$ and $K^0 = (2\pi/a)(2/3; 0)$, of the first Brillouin zone, where $a = 0.246$ nm is the lattice constant. Near the points K and K^0 the electrons have a linear Dirac-Weyl ("relativistic") type dispersion relation [1, 2]. Finally, in the continuum limit the electron wave function is described by the 8-component spinor, $\psi_{s;k}$, where $s = \pm 1/2$ is the spin index, $k = K/K^0$ is the valley index, and $\alpha = A/B$ is the sublattice index. Without the spin-orbit interaction [11, 12, 13] the Hamiltonian is described by two 4×4 matrices for each component of the electron spin. In the presence of a magnetic field per-

pendicular to the graphene plane the Hamiltonian matrix has the form

$$H = \frac{B}{2} \begin{pmatrix} 0 & 0 & x + i y & 0 & 0 & 1 \\ x + i y & 0 & 0 & 0 & 0 & C \\ 0 & 0 & 0 & 0 & x + i y & A \\ 0 & 0 & 0 & x + i y & 0 & 0 \end{pmatrix}; \quad (1)$$

where $\vec{r} = \vec{p} + e\vec{A}/c$, \vec{p} is the two-dimensional momentum, \vec{A} is the vector potential, and C is the band parameter. The ordering used for the basis states in the non-interacting Hamiltonian is $(K; A; K; B; K^0; A; K^0; B)$. The eigenfunctions of the Hamiltonian are specified by the Landau-level index $n = 0; 1; 2; \dots$ and the intra-Landau level index m that is gauge dependent. Each Landau level is four-fold degenerate due to the spin and valley degrees of freedom. The corresponding wave functions for an electron in the two valleys K and K^0 are described by

$$\psi_{K;n} = C_n \begin{pmatrix} 0 & \text{sgn}(n) i^{n/2} j^{1/2} & 1 \\ i^{n/2} j^{1/2} & 0 & C \\ 0 & 0 & A \end{pmatrix}; \quad (2)$$

$$\psi_{K^0;n} = C_n \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & C \\ i^{n/2} j^{1/2} & \text{sgn}(n) i^{n/2} j^{1/2} & A \end{pmatrix}; \quad (3)$$

where $C_n = 1$ for $n = 0$ and $C_n = 1/\sqrt{2}$ for $n \neq 0$. Here ψ_n is the standard Landau wave function for a particle with non-relativistic parabolic dispersion relation in the n -th Landau level. From Eqs. (2)–(3) it is clear that a specific feature of the relativistic dispersion law is the mixing of the non-relativistic Landau levels. This mixture is present only for $n \neq 0$ and strongly modifies the inter-electron interaction within a single Landau level.

In what follows, we study the partially occupied Landau levels with fractional filling factors. Partial occupation of the Landau levels is realized by doping of the graphene layer. Experimentally, different filling factors of the Landau levels are achieved by varying the applied magnetic field at a fixed electron concentration. In this case the ground state of the system and the excitation spectrum are fully determined by the inter-electron interactions. For the non-relativistic case this results in the incompressible fractional quantum Hall effect (FQHE) states at the fractional filling factors [14, 15]. Properties of these states are completely described by Haldane's pseudopotentials, V_m [16], which are the energies of two electrons with relative angular momentum m . The pseudopotentials for the n -th Landau level can be presented as

$$V_m^{(n)} = \sum_{q=0}^{2m} \frac{1}{2} q V(q) [F_n(q)]^2 L_m(q^2) e^{-q^2}; \quad (4)$$

where $L_m(x)$ are the Laguerre polynomials, $V(q) = 2/e^2 = (lq)$ is the Coulomb interaction in the momentum space, l is the dielectric constant, l is the magnetic length, and $F_n(q)$ is the form factor corresponding to the n -th Landau level. For relativistic electrons the form factor is given by the expression [9]

$$F_0(q) = L_0 \frac{q^2}{2} \quad (5)$$

$$F_{n \neq 0}(q) = \frac{1}{2} L_n \frac{q^2}{2} + L_{n-1} \frac{q^2}{2}; \quad (6)$$

while for the non-relativistic particles the form factors in Eq. (4) are $F_n(q) = L_n \frac{q^2}{2}$. This means that the inter-electron interactions for the relativistic and non-relativistic electrons are the same for $n = 0$ and are different for $n > 0$ [9]. In what follows, all energies are expressed in units of the Coulomb energy $V_c = e^2/l$.

In Fig. 1 we compare the pseudopotentials calculated from Eq. (4) for the relativistic and the non-relativistic cases. For the relativistic electrons [Fig. 1 (a)] we notice a clear suppression of the pseudopotential for $n = 1$ as compared to that at $n = 0$, only for $m = 0$, i.e. when both electrons are at the same spatial point. For all other m , we have the inequality $V_m^{(1)} > V_m^{(0)}$. This is different from the non-relativistic case where the pseudopotential is suppressed also for $m = 1$, i.e. $V_1^{(1)} < V_1^{(0)}$. We also see in Fig. 1 (a) that although the relativistic wave functions at $n = 1$ is the "mixture" of the $n = 0$ and $n = 1$ non-relativistic wave functions, the relativistic pseudopotential is not the average of the corresponding non-relativistic pseudopotentials. This is clearly seen for $m = 1$ where the relativistic pseudopotential at $n = 1$ is larger than the non-relativistic one for both $n = 0$ and $n = 1$. In Fig. 1 (b) the relativistic pseudopotentials are shown for different Landau levels. Here the special case is $m = 1$ where the dependence of the pseudopotential on the Landau-level index is non-monotonic, viz., the pseudopotential has the maximum value at $n = 1$.




FIG. 1: The pseudopotentials [Eq. (4)] as a function of the relative angular momentum (a) for relativistic and for non-relativistic 2D electrons in the first two Landau levels, and (b) for relativistic electrons in various Landau levels.

At all the other m values the trend is the same as for the non-relativistic case, i.e., for $m = 0$ the pseudopotential decreases with increasing n , while at $m > 1$ the pseudopotentials increase with n .

With the pseudopotentials for Dirac electrons at hand, we now evaluate the energy spectra of the many-electron states at fractional fillings of the Landau level. The calculations have been done in the spherical geometry [16] with the pseudopotentials given by Eq. (4). The radius of the sphere R is related to 2S of magnetic fluxes through the sphere in units of the flux quanta as $R = \sqrt{2S} l$. The single-electron states are characterized by the angular momentum S , and its z component S_z . For a given number of electrons N , the parameter S determines the filling factor of the Landau level. Due to the spherical symmetry of the problem, the many-particle states are described by the total angular momentum L , and its z component, while the energy depends only on L . At first we study the system with the fractional filling factor $\nu = 1/3$, which corresponds to the $\nu = 1/3$ FQHE. In the spherical geometry, the $\nu = 1/3$ FQHE state is realized at $S = (3/2)(N - 1)$. If the electron system is fully spin and valley polarized then we should expect the ground state to be in the Laughlin state [17] which is separated from the excited states by a finite gap. We calculated the energy spectra of a finite-size system by finding the lowest eigenvalues and eigenvectors of the interaction Hamiltonian matrix [18]. In these calculations we take into account the interaction between all the electrons of the partially occupied Landau levels. We have also addressed the question of polarization of the many-particle state. We assume that at a high magnetic field the system is always spin-polarized.




FIG. 2: Energy spectra of the eight-electron $\nu = 1/3$ -FQHE system, shown for different Landau levels: (a) $n = 0$ (stars) and $n = 1$ (dots), and (b) $n = 2$. The system is fully spin and valley polarized. The flux quanta is $2S = 21$.

In this case the system can be either valley-polarized or valley-unpolarized. Similar to the spin polarization of the standard FQHE states the valley polarization of the graphene system depends on the inter-valley asymmetry (same as the Zeeman splitting for the non-relativistic system) and filling factor of the Landau level. Namely, if the inter-valley splitting due to the inter-valley asymmetry is large then the system is always valley-polarized, while otherwise the valley polarization of the system depends on the filling factor of the Landau level.

In Fig. 2 we show the calculated energy spectra for the $1/3$ -FQHE state at different Landau levels. Here, the $1/3$ -FQHE state at the n -th Landau level is defined as the state corresponding to the $1/3$ filling factor (single valley, single spin) of the n -th Landau level, while all the lower energy Landau levels are completely occupied. Since the relativistic pseudopotential $V_m^{(0)}$ for $n = 0$ is the same as for the non-relativistic one, the $1/3$ state and the corresponding energy gap (in units of ε_c) will be the same in both cases. The deviation from the non-relativistic system occurs only at higher Landau levels. In Fig. 2 (a) the energy gap of the $1/3$ -state at $n = 1$ is noticeably enhanced compared to that at $n = 0$. This is a direct manifestation of the specific dependence of pseudopotentials $V_m^{(n)}$ on the Landau-level index. Due to the asymmetry of the electron wave functions, the spectra of the $1/3$ -FQHE state is mainly determined by the relative value of $V_1^{(n)}$ and $V_3^{(n)}$ pseudopotentials, which have the highest value at $n = 1$. The energy spectra of the $1/3$ -FQHE state at $n = 2$, shown in Fig. 2 (b), demonstrate a strong suppression of the gap when compared to the $n = 1$ and $n = 0$ FQHE states. We therefore conclude




FIG. 3: Energy spectra of the eight-electron $\nu = 2/3$ -FQHE system, shown for different Landau levels: $n = 0$ (stars) and $n = 1$ (dots). The system is valley-unpolarized and fully spin-polarized. The flux quanta is $2S = 11$.

that the $1/3$ -FQHE state in graphene is most stable at $n = 1$. Hence the inter-electron interaction effects are more pronounced at $n = 1$. Interestingly, this tendency is just the opposite to that of the non-relativistic system, where the excitation gap decreases monotonically with increasing Landau-level index [14].

The results in Fig. 2 correspond to a completely spin and valley polarized system. This polarization is achieved at a high magnetic field due to the Zeeman splitting and the valley asymmetry. The inter-valley asymmetry has two sources: the first one is due to interaction-induced "backscattering" between different valleys [9] while the second one is due to the asymmetry in the lattice-scale interactions within the two sublattices of graphene [10]. Since the positions of the electrons in two sublattices are shifted, the interaction between the electrons in the different sublattices is weaker than the interaction between the electrons in the same sublattice. Both effects vary as (a/l) , so they become more relevant at higher magnetic fields or at a smaller magnetic length.

The same picture holds for the other FQHE states of the type $1/m$, i.e., the state is most stable at $n = 1$. The new aspects of the interaction physics arise at other filling factors as well, when the lowest energy states are spin-unpolarized for the non-relativistic single-valley systems. The simplest example is $\nu = 2/3$. In this case the ground state of the non-relativistic electrons is spin-unpolarized at a small Zeeman splitting. The transition from the spin-polarized to the spin-unpolarized ground states in a tilted magnetic field is well established both experimentally and theoretically for non-relativistic electrons [15].

Here we address the problem of the polarized and unpolarized states in the graphene system. We assume that the Zeeman splitting is large enough so that all the states are spin-polarized. At the same time the inter-valley asymmetry is small and electrons can occupy both the valleys. In this case the valley-polarized and valley-unpolarized $2/3$ -FQHE states become relevant. For

a valley-polarized system the excitation spectra and the ground state properties of the $2=3$ -state are the same as those for $=1=3$ due to the particle-hole symmetry. Similar to the $1=3$ case we obtain numerically an enhancement of the excitation gap at $n = 1$ as compared to that at $n = 0$. The more complicated situation occurs for the valley-unpolarized system. First we compare, just as for the non-relativistic system [19], the energy of the ground states of the polarized and the unpolarized systems. In the spherical geometry the polarized and the unpolarized states are realized for different sizes of the sphere, i.e., for different flux quanta through the sphere. For the $2=3$ -FQHE system the polarized state occurs at $2S = 3N=2$, while the unpolarized one is at $2S = 3N=2 + 1$. Due to different size of the sphere in these two systems the finite-size corrections to the magnetic length should be introduced, $l^0 = (2S=N)^{1/2}l$ [19, 20].

We have calculated the ground state energies for the valley-polarized and the valley-unpolarized graphene system in the $n = 1$ and the $n = 0$ Landau levels for a eight-electron system in the spherical geometry. We found that for both $n = 0$ and $n = 1$ the valley-unpolarized state has the lower energy. The ground state energy per particle in the unpolarized system is lower than that for the polarized system by 0.073% for $n = 0$ and by 0.053% for $n = 1$. Therefore, the valley-unpolarized state is more favorable for $n = 0$. Here the gap between the polarized and the unpolarized states is suppressed for $n = 1$ as compared to the $n = 0$ case. This is opposite to the completely polarized system (Fig. 2), where the effects of interaction is the strongest for $n = 1$. Suppression of the interaction effects in an unpolarized system for $n = 1$ is also illustrated in Fig. 3, where the excitation spectra of the valley-unpolarized system is shown for $n = 0$ and $n = 1$. A strong suppression of the excitation gaps for $n = 1$ is clearly visible here. The origin of this sup-

pression can be understood from the dependence of the pseudopotentials $V_m^{(n)}$ on the relative angular momentum m in different Landau levels. Due to the Pauli exclusion principle the energetic properties of the polarized state is determined only by the pseudopotentials with odd angular momenta, $m = 1; 3; 5; \dots$. These pseudopotentials have the largest values for $n = 1$, which results in the strongest interaction effects for $n = 1$, in a polarized system. For an unpolarized system the properties of the ground and excited states depend on all the pseudopotentials. Since $V_m^{(n)}$ at $m = 0$ is strongly suppressed for $n = 1$ as compared to the $n = 0$ case, we expect a suppression of interaction effects in an unpolarized system in the $n = 1$ Landau level.

In conclusion, the relativistic nature of the energy dispersion of electrons in the graphene plane modifies the inter-electron interactions significantly. This results in a unique dependence of the ground state energy and the energy gaps of the graphene systems on the Landau-level index. For the valley-polarized states, i.e. at $=1=m$, the FQHE gaps have the largest values for $n = 1$. Based on these studies we conclude that the FQHE at $=1=m$ should be observed experimentally for both $n = 0$ and $n = 1$, perhaps in a higher mobility system. For the valley-unpolarized states, e.g., for the $2=3$ -FQHE state, the energy gaps are suppressed at $n = 1$ as compared to that for the $n = 0$ level. For both $n = 1$ and $n = 0$ the ground state of the $2=3$ -FQHE system is fully unpolarized. The inter-valley asymmetry will result in transitions between the valley-polarized and the valley-unpolarized states.

The work has been supported by the Canada Research Chair Program and a Canadian Foundation for Innovation Grant.

[z] Electronic address: tapash@physics.umanitoba.ca

[1] P. R. Wallace, Phys. Rev. 71, 622 (1947).

[2] T. Ando, in Nano-Physics & Bio-Electronics: A New Odyssey, edited by T. Chakraborty, F. Peeters, and U. Sivan (Elsevier, Amsterdam, 2002), Chap. 1.

[3] J. W. McClure, Phys. Rev. 104, 666 (1956); R. R. Haering and P. R. Wallace, J. Phys. Chem. Solids 3, 253 (1957).

[4] Y. Zheng and T. Ando, Phys. Rev. B 65, 245420 (2002).

[5] K. S. Novoselov, et al., Nature 438, 197 (2005); Y. Zhang, Y.-W. Tan, H. L. Stormer, and P. Kim, ibid. 438, 201 (2005).

[6] Y. Zhang, et al., Phys. Rev. Lett. 96, 136806 (2006).

[7] V. P. Gusynin and S. G. Sharapov, Phys. Rev. Lett. 95, 146801 (2005); E. McCann and V. I. Fal'ko, ibid. 96, 086805 (2006).

[8] M. Wilson, Phys. Today 59 (1), 21 (2006).

[9] K. Nomura and A. H. MacDonald, Phys. Rev. Lett. 96, 256602 (2006); M. O. Goerbig, R. Moessner, and B. Douçot, cond-mat/0604554.

[10] J. Aliloea and M. P. A. Fisher, cond-mat/0604601.

[11] C. L. Kane and E. J. Mele, Phys. Rev. Lett. 95, 226801 (2005).

[12] X. F. Wang and T. Chakraborty, cond-mat/0605498.

[13] N. A. Sinitsyn et al., cond-mat/0602598.

[14] T. Chakraborty and P. Pietiläinen, The Quantum Hall Effects (Springer, Heidelberg, 1995), 2nd edition.

[15] T. Chakraborty, Adv. Phys. 49, 959 (2000).

[16] F. D. M. Haldane, Phys. Rev. Lett. 51, 605 (1983); F. D. M. Haldane and E. H. Rezayi, ibid. 54, 237 (1985).

[17] R. B. Laughlin, Phys. Rev. Lett. 50, 1395 (1983).

[18] G. Fano, F. Ortolani, and E. Colombo, Phys. Rev. B 34, 2670 (1986).

[19] X. C. Xie, Y. Guo, F. C. Zhang, Phys. Rev. B 40, 3487 (1989).

[20] R. Morf, N. d'Ambrunenti, and B. J. Halperin, Phys. Rev. B 34, 3037 (1986).