

Kondo screening of a high-spin Nagaoka state in a triangular quantum dot

Akira Oguri^{*}, Yunori Nishikawa, Yoshihide Tanaka, Takahide Numata

Department of Material Science, Osaka City University, Sumiyoshi-ku, Osaka 558-8585, Japan

Received 9 June 2006; revised 9 June 2006; accepted 9 June 2006

Abstract

We study transport through a triangular quantum dot connected to two noninteracting leads using the numerical renormalization group (NRG). The triangle has a high-spin ground state of $S = 1$ caused by a Nagaoka ferromagnetism, when it is isolated and has one extra electron introduced into a half-filling. The results show that the conduction electrons screen the local moment via two separate stages with different energy scales. The half of the $S = 1$ is screened first by one of the channel degrees, and then at very low temperature the remaining half is fully screened to form a Kondo singlet. The transport is determined by two phase shifts for quasi-particles with even and odd parities, and then a two-term in a conductance in the series configuration is suppressed ($g_{\text{series}} \rightarrow 0$), while plateau of a four-term in a parallel conductance reaches a unitary limit value ($g_{\text{parallel}} \rightarrow 4e^2/h$) of two conducting modes.

PACS: 72.10.-d; 72.10.Bg; 73.40.-c

Keywords: Quantum dot; Kondo effect; Nagaoka Ferromagnetism; Fermi liquid; Numerical Renormalization Group

The Kondo effect in quantum dots is an active field of current research, and recently triangle quantum dots with triangle have been examined intensively [1,2].

One interesting property expected to be seen in a quantum-dot array with closed paths is that some degenerate states could be lifted by circular orbital motions of electrons to form a high-spin ground state due to the Nagaoka mechanism. In this report for clarifying, i) how the Nagaoka ferromagnetism that could manifest in the isolated triangle for a particular charge filling is screened by the conduction electrons, and ii) how it affects the low-temperature transport below the Kondo energy scale T_K , we present the results using the NRG approach [3,4,5], which is applicable to low temperatures $T \ll T_K$.

We start with a three-site Hubbard model connected to two non-interacting leads on the left (L) and right (R), as shown in Fig. 1 (a): $H = H_D + H_{\text{mix}} + H_{\text{lead}}$,

$$H_D = \sum_{\langle ij \rangle} t_{ij}^D \sum_{\sigma} d_{i\sigma}^{\dagger} d_{j\sigma} + d_{i\sigma}^{\dagger} d_{j\sigma} + \sum_i \sum_{\sigma} d_{i\sigma}^{\dagger} d_{i\sigma} + U \sum_i \sum_{\sigma} d_{i\sigma}^{\dagger} d_{i\sigma} d_{i\sigma}^{\dagger} d_{i\sigma}; \quad (1)$$

$$H_{\text{mix}} = \sum_i \sum_{\sigma} v_{iL} d_{i\sigma}^{\dagger} c_{L\sigma} + v_{iR} d_{i\sigma}^{\dagger} c_{R\sigma} + H.c.; \quad (2)$$

$$H_{\text{lead}} = \sum_{k \in L, R} \sum_{\sigma} c_{k\sigma}^{\dagger} c_{k\sigma}; \quad (3)$$

where t is the hopping matrix element between the dots, d the onsite energy, U the Coulomb interaction, and $N_D = 3$. A linear combination of the conduction electrons $\sum_k c_k = N$ hybridizes with the electrons in the dots via v , or v^2 , where ρ is the density of states for each lead. The low-energy states of the whole system including the leads show a local Fermi-liquid behavior, which is characterized by two phase shifts ν_{even} and ν_{odd} for the quasi-particles with the even and odd parities. Then the dc conductance g_{series} and total number of electrons in the dots N_{el} can be expressed at $T = 0$ in the form [3,4],

$$g_{\text{series}} = \frac{2e^2}{h} \sin^2 \nu_{\text{even}} \sin^2 \nu_{\text{odd}}; \quad (4)$$




Fig. 1. Schematic picture of (a) series and (b) parallel connections

^{*}Corresponding author.

Email address: oguri@sci.osaka-cu.ac.jp (Akira Oguri).




Fig. 2. Series () g_{series} and parallel () g_{parallel} conductances are plotted together with local charge () N_{el} as functions of ε_d for $U/(2\pi t) = 1.0$ and $\Gamma/t = 0.12$. In NRG, $t/D = 0.1$ and $\beta = 6.0$.

$$N_{\text{el}} = \sum_{i=1}^D \langle n_i \rangle = \frac{2}{\pi} \int_{-\pi}^{\pi} d\phi \left(\frac{1}{2} + \frac{1}{2} \cos \phi \right) = \frac{2}{\pi} \left(\frac{\pi}{2} + \frac{\pi}{2} \right) = 2 \quad \text{even} + \quad \text{odd} : \quad (5)$$

Furthermore, because of some symmetrical reasons, the conductance for the current flowing in the horizontal direction in a four-terminal geometry, as shown in Fig. 1 (b), can be related to these two phase shifts defined with respect to the series connection [3,4],

$$g_{\text{parallel}} = \frac{2e^2}{h} \sin^2 \phi_{\text{even}} + \sin^2 \phi_{\text{odd}} : \quad (6)$$

We have deduced ϕ_{even} and ϕ_{odd} from the fixed-point eigenvalues of the discretized Hamiltonian H_N of NRG.

In Fig. 2, the results of g_{series} , g_{parallel} and N_{el} are plotted as functions of ε_d that corresponds to the gate voltage. The number of electrons inside the triangle increases with decreasing ε_d showing a staircase behavior at $N_{\text{el}} = 1, 2, 3, 4$; and 6. Due to a degeneracy arising in the ground state at $\varepsilon_d/U = -1.5$, the average charge changes directly from 4 to 6 without taking a step corresponding to $N_{\text{el}} = 5$. We see that the conductances for small $\varepsilon_d/U \approx 0.8$ show the typical Kondo behavior, which has also been seen in a linear chain of quantum dots [3,4]. Namely, for odd occupancies $N_{\text{el}} = 1$ and 3, the plateaus of g_{series} and g_{parallel} reach the unitary-limit value $2e^2/h$ of a single conduction mode. We also found a very narrow dip of g_{series} at $\varepsilon_d/U = 0.6$. For an even occupancy at $N_{\text{el}} = 2$, the conductances are suppressed. The irregular behavior at $N_{\text{el}} = 4$ is partly relating to the fact that, for $U = 0$, the circular motion along the triangle forms two degenerate orbitals at $\varepsilon_b = t$ and a single orbital at $\varepsilon_a = 2t$. As two of the four electrons occupy the ε_b orbitals, the Coulomb repulsion lifts the degeneracy and makes a $S = 1$ state to be a many-body ground state for $\beta = 0$. This corresponds to the Nagaoka state, and also links to a flat-band ferromagnetism. Then, the conduction electrons from the leads screen the moment. At the filling $N_{\text{el}} = 4$, the phase shifts take the values $\phi_{\text{even}} = 3\pi/2$ and $\phi_{\text{odd}} = \pi/2$. It suppresses the series conductance $g_{\text{series}} = 0$, while for the parallel conductance both of the two conducting modes contribute to the plateau




Fig. 3. Low energy eigenvalues of a discretized Hamiltonian $H_N = D$ as functions of odd NRG step N for $\varepsilon_d/U = -0.925$, where $N_{\text{el}} = 4$. Other parameter are taken to be the same as those used for Fig. 2.

of $g_{\text{parallel}} = 4e^2/h$ at $T = 0$. The difference between the peak values of the two conductances itself is caused by an interference effect.

In order to give some insights into the screening mechanism of the $S = 1$ moment, the low-lying energy levels of the NRG version of Hamiltonian H_N are plotted against odd N in Fig. 3. The trajectory of the levels shows how the system evolves from the high-energy regime to the low-energy Fermi-liquid regime as N increases, or equivalently with decreasing energy $\varepsilon_N \rightarrow D/(N+1)^2$, where D is the half-width of the conduction bands and β is the parameter for the logarithmic discretization. Our results indicate clearly that two crossovers occur separately at $N \approx 20$ and $N \approx 80$. This feature and additional information about the total spin and electron number of the eigenstates show that at high temperatures $T \gg \varepsilon_{N=20}$ the $S = 1$ moment is free from the screening. Then at the intermediate temperatures, $\varepsilon_{N=80} > T > \varepsilon_{N=20}$, the half of the moment is screened by the conduction electrons from one of the channel degrees, and thus in this region the local moment is still in an under-screened situation. The full screening is completed finally at very low temperatures $T \ll \varepsilon_{N=80}$.

The competition between the Kondo effect and Nagaoka ferromagnetism could happen in a wide class of quantum dots that have closed paths for the orbital motion.

This work was supported by the Grant-in-Aid for Scientific Research from JSPS. Numerical computation was partly carried out in Yukawa Institute Computer Facility.

References

- [1] A. Vidan, R. M. Westervelt, M. Stopa, M. Hanson and A. C. Gossard, Appl. Phys. Lett. 85, (2004), p. 3602.
- [2] T. Kuzmenko, K. Kikoin and Y. A. Vishai, Phys. Rev. Lett. 96, (2006), p. 046601.
- [3] Y. Nisikawa and A. Oguri, Phys. Rev. B 73, (2006), p. 125108.
- [4] A. Oguri, Y. Nisikawa and A. C. Hewson: J. Phys. Soc. Jpn. 74 (2005), p. 2554.
- [5] A. Oguri and A. C. Hewson: J. Phys. Soc. Jpn. 74 (2005), p. 988.