

High-temperature series for the bond-diluted Ising model in 3, 4 and 5 dimensions

Meik Hellmund^{1,3,*} and Wolfhard Janke^{2,3,†}

¹ *Mathematisches Institut, Universität Leipzig, Augustusplatz 10/11, D-04109 Leipzig, Germany*

² *Institut für Theoretische Physik, Universität Leipzig, Augustusplatz 10/11, D-04109 Leipzig, Germany*

³ *Centre for Theoretical Sciences (NTZ) of the Centre for Advanced Study (ZHS), Universität Leipzig, Emil-Fuchs-Str. 1, D-04105 Leipzig, Germany*

(Dated: March 23, 2022)

In order to study the influence of quenched disorder on second-order phase transitions, high-temperature series expansions of the susceptibility and the free energy are obtained for the quenched bond-diluted Ising model in $d = 3$ –5 dimensions. They are analysed using different extrapolation methods tailored to the expected singularity behaviours. In $d = 4$ and 5 dimensions we confirm that the critical behaviour is governed by the pure fixed point up to dilutions near the geometric bond percolation threshold. The existence and form of logarithmic corrections for the pure Ising model in $d = 4$ is confirmed and our results for the critical behaviour of the diluted system are in agreement with the type of singularity predicted by renormalization group considerations. In three dimensions we find large crossover effects between the pure Ising, percolation and random fixed point. We estimate the critical exponent of the susceptibility to be $\gamma = 1.305(5)$ at the random fixed point.

PACS numbers:

05.50.+q Lattice theory and statistics (Ising, Potts, etc.)

64.60.Fr Equilibrium properties near critical points, critical exponents

75.10.Hk Classical spin models

75.10.Nr Spin-glass and other random models

I. INTRODUCTION

Since many years random Ising models have served as paradigmatic systems in which the influence of quenched disorder may be studied through different techniques. In the present work we report results obtained by high-temperature series expansions. Systematic series expansions (Domb and Green, 1974) for statistical physics models defined on a lattice provide an useful complement to field-theoretical renormalization group studies and large-scale numerical Monte Carlo simulations. This is in particular true when studying phase transitions and critical phenomena of quenched, disordered systems.

Series expansions techniques treat the quenched disorder average exactly and the infinite-volume limit is implicitly implied. Therefore one can obtain exact results up to a certain order in the inverse temperature for many quantities. Moreover, one can keep the disorder strength p as well as the dimension d as symbolic parameters and therefore analyse large regions of the parameter space of disordered systems. The critical part of the series expansion approach lies in the extrapolation techniques which are used in order to obtain information on the phase transition behaviour from the finite number of known coefficients. While for pure systems this usually works quite well, one can question the use of these extrapolation techniques in disordered systems, where the singularity structure of the free energy or susceptibility may be very complicated, involving Griffiths-type singularities or logarithmic corrections (Cardy, 1999). Yet our work indicates that at least for the model we consider here, the extrapolation techniques are of comparable quality as for the case of the pure (no disorder) Ising model.

We consider in this paper the Ising model on a hypercubic d -dimensional lattice \mathbb{Z}^d with bond dilution as a realization of quenched and uncorrelated disorder. The pure model has a second-order phase transition for $d \geq 2$, and the upper critical dimension, where mean-field behaviour sets in, is $d_u = 4$. The influence of quenched disorder can be estimated by the Harris criterion (Harris, 1974): For $d > d_u = 4$, the disorder is expected to change only non-universal quantities such as the transition temperature T_c . At the upper critical dimension $d = 4$, logarithmic or

*Electronic address: Meik.Hellmund@math.uni-leipzig.de

†Electronic address: Wolfhard.Janke@itp.uni-leipzig.de

even more subtle correction terms should appear, and in $d = 3$, the phase transition of the disordered system should be governed by a new “random” fixed point since for the pure model the critical exponent α of the specific heat is positive and disorder should hence be a relevant perturbation. Our high-temperature series analyses presented in this paper affirm this picture.

The rest of the paper is organized as follows: In Sect. II we first briefly recall the model and some of its properties. Section III is devoted to a description of the methods used by us for generating the series expansions, and Sect. IV first starts out with some remarks on the analysis techniques used. The main results are presented in the followings subsections, where we discuss our results for the random-bond Ising model in five (Sect. IV.A), four (Sect. IV.B) and three (Sect. IV.C) dimensions. Finally, Sect. V contains our conclusions.

II. THE MODEL

The ferromagnetic disordered Ising model on hypercubic lattices \mathbb{Z}^d is defined by the partition function

$$Z(\{J_{ij}\}) = \sum_{\{s_i\}} \exp \left(\beta \sum_{\langle ij \rangle} J_{ij} s_i s_j \right), \quad (1)$$

where $\beta = 1/k_B T$ is the inverse temperature, J_{ij} are quenched (non-negative) nearest-neighbour coupling constants, and the spins s_i can take on the two different values ± 1 . In our series expansion the combination $v_{ij} = \tanh(\beta J_{ij})$ will be the relevant expansion parameter. Quenched disorder averages $[\dots]_{P(J)}$ such as the free energy

$$-\beta F = [\ln Z]_{P(J)} \quad (2)$$

are taken over an uncorrelated bimodal distribution of the form

$$P(J_{ij}) = (1 - p)\delta(J_{ij} - J_0) + p\delta(J_{ij}), \quad (3)$$

which corresponds to bond dilution: With probability p , bonds are effectively absent from the lattice, so that $p = 0$ represents the pure system. The expansion parameter for averaged quantities is usually taken as $v = \tanh(\beta J_0)$.

A. Thermal phase transitions

The pure ($p = 0$) model has for $d \geq 2$ a second-order phase transition from the disordered high-temperature phase to a ferromagnetic low-temperature phase. Bond dilution decreases T_c . Approaching the geometric bond percolation threshold $p = p_c$ the critical temperature decreases to zero ($v_c \rightarrow 1$) since above this value only finite clusters of bonds are present and therefore no global ferromagnetic order is possible. Figure 1 gives an overview of the phase diagrams as calculated in this work from high-temperature series of the susceptibility. The influence of quenched disorder on the thermal fixed point is different above and below the upper critical dimension $d_u = 4$. For $d \geq 4$ the pure fixed point is, according to the Harris criterion (Harris, 1974), stable against the influence of disorder and one expects that the renormalization group (RG) flow goes along the critical line from the unstable percolation fixed point straight to the stable pure fixed point. The critical behaviour along this line of phase transitions is governed by the pure fixed point. For $d = 3$ the pure fixed point is unstable and one expects a new random fixed point governing the critical behaviour in the region $0 < p < p_c$, attracting RG flow from both sides (pure and percolation fixed point). More specifically, for $d = 4$ random disorder changes the form of the logarithmic corrections to the Gaussian fixed point (Aharony, 1976), whereas for $d = 5$ Gaussian behaviour is expected for $p = 0$ and $p > 0$ as well.

B. Crossover to the percolation point

The crossover behaviour of the diluted system near the percolation threshold of the underlying lattice is well-understood (Aizenman *et al.*, 1987; Coniglio, 1981; Stauffer and Aharony, 1992) and independent of the dimension of the lattice. The percolation critical point $p = p_c$, $T_c = 0$ ($v_c = 1$) is always unstable against thermal fluctuations (Coniglio, 1981) and plays for our system the role of the strong disorder fixed point (Iglói and Monthus, 2005).




FIG. 1 Critical coupling v_c as function of the dilution p for $d = 3, 4$, and 5 dimensions. The arrows on the top axis indicate the percolation thresholds.

The crossover exponent ϕ is exactly 1 and critical exponents approach their percolation values with a scaling behaviour of the form

$$\chi \sim (p_c - p)^{-\gamma_p} f_\chi \left(\frac{e^{-2J_0/T}}{p_c - p} \right) \quad (4)$$

$$\sim (p_c - p)^{-\gamma_p} f_\chi \left(\frac{1}{p_c - p} \times \frac{1 - v}{1 + v} \right), \quad (5)$$

where f_χ is a crossover-scaling function, and the critical line near $v = 1$, $p = p_c$ has the form

$$\frac{1 - v}{1 + v} \propto (p_c - p). \quad (6)$$

III. STAR-GRAPH EXPANSION

The star-graph expansion method has been reviewed many times in the literature (Singh and Chakravarty, 1987) as well as in our previous papers (Hellmund and Janke, 2003, 2005). We will only present some algorithmic details of our approach and make some remarks comparing different techniques for series generation.

“Star graph” is an old-fashioned name for 1-vertex-irreducible graphs. For observables F which allow a star-graph expansion (like the free energy and various inverse susceptibilities) this technique delivers a representation of F on an infinite (e.g. hypercubic) lattice,

$$F(\mathbb{Z}^d) = \sum_G E(G; \mathbb{Z}^d) W_F(G), \quad (7)$$

as sum over star graphs (1-vertex irreducible graphs) only. Here, $E(G; \mathbb{Z}^d)$ denotes the weak embedding number of the graph G in the given lattice structure (Martin, 1974) and $W_F(G)$ can be recursively calculated from the knowledge of $F(G)$ and $F(G')$ for all star subgraphs $G' \subset G$.

A property of eq. (7) important for us is that the quenched disorder average can be calculated on the level of single graphs as long as the disorder is uncorrelated. This is related to the fact that the method considers and counts weak embeddings of the graph into the lattice, i.e., embeddings where different bonds of a graph are always mapped to different bonds of the lattice. This is in contrast to the free embeddings used in linked-cluster expansions. Here, different vertices and bonds of the graph may correspond to the same vertex or bond of the lattice. Therefore in order to calculate the disorder average one has to classify the free embeddings of a graph according to their correspondence to embeddings of a collapsed graph (where some edges of G are identified). This is essentially the algorithm proposed in (Meyer-Ortmanns and Reisz, 1999) where the collapsed graphs are called “multiple-line graphs”. This method has not yet been used for actual calculations of disordered systems, presumably due to its combinatorial complexity.

TABLE I Number of star graphs with $N \geq 8$ links and non-vanishing embedding numbers on \mathbb{Z}^d . For $N = 1, 4, 6$, and 7 only a single star graph exists.

order N	8	9	10	11	12	13	14	15	16	17	18	19	20	21
# in $d = 3$	2	3	8	9	29	51	136	306	856	2237	6431	18487	55302	165730
# in $d = 4$ and 5	2	3	8	9	29	51	142	330	951	2561	7622	22688		

For other series generation methods as, e.g., the finite lattice method which holds the world record for the pure 3D Ising model (Arisue *et al.*, 2004) or methods using Schwinger-Dyson equations (Butera and Comi, 1999), no practical generalization to disordered systems is known.

For the generation of graphs we employed the *nauty* package by McKay (McKay, 1981) which makes very fast isomorphism tests by calculating a canonical representation of the automorphism group of the graphs. By this means, we classified all star graphs up to order 21 that can be embedded in hypercubic lattices, see Table I. For each of these graphs we calculated their weak embedding numbers for d -dimensional hypercubic lattices (up to order 17 for arbitrary d , order 19 for dimensions $d \leq 5$ and order 21 for $d \leq 3$). For this embedding count we implemented a refined version of the backtracing algorithm by Martin (Martin, 1974), making use of a couple of simplifications for bipartite hypercubic lattices \mathbb{Z}^d . After extensive tests to find the optimal algorithm for the innermost loop, the test for collisions in the embedding, we ended up using optimized hash tables.

In order to calculate the contribution $W(G)$ of a graph G of order N to the susceptibility series, one needs the partition sum $Z(\{J_{ij}\}|G)$ and the matrix of all spin-spin correlation functions $M_{kl}(\{J_{ij}\}|G) = \text{Tr} \delta_{s_k, s_l} e^{-\beta H(J_{ij})}$ as polynomials in the N variables J_{ij} . This is achieved by using the cluster representation

$$Z \propto \sum_C 2^{c(C)} \prod_{\langle ij \rangle \in C} \frac{2v_{ij}}{1 - v_{ij}}, \quad (8)$$

$$M_{kl} \propto \sum_{C_{kl}} 2^{c(C)} \prod_{\langle ij \rangle \in C} \frac{2v_{ij}}{1 - v_{ij}}, \quad (9)$$

where the sum in eq. (8) goes over all clusters $C \subseteq G$ which have only vertices with an even number of bonds and in eq. (9) the sum is restricted to all clusters $C_{kl} \subseteq G$ in which the vertices k and l have an odd number of bonds and all other have an even number of bonds. The exponent $c(C)$ counts the number of connected components of C . A graph with N bonds gives rise to 2^N clusters: every bond may be present or absent. The clusters can therefore be numerated by N -bit integers. These 2^N integers can be sorted in such a way that two consecutive numbers differ by exactly one bit (corresponding to the addition or deletion of one bond), an algorithm known as “Gray code” (Press *et al.*, 1992). This gives a simple algorithm for the calculation of the cluster sums in eqs. (8) and (9):

- Start with the full graph $C = G$, store the coordination number modulo 2 (“coordination bit”) for every vertex and count the total number r of odd vertices. Execute steps 2 and 3 from below for this cluster configuration.
- Iterate over all other $2^N - 1$ Gray codes C :
 1. Calculate the next Gray code (cluster configuration). Compared to the previous code, exactly one bond was added or deleted. Invert the coordination bit of the two involved vertices and calculate the change in r .
 2. If $r = 0$ then add $\prod_{n \in C} w_n$ to Z . The product has a factor $w_n = 2v_{ij}/(1 - v_{ij})$ for every non-zero bit n in C .
 3. If $r = 2$ then add $\prod_{n \in C} w_n$ to M_{kl} where k, l are the two odd vertices.

The prefactor $2^{c(C)}$ is taken into account by monitoring the change in the number of connected components $c(C)$ in each iteration step.

The further steps for the calculation of the susceptibility series are:

- Inversion of the Z polynomial as a series in the $\{v_{ij}\}$ up to the desired order.
- Averaging over quenched disorder,
 $N_{kl}(G) = [M_{kl}/Z]_{P(J)}$,
resulting in a matrix of polynomials in p and v .

- Inversion of the matrix N_{kl} and subgraph subtraction,
 $W_\chi(G) = \sum_{k,l} (N^{-1})_{kl} - \sum_{G' \subset G} W_\chi(G').$
- Collecting the results from all graphs,
 $1/\chi = \sum_G E(G; \mathbb{Z}^d) W_\chi(G).$

All calculations are done in arbitrary-precision integer arithmetic using the open source library GMP. For the polynomial arithmetic we developed our own optimized C++ template library using a degree-sparse representation of polynomials as linked lists (Knuth, 1998). The calculations took around two CPU years on an Opteron Linux Cluster.

IV. SERIES ANALYSES

The estimate of critical parameters from a high-temperature series involves extrapolation from a finite number of exactly known coefficients to the asymptotic form of the function. Many such extrapolation techniques have been developed and tested for different series and are comprehensively reviewed in (Guttmann, 1989). These extrapolation techniques are not rigorous. They make some assumptions about the expected form of the singularity at the critical temperature. Field theoretic techniques like the ϵ -expansion have similar problems.

Usually, error estimates rely on the scatter of the results of extrapolations with different parameters (like $[N/M]$ Padé approximants for different values of N and M). This may seriously underestimate systematic errors coming from wrong assumptions about the structure of the singularity.

In order to get a reliable picture, we will take into account several criteria, such as

- convergence of the analysis,
- scatter of different approximants,
- number of defective approximants,
- agreement between different extrapolation methods.

The basic methods we use are DLog-Padé approximation and inhomogeneous differential approximants (IDA) (Guttmann, 1989). In order to analyze confluent nonanalytic and logarithmic corrections, these methods are applied to suitably transformed forms of the series. The parameters of these transformations are fine-tuned according to the criteria listed above, a technique pioneered in (Adler *et al.*, 1991).

We obtain from the star-graph technique series in two variables v and p . Since the algorithm involves inversion of polynomials (and of matrices of polynomials) one has to find a consistent truncation criterion for the resulting series. The crucial observations are that a) the coefficient matrix a_{mn} of the resulting series is triangular (cf. App. A), i.e., the series is of the form

$$\chi(p, v) = \sum_{n=0}^{\infty} \sum_{m=0}^n a_{mn} q^m v^n \quad \text{where} \quad q = 1 - p, \quad (10)$$

and b) the contribution of a graph of order N starts with terms of order $q^N v^N$. This allows different consistent truncation schemes. In scheme A we calculate series which are correct up to a given order N in v , i.e., we calculate all non-vanishing coefficients a_{mn} with $n \leq N$. In scheme B, we calculate all non-vanishing coefficients a_{mn} which satisfy $m + n \leq 2N$.

Both schemes are useful in different regions of the parameter space. At small and medium disorder p we approach the critical point by extrapolating in v for constant values of p . Therefore we use the series calculated according to truncation scheme A in the form

$$\chi(v|p = \text{const.}) \propto \sum_{n=0}^N \left(\sum_{m=0}^n a_{mn} q^m \right) v^n = \sum_{n=0}^N c_n v^n. \quad (11)$$

Near the percolation threshold $p \lesssim p_c$ this gives less and less satisfactory results. As eq. (6) and Fig. 1 show, $p = \text{const.}$ -lines are reaching the critical line at a smaller and smaller angle. In this region, it is better to extrapolate along the locus $w = q/v = \text{const.}$ This is achieved by taking the series coefficients calculated with scheme B:

$$\chi(v|w = \text{const.}) \propto \sum_{m+n \leq 2N} a_{mn} w^m v^{n+m} = \sum_{n=0}^{2N} d_n v^n. \quad (12)$$

The analysis of scheme B series brings much better results than scheme A in the highly diluted region $p \lesssim p_c$. But the quality of the analysis is still lower (e.g., the scatter of different Padé approximants is larger) than in the regions with small or medium dilution. This is not surprising since we can not construct from our data a series consistently truncated in p . This would be necessary in order to extrapolate along the locus $v = \text{const.}$ and study the percolation transition directly. Nonetheless, a reliable estimate of the transition temperature v_c is possible up to $p \approx p_c$, giving the phase diagram in Fig. 1.

A. Five dimensions

The analysis of the 19th order series for the susceptibility of the 5D bond-diluted Ising model is a nice warmup in order to get an impression of the quality of different analysis methods. Unbiased DLog-Padé and inhomogeneous differential approximants (IDA) along different $p = \text{const.}$ loci give values of γ between 1.02 and 1.04 for a wide range of dilutions $p = 0 \dots 0.7$ with a slightly increasing tendency.

This effect – an overestimate of γ in a naïve analysis not taking into account confluent corrections – is well-known from the 3D pure Ising model. A good illustration is Fig. 2 in (Campostrini *et al.*, 2002): whereas the results for many different theories in the Ising universality class (ϕ^4 etc.) converge to $\gamma \approx 1.237$, even 25th order series for the Ising model give $\gamma \approx 1.244$. The remedy is the use of analysis methods tailored for confluent nonanalytic corrections,

$$f(v) \sim A(v_c - v)^{-\gamma} [1 + A_1(v_c - v)^{\Delta_1} + \dots], \quad (13)$$

such as the methods from (Adler *et al.*, 1991) called M1 and M2. The method M1 uses DLog-Padé approximants to

$$F(v) = (v_c - v) \frac{df}{dv} - \gamma f(v), \quad (14)$$

which has a pole at v_c with residue $\Delta_1 - \gamma$. For a given trial value of v_c the graphs of Δ_1 versus input γ are plotted for different Padé approximants and by adjusting v_c a point of optimal convergence is searched.

The M2 method starts with a transformation of the series in v into a series in $y = 1 - (1 - \frac{v}{v_c})^{\Delta_1}$, and then Padé approximants to

$$G(y) = \Delta_1(1 - y) \frac{d}{dy} \ln F(y) \quad (15)$$

are calculated which should converge to γ as $y \rightarrow 1$, i.e. $v \rightarrow v_c$. These methods are especially useful when taken as biased approximants with a given value of γ or Δ_1 as input.

Using the M2 method we find over a large range of the disorder strength $p = 0 \dots 0.7$ excellent convergence of the Padé approximants with a nearly p -independent value of Δ_1 . As an example, Fig. 2 shows a M2 plot at dilution $p = 0.5$ from which we read off $\gamma = 1.001(1)$, $\Delta_1 = 0.51(2)$ and determine $v_c = 0.227498$. Our overall estimate for Δ_1 for all disorder strengths $p = 0 \dots 0.7$ is $\Delta_1 = 0.50(5)$. The results of the M1 method are compatible with this but have poorer convergence.

For the pure ($p = 0$) 5D Ising model our critical values obtained from the 19th order series with the M2 analysis biased with $\gamma = 1$ are $v_c = 0.113425(3)$, $\Delta_1 = 0.50(2)$. As usual, the errors are estimates of the scatter of different Padé approximants. This should be compared with the value from Monte Carlo (MC) simulations (Luijten *et al.*, 1999) of $\beta_c = 0.1139152(4)$, i.e. $v_c = 0.1134250(4)$, which is in perfect agreement.

B. Four dimensions

Four is the upper critical dimension of the Ising model. Without impurities the scaling behaviour of the susceptibility and specific heat is thus expected to exhibit at the critical point logarithmic corrections. With $t \equiv |1 - T/T_c|$, the precise form reads as (Ballesteros *et al.*, 1998b; Kenna, 2004; Kenna and Lang, 1991, 1993)

$$\chi \sim t^{-1} |\ln t|^{1/3}, \quad (16)$$

$$C \sim |\ln t|^{1/3}, \quad (17)$$

where a non-singular background term in C has been omitted. For the disordered case it was shown long ago (Aharony, 1976) by a RG analysis that the critical behaviour is modified to take the form

$$\chi \sim t^{-1} \exp \left[D |\ln t|^{1/2} \right], \quad (18)$$

$$C \sim -|\ln t|^{1/2} \exp \left[-2D |\ln t|^{1/2} \right], \quad (19)$$




FIG. 2 M2 analysis (γ as function of Δ_1) for the $p = 0.5$ diluted 5D Ising model at $v_c = 0.227\,498$ based on 15 different $[N/M]$ Padé approximants.

where $D = \sqrt{6/53} \approx 0.336$. The crossover to mean-field behaviour as well as higher loop corrections (Geldart and De'Bell, 1993) modify this unusual exponential term by factors of the form $|\ln t|^{\Delta_1}$. Notice that the expression in eq. (19) approaches zero for $t \rightarrow 0$ such that C is no longer logarithmic divergent but stays finite in the presence of disorder.

1. Pure Ising model

Due to the logarithmic corrections in eqs. (16) and (17) a special treatment of the series is needed. The analysis of logarithmic corrections of the general form

$$f(v) \sim (v_c - v)^{-\gamma} |\ln(v_c - v)|^\delta \quad (20)$$

is possible by (Adler and Privman, 1981) calculating approximants for

$$F(v) = (v_c - v) \ln(v_c - v) \left[\frac{f'(v)}{f(v)} - \frac{\gamma}{v_c - v} \right], \quad (21)$$

where one expects for singularities of the form (20) that $\lim_{v \rightarrow v_c} F(v) = \delta$. In what follows we call this the M3 method.

The estimates for the pure 4D Ising model based on our 19th order series are: a) unbiased estimate, Fig. 3: $\gamma = 1.019(1)$, $\delta = 0.34(1)$, $v_c = 0.148\,607(10)$ and b) biased to $\gamma = 1$, Fig. 4: $\delta = 0.429$, $v_c = 0.148\,583(3)$. This should be compared to the analysis in (Stauffer and Adler, 1997) of a 17th order series (Gaunt *et al.*, 1979) which gave $v_c = 0.148\,588$ and to MC simulations (Bittner *et al.*, 2002) which found $\beta_c = 0.149\,697(2)$, i.e. $v_c = 0.148\,589(2)$. While the agreement of the estimates for v_c is excellent, this analysis also shows how difficult it is to obtain reliable estimates for the critical exponent δ of the logarithmic correction term.

2. Bond dilution

An extensive MC study of the *site*-diluted 4D model was done in (Ballesteros *et al.*, 1998b). The authors found data compatible with the theoretical prediction of a Gaussian fixed point with logarithmic corrections, but a precise fit of the logarithmic corrections was not possible.

A modification of the M3 analysis method which we refer to as M3a is able to take into account the specific form of eq. (18): For

$$f(v) \sim (v_c - v)^{-\gamma} \exp \left[D |\ln(v_c - v)|^{1/2} \right] \quad (22)$$




FIG. 3 Logarithmic exponent δ as function of the leading exponent γ for the pure 4D Ising model via unbiased M3 approximants at $v_c = 0.148\,607$. A pronounced clustering of the different $[N/M]$ Padé approximants around the point $\gamma \approx 1.019$, $\delta \approx 0.34$ is clearly observed.




FIG. 4 Logarithmic exponent δ as function of v_c for the pure 4D Ising model via M3 approximants biased to $\gamma = 1$. From the clustering of the $[N/M]$ Padé approximants one reads off $v_c \approx 0.148\,583$, $\delta \approx 0.429$. When assessing the clustering property notice that the y -scale is 10 times finer than in Fig. 3.

we calculate approximants to

$$F(v) \sim (v_c - v)(-\ln(v_c - v))^{1/2} \left[\frac{f'(v)}{f(v)} - \frac{\gamma}{v_c - v} \right] \quad (23)$$

which should behave as $\lim_{v \rightarrow v_c} F(v) = D/2$. We made an extensive analysis of our 19th order susceptibility series assuming the three forms eqs. (13), (20) resp. (16) and (22) resp. (18). It turns out that, while form (13) is not acceptable, both forms of logarithmic corrections allow a fit of the series data, both with a disorder dependent exponent δ or D , respectively. The quality of the M3a fits is, however, better than for the M3 fits. This is demonstrated in Figs. 5 and 6 where equal ranges of v_c ($\Delta v_c = 0.0003$) and of the exponent ($\Delta\delta = \Delta D = 0.1$) are shown.

We interpret this as an indication for the validity of the RG prediction eq. (18). Figure 7 shows the dilution dependency of our estimates for the critical parameter D . It illustrates the difficulty in differentiating between a behaviour of the form eq. (20) and the form eq. (22) by giving a non-zero result for D at $p = 0$. It is nevertheless




FIG. 5 Logarithmic exponent δ as function of v_c for the $p = 0.4$ diluted 4D model, results of a M3 analysis.




FIG. 6 Parameter D as function of v_c for the $p = 0.4$ diluted 4D model, results of a M3a analysis.

quite impressive that we can see at all such weak correction terms to the leading singularity and even estimate the parameters.

For the specific heat, one expects the weak logarithmic singularity in the pure case [eq. (17)] to be washed out by the disorder. An analysis of the free energy series $[F(v)]_{P(J)} = \sum a_i(p)v^i$ is usually more difficult and gives less satisfactory results compared to the susceptibility series. One loses two orders in v by calculating $C = \beta^2 \frac{d^2 F}{dv^2}$ and, more importantly, on bipartite lattices the series for $F(v)$ includes only even powers of v . So it has to be considered as a much shorter series in v^2 . In four dimensions we calculated $F(v, p)$ up to order 18, which gives a series for $C(v^2)$ of order 8 in v^2 .

Another difficulty is that the non-singular background terms are more influential in this case. Therefore DLog-Padé approximants show very poor convergence and one has to use inhomogeneous differential approximants (IDA) which are able to take background contributions into account. By this method, polynomials $P_N(v)$, $Q_M(v)$ and $R_L(v)$ (of order N , M and L , respectively) are determined such that

$$P_N(v)\hat{D}f(v) + Q_M(v)f(v) + R_L(v) = o(v^{(N+M+L)}), \quad (24)$$

where many different triples (N, M, L) with $N + M + L \leq (\text{order of } f)$ and two different variants of the differential operator \hat{D} , either $\hat{D}_1 = v d/dv$ or $\hat{D}_2 = d/dv$, are used. The critical point v_c is then given by the smallest positive




FIG. 7 Parameter D as function of p for the diluted 4D model, results of a M3a analysis. The horizontal line at $D = 0.336 \dots$ shows the predicted value $\sqrt{6/53}$.




FIG. 8 Critical exponent of $\partial^4 f(v)/\partial v^4$ as function of p for the diluted 4D model.

real root of $P_N(v)$ and the critical exponent by $Q_M(v_c)/P'_N(v_c)$.

We applied IDAs tailored to power-like singularities $\sim t^{-\alpha-2}$ to the fourth derivative $\partial^4 f(v)/\partial v^4$. The results are shown in Fig. 8. In the pure case we find a result consistent with $\partial^4 f(v)/\partial v^4 \sim (v_c - v)^{-2}$ indicating a logarithmic singularity in $C \sim \partial^2 f(v)/\partial v^2$. At nonzero values of the dilution parameter p the singularity of the fourth derivative of f is clearly weaker than $(v_c - v)^{-2}$, indicating the absence of a divergence in the specific heat in the disordered case.

C. Three dimensions

Let us finally turn to the physically most important case of three dimensions. The quest for a determination of the properties of the expected random fixed point in the 3D disordered Ising model is already rather long. A comprehensive compilation of results can be found in (Folk *et al.*, 2000, 2003) and (Pelissetto and Vicari, 2002), showing a wide scatter in the critical exponents of different groups, presumably due to large crossover effects. Recent MC simulations (Ballesteros *et al.*, 1998a; Berche *et al.*, 2005, 2004; Ivaneyko *et al.*, 2005) provide evidence for the random fixed point




FIG. 9 Critical exponent γ as function of p for the diluted 3D model obtained from about 300 DLog-Padé and inhomogeneous differential approximants for each dilution value.




FIG. 10 Critical exponent γ as function of p for the diluted 3D model obtained from series analyses with methods M1 and M2 (see text for details). The data of Fig. 9 (Pade/IDA) are also shown for comparison.

but also show large crossover effects due to the interference with the pure fixed point for $p \rightarrow 0$ and with the percolation fixed point for $p \rightarrow p_c$ (recall the discussion in Sect. II.B).

Early attempts using series expansions (Rapaport, 1972a,b) already indicated that the series are slower converging and more difficult to analyse than in the pure case. A review of earlier work can be found in (Stinchcombe, 1983). Series analysis in crossover situations is, in fact, extremely difficult. If the parameter p interpolates between regions governed by different fixed points, the exponent obtained from a finite number of terms of a series expansion must cross somehow between the two universal values, and does this usually quite slowly (Janke *et al.*, 2005). The mere existence of a plateau in $\gamma(p)$, however, is an indication that here a truly different critical behaviour is seen.

Our results are obtained from a large number of DLog-Padé and inhomogeneous differential approximants (around 300 for every dilution value) applied to the 21th order susceptibility series $\chi(p, v)$ compiled in App. A. The resulting estimates for the critical exponent γ shown in Fig. 9, however, do not exhibit any sign for a plateau.

Since confluent corrections are essential to understand crossover situations, we also performed a careful analysis using the M1 and M2 methods which both take such corrections explicitly into account, see Fig. 10. This figure shows results using both $v = \tanh(\beta J_0)$ as well as the coupling βJ_0 as expansion variables (denoted by, e.g., M1/ v and M1/ J). They give an indication of a plateau around $p = 0.3 \dots 0.4$, suggesting the presence of the random fixed

point with corresponding value of the critical exponent $\gamma = 1.305(5)$. Here, as usual in extrapolation techniques, the error is estimated from the scatter of different approximants and we are unable to give an estimate of the systematic error of the extrapolation. Our high-temperature series estimate is at least compatible with MC results for site and bond dilution (Ballesteros *et al.*, 1998a; Berche *et al.*, 2004; Calabrese *et al.*, 2003) which cluster quite sharply around $\gamma_{\text{MC}} = 1.34(1)$. Field-theoretic RG estimates (Folk *et al.*, 2000; Pelissetto and Vicari, 2000) favor slightly smaller exponents of $\gamma_{\text{RG}} = 1.32 - 1.33$, while experiments (Belanger *et al.*, 1986; Birgeneau *et al.*, 1983; Mitchell *et al.*, 1986) report values between $\gamma_{\text{exp}} = 1.31 - 1.44$; for a more detailed compilation, see e.g., Table 1 in Ref. (Berche *et al.*, 2005). Since it would be extremely demanding to further extend the series expansions in the disordered case, better series analysis methods for the case of crossover situations would be clearly desirable.

V. SUMMARY

We successfully applied the method of high-temperature series expansion to the bond-diluted Ising model in several dimensions. The computational effort is vast and increases faster than exponentially with the order of the expansion. But the extensive set of combinatorial data we generated on the way, such as the list of star graphs and their embedding numbers into hypercubic lattices, has a large number of potential applications, for example to models with other kinds of uncorrelated disorder like spin glasses. On the other hand, Monte Carlo simulations of systems with quenched disorder require an enormous amount of computing time because many realizations have to be simulated for the quenched average. For this reason it is hardly possible to scan a whole parameter range. Using the method of star-graph expansion we obtain this average exactly. Since the relevant parameters (degree of disorder p , spatial dimension d , etc.) are kept as symbolic variables, we can easily study large regions of the phase diagram.

Table II summarizes the main results of this work. Our estimates of the critical temperature for the 4- and 5-dimensional model are comparable in precision (as well as consistent) with the available Monte Carlo data for the pure case. For the bond-diluted model in 4 and 5 dimensions we demonstrated universality. The critical exponent γ keeps its Gaussian value $\gamma = 1$ up to the percolation threshold of the underlying lattice, despite the fact that the upper critical dimension of the percolation transition is six. In 5 dimensions we also show that the exponent Δ_1 of the confluent correction is universal for a large range of the dilution parameter p . In 4 dimensions we confirm the RG prediction that the weak logarithmic divergence of the specific heat of the pure model disappears in the disordered case due to logarithmic corrections of a special kind. We confirm the presence of these special corrections also in the case of the susceptibility. We furthermore determine the exponent of the logarithmic correction in the pure case with satisfying precision.

The case of 3 dimensions is, in a sense, the most difficult to analyze. We clearly see that the pure fixed point is unstable and does not describe the random system. Extrapolation methods not taking into account confluent corrections give a value for γ which changes monotonously with p . By using methods tailored to the consideration of confluent corrections we identify a plateau in the curve $\gamma(p)$ with critical exponent $\gamma = 1.305(5)$ which we see as evidence for the random critical point.

model		v_c	γ	confluent correction
5D	pure	0.113 425(3)	1.001(1)	power-like, $\Delta_1 = 0.50(2)$
	disordered		1.001(1)	power-like, $\Delta_1 = 0.50(5)$
4D	pure	0.148 607(3)	1.019(20)	logarithmic, $\delta = 0.34(2)$
	disordered		form eq. (18) confirmed	
3D	disordered		1.305(5)	

TABLE II Summary of results. In three dimensions, longer series for the pure Ising model exist, therefore we do not quote here our results for v_c or γ .

Acknowledgments

We thank Joan Adler for introducing us into the art and science of series extrapolation techniques.

Support by DFG grant No. JA 483/17-3 and partial support from the German-Israel-Foundation under grant No. I-653-181.14/1999 is gratefully acknowledged.

APPENDIX A: Coefficients of the 3D susceptibility

The following table of coefficients gives the complete information for calculating the susceptibility of the 3D bond-diluted Ising model up to the order v^{21} for any dilution p .

References

- Adler, J., A. Aharony, A. B. Harris, L. Klein, and Y. Meir, 1991, Series expansions for the Ising spin glass in general dimension, *Phys. Rev. B* **43**, 11249.
- Adler, J., and V. Privman, 1981, Two-dimensional percolation: Logarithmic corrections to the critical behaviour from series expansions, *J. Phys. A* **14**(11), L463.
- Aharony, A., 1976, New singularities in the critical behaviour of random Ising models at marginal dimensionalities, *Phys. Rev.* **13**(5), 2092.
- Aizenman, M., J. Chayes, L. Chayes, and C. Newman, 1987, The phase boundary in dilute and random Ising and Potts ferromagnets, *J. Phys. A* **20**, L313.
- Arisue, H., T. Fujiwara, and K. Tabata, 2004, Higher orders of the high-temperature expansion for the Ising model in three dimensions, *Nucl. Phys. (Proc. Suppl.)* **129**, 774 [hep-lat/0309158].
- Ballesteros, H. G., L. A. Fernández, V. Martín-Mayor, A. Muñoz Sudupe, G. Parisi, and J. J. Ruiz-Lorenzo, 1998a, Critical exponents of the three dimensional diluted Ising model, *Phys. Rev. B* **58**, 2740.
- Ballesteros, H. G., L. A. Fernández, V. Martín-Mayor, A. M. Sudupe, G. Parisi, and J. J. Ruiz-Lorenzo, 1998b, The four dimensional site-diluted Ising model: A finite-size scaling study, *Nucl. Phys. B* **512**, 681 [hep-lat/9707017].
- Belanger, D. P., A. R. King, and V. Jaccarino, 1986, Crossover from random-exchange to random-field critical behavior in $Fe_xZn_{1-x}F_2$, *Phys. Rev. B* **34**(1), 452.
- Berche, B., P. E. Berche, C. Chatelain, B. Berche, and W. Janke, 2005, Random Ising model in three dimensions: Theory, experiment and simulation – a difficult coexistence, *Condens. Matter Phys.* **8**(1), 47 [cond-mat/0411255].
- Berche, P. E., C. Chatelain, B. Berche, and W. Janke, 2004, Bond dilution in the 3D Ising model: A Monte Carlo study, *Eur. Phys. J. B* **38**, 463 [cond-mat/0402596].
- Birgeneau, R. J., R. A. Cowley, G. Shirane, , H. Yoshizawa, D. P. Belanger, A. R. King, and V. Jaccarino, 1983, Critical behavior of a site-diluted three-dimensional Ising magnet, *Phys. Rev. B* **27**(11), 6747.
- Bittner, E., W. Janke, and H. Markum, 2002, Ising spins coupled to a four-dimensional discrete Regge skeleton, *Phys. Rev. D* **66**, 024008 [hep-lat/0205023].
- Butera, P., and M. Comi, 1999, Enumeration of the self-avoiding polygons on a lattice by the Schwinger-Dyson equations, *Ann. Comb.* **3**, 277 [cond-mat/9903297].
- Calabrese, P., V. Martin-Mayor, A. Pelissetto, and E. Vicari, 2003, The three-dimensional randomly dilute Ising model: Monte Carlo results, *Phys. Rev. E* **68**, 036136.
- Campostrini, M., A. Pelissetto, P. Rossi, and E. Vicari, 2002, 25th-order high temperature expansion results for three-dimensional Ising-like systems on the simple cubic lattice, *Phys. Rev. E* **65**, 066127 [cond-mat/0201180].
- Cardy, J., 1999, Logarithmic correlations in quenched random magnets and polymers, *cond-mat/9911024*.
- Coniglio, A., 1981, Thermal phase transition of the dilute s -state Potts and n -vector models at the percolation threshold, *Phys. Rev. Lett.* **46**(4), 250.
- Domb, C., and M. S. Green (eds.), 1974, *Series Expansions for Lattice Models*, volume 3 of *Phase Transitions and Critical Phenomena* (Academic Press).
- Folk, R., Y. Holovatch, and T. Yavors'kii, 2000, Effective and asymptotic critical exponents of weakly diluted quenched Ising model: 3d approach versus $\sqrt{\epsilon}$ -expansion, *Phys. Rev. B* **61**(22), 15114 [cond-mat/9909121].
- Folk, R., Y. Holovatch, and T. Yavors'kii, 2003, Critical exponents of a three dimensional weakly diluted quenched Ising model, *Phys. Usp.* **46**, 169 [cond-mat/0106468].
- Gaunt, D. S., M. F. Sykes, and S. McKenzie, 1979, Susceptibility and fourth-field derivative of the spin-1/2 Ising model for $t > t_c$ and $d = 4$, *J. Phys. A* **12**(6), 871.
- Geldart, D. J. W., and K. De'Bell, 1993, Logarithmic corrections for dilute uniaxial ferromagnets at the critical dimension, *J. Stat. Phys.* **73**(1/2), 409.
- Guttmann, A. J., 1989, Asymptotic analysis of power-series expansions, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L. Lebowitz (Academic Press, New York), volume 13 of *Phase Transitions and Critical Phenomena*, pp. 1–234.
- Harris, A. B., 1974, Effect of random defects on the critical behaviour of Ising models, *Journal of Physics C: Solid State Physics* **7**(9), 1671.
- Hellmund, M., and W. Janke, 2003, Star graph expansions for bond diluted Potts models, *Phys. Rev. E* **67**, 026118.
- Hellmund, M., and W. Janke, 2005, High-temperature series expansions for random Potts models, *Condens. Matter Phys.* **8**(1), 59 [cond-mat/0502152].
- Iglói, F., and C. Monthus, 2005, Strong disorder RG approach of random systems, *Phys. Rep.* **412**, 277 [cond-mat/0502448].
- Ivaneyko, D., J. Ilnytskyi, B. Berche, and Y. Holovatch, 2005, Criticality of the random site Ising model: Metropolis, Swendsen-Wang and Wolff Monte Carlo algorithms, *Condens. Matter Phys.* **8**(1), 149 [cond-mat/0501291].
- Janke, W., B. Berche, C. Chatelain, P.-E. Berche, and M. Hellmund, 2005, Quenched disordered ferromagnets, in *Proceedings of "Lattice 2005 Dublin"*, PoS, p. 018, PoS(LAT2005)018.
- Kenna, R., 2004, Finite size scaling for $O(N)$ ϕ^4 -theory at the upper critical dimension, *Nucl. Phys. B* **691**, 292 [hep-lat/0405023].

- Kenna, R., and C. Lang, 1991, Finite-size scaling and the zeroes of the partition function in the ϕ_4^4 model, Phys. Lett. B **264**, 396.
- Kenna, R., and C. Lang, 1993, Renormalization group analysis of finite size scaling in the ϕ_4^4 model, Nucl. Phys. B **393**, 461 [hep-lat/9210009].
- Knuth, D. E., 1998, *The Art of Computer Programming*, volume 2 (Addison-Wesley), 2nd edition.
- Luijten, E., K. Binder, and H. Blöte, 1999, Finite-size scaling above the upper critical dimension revisited: The case of the five-dimensional Ising model, Eur. Phys. J. B **9**, 289 [cond-mat/9901042].
- Martin, J. L., 1974, Computer techniques for evaluating lattice constants, in (Domb and Green, 1974), pp. 97–112.
- McKay, B. D., 1981, Practical graph isomorphism, Congressus Numerantium **30**, 45, <http://cs.anu.edu.au/~bdm/nauty/>.
- Meyer-Ortmanns, H., and T. Reisz, 1999, Dynamical linked cluster expansions: A novel expansion scheme for point-link-point interactions, Int. J. Mod. Phys. A **14**, 947 [hep-lat/9804018].
- Mitchell, P. W., R. A. Cowley, H. Yoshizawa, P. Boni, Y. J. Uemura, and R. J. Birgeneau, 1986, Critical behavior of the three-dimensional site-random Ising magnet: $\text{Mn}_x\text{Zn}_{1-x}\text{F}_2$, Phys. Rev. B **34**(7), 4719.
- Pelissetto, A., and E. Vicari, 2002, Critical phenomena and renormalization-group theory, Phys. Rep. **368**, 549 [cond-mat/0012164].
- Pelissetto, A., and E. Vicari, 2000, Randomly dilute spin models: A six-loop field-theoretic study, Phys. Rev. B **62**, 6393.
- Press, W. H., S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, 1992, *Numerical Recipes in C* (Cambridge University Press).
- Rapaport, D. C., 1972a, The Ising ferromagnet with impurities: A series expansion approach. I, Journal of Physics C: Solid State Physics **5**(14), 1830.
- Rapaport, D. C., 1972b, The Ising ferromagnet with impurities: A series expansion approach: II, Journal of Physics C: Solid State Physics **5**(19), 2813.
- Singh, R. R. P., and S. Chakravarty, 1987, High-temperature series expansion for spin glasses. I. derivation of the series, Phys. Rev. B **36**(1), 546.
- Stauffer, D., and J. Adler, 1997, Logarithmic factors, critical temperature, and zero temperature flipping in the 4D kinetic Ising model, Int. J. Mod. Phys. C **8**(2), 263.
- Stauffer, D., and A. Aharony, 1992, *Introduction to Percolation Theory* (Taylor & Francis London), 2nd edition.
- Stinchcombe, R. B., 1983, Dilute magnetism, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L. Lebowitz (Academic Press, New York), volume 7, pp. 151–280.