

Quantum dynamics of Bose-Hubbard Hamiltonians beyond Hartree-Fock-Bogoliubov: The Bogoliubov backreaction approximation

I. Tikhonenkov,¹ J. R. Anglin,² and A. Vardi[†]

¹Department of Chemistry, Ben-Gurion University of the Negev, P.O.B. 653, Beer-Sheva 84105, Israel

²Fachbereich Physik, Technische Universität Kaiserslautern, D 67663, Kaiserslautern, Germany

We formulate a method for studying the quantum field dynamics of ultracold Bose gases confined within optical lattice potentials, within the lowest Bloch-band Bose-Hubbard model. Our formalism extends the two-sites results of Phys. Rev. Lett. 86, 000568 (2001) to the general case of M lattice sites. The methodology is based on mapping the Bose-Hubbard Hamiltonian to an $SU(M)$ pseudospin problem and truncating the resulting hierarchy of dynamical equations for correlation functions, up to pair-correlations between $SU(M)$ generators. Agreement with few-site exact many-particle calculations is consistently better than the corresponding Hartree-Fock-Bogoliubov approximation. Moreover, our approximation compares favorably with a more elaborate two-particle irreducible effective action formalism, at a fraction of the analytic and numerical effort.

PACS numbers: 3.75.Kk

I. INTRODUCTION

Strong correlation effects, which imply enhanced quantum fluctuations around mean field order parameters, are playing an increasingly important role in recent experiments on dilute quantum gases. One strategy for boosting the importance of correlations and fluctuations involves the control of coupling parameters. Interatomic interactions can be effectively tuned by means of magnetic Feshbach resonances [1, 2, 3, 4, 5, 6], allowing for a controlled transition into the non-unitary regime $n^{1/3}a_s > 1$, where the effective s -wave scattering length a_s is larger than the average distance between particles $n^{1/3}$ with n being the number-density of the gas. Quantum fluctuations also dominate quasi-one-dimensional systems [7, 8, 9, 10, 11, 12, 13, 14, 15] where transverse confinement may be used to increase the effective coupling strength $g_{1D} = 2\hbar^2 a_s / (m l_\perp^2)$ without explicit control of the three-dimensional s -wave scattering length. In the extreme Tonks-Girardeau strong-coupling regime $g_{1D} m = (\hbar^2 n) \gg 1$, spatial correlations dictate the inpenetrability of bosons, leading to ideal fermion like density distributions [8, 9].

An alternative to increasing effective interaction strengths, is to decrease other (e.g. kinetic) terms in the many-body Hamiltonian. In a Bose gas confined by an optical lattice, an effective momentum cutoff is introduced by controlling the barrier heights, thus suppressing the hopping frequency J between adjacent sites. Given N particles interacting with strength U , the strong-interaction regime is achieved for $UN = J > 1$, as manifested in the quantum transition from a superfluid to a Mott-insulator phase [16, 17, 18].

Considerable theoretical effort is currently aimed at developing efficient methods for the description of correlated quantum gases far from equilibrium. One approach relies on perturbations of the lowest-order mean-field theory given by the Gross-Pitaevskii (GP) equation. The result is a family of mean-field pairing theories. The

standard zero-temperature Bogoliubov prescription [19] gives the natural small-oscillation modes by linearization about the GP ground-state. However, this linear response theory does not account for the backreaction of excitations on the condensate order-parameter and is thus limited to small perturbations and short timescales. Backreaction is accounted for within the Hartree-Fock-Bogoliubov (HFB) theory, which prescribes a set of coupled equations for the condensate order-parameter and pair correlation functions [20, 21, 22, 23]. Since both normal and anomalous correlations are included, this approach comes at the cost of ultraviolet divergences of anomalous quantities. While this problem is relatively easy to deal with by renormalization of the coupling parameters, a more serious issue, also related to the inclusion of anomalous correlations, is the HFB spectral gap [20]. This unphysical gap in the excitation spectrum results in from the breaking of $U(1)$ gauge symmetry and the consequent elimination of the Goldstone modes corresponding to gauge transformations of the broken symmetry solution. An intermediate form between Bogoliubov and HFB is the HFB-Popov (HFB-P) approximation [20, 24] where $U(1)$ symmetry is restored by elimination of noncondensate anomalous terms only. While the resulting theory is gapless, it does not conserve the total number of particles and is thus inadequate for describing dynamical condensate depletion. Finally, if all anomalous quantities are neglected, one obtains the bosonic Hartree-Fock (HF) theory [20] which is both gapless and conserving, but does not allow for any dynamical depletion, since the populations of condensed and non-condensed particles are conserved separately. It is thus highly desirable to develop a theoretical description that (a) is $U(1)$ invariant and hence gapless, (b) conserves the total number of particles, yet (c) allows for dynamical depletion of the condensate.

Recently, a perturbative approximation scheme based on a two-particle irreducible (2PI) effective action expansion, has been used to study the nonequilibrium dynamics of condensates in optical lattices [23] within the

lowest Bloch-band Bose-Hubbard model. Within the 2PI effective action expansion, the Bogoliubov and HFB theories emerge as one-loop and a single two-loop correction respectively, to the classical GP action. Higher-order approximations, obtained by including two-vertex terms in the diagrammatic expansion of the effective action (denoted as in Ref. [23] by '2nd') and by a $1=N$ expansion up to second-order in the coupling strength (denoted henceforth by ' $1=N$ ') with N being the number of auxiliary classical fields used to approximate the quantum field, have been compared with HFB and exact few-sites numerical calculations. The results demonstrate some improvement of the higher-order approximations over HFB in predicting the exact many-body dynamics. However, at sufficiently long times all approximations fail due to interaction effects. A nonperturbative $1=N$ 2PI effective action expansion approach have also been developed and applied to the equilibration of a homogeneous Bose gas in 1D [25].

In this work we develop a mean-field theory for the description of quantum dynamics in the Bose-Hubbard model. The technique, referred to here as Bogoliubov Back Reaction (BBR), is a many-site extension of previous work on a two-site model [26, 27], based on the perturbation of equations of motion for the reduced single-particle density operator, instead of the usual field operator approach. The resulting equations involve the two-point reduced single-particle density matrix (SPDM) and the four-point correlation functions. They contain only normal (i.e. number conserving) quantities, and are thus $U(1)$ symmetric. The approximation conserves the total number of particles, yet it allows for population transfer from the condensate to the excitations, thus accounting for condensate depletion during the evolution. We compare BBR calculations with full many-body numerical results for up to a hundred particles and five lattice sites, as well as with HFB and 2PI effective action results. The BBR results give better, longer-time predictions than current rival approximations, at a small fraction of the theoretical effort.

In section II we present the Bose-Hubbard model and the standard HFB approach. In section III we transform the Bose-Hubbard Hamiltonian with M lattice sites into an $SU(M)$ pseudospin problem, derive dynamical equations for the $SU(M)$ generators spanning the single-particle density operator, and truncate the resulting hierarchy of dynamical equations for correlation functions to obtain the BBR equations of motion. Section IV contains numerical few-sites results and comparison with HFB as well as 2PI effective action approximation methods. Discussion, conclusions and prospects for future research are presented in section V.




FIG. 1: (color online) Population imbalance w in a two-site system as a function of rescaled time τ for $L = 2$ (a), $L = 4$ (b), $L = 5$ (c), and $L = 10$ (d). The total number of particles is set to $N = 100$. Solid blue lines, corresponding to exact many-body numerical results, are compared to the GP (dotted lines), HFB (dash-dotted lines), and BBR (dashed lines) approximations.

II. CONVENTIONAL MEAN-FIELD THEORIES: GROSS-PITAEVSKII AND HARTREE-FOCK-BOGOLIUBOV

We begin with the standard Bose-Hubbard model Hamiltonian for an ultracold gas in a one-dimensional periodic optical lattice

$$\hat{H} = J \sum_i \hat{a}_{i+1}^\dagger \hat{a}_i + \hat{a}_i^\dagger \hat{a}_{i+1} + \frac{U}{2} \sum_i \hat{a}_i^\dagger \hat{a}_i \hat{a}_i \hat{a}_i \quad (1)$$

where $\hat{a}_i$ and $\hat{a}_i^\dagger$ are annihilation and creation operators respectively, for a particle in site i . We consider only on-site interactions with strength U and nearest-neighbor tunneling with hopping rate J . These approximations are justified because adjacent site interactions and next-to-nearest-neighbor tunneling amplitudes are characteristically at least two orders of magnitude smaller than on-site interactions and nearest-neighbor hopping [16]. The Bose-Hubbard model is viable as long as there are no transitions into excited Bloch bands.

Using the Hamiltonian (1) we write the Heisenberg equations of motion for the field operators $\hat{a}_j$,

$$i \frac{d}{dt} \hat{a}_j = J (\hat{a}_{j-1} + \hat{a}_{j+1}) + U \hat{a}_j^\dagger \hat{a}_j \hat{a}_j : \quad (2)$$

The lowest order mean-field theory for the Bose-Hubbard model is obtained by replacing the field operators $\hat{a}_j$, and $\hat{a}_j^\dagger$ by c -numbers a_j and a_j^* . This approximation is tantamount to assuming coherent many-body states with a well-defined phase between sites. Rescaling $a_j \rightarrow a_j / \sqrt{N}$




FIG. 3: (color online) Evolution of the leading eigenvalue (above) and single-particle entropy (below) for a two-site system with $N = 20; 40; 80$ and $L = 2$. Exact many-body numerics (solid blue line) is compared with the HFB approximation (green dotted line) and the BBR approximation (red dashed line).

III. THE BOGOLIUBOV BACKREACTION EQUATIONS

Instead of the conventional mean-field approaches, based on the site field operators $\hat{a}_j$, we construct a mean-field formalism using the reduced single-particle density operator $\hat{a}_j^y \hat{a}_k$, treating it as the fundamental quantity. We have previously applied this approach to the case of a two-site model [26, 27]. Here we extend it to the general M -site case. It is convenient to rewrite the Hamiltonian (1) in terms of the $M^2 - 1$ traceless operators which generate $SU(M)$:

$$\begin{aligned} \hat{u}_{j;k} &= \hat{a}_j^y \hat{a}_k + \hat{a}_k^y \hat{a}_j; 1 \leq k < j \leq M \\ \hat{v}_{j;k} &= i \frac{\hat{a}_j^y \hat{a}_k - \hat{a}_k^y \hat{a}_j}{2}; 1 \leq k < j \leq M \\ \hat{w}_1 &= \frac{2}{1(1+1)} \sum_{j=1}^{M-1} \hat{n}_j - \frac{1}{M} \hat{n}^2; 1 \leq j \leq M-1 \end{aligned} \quad (13)$$

Since it is easily verified that

$$\frac{1}{2} \sum_{j=1}^{M-1} \hat{w}_j^2 + \frac{1}{M} \hat{n}^2 = \sum_{j=1}^{M-1} \hat{n}_j^2 \quad (14)$$

where $\hat{n} = \sum_{j=1}^M \hat{n}_j$ is the total particle number, equation (1) can be rewritten, eliminating c-number terms, as:

$$\hat{H} = J \sum_{j=1}^{M-1} \hat{u}_{j+1;j} + \frac{U}{4} \sum_{j=1}^{M-1} \hat{w}_j^2; \quad (15)$$

Using the $SU(M)$ generators we construct a pseudo-

spin vector operator,

$$\hat{S} = (\hat{u}_{21}; \hat{u}_{32}; \dots; \hat{u}_{31}; \hat{u}_{42}; \dots; \hat{v}_{21}; \hat{v}_{32}; \dots; \hat{v}_{31}; \hat{v}_{42}; \dots; \hat{w}_1; \hat{w}_2; \dots; \hat{w}_{M-1}); \quad (16)$$

so that the Hamiltonian (15) takes the form:

$$\hat{H} = J \sum_{j=1}^{M-1} \hat{S}_j + \frac{U}{4} \sum_{j=M^2-M}^{M^2-1} \hat{S}_j^2; \quad (17)$$

The Heisenberg equations of motion for the operators $\hat{S}_i$ and their products $\hat{S}_i \hat{S}_1$ then read:

$$i \frac{d}{dt} \hat{S}_i = J \sum_{j=1}^{M-1} \sum_{k=1}^{M^2-1} c_{ij}^k \hat{S}_k \quad (18)$$

$$+ \frac{U}{4} \sum_{j=M^2-M+1}^{M^2-1} \sum_{k=1}^{M^2-1} c_{ij}^k (\hat{S}_k \hat{S}_j + \hat{S}_j \hat{S}_k);$$

$$i \frac{d}{dt} \hat{S}_i \hat{S}_1 = J \sum_{j=1}^{M-1} \sum_{k=1}^{M^2-1} (c_{ij}^k \hat{S}_k \hat{S}_1 + c_{1j}^k \hat{S}_i \hat{S}_k) \quad (19)$$

$$+ \frac{U}{4} \sum_{j=M^2-M+1}^{M^2-1} \sum_{k=1}^{M^2-1} c_{ij}^k (\hat{S}_k \hat{S}_j + \hat{S}_j \hat{S}_k) \hat{S}_1$$

$$+ \frac{U}{4} \sum_{j=M^2-M+1}^{M^2-1} \sum_{k=1}^{M^2-1} c_{1j}^k \hat{S}_i (\hat{S}_k \hat{S}_j + \hat{S}_j \hat{S}_k);$$

where the coefficients c_{ij}^k are the structure constants of the $SU(M)$ group. We note that for $M = 2$ the Hamiltonian (15) and the dynamical equations (18)–(19) reduce to the familiar Bloch forms used in Refs. [26, 27]. The M -site system is a direct extension of the two-mode case, in that hopping terms induce linear Rabi-like oscillations in the vw subspace, whereas on-site interactions lead to nonlinear phase precession in the uv subspace.

The reduced single-particle density matrix is obtained from the expectation value of $\hat{S}$, according to:

$$= \frac{N}{2} I + \frac{1}{2} \sum_{j=1}^{M^2-1} \langle \hat{S}_j \rangle \hat{S}_j; \quad (20)$$

where I is a unit matrix of order M and $\hat{S}_j$ are the $M^2 - M$ irreducible representations of the $SU(M)$ generators (e.g. Pauli matrices for $M = 2$, Schwinger matrices for $M = 3$ etc.). We will therefore focus on the dynamics of the ‘hyper-Bloch-vector’ $\mathbf{S} = \langle \hat{S} \rangle$ with $i=2N$. The lowest-order mean-field approximation replaces the vector of operators $\hat{S}$ by the vector of their expectation values $\mathbf{S}$, thus truncating $\langle \hat{S}_i \hat{S}_j \rangle$ to $\langle \hat{S}_i \rangle \langle \hat{S}_j \rangle$. This results in the nonlinear pseudospin-precession form of the GP equations,

$$\frac{d}{dt} \mathbf{S} = \mathbf{B}(\mathbf{S}) \times \mathbf{S} \quad (21)$$




FIG. 4: (color online) Site-populations in a three-site system as a function of rescaled time τ for $N = 20; 40; 80$ and fixed $L = 2$. Blue, green, and red lines correspond to 1st, 2nd, and 3rd site populations, respectively. Solid lines depict the full many-body dynamics, whereas dotted and dashed lines correspond to the HFB approximation and the BBR approximation, respectively.

where

$$B(S) = (B_1; B_2; \dots; B_{M^2-1}) ; \quad (22)$$

with

$$B_j = \begin{cases} 0 & j = 1; \dots; M(M-1)=2 \\ L S_j & j = M(M-1)+1; \dots; M^2-1 \end{cases}$$

It is readily verified that Eq. (21) is exactly equivalent to the discrete GP equation (3). In addition to the conservation of the total number $\text{Tr}(\rho)$ there exist, within GP theory, $M-1$ independent constants of the motion $\text{Tr}(\rho^m)$ with $m = 2; \dots; M-1$. For example, for $M = 2$ the GP mean-field theory also conserves the single-particle purity $\text{Tr}(\rho^2)$, which is just the length of the three-dimensional Bloch vector. Deviations from this classical field theory, due to interparticle entanglement and loss of single particle coherence, will show up as a reduction in these classically conserved quantities.

The BBR approximation is obtained by going one level deeper in the hierarchy of dynamical equations for expectation




FIG. 5: (color online) Leading eigenvalue of the reduced single-particle density matrix and single-particle entropy $-\text{Tr}(\rho \ln \rho)$, as a function of rescaled time τ in a three-site system with $N = 20; 40; 80$ and $L = 2$. Exact results (solid blue lines) are compared to HFB calculations (dotted green lines) and BBR calculations (dashed red lines).

tation values. Taking the expectation values of Eq. (18) and Eq. (19) and truncating

$$\begin{aligned} \hbar \hat{S}_i \hat{S}_j \hat{S}_k i &= \hbar \hat{S}_i \hbar \hat{S}_j \hat{S}_k i + \hbar \hat{S}_j \hbar \hat{S}_i \hat{S}_k i \\ &+ \hbar \hat{S}_k \hbar \hat{S}_i \hat{S}_j i - 2 \hbar \hat{S}_i \hbar \hat{S}_j \hbar \hat{S}_k i ; \end{aligned} \quad (23)$$

we obtain the BBR equations of motion:

$$\begin{aligned} i \frac{d}{dt} S_i &= \sum_{j=1}^{M^2-1} \sum_{k=1}^{M^2-1} c_{ij}^k S_k \\ &+ L \sum_{j=M^2-M+1}^{M^2-1} \sum_{k=1}^{M^2-1} c_{ij}^k (S_j S_k + S_{jk}) \end{aligned} \quad (24)$$

$$\begin{aligned} i \frac{d}{dt} S_{ij} &= \sum_{j=1}^{M^2-1} \sum_{k=1}^{M^2-1} c_{ij}^k S_{ik} + c_{ij}^k S_{ik} \\ &+ L \sum_{j=M^2-M+1}^{M^2-1} \sum_{k=1}^{M^2-1} c_{ij}^k (S_{ij} S_k + S_{ik} S_j) \\ &+ L \sum_{j=M^2-M+1}^{M^2-1} \sum_{k=1}^{M^2-1} c_{ij}^k (S_{ij} S_k + S_{ik} S_j) ; \end{aligned} \quad (25)$$

where $S_j = \frac{\hbar \hat{S}_j i}{2N}$ and $S_{jk} = \frac{\hbar \hat{S}_j \hat{S}_k + \hat{S}_k \hat{S}_j i}{4N^2}$. In the following section we compare the accuracy of the BBR approximation with respect to GP, HFB, and 2PI effective action.




FIG. 6: (color online) Site-populations in a four-site system as a function of rescaled time for $N = 20; 40; 80$ and fixed $L = 2$. Blue, green, red, and cyan lines correspond to 1st, 2nd, 3rd, and 4th site populations, respectively. Solid lines depict the full many-body dynamics, whereas dotted and dashed lines correspond to the HFB approximation and the BBR approximation, respectively.

IV. NUMERICAL RESULTS

In order to test the accuracy of the BBR approximation compared to other methods, we carried out exact numerical calculations for limited numbers of particles and sites (up to $N = 100$ particles and $M = 5$ lattice sites). The Hamiltonian (1) was represented in terms of site-number states and the N -body Schrödinger equation was solved numerically, as in Refs. [23, 26, 27]. These many-body results were then compared with BBR mean-field calculations, as well as with GP, HFB and variants of the 2PI effective action method.

In Fig. 1 the evolution of fractional population difference for a hundred particles in two-sites, is plotted for various values of the coupling parameter L . Within the GP mean-field theory, full-amplitude Rabi-like oscillations are predicted in the linear regime with $L < 2$ (Fig. 1(a)). As the transition is made to the strong-coupling regime, the oscillation becomes increasingly more non-linear, until when $L \geq 4$ macroscopic self-trapping is attained (Figs. 1(b)-1(d)). The value of $L = 4$ is par-

ticularly interesting because for this coupling a trajectory starting from a single-populated site becomes dynamically unstable when site-populations equilibrate. In previous work we have shown that this dynamical instability serves as a quantum-noise amplifier [26, 27], so that the growth of the deviation of a quantum trajectory from the corresponding GP prediction is initially exponential, leading to a $\log(1/N)$ slow convergence of the many-body quantum-field results to the classical GP prediction. Thus, while the naive expectation would be that quantum fluctuations would simply grow with the coupling parameter L , their role is in fact maximized for $L = 4$, as evident in Fig. 1(b). It is clear from Fig. 1 that the BBR approximation gives a better description of the ensuing quantum dynamics, for longer timescales, than HFB does.

Convergence of various approximations with increasing number of particles is demonstrated in Fig. 2, where the two-sites population dynamics is plotted for increasing particle numbers, keeping a fixed coupling value of $L = 2$. In addition to the exact, BBR, and HFB results, we also plot two calculations based on the 2PI effective action approach, taken from Fig. 5 of Ref. [23] (our exact and HFB results exactly coincide with the corresponding lines in that figure). Here too, the BBR approximation (red dashed lines) gives a more accurate description of the dynamics than any of the other methods, attaining a nearly perfect convergence in the given time-frame for $N = 80$ particles. In comparison, standard HFB fails to depict the damping of coherent oscillations, whereas the 2PI effective action methods tend to over-damp. We note, that in terms of formalistic complexity alone, the BBR approximation is far simpler than the non-instantaneous integro-differential equations used in the 2PI effective action methods [23]. In fact, it is even simpler than HFB, in that only normal quantities are involved, giving a total of nine equations for two sites, as opposed to fifteen in HFB.

Dynamical condensate depletion is also well-depicted by the BBR approximation. In Fig. 3 we plot the evolution of the leading eigenvalue of the reduced single-particle density matrix and the single-particle von-Neumann entropy $-\text{Tr}(\ln \rho)$, corresponding to the population dynamics of Fig. 2. While HFB calculations seem to give an abrupt deviation of the predicted condensate fraction from its exact value, the BBR results converge well, giving a reasonably accurate description of BEC depletion.

The same qualitative behavior carries over to systems with more than two sites. In Fig. 4 and Fig. 5, population dynamics and condensate depletion are shown for a three-sites system with $N = 20; 40; 80$ particles. Similarly to the two-sites case, the BBR approximation constitutes a significant improvement over the HFB approach, giving a better description of populations as well as coherences. The same is also true for the four-sites case shown in Fig. 6.

The faster convergence of BBR as compared with the




FIG. 7: (color online) Characteristic times at which the Cartesian distance between the exact Bloch vector and its HFB (green, x's) and BBR (blue, circles) approximations, reaches a predetermined threshold, as a function of N for $L = 1$ (a), $L = 4$ (b), $L = 6$ (c), and $L = 10$ (d). The break-threshold is set to 0.2 in (a)–(c), and to 0.05 in (d).

HFB approximation is illustrated in Fig. 7, where characteristic breaktimes of the two approximations in a two-site calculation, are plotted as a function of the total number of particles N . As anticipated, breaktimes grow as N when the classical dynamics is regular (7a, 7c, 7d) and as $\log N$ when the classical trajectory hits the dynamical instability (7b). The BBR calculations give consistently longer breaktimes, with a more regular convergence pattern.

V. DISCUSSION

The rich regime of strongly correlated many body physics, which ultracold atom experiments are now beginning to probe, will surely not be fully conquered by any simple hierarchy truncation scheme such as BBR. Nor does BBR offer anything like an exact solution even to the problems to which we have applied it in this paper; its improvements over its rivals are incremental rather than revolutionary. On the other hand it should be born in mind that incremental improvements in theory are more significant in the context of ultracold gases than in traditional condensed matter, because in the new atomic systems samples are precisely characterized, controlled, and measured, and relevant microphysics is clearly known. It is perfectly plausible in these systems that we may come to learn important qualitative principles from experimental discrepancies on the few percent

level.

The merits of BBR that we would like to emphasize, along with its very reasonable level of accuracy, are its simplicity and its direct relation to experimental reality. It involves only quantities which are directly observed in single- and two-particle number-conserving measurements, and it respects the fact that in current quantum gas laboratories atoms are neither created nor destroyed. And it is conceptually and computationally straightforward.

In one sense it is of course conceptually all too straightforward: like all hierarchy truncation schemes since Boltzmann's, it is an uncontrolled approximation, whose accuracy is therefore arguably as much a puzzle as it is a solution. Insofar as truncating a hierarchy at a deeper level is grounds for expecting higher accuracy, however, the advantage of BBR is clear: it is a truncation at fourth order in field operators, compared to only second order for HFB. Deeper level truncation often involves proliferation of terms, to the point of sharply diminishing returns in accuracy versus effort; but BBR avoids this, and manages to use fewer equations than HFB, because it eliminates all anomalous terms.

And this leads us to conclude by indicating some of the potential future applications of the results of this paper.

Why do hierarchy truncations often work as well as they do? What determines the best way of truncating a hierarchy? These are questions that have been raised ever since Boltzmann's stosszahlansatz produced the arrow of time, but they have yet to be fully answered. With current experimental capabilities for precise and controlled measurements on ultracold gases, introducing a physically motivated alternative truncation scheme, as this paper has done, may contribute to new progress on these questions.

Finally, another conceptual merit of BBR is that because it is based on the single particle density matrix, rather than just the macroscopic wave function, it makes such a conceptually important quantity as the single particle entropy (the entropy of Boltzmann) a basic ingredient in the theory, rather than a perturbative afterthought. Rethinking entropy, heretofore mainly in the context of quantum information and computation theory, is another major thrust of current physics; the alternative viewpoint offered by BBR may potentially be of some value in a broader conception of this project.

Acknowledgments

This work was supported in part by grants from the Minerva foundation for a junior research group and the Israel Science Foundation for a Center of Excellence (grant No. 8006/03).

[1] W. C. Stwalley, Phys. Rev. Lett. 37, 1628 (1976).

[2] E. Tiesinga et al., Phys. rev. A 46, R1167 (1992).

- [3] E. Tiesinga et al, Phys. Rev. A 47, 4114 (1993).
- [4] E. Timmermans, P. Tommasini, M. Hussein, and A. Kerman, Phys. Rep. 315, 199 (1999).
- [5] F. H. Mies et al, Phys. rev. A 61, 022721 (2000).
- [6] S. Inouye, M. R. Andrews, J. Stenger, H.-J. Miesner, D. M. Staper-Kum, and W. Ketterle, Nature 392, 151 (1998).
- [7] E. Lieb and W. Liniger, Phys. Rev. 130, 1605 (1963); E. Lieb, Phys. Rev. 130, 1616 (1963).
- [8] M. Girardeau, J. Math. Phys. 1, 516 (1960).
- [9] M. Girardeau and E. M. Wright, Phys. Rev. Lett. 84, 5691 (2000).
- [10] M. Olshanii, Phys. Rev. Lett. 81, 938 (1998).
- [11] D. S. Petrov, G. Schlyapnikov, and J. T. M. Valraven, Phys. Rev. Lett. 85 3745 (2000).
- [12] V. Dunjko, V. Lorent, and M. Olshanii, Phys. Rev. Lett. 86 5413 (2001).
- [13] F. Schreck et al, Phys. Rev. Lett. 87, 080403 (2001); A. Gorlitz et al, Phys. Rev. Lett. 87, 130402 (2001); M. Greiner et al, Phys. Rev. Lett. 87, 160405 (2001); H. Moritz et al, Phys. Rev. Lett. 91, 250402 (2003); B. L. Tolra et al, Phys. Rev. Lett. 92, 190401 (2004); T. Stoferle et al, Phys. Rev. Lett. 92, 130403 (2004).
- [14] B. Paredes et al, Nature (London) 429, 377 (2004).
- [15] T. Kinoshita, T. Wenger, and D. S. Weiss, Nature (London) 440, 900 (2006).
- [16] D. Jaksch et al, Phys. Rev. Lett. 81, 3108 (1998).
- [17] D. van Oosten, P. van der Straten, and H. T. C. Stoof, Phys. Rev. A 63, 053601 (2001).
- [18] M. Greiner et al, Nature (London) 415, 39 (2002).
- [19] N. N. Bogoliubov, J. Phys. (USSR) 11, 23 (1947), reprinted in D. Pines, The Many-Body Problem, (W. A. Benjamin, New York, 1961).
- [20] A. Griffin, Phys. Rev. B 53, 9341 (1996).
- [21] N. P. Proukakis and K. Burnett, J. Res. Natl. Inst. Stand. Technol. 101 457 (1996); N. P. Proukakis, K. Burnett, and H. T. C. Stoof, Phys. Rev. A 57, 1230 (1998).
- [22] M. Holland, J. Park, and R. Walsler, Phys. Rev. Lett. 86, 1915 (2001).
- [23] A. M. Rey et al, Phys. Rev. A 69 033610 (2004)
- [24] V. N. Popov, Functional Integrals and Collective Excitations, (Cambridge University Press, Cambridge, 1987).
- [25] T. Gasenzer et al, Phys. Rev. A 72 063604 (2005).
- [26] A. Vardi and J. R. Anglin, Phys. Rev. Lett. 86 568 (2001).
- [27] J. R. Anglin and A. Vardi Phys. Rev. A 64 013605 (2001).
- [28] A. Smorzi et al, Phys. Rev. Lett. 79 4950 (1997).
- [29] S. Raghavan et al, Phys. Rev. A 59 620 (1999).