Nonequilibrium Bose systems and nonground-state Bose-Einstein condensates

V.I. Yukalov^{1,2}

¹Bogolubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna 141980, Russia

> ² Theoretische Physik, Universität Essen, Universitätsstrasse 5, Essen 45117, Germany

Abstract

The theory of resonant generation of nonground-state Bose-Einstein condensates is extended to Bose-condensed systems at finite temperature. The generalization is based on the notion of representative statistical ensembles for Bose systems with broken global gauge symmetry. Self-consistent equations are derived describing an arbitrary nonequilibrium nonuniform Bose system. The notion of finite-temperature topological coherent modes, coexisting with a cloud of noncondensed atoms, is introduced. It is shown that resonant generation of these modes is feasible for a gas of trapped Bose atoms at finite temperature.

Key words: Bose systems; representative statistical ensembles; broken gauge symmetry; conserving and gapless theory; condensate wave function; Bose-Einstein condensates; topological coherent modes

PACS numbers: 03.75.Hh, 03.75.Kk, 03.75.Nt, 05.30.Ch, 05.30.Jp, 05.45.Yv, 05.70.Ce, 67.40.Db

1 Introduction

Statistical systems with Bose-Einstein condensate exhibit a variety of very interesting phenomena, both equilibrium as well as nonequilibrium (see, e.g., book [1] and review articles [2-5]). One such a nontrivial effect is the possibility of creating nonground-state Bose-Einstein condensates of trapped atoms, as was advanced in Ref. [6]. The properties of these nonground-state condensates have been studied in a number of papers, for instance in Refs. [6-19], and are recently summarized in survey [20]. All these works [6-20] deal solely with the case of trapped atoms forming a dilute gas at zero-temperature, when all atoms can be condensed, so that there is no noncondensed atoms. However the latter are always present in real experiments at finite temperature. Interactions between atoms also produce an admixture of noncondensed particles. Then the question arises whether the nonground-state condensates can be generated, when the trapped atoms do not form a purely condensed system, but there is also a cloud of noncondensed atoms. To answer the above question, it is necessary to be based on a reliable general theory of nonequilibrium nonuniform Bose-condensed systems. The description of such systems is commonly done by employing the Bogolubov ideas [21–24], when the global gauge symmetry is broken by means of the Bogolubov shift. However, Hohenberg and Martin [25], first, emphasized that all theories of Bose systems with broken gauge symmetry suffer from an internal inconsistency, being either nonconserving or displaying an unphysical gap in the collective spectrum. In both these cases, such theories do not correspond to stable statistical systems. A detailed discussion of this problem has recently been done by Andersen [3].

To overcome the standard deficiency of theories with broken gauge symmetry, it is necessary to invoke the notion of representative ensembles [26]. This notion is strictly formulated in Sec. 2 for arbitrary statistical systems. The appropriate representative ensemble for a Bose system with broken gauge symmetry is constructed in Sec. 3. The resulting evolution equations are derived from the general variational principle for the extremum of action functional, which makes all equations self-consistent and the theory conserving and gapless, as is shown in Sec. 4. The derived evolution equation for the condensate wave function is analysed in Sec. 5, where topological coherent modes at finite temperature are defined. Resonant generation of these coherent modes, corresponding to nonground-state condensates, is demonstrated to be feasible even in the presence of a substantial admixture of noncondensed atoms.

Throughout the paper, the system of units is used, where the Planck and Boltzmann constants are set to unity, $\hbar = 1$, $k_B = 1$.

2 Representative statistical ensembles

Statistical systems are characterized by statistical ensembles, which means the following. First, one has to specify a space \mathcal{F} of microstates spanning all admissible quantum or dynamical states of the given system. Second, a statistical operator $\hat{\rho}(0)$ at the initial time t = 0has to be fixed. Third, temporal evolution of either the statistical operator or of observable quantities is to be defined, which is given by defining the action of the time derivative $\partial/\partial t$. Thus, a general *nonequilibrium statistical ensemble* can be denoted as a triplet

$$\left\{ \mathcal{F}, \hat{\rho}(0), \frac{\partial}{\partial t} \right\} . \tag{1}$$

For stationary or, in particular, equilibrium statistical systems, when their evolution is either trivial or absent, an equilibrium statistical ensemble is a pair $\{\mathcal{F}, \hat{\rho}\}$, where $\hat{\rho}(t) = \hat{\rho}(0) \equiv \hat{\rho}$.

The temporal evolution equation of operators in the Heisenberg representation can be symbolized with the help of the evolution operator U(t), so that the evolution of an operator $\hat{C}(t)$ is given by the relation

$$\hat{C}(t) = \hat{U}^+(t)\hat{C}(0)\hat{U}(t)$$
.

Then the *statistical average* of this operator is

$$\langle \hat{C}(t) \rangle \equiv \operatorname{Tr} \hat{\rho}(0)\hat{C}(t) = \operatorname{Tr} \hat{\rho}(t)\hat{C}(0)$$

where the trace operator is accomplished over the given state of microstates \mathcal{F} .

Each statistical system is characterized by a set of dynamical variables. The latter, keeping in mind quantum statistical systems, can be called field variables, whose example is a set $\psi(x,t) = [\psi_j(x,t)]$ of field operators $\psi_j(x,t)$, with an index j enumerating the set members. Here t is time and x is a collection of all other variables, which, e.g., could be spatial coordinates or momenta.

The system energy operator is given by a Hamiltonian $H[\psi]$, which is a functional of the field variables ψ . The related Lagrangian is

$$\hat{L}[\psi] \equiv \int \psi^{\dagger}(x,t) \ i \ \frac{\partial}{\partial t} \ \psi(x,t) \ dx \ - \ \hat{H}[\psi] \ , \tag{2}$$

where the set $\psi = [\psi_j]$ can be treated as a column.

Strictly speaking, for correctly defining a statistical system, it is not always sufficient to fix just a Hamiltonian $\hat{H}[\psi]$ or a Lagrangian $\hat{L}[\psi]$, but it is necessary to formulate additional conditions or constraints for making the system uniquely defined. Suppose there is a family $\{\hat{C}_i[\psi]\}$ of self-adjoint operators $\hat{C}_i[\psi] = \hat{C}_i^+[\psi]$, which will be called *condition operators*, when the required additional constraints, imposed on the system, are formulated as *statistical conditions*

$$C_i = \langle \hat{C}_i[\psi] \rangle , \qquad (3)$$

defined as the statistical averages of the condition operators. A statistical ensemble can correctly represent the considered statistical system only when all appropriate statistical conditions are accurately taken into account.

Representative statistical ensemble is a statistical ensemble that correctly represents the given statistical system, uniquely defining all its physical properties. The construction of representative ensembles for equilibrium systems is described in Ref. [27]. Then the equilibrium statistical operators are obtained from the conditional minimization of information functionals [28]. Now, we shall generalize the definition of representative ensembles to arbitrary nonequilibrium statistical systems.

Let us consider a physical system, whose correct definition requires the validity of statistical conditions (3) for some condition operators. It is worth stressing that the latter are assumed to be self-adjoint, but they do not need to be compulsorily the integrals of motion. For instance, the normalization condition $N = \langle \hat{N} \rangle$ for the total number of particles Ninvolves the number-of-particle operator \hat{N} , which does not commute with the Hamiltonian, when the global gauge symmetry is broken.

The most general way of deriving the evolution equations is by extremizing an action functional [29]. For a system under constraints, imposed by the statistical conditions (3), this implies the conditional extremization of the *effective action*

$$A[\psi] \equiv \int \left\{ \hat{L}[\psi] - \sum_{i} \nu_i \hat{C}_i[\psi] \right\} dt , \qquad (4)$$

with the Lagrange multipliers ν_i guaranteeing the validity of constraints (3). Defining the grand Hamiltonian

$$H[\psi] \equiv \hat{H}[\psi] + \sum_{i} \nu_i \hat{C}_i[\psi] , \qquad (5)$$

and using Lagrangian (2), we have for the action functional (4) the form

$$A[\psi] = \int \left\{ \int \psi^{\dagger}(x,t) \ i \ \frac{\partial}{\partial t} \ \psi(x,t) - H[\psi] \right\} \ dt \ . \tag{6}$$

The extremization of the effective action, defined by the variation

$$\delta A[\psi] = 0 \tag{7}$$

with respect to the field variables ψ and ψ^{\dagger} , implies the validity of the variational equations

$$\frac{\delta A[\psi]}{\delta \psi_j^{\dagger}(x,t)} = 0 , \qquad \frac{\delta A[\psi]}{\delta \psi_j(x,t)} = 0 .$$
(8)

In view of action (6), this is equivalent to the equations

$$i \frac{\partial}{\partial t} \psi_j(x,t) = \frac{\delta H[\psi]}{\delta \psi_j^{\dagger}(x,t)} \tag{9}$$

and their Hermitian conjugate.

Thus, we obtain the evolution equations (9) for the field variables. As is seen, the evolution is governed by the grand Hamiltonian (5). In the Heisenberg representation, a field operator ψ_j satisfies the Heisenberg equation of motion

$$i \frac{\partial}{\partial t} \psi_j(x,t) = [\psi_j(x,t), H[\psi]] ,$$

which is equivalent to Eq. (9). Then the time evolution of the field operator is described by the relation

$$\psi_j(x,t) = \hat{U}^+(t)\psi_j(x,0)\hat{U}(t)$$

with the evolution operator satisfying the Schrödinger equation

$$i \frac{d}{dt} \hat{U}(t) = H[\psi(x,0)]\hat{U}(t)$$
 (10)

The evolution equations, either (9) or (10), is a necessary component for defining a nonequilibrium statistical ensemble. This ensemble is representative, since the evolution equations are derived with taking account of all statistical conditions uniquely characterizing the considered statistical system. The given additional constraints define the grand Hamiltonian (5) governing the system evolution. As is clear, the properties of a system under the given constraints can be essentially different from the properties of a system under other or without constraints. This is why the usage of representative ensembles is crucially important for correctly describing physical systems. The general nonequilibrium representative ensemble, in the case of an equilibrium system, reduces to the equilibrium representative ensemble with the same grand Hamiltonian (5).

3 Broken gauge symmetry

Now we shall use the notion of representative ensembles, formulated above, for developing a correct general theory for Bose systems with broken gauge symmetry, which is associated with the appearance of Bose-Einstein condensate. The standard way of breaking the global gauge symmetry is by means of the Bogolubov shift

$$\psi(\mathbf{r},t) \longrightarrow \hat{\psi}(\mathbf{r},t) \equiv \eta(\mathbf{r},t) + \psi_1(\mathbf{r},t) , \qquad (11)$$

in which $\eta(\mathbf{r}, t)$ is the condensate wave function and $\psi_1(\mathbf{r}, t)$ is the field operator of uncondensed atoms, with \mathbf{r} being the spatial coordinate. The operators ψ and ψ_1 satisfy the same Bose commutation relations. The passage from ψ to ψ_1 corresponds to a canonical transformation realizing nonequivalent operator representations [30]. The Fock space $\mathcal{F}(\psi)$, generated by the field operator ψ^{\dagger} , characterizes the space of microstates for a system without Bose-Einstein condensate, while the Fock space $\mathcal{F}(\psi_1)$, generated by the field operator ψ_1^{\dagger} , is the space of microstates for a system with Bose-Einstein condensate. The spaces $\mathcal{F}(\psi)$ and $\mathcal{F}(\psi_1)$ are mutually orthogonal [30]. The condensate wave function $\eta(\mathbf{r}, t)$ is the same as the coherent field. The field variables of condensed and uncondensed atoms are mutually orthogonal, such that

$$\int \eta^*(\mathbf{r}, t) \psi_1(\mathbf{r}, t) \, d\mathbf{r} = 0 \; . \tag{12}$$

The Bogolubov shift (11) is sufficient for breaking the global gauge symmetry. Note that the method of infinitesimal sources is not always able to break the gauge symmetry [26].

Having now two field variables, η and ψ_1 , instead of just one ψ , requires to have two normalization conditions. One is the normalization of the condensate wave function to the number of condensed atoms

$$N_0 \equiv \int |\eta(\mathbf{r}, t)|^2 d\mathbf{r} , \qquad (13)$$

where N_0 , in general, can be a function of time, which is not marked explicitly just for brevity. Defining the operator $\hat{N}_0 \equiv N_0 \hat{1}$, with $\hat{1}$ being a unity operator in $\mathcal{F}(\psi_1)$, we can represent normalization (13) in the standard form of a statistical condition

$$N_0 = \langle \hat{N}_0 \rangle , \qquad (14)$$

as in Eq. (3). Here and in what follows, all statistical averages are accomplished over the space of microstates $\mathcal{F}(\psi_1)$.

The second normalization condition is that one for the number of uncondensed atoms

$$N_1 = \langle \hat{N}_1 \rangle , \qquad (15)$$

where the operator for the number of uncondensed atoms is

$$\hat{N}_1 \equiv \int \psi_1^{\dagger}(\mathbf{r}, t) \psi_1(\mathbf{r}, t) \, d\mathbf{r} \,. \tag{16}$$

In general, N_1 can also be a function of time in a nonequilibrium system.

The total number of atoms is

$$N = N_0 + N_1 , (17)$$

which could be considered as a normalization condition, instead of Eq. (15). For two field variables, we have to fix not less and not more than two normalization conditions.

The atomic densities and the related atomic fractions are denoted as

$$\rho_0 \equiv \frac{N_0}{V} , \qquad \rho_1 \equiv \frac{N_1}{V} , \qquad \rho \equiv \frac{N}{V} ,$$
$$n_0 \equiv \frac{N_0}{N} = \frac{\rho_0}{\rho} , \qquad n_1 \equiv \frac{N_1}{N} = \frac{\rho_1}{\rho} , \qquad (18)$$

where V is the system volume. For these, one has

$$\rho = \rho_0 + \rho_1 , \qquad n_0 + n_1 = 1 . \tag{19}$$

It is assumed that ρ_0 and n_0 are not zero in the thermodynamic limit.

The number of condensed atoms N_0 is defined so that to make the system stable. In equilibrium, N_0 is to be found from the minimization of a thermodynamic potential. For a nonequilibrium system, N_0 must ensure the dynamic stability of the solution for the condensate function $\eta(\mathbf{r}, t)$ satisfying the related evolution equation.

For a system with broken gauge symmetry, the average $\langle \psi_1 \rangle$ may be nonzero. This, however, would mean that quantum numbers, such as spin or momentum, are not conserved. One, therefore, has to impose an additional constraint

$$\langle \psi_1(\mathbf{r},t) \rangle = 0$$
. (20)

By introducing the condition operator

$$\hat{\Lambda}[\hat{\psi}] \equiv \int \left[\lambda(\mathbf{r}, t) \psi_1^{\dagger}(\mathbf{r}, t) + \lambda^*(\mathbf{r}, t) \psi_1(\mathbf{r}, t) \right] dt , \qquad (21)$$

in which $\lambda(\mathbf{r}, t)$ is a complex function, constraint (20) can be rewritten in the standard form (3) as the quantum-number conservation condition

$$\langle \hat{\Lambda}[\hat{\psi}] \rangle = 0$$
. (22)

The grand Hamiltonian (5) has to be defined by taking into account the statistical conditions (14), (15), and (22), which yields

$$H[\eta, \psi_1] = \hat{H}[\hat{\psi}] - \mu_0 \hat{N}_0 - \mu_1 \hat{N}_1 - \Lambda[\hat{\psi}] .$$
(23)

The action functional (6), with the Bogolubov shift (11), takes the form

$$A[\eta,\psi_1] = \int \left\{ \int \left[\eta^*(\mathbf{r},t) \ i \ \frac{\partial}{\partial t} \ \eta(\mathbf{r},t) + \psi_1^{\dagger}(\mathbf{r},t) \ i \ \frac{\partial}{\partial t} \ \psi_1(\mathbf{r},t) \right] \ d\mathbf{r} - H[\eta,\psi_1] \right\} \ dt \ .$$
(24)

The extremization of the action functional (24) implies two variational equations, for the condensate function,

$$\frac{\delta A[\eta, \psi_1]}{\delta \eta^*(\mathbf{r}, t)} = 0 , \qquad (25)$$

and for the field operator of uncondensed atoms,

$$\frac{\delta A[\eta, \psi_1]}{\delta \psi_1^{\dagger}(\mathbf{r}, t)} = 0.$$
(26)

These equations, in view of the action functional (24), are equivalent to the evolution equations

$$i \frac{\partial}{\partial t} \eta(\mathbf{r}, t) = \frac{\delta H[\eta, \psi_1]}{\delta \eta^*(\mathbf{r}, t)}$$
(27)

and

$$i \frac{\partial}{\partial t} \psi_1(\mathbf{r}, t) = \frac{\delta H[\eta, \psi_1]}{\delta \psi_1^{\dagger}(\mathbf{r}, t)} .$$
(28)

Let us take the energy operator in the standard form

$$\hat{H}[\hat{\psi}] = \int \hat{\psi}^{\dagger}(\mathbf{r}, t) \left(-\frac{\nabla^2}{2m} + U\right) \hat{\psi}(\mathbf{r}, t) d\mathbf{r} + \frac{1}{2} \int \hat{\psi}^{\dagger}(\mathbf{r}, t) \hat{\psi}^{\dagger}(\mathbf{r}', t) \Phi(\mathbf{r} - \mathbf{r}') \hat{\psi}(\mathbf{r}', t) \hat{\psi}(\mathbf{r}, t) d\mathbf{r} d\mathbf{r}' , \qquad (29)$$

in which $U = U(\mathbf{r}, t)$ is an external field and $\Phi(\mathbf{r}) = \Phi(-\mathbf{r})$ is an interaction potential. To satisfy the conservation condition (22), the grand Hamiltonian (23) must have no linear in ψ_1 terms [31]. This is achieved by chosing the Lagrange multiplier

$$\lambda(\mathbf{r},t) = \left[-\frac{\nabla^2}{2m} + U + \int \Phi(\mathbf{r} - \mathbf{r}') |\eta(\mathbf{r}',t)|^2 d\mathbf{r}'\right] \eta(\mathbf{r},t) .$$
(30)

The evolution equations (27) and (28) yield

$$i \frac{\partial}{\partial t} \eta(\mathbf{r}, t) = \left(-\frac{\nabla^2}{2m} + U - \mu_0 \right) \eta(\mathbf{r}, t) + \int \Phi(\mathbf{r} - \mathbf{r}') \left[|\eta(\mathbf{r}', t)|^2 \eta(\mathbf{r}) + \hat{X}(\mathbf{r}, \mathbf{r}') \right] d\mathbf{r}'$$
(31)

and, respectively,

$$i \frac{\partial}{\partial t} \psi_1(\mathbf{r}, t) = \left(-\frac{\nabla^2}{2m} + U - \mu_1\right) \psi_1(\mathbf{r}, t) + \int \Phi(\mathbf{r} - \mathbf{r}') \left[|\eta(\mathbf{r}')|^2 \psi_1(\mathbf{r}) + \eta^*(\mathbf{r}')\eta(\mathbf{r})\psi_1(\mathbf{r}') + \eta(\mathbf{r}')\eta(\mathbf{r})\psi_1^{\dagger}(\mathbf{r}') + \hat{X}(\mathbf{r}, \mathbf{r}')\right] d\mathbf{r}', \qquad (32)$$

where, for brevity, the time-dependence is not explicitly shown in the right-hand sides of these equations, and the correlation operator

$$\hat{X}(\mathbf{r},\mathbf{r}') \equiv \psi_1^{\dagger}(\mathbf{r}')\psi_1(\mathbf{r}')\eta(\mathbf{r}) + \psi_1^{\dagger}(\mathbf{r}')\eta(\mathbf{r}')\psi_1(\mathbf{r}) + \eta^*(\mathbf{r}')\psi_1(\mathbf{r}')\psi_1(\mathbf{r}) + \psi_1^{\dagger}(\mathbf{r}')\psi_1(\mathbf{r}')\psi_1(\mathbf{r})$$
(33)

is introduced.

To obtain an equation for the condensate wave function, we need to average Eq. (31). For this purpose, let us define the normal density matrix

$$\rho_1(\mathbf{r}, \mathbf{r}') \equiv \langle \psi_1^{\dagger}(\mathbf{r}')\psi_1(\mathbf{r}) \rangle, \qquad (34)$$

the anomalous density matrix

$$\sigma_1(\mathbf{r}, \mathbf{r}') \equiv \langle \psi_1(\mathbf{r}')\psi_1(\mathbf{r}) \rangle, \qquad (35)$$

and the densities

$$\rho_0(\mathbf{r}) \equiv |\eta(\mathbf{r})|^2 , \qquad \rho_1(\mathbf{r}) \equiv \rho_1(\mathbf{r}, \mathbf{r}) = \langle \psi_1^{\dagger}(\mathbf{r})\psi_1(\mathbf{r}) \rangle ,$$

$$\sigma_1(\mathbf{r}) \equiv \sigma_1(\mathbf{r}, \mathbf{r}) = \langle \psi_1(\mathbf{r})\psi_1(\mathbf{r}) \rangle . \qquad (36)$$

The total density of atoms is the sum

$$\rho(\mathbf{r}) = \rho_0(\mathbf{r}) + \rho_1(\mathbf{r}) \tag{37}$$

of the condensate density $\rho_0(\mathbf{r})$ and of the density $\rho_1(\mathbf{r})$ of uncondensed atoms. Also, let us use the notation

$$\xi(\mathbf{r},\mathbf{r}') \equiv \langle \psi_1^{\dagger}(\mathbf{r}')\psi_1(\mathbf{r}')\psi_1(\mathbf{r}) \rangle \quad .$$
(38)

Then the average of the correlation operator (33) becomes

$$\langle \hat{X}(\mathbf{r},\mathbf{r}')\rangle = \rho_1(\mathbf{r}')\eta(\mathbf{r}) + \rho_1(\mathbf{r},\mathbf{r}')\eta(\mathbf{r}') + \sigma_1(\mathbf{r},\mathbf{r}')\eta^*(\mathbf{r}') + \xi(\mathbf{r},\mathbf{r}') .$$
(39)

Finally, averaging Eq. (31), we find the equation for the condensate function

$$i \frac{\partial}{\partial t} \eta(\mathbf{r}, t) = \left(-\frac{\nabla^2}{2m} + U - \mu_0\right) \eta(\mathbf{r}) + \int \Phi(\mathbf{r} - \mathbf{r}') \left[\rho(\mathbf{r}')\eta(\mathbf{r}) + \rho_1(\mathbf{r}, \mathbf{r}')\eta(\mathbf{r}') + \sigma_1(\mathbf{r}, \mathbf{r}')\eta^*(\mathbf{r}') + \xi(\mathbf{r}, \mathbf{r}')\right] d\mathbf{r}', \quad (40)$$

where again, for brevity, the temporal dependence in the right-hand side is not explicitly shown. This is a general equation valid for an arbitrary nonequilibrium Bose-condensed system.

It is worth noting that, contrary to Eq. (31), the average of Eq. (32) is not defined because of the following. The grand Hamiltonian does not change being complimented by the term

$$z[\psi_1] \equiv \int \langle \zeta(\mathbf{r},t)\psi_1^{\dagger}(\mathbf{r},t) + \zeta^*(\mathbf{r},t)\psi_1(\mathbf{r},t) \rangle d\mathbf{r} ,$$

which is identically zero owing to constraint (20) for any complex function $\zeta(\mathbf{r}, t)$. However, the variation of such a term results in

$$\frac{\delta z[\psi_1]}{\delta \psi_1^{\dagger}(\mathbf{r},t)} = \zeta(\mathbf{r},t)$$

which, generally, is not zero. Therefore, replacing $H[\eta, \psi_1]$ by $H[\eta, \psi_1] + z[\psi_1]$ would lead to the appearance in Eq. (28) of an additional undefined term $\zeta(\mathbf{r}, t)$. Equation (32) must be used for defining the evolution of the correlation functions $\langle \psi_1^{\dagger}\psi_1 \rangle$ and $\langle \psi_1\psi_1 \rangle$. But for these functions, an additive term ζ does not play any role, since $\langle \zeta\psi_1^{\dagger} \rangle = \langle \zeta\psi_1 \rangle = 0$ due to constraint (20).

4 Self-consistent equations

The normal and anomalous density matrices (34) and (35), entering the evolution equation (40) for the condensate function, can be expressed through Green functions. One needs the normal and anomalous Green functions [32–34], which can be assembled in a matrix $G(12) = [G_{\alpha\beta}(12)]$ with the elements

$$G_{11}(12) = -i < \hat{T}\psi_1(1)\psi_1^{\dagger}(2) > , \qquad G_{12}(12) = -i < \hat{T}\psi_1(1)\psi_1(2) > ,$$

$$G_{21}(12) = -i < \hat{T}\psi_1^{\dagger}(1)\psi_1^{\dagger}(2) > , \qquad G_{22}(12) = -i < \hat{T}\psi_1^{\dagger}(1)\psi_1(2) > , \qquad (41)$$

where \hat{T} is the chronological operator. Here and in what follows, the shorthand notation is used, denoting the set $\{\mathbf{r}_j, t_j\}$ just by the number j.

We shall also need the combination

$$\Psi(123) \equiv \psi_1(1)\psi_1(2)\psi_1^{\dagger}(3) + \eta(1)\psi_1(2)\psi_1^{\dagger}(3) + \psi_1(1)\eta(2)\psi_1^{\dagger}(3) + \psi_1(1)\psi_1(2)\eta^*(3) + \eta(1)\eta(2)\psi_1^{\dagger}(3) + \eta(1)\psi_1(2)\eta^*(3) + \psi_1(1)\eta(2)\eta^*(3) .$$
(42)

Using this, we define the binary Green function, which is a matrix $B(1234) = [B_{\alpha\beta}(1234)]$ with the elements

$$B_{11}(1234) = -\langle \hat{T}\Psi(123)\psi_1^{\dagger}(4) \rangle, \qquad B_{12}(1234) = -\langle \hat{T}\Psi(123)\psi_1(4) \rangle,$$

$$B_{21}(1234) = -\langle \hat{T}\Psi^{+}(123)\psi_{1}^{\dagger}(4) \rangle, \qquad B_{22}(1234) = -\langle \hat{T}\Psi^{+}(123)\psi_{1}(4) \rangle .$$
(43)

We may note that

$$\lim_{t_{12}\to -0} \langle \hat{T}\Psi(122) \rangle = \rho_1(2)\eta(1) + \rho_1(12)\eta(2) + \sigma_1(12)\eta^*(2) + \xi(12) = \langle \hat{X}(12) \rangle,$$

where $t_{12} = t_1 - t_2$.

Introducing the retarded interaction potential

$$\Phi(12) \equiv \Phi(\mathbf{r}_1 - \mathbf{r}_2)\delta(t_{12} + 0) , \qquad (44)$$

we define the self-energy $\Sigma(12)$ by the relation

$$\int \Sigma(13)G(32) \ d(3) = i \int \Phi(13)B(1332) \ d(3) \ . \tag{45}$$

Then, from Eq. (32), we obtain the matrix equation

$$\left(\hat{\tau}i \ \frac{\partial}{\partial t_1} + \frac{\nabla_1^2}{2m} \ - \ U(1) + \mu_1\right) G(12) - \int \Sigma(13) G(32) \ d(3) = \delta(12)\hat{1} \ , \tag{46}$$

in which

$$\hat{\tau} \equiv \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$
, $\hat{1} \equiv \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$.

Equation (46) for the Green function is to be complemented by equation (40) for the condensate function, which can be represented as

$$i \frac{\partial}{\partial t_1} \eta(1) = \left(-\frac{\nabla_1^2}{2m} + U(1) - \mu_0 \right) \eta(1) + \int \Phi(12) \left[\rho_0(2)\eta(1) + \langle \hat{T}\Psi(122) \rangle \right] d(2) , \quad (47)$$

where $\rho_0(1) \equiv |\eta(1)|^2$.

Equations (46) and (47) is a self-consistent set of equations derived from the extremization of the action functional (24). Hence, these equations respect all conservation laws of the Hamiltonian, as is should be for the equations derived from a variational procedure [3,35]. It is important to stress that the self-consistency of the equations is ensured by two, generally different, Lagrange multipliers μ_0 and μ_1 , playing the role of the chemical potentials for condensed and uncondensed atoms, respectively. The potential μ_0 guarantees the normalization condition (14), hence, $\mu_0 = \mu_0(N_0)$. The normalization condition (15) defines $\mu_1 = \mu_1(N_1)$. The number of condensed atoms is to be such that to provide the stability of the system of N atoms, so that $N_0 = N_0(N)$. Because of Eq. (17), $N_1 = N - N_0$. Therefore, $\mu_0 = \mu_0(N)$ and $\mu_1 = \mu_1(N)$. But there is no necessity that μ_0 be equal to μ_1 , though it may occasionally happen. Thus, this happens in the Bogolubov approximation [21–24], which is valid for asymptotically weak interactions. However, for more elaborate approximations, this is not so. To illustrate this fact, let us consider an equilibrium system, when $\eta(\mathbf{r}, t) = \eta(\mathbf{r})$ does not depend on time. Then Eq. (47) becomes

$$\left(-\frac{\nabla^2}{2m} + U\right)\eta(\mathbf{r}) + \int \Phi(\mathbf{r} - \mathbf{r}')\left[\rho(\mathbf{r}')\eta(\mathbf{r}) + \rho_1(\mathbf{r}, \mathbf{r}')\eta(\mathbf{r}') + \sigma_1(\mathbf{r}, \mathbf{r}')\eta^*(\mathbf{r}') + \xi(\mathbf{r}, \mathbf{r}')\right] d\mathbf{r}' = \mu_0\eta(\mathbf{r}) .$$
(48)

The grand thermodynamic potential is

$$\Omega = -T \ln \operatorname{Tr} \exp\left(-\beta H[\eta, \psi_1]\right) , \qquad (49)$$

and the free energy being

$$F = -T \ln \operatorname{Tr} \exp\left\{-\beta \left(\hat{H}[\hat{\psi}] - \hat{\Lambda}[\hat{\psi}]\right)\right\} , \qquad (50)$$

where $\beta = T^{-1}$ is inverse temperature. For potentials (49) and (50), we have the relation

$$\Omega = F - \mu_0 N_0 - \mu_1 N_1 . \tag{51}$$

The number of condensed atoms is defined by the stability condition implying the minimum of the grand potential (49),

$$\frac{\partial\Omega}{\partial N_0} = 0 , \qquad \frac{\partial^2\Omega}{\partial N_0^2} > 0 .$$
(52)

More generally, the condensate wave function, assuring the system stability, is defined by the condition

$$\frac{\delta\Omega}{\delta\eta(\mathbf{r})} = \langle \frac{\delta H[\eta,\psi_1]}{\delta\eta(\mathbf{r})} \rangle = 0.$$
(53)

The latter, owing to Eq. (27), gives Eq. (48).

Since $N_0 = N_0(N)$ and $N_1 = N_1(N)$, it is the total number of atoms N that can only be fixed for a system, which requires the relation

$$\Omega = F - \mu N , \qquad (54)$$

in which μ is the system chemical potential. Comparing Eqs. (51) and (54), immediately results in the definition of the system chemical potential

$$\mu = n_0 \mu_0 + n_1 \mu_1 . \tag{55}$$

For the latter, one has the standard thermodynamic equations

$$\mu = \left(\frac{\partial F}{\partial N}\right)_{TV} , \qquad N = -\left(\frac{\partial \Omega}{\partial \mu}\right)_{TV}$$

Let us emphasize that, since N_0 and N_1 are uniquely defined through the total number of atoms N, neither N_0 nor N_1 can be treated as independent thermodynamic variables, because of which, generally, $\mu_0 \neq \mu_1$.

To give a more explicit demonstration that $\mu_0 \neq \mu_1$, let us turn to a uniform system, when the density matrices $\rho_1(\mathbf{r}, \mathbf{r}')$, $\sigma_1(\mathbf{r}, \mathbf{r}')$, as well as $\xi(\mathbf{r}, \mathbf{r}')$, depend solely on the difference $\mathbf{r} - \mathbf{r}'$. Then the densities $\rho_1 = \rho_1(\mathbf{r}, \mathbf{r})$, $\sigma_1(\mathbf{r}, \mathbf{r})$, and $\xi_1(\mathbf{r}, \mathbf{r})$ are constants, together with the order parameter $\eta = \eta(\mathbf{r})$. Also, normalization (14) yields $\eta = \sqrt{\rho_0}$. From Eq. (48), we find

$$\mu_0 = \rho \Phi_0 + \int \left(n_k + \sigma_k + \frac{\xi_k}{\sqrt{\rho_0}} \right) \Phi_k \, \frac{d\mathbf{k}}{(2\pi)^3} \,, \tag{56}$$

where Φ_k , n_k , σ_k , and ξ_k are the Fourier transforms of the corresponding quantities $\Phi(\mathbf{r})$, $\rho_1(\mathbf{r}, 0)$, $\sigma_1(\mathbf{r}, 0)$, and $\xi(\mathbf{r}, 0)$, respectively. For the Fourier transforms $G_{\alpha\beta}(\mathbf{k}, \omega)$ of the Green functions $G_{\alpha\beta}(12)$, we find the former from Eq. (46). The poles of $G_{\alpha\beta}(\mathbf{k}, \omega)$ define the single-atom spectrum, which is gapless, provided that

$$\mu_1 = \Sigma_{11}(0,0) - \Sigma_{12}(0,0) , \qquad (57)$$

where $\Sigma_{\alpha\beta}(\mathbf{k},\omega)$ is the Fourier transform of the self-energy $\Sigma_{\alpha\beta}(12)$. Note that the gapless spectrum is a necessary requirement for the existence of a stable system with Bose-Einstein condensate [30]. Equation (57) is the Hugenholtz-Pines relation that could be derived either from thermodynamic equations [33] or from the Ward identities with respect to the variation of gauge [24].

One often considers Bose-condensed systems in the contact-potential approximation, when

$$\Phi(\mathbf{r}) = \Phi_0 \delta(\mathbf{r}) \qquad \left(\Phi_0 \equiv 4\pi \; \frac{a_s}{m}\right) \; ,$$

with a_s being a scattering length. Then $\Phi_k = \Phi_0$, and Eq. (56) gives

$$\mu_0 = \left(\rho + \rho_1 + \sigma_1 + \frac{\xi}{\sqrt{\rho_0}}\right) \Phi_0 , \qquad (58)$$

where

$$\rho_1 = \int n_k \frac{d\mathbf{k}}{(2\pi)^3}, \qquad \sigma_1 = \int \sigma_k \frac{d\mathbf{k}}{(2\pi)^3}, \qquad \xi = \int \xi_k \frac{d\mathbf{k}}{(2\pi)^3}.$$

Clearly, there is no such a general law that would require the identity of μ_0 and μ_1 .

To be even more specific, let us consider the Hartree-Fock-Bogolubov approximation, when $\xi(\mathbf{r}, \mathbf{r}') = 0$. Then for Eq. (57), we find

$$\mu_1 = \rho \Phi_0 + \int (n_k - \sigma_k) \Phi_k \, \frac{d\mathbf{k}}{(2\pi)^3} \,. \tag{59}$$

In the case of the contact potential, we obtain

$$\mu_0 = (\rho + \rho_1 + \sigma_1)\Phi_0 , \qquad \mu_1 = (\rho + \rho_1 - \sigma_1)\Phi_0 .$$
(60)

As is evident, $\mu_0 \neq \mu_1$.

In the standard approach, without using the representative ensemble, one assumes that μ_0 equals μ_1 , hence, according to Eq. (55), equals μ . But then one comes to the explicit inconsistency in Eqs. (60). A common way of trying to treat this inconsistency is by neglecting the anomalous average σ_1 , calling this the "Popov approximation". However, this unjustified trick has nothing to do with Popov, as is easy to infer from his original works [36–38]. Moreover, it is easy to show that at low temperatures the anomalous average σ_1 can be much larger than the normal average ρ_1 , that is, neglecting σ_1 has nothing to do with a reasonable approximation [30,39].

By employing the representative ensemble, as is done in the present paper, we introduce two Lagrange multipliers, μ_0 and μ_1 , for two normalization conditions (14) and (15). These multipliers are not obliged to be equal, but are to be such that to render the whole theory completely self-consistent.

5 Topological coherent modes

Equation (48), defining the condensate function $\eta(\mathbf{r})$, corresponds to a stable equilibrium system, when condensing atoms pile down to the ground-state quantum level. This equation can be generalized for describing arbitrary stationary states $\eta_n(\mathbf{r})$, enumerated by a quantum multi-index n, so that, for the contact potential, we have

$$\left[-\frac{\nabla^2}{2m} + U(\mathbf{r})\right]\eta_n(\mathbf{r}) + \Phi_0\left\{\left[|\eta_n(\mathbf{r})|^2 + 2\rho_1^{(n)}(\mathbf{r})\right]\eta_n(\mathbf{r}) + \sigma_1^{(n)}(\mathbf{r})\eta_n^*(\mathbf{r}) + \xi^{(n)}(\mathbf{r})\right\} = E_n\eta_n(\mathbf{r}) , \qquad (61)$$

where $\rho_1^{(n)}(\mathbf{r})$, $\sigma_1^{(n)}(\mathbf{r})$, and $\xi^{(n)}(\mathbf{r})$ are the solutions to the self-consistent system of equations for $\rho_1(\mathbf{r})$, $\sigma_1(\mathbf{r})$, and $\xi(\mathbf{r}) \equiv \xi(\mathbf{r}, \mathbf{r})$, respectively, under the condition that $\eta(\mathbf{r})$ is replaced by $\eta_n(\mathbf{r})$. The ground-state energy level corresponds to

$$E_0 \equiv \min_n E_n = \mu_0 , \qquad (62)$$

with $\eta_0(\mathbf{r}) \equiv \eta(\mathbf{r})$. But, in general, there can exist a whole set of coherent states $\eta_n(\mathbf{r})$, with a spectrum of energies E_n . The condensate functions $\eta_n(\mathbf{r})$ will be called the *topological* coherent modes.

It is clear that, for a system at absolute equilibrium, the sole pertinent mode is the ground state $\eta(\mathbf{r})$. In order to generate excited coherent modes, it is necessary to deal with a nonequilibrium system. Then one has to consider the time-dependent equation (40) for the condensate function, which, for the contact potential, reads as

$$i \frac{\partial}{\partial t} \eta(\mathbf{r}, t) = \left(-\frac{\nabla^2}{2m} + U - \mu_0 \right) \eta(\mathbf{r}, t) + \Phi_0 \left\{ \left[\rho_0(\mathbf{r}, t) + 2\rho_1(\mathbf{r}, t) \right] \eta(\mathbf{r}, t) + \sigma_1(\mathbf{r}, t) \eta^*(\mathbf{r}, t) + \xi(\mathbf{r}, t) \right\} , \qquad (63)$$

where $U = U(\mathbf{r}, t)$ is a time-dependent external potential. The latter can be represented as a sum

$$U(\mathbf{r},t) = U(\mathbf{r}) + V(\mathbf{r},t) \tag{64}$$

of a trapping potential $U(\mathbf{r})$ and a modulating potential $V(\mathbf{r}, t)$.

If in Eq. (63) we set $\rho_1(\mathbf{r},t) = 0$, $\sigma_1(\mathbf{r},t) = 0$, and $\xi(\mathbf{r},t) = 0$, then we come to the usual temporal Gross-Pitaevskii equation [1]. The generation of various topological coherent modes on the basis of the latter equation has been investigated earlier [6–20,40]. However, in general, we have to analyse the full Eqs. (61) and (63).

Suppose that at the initial time t = 0, the atomic system is in equilibrium, so that

$$\eta(\mathbf{r},0) = \eta_0(\mathbf{r}) \equiv \eta(\mathbf{r}) . \tag{65}$$

Assume that we wish to generate a nonground-state condensate, corresponding to a topological coherent mode labelled by the index $n = n_1$, with the energy $E_1 \equiv E_{n_1}$. For this purpose, we impose an alternating modulating potential

$$V(\mathbf{r},t) = V_1(\mathbf{r})\cos\omega t + V_2(\mathbf{r})\sin\omega t , \qquad (66)$$

with the frequency ω tuned close to the transition frequency $\omega_1 \equiv E_1 - E_0$, which implies the resonance condition

$$\left|\frac{\Delta\omega}{\omega}\right| \ll 1 \qquad (\Delta\omega \equiv \omega - \omega_1) .$$
 (67)

Imposing an external alternating field will, of course, destroy the ground-state condensate by means of two processes. One is the resonant process of transferring condensed atoms from the ground state mode, with the energy $E_0 = \mu_0$, to the chosen excited mode, with the energy E_1 . And also, there will be nonresonant processes of taking atoms to other nonresonant coherent modes, as well as the process of transferring atoms from the condensate to the cloud of noncondensed atoms. Estimates show [15] that it is feasible to arrange such a setup that the resonant generation would occur much faster than other nonresonant processes. Assuming this, we may consider the solution of Eq. (63) at times shorter than the critical time t_c , when the resonant generation prevails, and the number of condensed atoms stays practically constant,

$$\int |\eta(\mathbf{r},t)|^2 d\mathbf{r} = \int |\eta(\mathbf{r})|^2 d\mathbf{r} = N_0 .$$
(68)

The topological coherent modes are also normalized to N_0 , because of which we can set

$$\eta_n(\mathbf{r}) = \sqrt{N_0} \,\varphi_n(\mathbf{r}) \,, \qquad \int |\varphi_n(\mathbf{r})|^2 d\mathbf{r} = 1 \,. \tag{69}$$

Then we look for the solution of Eq. (63) in the form of the mode expansion

$$\eta(\mathbf{r},t) = \sum_{n} C_n(t)\eta_n(\mathbf{r})e^{-i\omega_n t} , \qquad (70)$$

in which $\omega_n \equiv E_n - E_0$ and the coefficient functions $C_n(t)$ are assumed to be slow functions of time, as compared to the exponential $\exp(-i\omega_n t)$, which means the condition

$$\frac{1}{\omega_n} \left| \frac{dC_n}{dt} \right| \ll 1 . \tag{71}$$

From the normalization condition (68), with expansion (70), after averaging over time, while keeping the quasi-invariants $C_n(t)$ fixed, we get

$$\sum_{n} |C_n(t)|^2 = 1 .$$
(72)

Substituting expansion (70) into Eq. (63), we multiply the latter by $\eta_n^*(\mathbf{r}) \exp(i\omega_n t)$, integrate over \mathbf{r} , and average over time, following the averaging technique [6,14,15]. To this end, we need the following integrals

$$\begin{split} \alpha_{mn} &\equiv \Phi_0 N_0 \int |\varphi_m(\mathbf{r})|^2 \left[2|\varphi_n(\mathbf{r})|^2 - |\varphi_m(\mathbf{r})|^2 \right] \, d\mathbf{r} \;, \\ \beta_{mn} &\equiv \int \varphi_m^*(\mathbf{r}) \left[V_1(\mathbf{r}) - iV_2(\mathbf{r}) \right] \varphi_n(\mathbf{r}) \, d\mathbf{r} \;, \\ \overline{\alpha}_{nn}(t) &\equiv \alpha_{nn} - \Phi_0 \int \varphi_n^*(\mathbf{r}) \left\{ 2 \left[\rho_1^{(n)}(\mathbf{r}) - \rho_1(\mathbf{r}, t) \right] \varphi_n(\mathbf{r}) + \sigma_1^{(n)}(\mathbf{r}) \varphi_n^*(\mathbf{r}) + \frac{\xi^{(n)}(\mathbf{r})}{\sqrt{N_0}} \right\} \, d\mathbf{r} \;, \\ \alpha_{nn} &= \Phi_0 N_0 \int |\varphi_n(\mathbf{r})|^4 d\mathbf{r} \;, \qquad \Delta_{mn} \equiv \Delta \omega + \alpha_{00} - \alpha_{11} \;. \end{split}$$

We presume that, at the resonant stage, when $t \ll t_c$, the quantity $\overline{\alpha}_{nn}(t)$ is a slow function of time, such that

$$\frac{t_c}{|\overline{\alpha}_{nn}|} \left| \frac{d\overline{\alpha}_{nn}}{dt} \right| \ll 1 .$$
(73)

Then, introducing the functions

$$c_n(t) \equiv C_n(t) \exp\left\{i\overline{\alpha}_{nn}(t)t\right\} , \qquad (74)$$

we, finally, arrive at the system of two equations

$$i \frac{dc_0}{dt} = \alpha_{01} |c_1|^2 c_0 + \frac{1}{2} \beta_{01} c_1 \exp(i\Delta_{01}t) ,$$

$$i \frac{dc_0}{dt} = \alpha_{10} |c_0|^2 c_1 + \frac{1}{2} \beta_{01}^* c_0 \exp(-i\Delta_{01}t) .$$
(75)

Equations (75) have the same form as studied in the previous publications [6,8,10,14–16,20] describing the resonant generation of topological coherent modes on the basis of the Gross-Pitaevskii equation.

The reduction of the general equation for the condensate function (63) to the system of Eqs. (75) demonstrates that the topological coherent modes, corresponding to nongroundstate condensates, can be resonantly generated also in the presence of the cloud of uncondensed atoms. The procedure of the resonant generation requires the validity of several conditions discussed above. In particular, the resonant process must be much faster than nonresonant ones. After the critical time t_c , power broadening destroys the resonant procedure and nonresonant processes become prevailing. According to estimates [15], the critical time is of the order of $t_c \sim \omega/(\alpha^2 + \beta^2)$, where $\alpha \equiv (\alpha_{01} + \alpha_{10})/2$ and $\beta \equiv |\beta_{01}|$. At the initial stage of time $t \ll t_c$, the resonant generation is feasible. The critical time t_c can be made comparable to the lifetime of atoms in a trap [15].

References

- L. Pitaevskii and S. Stringari, Bose-Einstein Condensation in Dilute Gases (Clarendon, Oxford, 2003).
- [2] P.W. Courteille, V.S. Bagnato, and V.I. Yukalov, Laser Phys. 11, 659 (2001).
- [3] J.O. Andersen, Rev. Mod. Phys. **76**, 599 (2004).
- [4] K. Bongs and K. Sengstock, Rep. Prog. Phys. 67, 907 (2004).
- [5] V.I. Yukalov, Laser Phys. Lett. 1, 435 (2004).
- [6] V.I. Yukalov, E.P. Yukalova, and V.S. Bagnato, Phys. Rev. A 56, 4845 (1997).
- [7] K.P. Marzlin and W. Zhang, Phys. Rev. A 57, 4761 (1998).
- [8] V.I. Yukalov, E.P. Yukalova, and V.S. Bagnato, Laser Phys. 10, 26 (2000).
- [9] Y.S. Kivshar, T.J. Alexander, and S.K. Turitsyn, Phys. Lett. A 278, 225 (2001).
- [10] V.I. Yukalov, E.P. Yukalova, and V.S. Bagnato, Laser Phys. 11, 455 (2001).
- [11] R. D'Agosta, B.A. Malomed, and C. Presilla, Laser Phys. **12**, 37 (2002).
- [12] R. D'Agosta and C. Presilla, Phys. Rev. A 65, 043609 (2002).
- [13] N.P. Proukakis and L. Lambropoulos, Eur. Phys. J. D **19**, 355 (2002).
- [14] V.I. Yukalov and E.P. Yukalova, J. Phys. A **35**, 8603 (2002).
- [15] V.I. Yukalov, E.P. Yukalova, and V.S. Bagnato, Phys. Rev. A 66, 043602 (2002).
- [16] V.I. Yukalov, E.P. Yukalova, and V.S. Bagnato, Laser Phys. **13**, 861 (2003).
- [17] S.K. Adhikari, Phys. Lett. A **308**, 302 (2003).

- [18] V.I. Yukalov, K.P. Marzlin, and E.P. Yukalova, Phys. Rev. A 69, 023620 (2004).
- [19] S.K. Adhikari, Phys.Rev. A **69**, 063613 (2004).
- [20] V.I. Yukalov, K.P. Marzlin, E.P. Yukalova, and V.S. Bagnato, Am. Inst. Phys. Conf. Proc. 770, 218 (2005).
- [21] N.N. Bogolubov, J. Phys. (Moscow) **11**, 23 (1947).
- [22] N.N. Bogolubov, Moscow Univ. Phys. Bull. 7, 43 (1947).
- [23] N.N. Bogolubov, Lectures on Quantum Statistics (Gordon and Breach, New York, 1967), Vol. 1.
- [24] N.N. Bogolubov, Lectures on Quantum Statistics (Gordon and Breach, New York, 1970), Vol. 2.
- [25] P.C. Hohenberg and P.C. Martin, Ann. Phys. **34**, 291 (1965).
- [26] V.I. Yukalov, Phys. Rev. E **72**, 066119 (2005).
- [27] V.I. Yukalov, Phys. Rep. **208**, 395 (1991).
- [28] V.I. Yukalov, Statistical Green's Functions (Queen's University, Kingston, 1998).
- [29] H. Kleinert, Path Integrals (World Scientific, Singapore, 2004).
- [30] V.I. Yukalov, Laser Phys. **16**, 511 (2006).
- [31] N.N. Bogolubov and N.N. Bogolubov Jr., Introduction to Quantum Statistical Mechanics (Gordon and Breach, Lausanne, 1994).
- [32] S.T. Beliaev, J. Exp. Theor. Phys. 7, 289 (1958).
- [33] N.M. Hugenholtz and D. Pines, Phys. Rev. **116**, 489 (1959).
- [34] P.O. Fedichev and G.V. Shlyapnikov, Phys. Rev. A 58, 3146 (1998).
- [35] G. Baym and G. Grinstein, Phys. Rev. D 15, 2897 (1977).
- [36] V.N. Popov, J. Exp. Theor. Phys. **20**, 1185 (1965).
- [37] V.N. Popov, Functional Integrals in Quantum Field Theory and Statistical Physics (Reidel, Dordrecht, 1983).
- [38] V.N. Popov, Functional Integrals and Collective Modes (Cambridge University, New York, 1987).
- [39] V.I. Yukalov and E.P. Yukalova, Laser Phys. Lett. 2, 506 (2005).
- [40] C. Weiss and T. Jinasundra, Phys. Rev. A **72**, 053626 (2005).