Magnetic Field Dependence of T_{AB} in Super uid ³H e

GA.Baramidze and GA.Kharadze

Andronikashvili Institute of Physics, Tamarashvili str. 6, 0177 Tbilisi, Georgia (e-mail: gogi@iphac.ge)

April 15, 2024

A bstract

The magnetic eld dependence of the A ! B transition temperature T_{AB} in the super uid ³H e is reconsidered in order to take into account the linear-in-eld contribution beyond the approximation used in Ref.[7]. In the high eld region, where the quadratic-in-eld contribution prevails, the well known answer is restored. On the other hand, it is pointed out that the Ferm i liquid e ects shift the observability of the linear-in-eld region to rather low magnetic elds.

Soon after the discovery of the super uidity of liquid ³H e it was established that near critical temperature T_c (P) the phase diagram of super uid state experiences a profound m odi cation under the action of even sm all external magnetic elds [1, 2, 3]. This modi cation shows up in elimination of the direct norm alto B phase transition over entire phase diagram in favour of an anisotropic A phase.

The A ! B transition temperature $T_{A\,B}$ (H) < $T_{\rm c}$ can be established by equating the free energies F_A (H) and F_B (H). Near the zero- eld transition temperature T_c (P) the free energy can be found by m inim izing the G inzburg-Landau functional

$$F_{s} = 3 < > + F_{s}^{(4)}(~);$$
 (1)

where the fourth order contribution in the order-parameter $\widetilde{\ }(\,\hat{k})$ reads as

$$F_{s}^{(4)} = 9f_{1} < > < > + _{2} < j^{2}j^{2} > ^{2} + _{3} < > < >$$

$$(2)$$

$$+ _{4} < > < > + _{5} < > < > g:$$

In Eqs. (1) and (2) the spin-vector ~ is de ned according to the general expression for the order-parameter A_i of the spin-triplet p-wave Cooper condensate: $(\hat{k}) = A_i \hat{k}_i$, and the angular brackets < > stand for an average over the momentum direction \hat{k} on the Fermi surface. In presence of a magnetic eld H the tensor coe cient of the second order term in Eq.(1) reads as

=

$$+ ig_1$$
" H + g_2 H H ; (3)

where

$$= \frac{1}{3} N_{\rm F} \ln \frac{T}{T_{\rm c}} ; \qquad (4a)$$

$$g_1 H = \frac{1}{3} N_F h;$$
 (4b)

$$g_2 H^2 = \frac{1}{3} N_F h^2;$$
 (4c)

with $h = h H = (2k_B T_c) = H = H_o$. The dimensionless coe cients and could be considered as the phenom enological quantities although their values can be estimated according to the microscopic calculations. In the weak-coupling approximation [4]

$$_{NC} = \frac{N_{F}^{0}}{N_{F}} k_{B} T_{C} \ln \frac{2_{E} !_{C}}{T_{C}}$$
(5)

where N_F^0 stands for the derivative of the quasiparticle DOS with respect to the energy at the Ferm i level. A detailed calculations which take into account the linear-in-eld corrections to the Ferm i liquid parameters are performed in Ref. [5].

As to the -coe cient, it stems from the free energy part

$$F_{\rm H}^{(2)} = \frac{1}{2} {}_{\rm s} {\rm H}^{2}; {}_{\rm s} = {}_{\rm s} {}_{\rm N};$$
 (6)

where $_{\rm s}$ ($_{\rm N}$) stands for the m agnetic susceptibility of the super uid (norm al) phase. For the B phase near T_c, $F_{\rm H}^{(2)} = g_2 {\rm H}^2 {}^2$ where $g_2 {\rm H}^2$ is given according to Eq. (4c) with

$$= \frac{7 (3)}{4^{2}} \frac{1}{(1 + F_{0}^{a})^{2}}$$
(7)

Here the Ferm i liquid parameter $F_o^a = -3=4$ and is weakly pressure dependent.

As it is well known, the linear-in-eld term (4b) is the origin of a tiny splitting of the A phase (due to a sm all asymmetry of the density of quasiparticle states at the Ferm i level). The quadratic-in-eld contribution (4c) gives rise to the magnetic eld suppression of " $_{\#}$ component of the energy gap of the B phase.

Based on the argument that the term (4b) is rather small, in the majority of considerations of $T_{AB} = T_{AB}$ (H) this term is usually discarded, and in a standard way one starts from the following expressions for the equilibrium free energies of the A and B phases:

$$F_{A} = \frac{1}{4_{245}}^{2};$$
 (8a)

$$F_{\rm B} = \frac{1}{2(3_{12} + 3_{45})} \frac{3}{2}^2 + g_2 H^2 + \frac{2_{12} + 3_{45}}{2_{345}} (g_2 H^2)^2 :$$
(8b)

Since here the linear-in-eld contribution (proportional to $g_1)$ is dropped, the action of the magnetic $\,$ eld appears only in F $_B$.

Equating (8a) and (8b) it is found that $T_{A\,B}\ (\!H\!$) is to be obtained from the equation

1

$$= {}^{2} + a + b = 0$$
 (9)

with

$$a = 2P_1 g_2 H^2;$$

$$b = P_1^2 (1 - q_1) (g_2 H^2)^2$$
(10)

where the coe cients P₁ and q_1 are dened in the Appendix. From Eq.(9) it follows the answer for $_{AB} = 1$ $T_{AB} = T_c$:

$$AB = P_1 \left(1 + \frac{p_{\overline{q_1}}}{q_1}\right) h^2$$
(11)

which reproduces a well known result obtained in Ref.[6].

Now we turn to the role of the linear-in-eld contribution to T_{AB} (H). This question was rst posed in Ref.[7]. The starting point is the construction of the expressions for F_{A2} (H) and F_B (H), the contribution (4b) being taken into account. In a standard way it is established that

$$F_{A2} = \frac{1}{4_{245}} \frac{2}{5} \frac{-\frac{245}{5}}{5} g_1^2 H^2 ; \qquad (12)$$

$$F_{B} = \frac{1}{2(3_{12} + 345)} \left(\frac{3}{2}^{2} + g_{2}H^{2} + \frac{3_{12} + 345}{4(3_{1} + 35)}g_{1}^{2}H^{2} + \frac{2_{12} + 345}{2(3_{1} + 35)}g_{1}^{2}H^{2} + \frac{2_{12} + 345}{2(3_{1} + 35)}(g_{2}H^{2})^{2} + \frac{(3_{12} + 345)^{2}}{(3_{1} + 35)} \left(\frac{1}{345}g_{1}^{2}g_{2}H^{4} + \frac{1(3_{12} + 345)}{4(3_{1} + 35)}g_{1}^{2}H^{4}\right)^{2} :$$

$$(13)$$

 $C\,\text{om}\,parison\,\,of\,F_{A\,2}$ and $F_B\,$ gives an equation for $T_{A\,B}$ (H):

'
$$(T_{AB} = T_c) = {}^4 + a {}^3 + b {}^2 + c + d = 0$$
 (14)

where now

$$a = 2P_{1}g_{2}H^{2};$$

$$b = P_{1}P_{2}P_{3}g_{1}^{2}H^{2} + P_{1}^{2}(1 - q_{1})(g_{2}H^{2})^{2};$$

$$c = P_{1}P_{2}^{2}P_{4}g_{1}^{2}g_{2}H^{4};$$

$$d = P_{1}P_{2}^{3}P_{5}(g_{1}H^{2})^{4};$$

(15)

The de nition of all the coe cients P $_{\rm a}$ is given in the Appendix.

In Ref.[7] the second and third lines in Eq.(13), containing terms on the order H ⁴ were neglected. This has the following in uence on the answer for T_{AB} (H): i) neglection of the contribution proportional to $(g_2 H^2)^2$ makes it in possible one to reproduce the correct answer given by Eq.(11) (the term

 $p = \overline{q_{1}}$ will be lost), ii) neglection of the contribution collected in the square brackets of the third line of Eq.(13) changes the answer for the linear-in-eld contribution to T_{AB} (H). To avoid this drawbacks we address Eq.(14) and, as a rst step, perform the variable transformation $! x = \frac{1}{4}a$, after which an equation for x ($T_{AB} = T_{c}$) is obtained:

'
$$(T_{AB} = T_c) = x^4 + Ax^2 + Bx + C = 0$$
 (16)

with the coe cients

$$A = P_{1}P_{2}P_{3} (g_{1}H)^{2} + \frac{1}{2}P_{1}^{2} (1 + 2q_{1}) (g_{2}H^{2})^{2} ;$$

$$B = q_{1}P_{1}^{3} (g_{2}H^{2})^{3} + q_{2}P_{1}P_{2}g_{1}^{2}g_{2}H^{4};$$

$$C = P_{1}P_{2}^{3}P_{5} (g_{1}H)^{4} + \frac{1}{16}P_{1}^{4} (1 - 4q_{1}) (g_{2}H^{2})^{4} + \frac{1}{4}P_{1}^{2}P_{2} (P_{1}P_{3} - 2q_{2}) (g_{1}g_{2}H^{3})^{2}:$$

$$(17)$$

It is to be noticed, that as a result of the variable transform ation used, in Eq. (16) the cubic term is absent. At the same time the coe cient of the linear term B is zero in the weak-coupling approximation.

In order to solve Eq. (16) we use the decomposition $x = x_0^{p} \overline{q_1} P_1 q_2 H^2$ which generates the decomposition $' = '_0 + '$ with $'_0$ being the solution of the equation

$$'_{o} = x_{o}^{4} + A_{o}x_{o}^{2} + C_{o} = 0;$$
 (18)

where A_{\circ} and C_{\circ} are the coe cients A and C taken at $q_1 = q_2 = 0$. From Eq. (18) it is found that

$$x_{o}(T_{AB} = T_{c}) = P_{o}^{2}(g_{1}H)^{2} + \frac{1}{4}P_{1}^{2}(g_{2}H^{2})^{2}$$
(19)

where

$$P_{o}^{2} = \frac{1}{2} P_{1} P_{2} P_{3} [1 + \frac{q}{1 + 4P_{2}P_{5} = P_{1} P_{3}^{2}}]:$$
(20)

The direct inspection shows that ' is the sum of terms proportional to the powers of q and the minim alpower in H contained in ' is H^5 . For this

reason we can use an approximation with ' disregarded, and as a result it is found that

$$(T_{AB} = T_{c}) = x \quad \frac{1}{4}a = x_{o} \quad P = \frac{1}{q_{1}}P_{1}g_{2}H^{2} \quad \frac{1}{2}P_{1}g_{2}H^{2} =$$

$$= P_{1} \quad \frac{1}{2} + P = \frac{1}{q_{1}}g_{2}H^{2} \quad S = \frac{1}{P_{o}^{2}}(g_{1}H^{2})^{2} + \frac{1}{4}P_{1}^{2}(g_{2}H^{2})^{2}; \qquad (21)$$

U sing this result we nally have the following simple answer for $_{AB}$:

c

$${}_{AB} = P_1 \frac{1}{2} + \frac{p_{\overline{q_1}}}{p_1} h^2 + \frac{p_{\overline{q_2}}}{p_{\overline{q_1}}} (h^2)^2 + \frac{1}{4} P_1^2 (h^2)^2$$
(22)

By introducing a characteristic magnetic eld H $=2\,(P_{\circ}{=}P_{1})\,($ $=)\,H_{\circ}$ the asymptotic regions where the quadratic-in-eld (H >> H) and the linear-in-eld (H << H) contributions to $T_{A\,B}$ prevail are found:

$${}_{AB} = \begin{pmatrix} P_{1} (1 + {}^{p} \overline{q_{1}}) & h^{2}; & H >> H \\ P_{0} & h; & H << H : \end{pmatrix}$$
(23)

For H >> H the well known result [4,8] is reproduced.

In order to isolate the role of the linear-in-eld contribution to $T_{A\,B}\,$ it is convenient to consider

$$_{AB}(h) = _{AB}(h) (1 + {}^{P}\overline{q_{I}})P_{1} h^{2};$$
 (24)

and construct graphically

$$f(h) = {}_{AB}(h) = h = P_{o} (1 + (h=h)^{2} h=h);$$
 (25)

where the scaling value $h = 2(P_0=P_1)(=)$.

Below f (h) is plotted for the pressure P = 10 bar (P_o ' 5;4 1 $\hat{\sigma}$;h ' 1;6 1 $\hat{\sigma}$).

A cknow ledgem ents

The authors highly appreciate the valuable discussions with late Dr. Sh N ikolaishvili.



Figure 1: The bold curve corresponds to the Landau parameter $F_o^a = 3=4$ and the thin curve is constructed at $F_o^a = 0$. It is obvious that the Ferm i liquid e ect shifts the linear-in-eld contribution to rather small values of $H << H = h H_o'$ 420 G.

Appendix

The coe cients P $_a$ (a = 1;2;3;4;5) and q_a (a = 1;2) introduced in the main text are de ned as follows:

$$P_{1} = \frac{245}{2_{345}};$$

$$P_{2} = \frac{3_{12} + 345}{4_{31}};$$

$$P_{3} = \frac{45_{31}}{5};$$

$$P_{4} = \frac{2_{1}}{345};$$

$$P_{5} = \frac{1}{2(4 \ 3_{1} \ 3_{5})};$$

$$q_{1} = \frac{(3_{12} + 3_{45})(2_{13} \ 3_{45})}{245 \ 345};$$

$$q_{2} = P_{1}P_{3} \ P_{2}P_{4};$$

where $_{ij} = _i + _j +$

The coe cients q_1 and q_2 contain only the strong-coupling corrections $_{\rm i}$ de ned according to the decom position

$$1 = 0 + 1;$$

$$2 = 2 + 2;$$

$$3 = 2 + 3;$$

$$4 = 2 + 4;$$

$$5 = 2 + 5;$$

$$0 = \frac{7 + 3}{240} + 5;$$

References

- [1] W J.Gully et al, Phys. Rev. A 8, 1633 (1973).
- [2] D N. Paulson et al, Phys. Rev. Lett. 32, 1098 (1974).
- [3] J.D. Feder et al, Phys. Rev. Lett. 47, 428 (1981).
- [4] V.Ambegaonar and D.Mermin, Phys. Rev. Lett. 30, 81 (1973).
- [5] K.Bedell and K.Quadar, Phys. Rev. B 30, 2894 (1984).

- [6] A L. Fetter, in "Quantum Statistics and the Many-Body Problem", Plenum, NY, 1975; J.Low Temp. Phys. 23, 245 (1976).
- [7] K.Levin and O.W alls, Phys. Rev. B 15, 4256 (1977).
- [8] Y H. Tang et al, Phys. Rev. Lett. 67, 1775 (1991).