

High-temperature series expansions for the q -state Potts model on a hypercubic lattice and critical properties of percolation

Meik Hellmund^{1,2,*} and Wolfhard Janke^{3,2,†}

¹*Mathematisches Institut, Universität Leipzig, Augustusplatz 10/11, D-04109 Leipzig, Germany*

²*Centre for Theoretical Sciences (NTZ) of the Centre for Advanced Study (ZHS), Universität Leipzig, Emil-Fuchs-Str. 1, D-04105 Leipzig, Germany*

³*Institut für Theoretische Physik, Universität Leipzig, Augustusplatz 10/11, D-04109 Leipzig, Germany*

(Dated: March 23, 2022)

We present results for the high-temperature series expansions of the susceptibility and free energy of the q -state Potts model on a D -dimensional hypercubic lattice \mathbb{Z}^D for arbitrary values of q . The series are up to order 20 for dimension $D \leq 3$, order 19 for $D \leq 5$ and up to order 17 for arbitrary D . Using the $q \rightarrow 1$ limit of these series, we estimate the percolation threshold p_c and critical exponent γ for bond percolation in different dimensions. We also extend the $1/D$ expansion of the critical coupling for arbitrary values of q up to order D^{-9} .

PACS numbers:

05.50.+q Lattice theory and statistics (Ising, Potts, etc.)

05.70.Jk Critical point phenomena

64.60.Fr Equilibrium properties near critical points, critical exponents

64.60.Ak Renormalization-group, fractal, and percolation studies of phase transitions

I. INTRODUCTION

Since many years the q -state Potts model (Wu, 1982) has served as paradigmatic system for the study of phase transitions and critical phenomena. Fortuin and Kasteleyn reformulated it as a random cluster model in which q appears as a general parameter no longer restricted to $q = 2, 3, \dots$ and where bond percolation appears as the special case $q = 1$.

Systematic series expansions have a long history (Domb and Green, 1974) as a method to analyse lattice models of phase transitions. In the case of Potts models, high-temperature series for the susceptibility and free energy with q as a free parameter have been obtained for D -dimensional hypercubic lattices in (Moraal, 1993) (free energy only) and (Schreider and Reger, 1994). We extend these series considerably and use them for a scan of the phase diagram in (q, D) -space. A detailed analysis is presented for the case of bond percolation ($q = 1$) on higher-dimensional hypercubic lattices, extending the analysis of 15th order percolation series in arbitrary dimensions of (Adler *et al.*, 1990).

We shortly review the model and its cluster formulation in Sect. II and the star-graph expansion technique in Sect. III. In the following sections we present results for general parameter q in two (Sect. IV) and higher (Sect. V) dimensions. Section VI treats the large-dimensionality expansion of the critical coupling, and in Sect. VII the case of bond percolation, i.e., the limit $q = 1$ is analyzed. We end with a short summary in Sect. VIII.

II. THE q -STATE POTTS MODEL

Let $G = (V, E)$ be an undirected graph, that is a set V of vertices together with a set E of edges defining which pairs of vertices are considered “nearest neighbors”. Put a spin s_i on each vertex $i \in V$ which can take q different values. The q -state Potts model (see, e.g., (Wu, 1982) for a review) on this graph is then defined by the Hamiltonian

$$H = -J \sum_{(ij) \in E} \delta(s_i, s_j), \quad (1)$$

*Electronic address: Meik.Hellmund@math.uni-leipzig.de

†Electronic address: Wolfhard.Janke@itp.uni-leipzig.de

where $\delta(.,.)$ is the Kronecker delta symbol. The Fortuin-Kasteleyn representation shows that the partition function $Z_G = \sum_{\{s_i\}} e^{-\beta H}$ for a finite graph G is actually a polynomial in q and $w = e^{\beta J} - 1$:

$$Z_G(q, w) = \sum_{C \subseteq E} q^{N_c(C)} w^{|C|}. \quad (2)$$

Here the sum is over all $2^{|E|}$ clusters, i.e. subsets C of the set E of edges; $N_c(C)$ is the number of connected components where each isolated vertex is counted as one component and $|C|$ is the number of edges of C . This allows one to study the model for arbitrary values of $q \notin \{2, 3, \dots\}$ which can no longer be represented by a local interaction of spin degrees of freedom of the original Hamiltonian (1). Of special interest in statistical physics are the limits $q \rightarrow 1$ and $q \rightarrow 0$ which can be shown to be related to percolation and tree percolation on the graph G , respectively.

A two-point correlation function can be defined in the cluster representation simply by

$$D(i, j) = \frac{1}{Z_G} \sum_{C_{ij} \subseteq E} q^{N_c(C)} w^{|C|}, \quad (3)$$

where the cluster sum is restricted to those clusters C_{ij} in which the vertices i and j belong to the same connected component. The susceptibility related to this correlator,

$$\chi_G(q, w) = \frac{1}{|V|} \sum_{i, j \in V} D(i, j), \quad (4)$$

equals for $q = 2, 3, \dots$ the usual magnetic susceptibility of the q -state Potts model in the high-temperature phase (i.e., as long as $\langle s_i \rangle = 0$).

In the case $q = 1$ the weight of a cluster in the partition sum depends only on its number of edges. This case therefore describes bond percolation where the local bond probability is given by $p = w/(w + 1)$. It easily follows that here the susceptibility $\chi(q = 1, w)$ measures the average size of the percolation cluster.

The Potts partition function $Z_G(q, w)$ is essentially equivalent to the Tutte or dichromatic polynomial of the graph G , see, e.g., (Sokal, 2005) and references therein. It encodes a lot of combinatorial properties of the graph such as the number of spanning trees, the number of spanning forests, the number of possible acyclic orientations, or the number of vertex colourings with q different colours using different colours on neighbouring vertices. The last mentioned number, for example, equals $\lim_{\beta J \rightarrow -\infty} Z_G(q, e^{\beta J} - 1) = Z_G(q, -1)$, which is the number of ground states of the antiferromagnetic q -state Potts model on the graph G .

In statistical physics, on the other hand, the interest in these models is related to the existence of a phase transition between a disordered high-temperature (low-density in case of percolation clusters) and an ordered low-temperature (large w) phase if we go to infinite graphs like the D -dimensional hypercubic lattice $G = \mathbb{Z}^D$. Depending on q and D , these transitions can be first or second order. Whereas in $D = 2$ dimensions many exact results are available showing that the second-order nature changes to first-order for $q > 4$, in $D = 3$ and higher dimensions there is strong numerical evidence that the transition is of first order for all integer values $q > 2$. In the case of a second-order phase transition at β_c , the behavior of the susceptibility (4) is characterized by the critical exponent γ : $\chi_G \sim |\beta - \beta_c|^{-\gamma}$.

III. HIGH-TEMPERATURE SERIES EXPANSIONS

Series expansions are an important tool for the extraction of information on statistical systems which is exact (up to the order of the expansion) in the thermodynamic limit. There exist different well-established methods (Domb and Green, 1974) for the systematic generation of high-temperature series expansions, as for example the linked cluster and the star-graph method. The latter one got its name from the fact that only 1-vertex-irreducible graphs contribute to the series. In an attempt to study systems with quenched disorder via series expansions (Hellmund and Janke, 2002a,b, 2003, 2005; Janke *et al.*, 2005) we used this method and developed a comprehensive software toolbox for generating and enumerating star graphs which is re-used in this work.

The basic idea of the star-graph method is to assemble the value of some extensive thermodynamic quantity F on a large or even infinite graph from its values on subgraphs: Graphs constitute a partially ordered set under the “subgraph” relation. Therefore, for every function $F(G)$ defined on the set of graphs exists another function $W_F(G)$ such that for all graphs G

$$F(G) = \sum_{g \subseteq G} W_F(g), \quad (5)$$




FIG. 1 Two star graphs of order 19 (left) and 17 (right) and their weak embedding numbers. Due to its topology, the result is complete (valid for arbitrary dimension D) for the graph of order 17. For the graph of order 19 it is valid only up to $D = 6$.

and this function can be calculated recursively via

$$W_F(G) = F(G) - \sum_{g \subset G} W_F(g). \quad (6)$$

This gives for an infinite (e.g. hypercubic) lattice

$$F(\mathbb{Z}^D) = \sum_G (G : \mathbb{Z}^D) W_F(G), \quad (7)$$

where $(G : \mathbb{Z}^D)$ denotes the weak embedding number of the graph G in the given lattice structure (Martin, 1974).

The following observation makes this a useful method: Let G be a graph with an articulation vertex where two star subgraphs $G_{1,2}$ are glued together. Then $W_F(G)$ vanishes if

$$F(G) = F(G_1) + F(G_2). \quad (8)$$

An observable F for which eq. (8) is true on arbitrary graphs with articulation points allows a star-graph expansion. Then all non-star graphs have zero weight W_F in the sum eq. (7). It is easy to see that the (properly normalized) free energy $\ln Z$ has this property and it can be shown that the inverse susceptibility $1/\chi$ has it, too.

The weak embedding numbers $(G : \mathbb{Z}^D)$ of a graph G into the hypercubic lattice are counted using a refined version of the backtracing algorithm by Martin (Martin, 1974), making use of a couple of simplifications for bipartite hypercubic lattices \mathbb{Z}^D . Figure 1 shows two typical results. A star graph of order n can use at most $\lfloor \frac{n}{2} \rfloor$ dimensions of the lattice and every embedding using m dimensions appears in \mathbb{Z}^D with multiplicity $\binom{D}{m}$.

For the symbolic calculations of the partition sum and susceptibility on every star graph up to a given order (ca. 80 000 up to order 20) as polynomials in q and w , we developed a C++ template library using an expanded degree-sparse representation of polynomials and series in many variables. The open source library GMP is used for the arbitrary-precision arithmetics. Our series for the free energy $\ln Z$ and the susceptibility χ with q as a free parameter are up to order 20 for dimension $D \leq 3$, order 19 for $D \leq 5$ and up to order 17 for arbitrary D .

In order to check our algorithms we did a careful comparison with older published series data. Our series agree with the 10th order susceptibility and free energy series for arbitrary q and D of (Schreider and Reger, 1994) (modulo some misprints in their expression for $\ln Z$) as well as with the 16th order $D = 2$ free energy series of (Moraal, 1993) but disagree in several places with the 11th order, arbitrary dimension free energy series in the last mentioned paper. For the special case of bond percolation $q = 1$ they agree with the 15th order arbitrary-dimension series of (Adler *et al.*, 1990). In the special case of the three-dimensional Ising model ($q = 2$, $D = 3$) our 20th order series also conform with the 25th order series of (Butera and Comi, 2002) generated by linked-cluster expansion. They also agree with the 21th order $q = 3$, $D = 3$ partition function series of (Guttmann and Enting, 1994) obtained by finite lattice methods. In two dimensions, much longer series exist for special models. Our 20th order susceptibility series agree with the $q = 3$ and 4, $D = 2$ series of (Enting and Guttmann, 2003).

We close this section with some remarks about the analysis of the thus generated expansions. The estimate of critical parameters from a high-temperature series involves extrapolation from a finite number of exactly known coefficients to the asymptotic form of the function. Many such extrapolation techniques have been developed and tested for different series and are comprehensively reviewed in (Guttmann, 1989). These extrapolation techniques are not rigorous. They make some assumptions about the expected form of the singularity at the critical temperature. Usually, error estimates rely on the scatter of the results of extrapolations with different parameters (like $[N/M]$ Padé approximants for different values of N and M). This may seriously underestimate systematic errors coming from wrong assumptions about the structure of the singularity.

In order to get a reliable picture, we will take into account several criteria, such as




FIG. 2 Critical coupling w_c as function of q in 2 dimensions.

- convergence of the analysis,
- scatter of different approximants,
- number of defective approximants,
- agreement between different extrapolation methods.

The basic methods we use are DLog-Padé approximation and inhomogeneous differential approximants (IDA) (Guttman, 1989). In order to analyze confluent nonanalytic and logarithmic corrections, these methods are applied to suitably transformed forms of the series. The parameters of these transformations are fine-tuned according to the criteria listed above, a technique pioneered in (Adler *et al.*, 1990).

IV. TEST: TWO DIMENSIONS

The Potts model in two dimensions is exhaustively reviewed in (Jacobsen *et al.*, 2005). We repeat here the principal facts: Baxter has determined the exact free energy on \mathbb{Z}^2 along the two curves

$$w_F = \sqrt{q} \quad (9)$$

$$w_A = -2 + \sqrt{4 - q} \quad (10)$$

in (q, w) parameter space. The curve (9) gives the locus of the ferromagnetic phase transition and is perfectly understood in terms of conformal field theory. Critical exponents are exactly known, e.g.

$$\gamma = \frac{4 + 3k^2}{12k - 6k^2} \quad \text{where} \quad \sqrt{q} = -2 \cos(\pi/k). \quad (11)$$

The curve (10) is conjectured to be the locus of the antiferromagnetic transition which agrees with the known facts for $q = 2, 3, 4$.

An analysis of the susceptibility series of order 20 using Padé approximants to the logarithmic derivative of the susceptibility (DLog-Padé approximants) is shown for the ferromagnetic case in Figs. 2 and 3 where around 15–25 different $[N, M]$ approximants are plotted for each value of q . The continuous lines present the exact results (9) and (11). We find a quite satisfactory agreement also for noninteger values of q , however, the scatter of different approximants and the number of defective approximants increases for smaller values of q . This may be related to the fact that the $q \rightarrow 0$ limit is a quite intricate multicritical point (Jacobsen *et al.*, 2005).




FIG. 3 Critical exponent γ as function of q in 2 dimensions.




FIG. 4 Critical coupling v_c as function of q in different dimensions.

V. HIGHER DIMENSIONS: OVERVIEW

The explicit q - and D -dependence of our series allows us to get a fast overview over the phase structure in a large parameter range. Figures 4 and 5 show the results of a DLog-Padé analysis of the susceptibility series for the critical coupling $v = (e^{\beta J} - 1)/(e^{\beta J} - 1 + q) = w/(w + q)$ and exponent γ .

Several remarks are in order:

- At the upper critical dimension ($D = 4$ for $q = 2$, $D = 6$ for $q = 1$) the DLog-Padé analysis finds a value $\gamma \approx 1.08$ slightly larger than 1. This effect is well-known and due to the existence of logarithmic corrections.
- For $D \geq 3, q \geq 3$ the phase transition is of first order and the correlation length remains finite. In analysing high-temperature series by Padé approximants or similar techniques, the approximant will provide an analytic continuation beyond the first-order transition temperature T_0 into a metastable region on a pseudo-spinodal line with a singularity $T_c^* < T_0$ and effective exponents at T_c^* . Our v_c in this region is therefore an upper bound of the real transition point.




FIG. 5 Critical exponent γ as function of q in different dimensions.

- For $D = 3$, the curve $\gamma(q)$ passes $\gamma = 1$ at $q \approx 2.7$, marking the onset of a first-order transition. This is consistent with slightly smaller estimates in the literature coming from Monte Carlo computer simulations and field theoretic calculations (see (Grollau *et al.*, 2001) and references therein). It would be interesting to see if the crossover point can be understood as a kind of tricritical point in the (q, v) parameter space.

Since high-temperature series alone are not sufficient to analyse first-order transitions (and much longer series exist for the special case of the three-dimensional Ising model $q = 2, D = 3$), we concentrate our analysis in the second part of the paper on the $q = 1$ case of bond percolation.

VI. LARGE-DIMENSIONALITY EXPANSION

The dimension of the lattice enters the star-graph series expansion only through the embedding numbers (eq. (7)) and we keep it as a free parameter in our series (see Fig. 1). This allows one (Fisher and Singh, 1990) to extract a large- D expansion for the critical coupling from the susceptibility series by iteratively solving the critical point equation $1/\chi(D, v_c) = 0$ in terms of v_c as series in $1/D$. This gives the large- D expansion of v_c up to order D^{-n} if χ is known as series in v up to order $2n$. For the $O(N)$ model the first terms of the $1/D$ expansion of the critical coupling for arbitrary N were obtained by Gerber and Fisher (Gerber and Fisher, 1974), who also demonstrated by solving the large N limit that this expansion is only asymptotic.

As far as we know, our analogous series for the q -state Potts model with arbitrary q is new. For the percolation ($q = 1$) case the first 5 orders were calculated in (Gaunt and Ruskin, 1978). We give the series for v_c in terms of the branching number σ which is one less than the coordination number of the lattice and therefore for the hypercubic lattice $\sigma = 2D - 1$,

$$v_c(q, \sigma) = \frac{1}{\sigma} \left[1 + \frac{8-3q}{2\sigma^2} + \frac{3(8-3q)}{2\sigma^3} + \frac{3(68-31q+q^2)}{2\sigma^4} + \frac{8664-3798q-11q^2}{12\sigma^5} \right. \\ \left. + \frac{78768-36714q+405q^2-50q^3}{12\sigma^6} + \frac{1476192-685680q-2760q^2-551q^3}{24\sigma^7} \right. \\ \left. + \frac{7446864-3524352q-11204q^2-6588q^3-9q^4}{12\sigma^8} + \dots \right]. \quad (12)$$

By using σ as expansion variable we see that the first correction to the Bethe lattice result $v_c = 1/\sigma$ is absent for all values of q . Figure 6 shows plots of the critical coupling as obtained from eq. (12) by simply summing up all terms. Since these expansions are asymptotic series, for relatively small D a more careful treatment with appropriately adapted maximal order would actually be necessary, as discussed below for the percolation case $q = 1$. Of course,




FIG. 6 Plots of eq. (12) (using all available terms, cf. text) for different values of q .

in the case of first-order phase transitions, the expansion (12) yields the pseudo-spinodal transition point in the meta-stable region.

VII. $q \rightarrow 1$: PERCOLATION

The $q \rightarrow 1$ limit of the Potts model describes bond percolation with local bond probability $p = w/(w+1) = v$. Standard scaling theory (Stauffer and Aharony, 1994) describes the number of clusters of size s for large s near the percolation threshold p_c by

$$n(s, p) \sim s^{-\tau} f((p_c - p)s^\sigma), \quad (13)$$

introducing the critical exponents τ and σ . The $q \rightarrow 1$ limit of the Potts model susceptibility describes the average cluster size and $\frac{\partial \ln Z}{\partial q}$ the number of clusters. The respective critical exponents are related to the scaling law eq. (13) by

$$\gamma = \frac{3 - \tau}{\sigma}, \quad (14)$$

$$2 - \alpha = \frac{\tau - 1}{\sigma}. \quad (15)$$

A throughout analysis of percolation series in higher dimensions was done some years ago by (Adler *et al.*, 1990). Since we extended the series from 15 to 17 (for $D \geq 6$), 19 (for $D = 4, 5$) and 20 (in $D = 3$) terms, a careful re-analysis seems to be in place (for extensive percolation series expansions in two dimensions, see (Conway and Guttmann, 1995) and, recently, (Jensen and Ziff, 2006)).

A. Up to the upper critical dimension

The divergence of the susceptibility (mean finite cluster size) is expected to be of the form

$$\chi(p) = A(p_c - p)^{-\gamma} [1 + a_1(p_c - p)^{\Delta_1} + \dots] \quad (16)$$

for $D \neq 6$ and

$$\chi(p) = A(p_c - p)^{-\gamma} |\ln(p_c - p)|^\delta + \dots \quad (17)$$

at the upper critical dimension $D = 6$ with $\gamma = 1$ and $\delta = 2/7$. For $D > 6$, one expects a Gaussian fixed point with $\gamma = 1$.

TABLE I Percolation thresholds (bond percolation on \mathbb{Z}^D) and critical exponents.

D	Methods	p_c	σ	τ	$\gamma = \frac{3-\tau}{\sigma}$
2	exact, CFT	$\frac{1}{2}$	$\frac{36}{91}$	$\frac{187}{91}$	$\frac{43}{18}$
3	MC (Ziff and Stell, 1988)	0.248 812(2)	0.453(1)	2.189(1)	1.795(5)
	HTS (Adler <i>et al.</i> , 1990)	0.248 8(2)			1.805(20)
	MC (Lorenz and Ziff, 1998a)	0.248 812 6(5)	0.4522(8) 0.4539(3)	2.18906(6) 2.18925(5)	1.7933(85)
	MCS (Ballesteros <i>et al.</i> , 1999)				1.7862(30)
	MCS (Deng and Blöte, 2005)				1.804(5)
	present work	0.248 91(10)			
4	HTS (Adler <i>et al.</i> , 1990)	0.160 05(15)	2.313(3)		1.435(15)
	MCS (Ballesteros <i>et al.</i> , 1997)				1.44(2)
	MC (Paul <i>et al.</i> , 2001)	0.160 130(3)			
	MC (Grassberger, 2003)	0.160 131 4(13)			
	present work	0.160 08(10)			1.435(5)
5	HTS (Adler <i>et al.</i> , 1990)	0.118 19(4)	2.412(4)		1.185(5)
	MC (Paul <i>et al.</i> , 2001)	0.118 174(4)			
	MC (Grassberger, 2003)	0.118 172(1)			
	present work	0.118 170(5)			1.178(2)
6	RG (Essam <i>et al.</i> , 1978)		$\frac{1}{2}$	$\frac{5}{2}$	$\chi \sim t^{-1} \ln t ^\delta$, $\delta = \frac{2}{7}$
	HTS (Adler <i>et al.</i> , 1990)	0.094 20(10)			
	MC (Grassberger, 2003)	0.094 201 9(6)			
	present work	0.094 202 0(10)			$\delta = 0.40(2)$
> 6	RG		$\frac{1}{2}$	$\frac{5}{2}$	1

MC = Monte Carlo, MCS = Monte Carlo, **site** percolation, RG = Renormalization group, HTS = High-temperature series

We analyse the series using DLog-Padé approximants, inhomogeneous differential approximants and the methods termed M1 and M2 from (Adler *et al.*, 1991, 1990) especially tailored to take confluent nonanalytic correction terms into account. The method M1 uses DLog-Padé approximants to

$$F(p) = (p_c - p) \frac{d\chi}{dp} - \gamma \chi(p), \quad (18)$$

which has a pole at p_c with residue $\Delta_1 - \gamma$. For a given trial value of p_c the graphs of Δ_1 versus input γ are plotted for different Padé approximants and by adjusting p_c a point of optimal convergence is searched.

The M2 method starts with a transformation of the series in p into a series in $y = 1 - (1 - p/p_c)^{\Delta_1}$, and then Padé approximants to

$$G(y) = \Delta_1(1 - y) \frac{d}{dy} \ln F(y) \quad (19)$$

are calculated which should converge to γ as $y \rightarrow 1$, i.e. $p \rightarrow p_c$. These methods are especially useful when taken as biased approximants with a given value of γ or Δ_1 as input. Our results are compiled in Table I, where we quote for comparison also previous estimates based on Monte Carlo simulations and analyses of shorter series expansions.

Generally the series analysis appears to be more difficult for lower dimensions. The p_c we quote for $D = 3$ is an average over a large number (ca. 200) of Padé and IDA approximants. The general pattern in the scatter of data is that larger values of p_c corresponds with larger values of the critical exponent γ . The value for γ we quote as our result in three dimensions is obtained by biasing the M1 and M2 analyses with the more precise p_c value from Monte Carlo simulations. Moving on to $D = 4$ and $D = 5$ dimensions, nothing spectacular happens and in particular we do not observe any indication of logarithmic corrections to the critical behavior in $D = 5$, in accordance with renormalization group expectations and also with newer simulations, cf. the careful discussion of five-dimensional percolation in (Fortunato *et al.*, 2004).

Due to the theoretically expected multiplicative logarithmic corrections (17) at the upper critical dimension a special treatment of the case $D = 6$ is needed. The analysis of logarithmic corrections of the general form

$$f(p) \sim (p_c - p)^{-\gamma} |\ln(p_c - p)|^\delta \quad (20)$$

is possible by (Adler and Privman, 1981) calculating approximants for

$$F(p) = (p_c - p) \ln(p_c - p) \left[\frac{f'(p)}{f(p)} - \frac{\gamma}{p_c - p} \right], \quad (21)$$

TABLE II Bond percolation thresholds for hypercubic lattices \mathbb{Z}^D for dimensions $D \geq 5$.

D	Present HT series	MC (Grassberger, 2003)	Present $1/D$ -expansion eq. (12)
5	0.118 170(5)	0.118 172(1)	0.118 149
6	0.094 202 0(10)	0.094 201 9(6)	0.094 354 3
7	0.078 682(2)	0.078 675 2(3)	0.078 688 1
8	0.067 712(1)	0.067 708 39(7)	0.067 708 0
9	0.059 497(1)	0.059 496 01(5)	0.059 495 1
10	0.053 093 5(5)	0.053 092 58(4)	0.053 092 13
11	0.047 950 3(1)	0.047 949 69(1)	0.047 949 47
12	0.043 724 1(1)	0.043 723 86(1)	0.043 723 76
13	0.040 187 7(1)	0.040 187 62(1)	0.040 187 57
14	0.037 183 8(1)		0.037 183 68

where one expects for singularities of the form (20) that $\lim_{p \rightarrow p_c} F(p) = \delta$. Using this method biased to $\gamma = 1$ we find the values for p_c and δ listed in Table I.

B. The percolation threshold in large dimensions

In (Grassberger, 2003), Grassberger compared his Monte Carlo results for p_c with a large- D expansion up to order 5 in $1/D$ (Gaunt and Ruskin, 1978). Since we extended this expansion up to order 9, a new comparison may be in order. This expansion is believed to be only asymptotic. Therefore we stop summing up terms of the expansion if the next term is larger than the current one. This recipe results in using only the terms up to order 7 in $D = 5$ and using all available terms for $D > 5$. We also analyzed our susceptibility series at fixed D using differential approximants and the M1 and M2 methods. Above the critical dimension, the M1 and M2 methods are used in a variant biased to $\gamma = 1$. Our results are collected in Table II.

For higher dimensions our large-dimensionality expansion is in perfect agreement with Grassberger's recent Monte Carlo (MC) data (Grassberger, 2003), whereas direct analyses of the susceptibility series in the respective dimension give slightly larger estimates for p_c . An obvious outlier is $D = 6$ where the cut-off criterion for the $1/D$ -expansion apparently does not work (taking artificially one term less would significantly improve the estimate).

C. Mean cluster number and moments of the cluster number

Our series for the free energy $\ln Z$ do not allow a reliable estimation of the critical exponent α . This phenomenon is already known from, e.g., Ising model series. One needs higher derivatives $\frac{\partial^n \ln Z}{\partial p^n}$ in order to see a singular contribution which shortens the series and due to the nature of the non-singular background the usual extrapolation methods give much worse results than, e.g., for the susceptibility series.

On the other hand, we can determine some interesting non-universal quantities such as the mean number of clusters per site,

$$\langle n_c \rangle = \lim_{||G|| \rightarrow \infty} \langle N_c \rangle / ||G||, \quad (22)$$

and its n th moments. Here, $||G||$ is the number of sites of the graph G . Using $w = q \frac{p}{1-p}$ and denoting the number of edges of the graph by $|G|$, we rewrite the partition sum (2) as

$$Z_G(q, p) = \frac{1}{(1-p)^{|G|}} \sum_C q^{N_c(C)+|C|} p^{|C|} (1-p)^{|G|-|C|}, \quad (23)$$

implying that

$$\langle (N_c + |C|)^n \rangle = \lim_{q \rightarrow 1} \frac{1}{Z} \left(q \frac{\partial}{\partial q} \right)^n Z. \quad (24)$$

The mean number of active bonds per side $n_b = |C|/||G||$ is for a D -dimensional hypercubic lattice given by $\langle n_b \rangle = pD$ and we get

$$\langle n_c \rangle = \lim_{q \rightarrow 1} \frac{\partial f}{\partial q} - pD \quad \text{and} \quad ||G|| [\langle (n_c + n_b)^2 \rangle - \langle n_c + n_b \rangle^2] = \lim_{q \rightarrow 1} \frac{\partial}{\partial q} \left(q \frac{\partial f}{\partial q} \right), \quad (25)$$

TABLE III Mean cluster number and fluctuation.

Dim.	$\langle n_c \rangle$	$\ G\ [\langle (n_c + n_b)^2 \rangle - \langle n_c + n_b \rangle^2]$
2	0.097 9(1)	0.161(3)
3	0.272 89(3)	0.029 7(5)
4	0.365 494(2)	0.007 56(2)
5	0.411 852(5)	0.003 04(3)
6	0.436 327(5)	0.001 64(1)

where f is the free energy $\ln Z$ per site.

By using Padé extrapolation to estimate $\partial f / \partial q$ and $\partial^2 f / \partial q^2$ at the percolation threshold p_c , we obtain the estimates for the mean cluster numbers and the variances eq. 25 given in Table III. As usual, the error estimates in Table III characterize the scatter of Padé approximants and do not include systematic errors which are presumably much larger.

For the first number in Table III, the number of clusters per site $\langle n_c \rangle$ in critical $D = 2$ bond percolation, Temperley and Lieb derived an exact expression in 1971. This expression was simplified in (Ziff *et al.*, 1997) to $\langle n_c \rangle = \frac{3\sqrt{3}-5}{2} \approx 0.098076$ which affirms our analysis. For bond percolation in three dimensions, Monte Carlo simulations (Lorenz and Ziff, 1998b) obtained $\langle n_c \rangle = 0.272\,931\,0(5)$. The fluctuations were studied in a recent Monte Carlo simulation (Deng and Yang, 2006). Their results given for $\|G\| [\langle n_c^2 \rangle - \langle n_c \rangle^2]$ and $\|G\| [\langle n_c n_b \rangle - \langle n_c \rangle \langle n_b \rangle]$ together with $\|G\| [\langle n_b^2 \rangle - \langle n_b \rangle^2] = \frac{z}{2} p(1-p)$ where $z = 2D$ is the coordination number of the lattice and p equals p_c imply $\|G\| [\langle (n_c + n_b)^2 \rangle - \langle n_c + n_b \rangle^2] = 0.164\,45(8)$ for two dimensions and $0.030\,71(45)$ for three dimensions, slightly larger but still consistent with our data.

VIII. SUMMARY

We successfully applied the method of high-temperature series expansion to q -state Potts model on hypercubic lattices. Modern computer facilities enable the calculation of such series while keeping some parameters (like q or even D) symbolic. This allows one to scan a whole parameter range. Even for the special case $q = 1$ we extended the known series by several terms and the results of the singularity analysis are comparable to and in good agreement with Monte Carlo data. By reverting the inverse susceptibility series, we derived the large- D expansion for the (pseudo-) transition point of general q -state Potts models up to order D^{-9} . Further applications such as the study of the tree percolation limit $q \rightarrow 0$ are conceivable, too.

Acknowledgments

We thank Joan Adler for discussions and help with the series analysis and Robert Ziff for a discussion on cluster cumulants. Support by DFG grant No. JA 483/17-3 and partial support from the German-Israeli-Foundation under grant No. I-653-181.14/1999 is gratefully acknowledged.

References

- Adler, J., A. Aharony, A. B. Harris, L. Klein, and Y. Meir, 1991, Series expansions for the Ising spin glass in general dimension, Phys. Rev. B **43**, 11249.
- Adler, J., Y. Meir, A. Aharony, and A. B. Harris, 1990, Series study of percolation moments in general dimension, Phys. Rev. B **41**, 9183.
- Adler, J., and V. Privman, 1981, Two-dimensional percolation: Logarithmic corrections to the critical behaviour from series expansions, J. Phys. A **14**(11), L463.
- Ballesteros, H. G., L. A. Fernández, V. Martín-Mayor, G. Parisi, and J. J. Ruiz-Lorenzo, 1999, Scaling corrections: site-percolation and Ising model in three dimensions, J. Phys. A **32**, 1 [cond-mat/9805125].
- Ballesteros, H. G., L. A. Fernández, V. Martín-Mayor, A. M. Sudupe, G. Parisi, and J. J. Ruiz-Lorenzo, 1997, Measures of critical exponents in the four dimensional site percolation, Phys. Lett. B **400**, 346 [hep-lat/9612024].
- Butera, P., and M. Comi, 2002, A library of extended high-temperature expansions of basic observables for the spin S Ising models on two- and three-dimensional lattices, J. Stat. Phys. **109**, 311 [hep-lat/0204007].
- Conway, A. R., and A. J. Guttmann, 1995, On two-dimensional percolation, J. Phys. A **28**(4), 891.
- Deng, Y., and H. W. J. Blöte, 2005, Monte Carlo study of the site-percolation model in two and three dimensions, Phys. Rev. E **72**, 016126.
- Deng, Y., and X. Yang, 2006, Finite-size scaling of energylike quantities in percolation, Phys. Rev. E **73**, 066116.

- Domb, C., and M. S. Green (eds.), 1974, *Series Expansions for Lattice Models*, volume 3 of *Phase Transitions and Critical Phenomena* (Academic Press, New York).
- Enting, I. G., and A. J. Guttmann, 2003, Susceptibility amplitudes for the three- and four-state Potts models, *Physica A* **321**, 90.
- Essam, I. W., D. S. Gaunt, and A. J. Guttmann, 1978, Percolation theory at the critical dimension, *J. Phys. A* **11**(10), 1983.
- Fisher, M. E., and R. R. Singh, 1990, Critical points, large-dimensionality expansions, and the Ising spin glass, in *Disorder in Physical Systems*, edited by G. Grimmett and D. Welsh (Oxford University Press, Oxford), pp. 87–111.
- Fortunato, S., A. Aharony, A. Coniglio, and D. Stauffer, 2004, Number of spanning clusters at the high-dimensional percolation thresholds, *Phys. Rev. E* **70**, 056116 [cond-mat/0407276].
- Gaunt, D. S., and H. Ruskin, 1978, Bond percolation processes in d dimensions, *J. Phys. A* **11**(7), 1369.
- Gerber, P. R., and M. E. Fisher, 1974, Critical temperatures of classical n -vector models on hypercubic lattices, *Phys. Rev. B* **10**(11), 4697.
- Grassberger, P., 2003, Critical percolation in high dimensions, *Phys. Rev. E* **67**, 036101 [cond-mat/0202144].
- Grollau, S., M. L. Rosinberg, and G. Tarjus, 2001, The ferromagnetic q -states Potts model on three-dimensional lattices: an analytical study for real values of q , *Physica A* **286**, 460 [cond-mat/0011483].
- Guttmann, A. J., 1989, Asymptotic analysis of power-series expansions, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L. Lebowitz (Academic Press, New York), volume 13, pp. 1–234.
- Guttmann, A. J., and I. G. Enting, 1994, Series studies of the Potts model: III. The 3-state model on the simple cubic lattice, *J. Phys. A* **27**, 5801 [hep-lat/9312083].
- Hellmund, M., and W. Janke, 2002a, High-temperature series expansions for random-bond Potts models on \mathbb{Z}^D , *Comp. Phys. Comm.* **147**(1–2), 435.
- Hellmund, M., and W. Janke, 2002b, Random-bond Potts models on hypercubic lattices: high-temperature series expansions, *Nucl. Phys. B (Proc. Suppl.)* **106–107**, 923.
- Hellmund, M., and W. Janke, 2003, Star-graph expansions for bond diluted Potts models, *Phys. Rev. E* **67**, 026118.
- Hellmund, M., and W. Janke, 2005, High-temperature series expansions for random Potts models, *Condens. Matter Phys.* **8**(1), 59 [cond-mat/0502152].
- Jacobsen, J. L., J. Salas, and A. D. Sokal, 2005, Spanning forests and the q -state Potts model in the limit $q \rightarrow 0$, *J. Stat. Phys.* **119**, 1153 [cond-mat/0401026].
- Janke, W., B. Berche, C. Chatelain, P.-E. Berche, and M. Hellmund, 2005, in *Proceedings of “Lattice 2005 Dublin”*, PoS, p. 018, PoS(LAT2005)018.
- Jensen, I., and R. M. Ziff, 2006, The universal amplitude ratio Γ^-/Γ^+ for two-dimensional percolation, eprint cond-mat/0607146.
- Lorenz, C. D., and R. M. Ziff, 1998a, Precise determination of the bond percolation thresholds and finite-size scaling corrections for the sc, fcc, and bcc lattices, *Phys. Rev. E* **57**(1), 230 [cond-mat/9710044].
- Lorenz, C. D., and R. M. Ziff, 1998b, Universality of the excess number of clusters and the crossing probability function in three-dimensional percolation, *J. Phys. A* **31**(40), 8147.
- Martin, J. L., 1974, Computer techniques for evaluating lattice constants, in (Domb and Green, 1974), pp. 97–112.
- Moraal, H., 1993, High-temperature expansions for classical spin models, *Physica A* **197**, 457.
- Paul, G., R. M. Ziff, and H. E. Stanley, 2001, Percolation threshold, Fisher exponent, and shortest path exponent for 4 and 5 dimensions, *Phys. Rev. E* **64**, 026115 [cond-mat/0101136].
- Schreider, G., and J. D. Reger, 1994, The q -state Potts-ferromagnet on d -dimensional hypercubic lattices, *J. Phys. A* **27**(4), 1071.
- Sokal, A. D., 2005, The multivariate Tutte polynomial (alias Potts model) for graphs and matroids, in *Surveys in Combinatorics, 2005*, edited by B. S. Webb (Cambridge University Press, Cambridge), number 327 in London Mathematical Society Lecture Note Series, pp. 173–226 [math/0503607].
- Stauffer, D., and A. Aharony, 1994, *Introduction to Percolation Theory* (Taylor & Francis, London), revised 2nd edition.
- Wu, F. Y., 1982, The Potts model, *Rev. Mod. Phys.* **54**(1), 235.
- Ziff, R. M., S. R. Finch, and V. S. Adamchik, 1997, Universality of finite-size corrections to the number of critical percolation clusters, *Phys. Rev. Lett.* **79**(18), 3447 [cond-mat/9707168].
- Ziff, R. M., and G. Stell, 1988, University of Michigan Report No. 88-4.