

Scaling property and the generalized entropy uniquely determined by a fundamental nonlinear differential equation

Hiroki Suyari

Department of Information and Image Sciences,
Chiba University, Chiba 263-8522, Japan

Tatsuaki Wada^y

Department of Electrical and Electronic Engineering,
Ibaraki University, Hitachi, Ibaraki 316-8511, Japan

(Dated: November 18, 2021)

Abstract

We derive a scaling property from a fundamental nonlinear differential equation whose solution is the so-called q -exponential function. A scaling property has been believed to be given by a power function only, but actually more general expression for the scaling property is found to be a solution of the above fundamental nonlinear differential equation. In fact, any power function is obtained by restricting the domain of the q -exponential function appropriately. As similarly as the correspondence between the exponential function and Shannon entropy, an appropriate generalization of Shannon entropy is expected for the scaling property. Although the q -exponential function is often appeared in the optimal distributions of some one-parameter generalized entropies such as Renyi entropy, only Tsallis entropy is uniquely derived from the algebra of the q -exponential function, whose uniqueness is shown in the two ways in this paper.

Keywords: q -exponential function, scaling property, q -product, q -multinomial coefficient, q -Stirling's formula, Tsallis entropy, generalized Shannon additivity

Electronic address: suyari@faculty.chiba-u.jp, suyari@ieee.org

^yElectronic address: wada@ee.ibaraki.ac.jp

I. SCALING PROPERTY DERIVED FROM A FUNDAMENTAL NONLINEAR DIFFERENTIAL EQUATION

The exponential function is often appeared in every scientific field. Among many properties of the exponential function, the linear differential function $dy/dx = y$ is the most important characterization of the exponential function. A slightly nonlinear generalization of this linear differential equation is given by

$$\frac{dy}{dx} = y^q \quad (q \in \mathbb{R}) : \quad (1)$$

(See the equation (17) at page 5 of [1] and the equations (22)–(23) at page 8 of [2].) This nonlinear differential equation is equivalent to

$$\int \frac{1}{y^q} dy = \int dx : \quad (2)$$

Then we define the so-called q -logarithm $\ln_q x$:

$$\ln_q x := \frac{x^{1-q} - 1}{1-q} \quad (3)$$

as a generalization of $\ln x$. Applying the property:

$$\frac{d}{dx} \ln_q x = \frac{1}{x^q}; \quad (4)$$

to (2), we obtain

$$\ln_q y = x + C \quad (5)$$

where C is any constant [3]. Then we define the so-called q -exponential $\exp_q(x)$ as the inverse function of $\ln_q x$ as follows:

$$\exp_q(x) := \begin{cases} [1 + (1-q)x]^{\frac{1}{1-q}} & \text{if } 1 + (1-q)x > 0; \\ 0 & \text{otherwise.} \end{cases} \quad (6)$$

Note that the q -logarithm and q -exponential recover the usual logarithm and exponential when $q \rightarrow 1$, respectively (See the pages 84–87 of [4] for the detail properties of these generalized functions $\ln_q x$ and $\exp_q(x)$). Thus, the general solution to the nonlinear differential equation (1) becomes

$$y = \exp_q(x + C) = \exp_q(C) \exp_q\left(\frac{x}{\exp_q(C)^{1-q}}\right) \quad (7)$$

where C is any constant satisfying $1 + (1 - q)C > 0$: Dividing the both sides by $\exp_q(C)$ of the above solution, we obtain

$$\frac{y}{\exp_q(C)} = \exp_q \left(\frac{x}{\exp_q(C)^{1-q}} \right) \quad (8)$$

Under the following scaling:

$$y_0 := \frac{y}{\exp_q(C)}; \quad x_0 := \frac{x}{\exp_q(C)^{1-q}}; \quad (9)$$

we obtain

$$y_0 = \exp_q(x_0) \quad (10)$$

This means that the solution of the nonlinear differential equation (1) obtained above is "scale-invariant" under the above scaling (9). Moreover, we can choose any constant C satisfying $1 + (1 - q)C > 0$ because C is an integration constant of (2).

Note that the above scaling (9) with respect to both variables x and y can be observed only when $q \neq 1$ and $C \neq 0$. In fact, when $q = 1$; i.e., $y = \exp(x + C)$; (9) reduces to the scaling with respect to only y ; i.e., $x_0 = x$, and when $C = 0$; both scalings in (9) disappears [3].

The above scaling property of the nonlinear differential equation (1) is very significant in the fundamental formulations for every generalization based on (1). We summarize the important points in the above fundamental result.

1. C in the scaling (9) is any constant satisfying $1 + (1 - q)C > 0$ because C is an integration constant of (2). This means that the scaling (9) is arbitrary for any q and C if q and C satisfies $1 + (1 - q)C > 0$ ($q \neq 1$ and $C \neq 0$).
2. In general studies of differential equation, C is determined by the initial condition of the nonlinear differential equation (1). This means, when an observable in a dynamics grows according to the nonlinear differential equation (1), the initial condition determines the scaling of the dynamics. This is applicable to the analysis of the chaotic dynamics [5][6][7][8][9][10][11].
3. In general, for a mapping $f : X \rightarrow Y$, $f|_{X_0} : X_0 \rightarrow Y$ is called a restriction of a mapping f to X_0 which is denoted by $f|_{X_0}$. Let f_q be a q -exponential function (6)

from $\mathbb{R} \rightarrow \mathbb{R}^+$. For the restricted domain RQ_q defined by

$$RQ_q := \{x \in \mathbb{R} \mid (1-q)x \leq 1, q \in \mathbb{R}\}; \quad (11)$$

a restriction of a mapping f_q to RQ_q is denoted by

$$fQ_q := f_q|_{RQ_q} : RQ_q \rightarrow \mathbb{R}^+; \quad (12)$$

Then fQ_q becomes a power function:

$$fQ_q(x) = ((1-q)x)^{\frac{1}{1-q}} = x^{\frac{1}{1-q}}; \quad (13)$$

In the above formulations the only case $q < 1$ is discussed. As shown in the section IV, the q -generalizations along the line of (1) has a symmetry $q \leftrightarrow 2-q$, i.e., $1 - (1-q) \leftrightarrow 1 + (1-q)$. Therefore, the above $fQ_q(x)$ can be replaced by a restriction of a mapping f_{2-q} to RQ_{2-q} :

$$fQ_{2-q}(x) = ((q-1)x)^{\frac{1}{q-1}} = x^{\frac{1}{q-1}} \quad (14)$$

in accordance with the symmetry, which implies that the case $q > 1$ can be discussed. In this way, the restriction of the q -exponential function to the domain RQ_q ($q < 1$) or RQ_{2-q} ($q > 1$) coincide with a power function, which has been often appeared and discussed in science. In general, the restricted domain RQ_q or RQ_{2-q} is called "scaling domain" and its corresponding range is called "scaling range".

4. A power function $f : \mathbb{R} \rightarrow \mathbb{R}$ is known to be characterized by the following functional equation, i.e., there exists a function $g : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$f(bx) = g(b)f(x) \quad (15)$$

holds for any $b; x \in \mathbb{R}$. The above functional equation uniquely determines a power function

$$f(x) = f(1)x \quad (16)$$

for choosing $g(b) = b$. See some references such as [12] for the proof. On the other hand, a q -exponential function is characterized by the nonlinear differential equation (1) as similarly as an exponential function. Moreover, the solutions of (1) are scale-invariant under the scaling (9) and reduce to power functions when the domain is restricted to RQ_q or RQ_{2-q} as shown above. In fact, by restricting the domain of (8) to

RQ ; the general solution (8) of the nonlinear differential equation (1) reduces to the following power function according to (13).

$$\frac{y}{\exp_q(C)} = (1 - q) \frac{x}{\exp_q(C)^{1-q}} = \frac{(1 - q)^{\frac{1}{1-q}}}{\exp_q(C)} x^{\frac{1}{1-q}}; \quad (17)$$

that is,

$$y = (1 - q)^{\frac{1}{1-q}} x^{\frac{1}{1-q}}; \quad (18)$$

which is equivalent to (16) if

$$= \frac{1}{q - 1}; \quad (19)$$

Therefore, many discussions on "exponential versus power-law", i.e., " $\frac{dy}{dx} = y$ versus $f(x) = g(b)f(x)$ " should be replaced by "exponential versus q -exponential", i.e., " $\frac{dy}{dx} = y$ versus $\frac{dy}{dx} = y^q$ ", which is more natural from mathematical point of view.

As shown in these discussions, the fundamental nonlinear differential equation (1) provides us with not only the characterization of the q -exponential function but also the scaling property in its solution.

As similarly as the relation between the exponential function $\exp(x)$ and Shannon entropy, we expect the corresponding information measure to the q -exponential function $\exp_q(x)$. There exist some candidates such as Renyi entropy, Tsallis entropy and so on. But the algebra derived from the q -exponential function uniquely determines Tsallis entropy as the corresponding information measure. In the following sections of this paper, we present the two mathematical results to uniquely determine Tsallis entropy by means of the already established formulations such as the q -exponential law, the q -multiplicative coefficient and q -Stirling's formula.

II. q -EXPONENTIAL LAW

The exponential law plays an important role in mathematics, so this law is also expected to be generalized based on the q -exponential function $\exp_q(x)$. For this purpose, the new multiplication operation \otimes_q is introduced in [13] and [14] for satisfying the following identities:

$$\ln_q(x \otimes_q y) = \ln_q x + \ln_q y; \quad (20)$$

$$\exp_q(x) \otimes_q \exp_q(y) = \exp_q(x + y); \quad (21)$$

The concrete form of the q -logarithm or q -exponential has been already given in the previous section, so that the above requirements as q -exponential law leads us to the definition of \cdot_q between two positive numbers.

Definition 1 For two positive numbers x and y , the q -product \cdot_q is defined by

$$x \cdot_q y := \begin{cases} [x^{1-q} + y^{1-q} - 1]^{\frac{1}{1-q}}; & \text{if } x > 0; y > 0; x^{1-q} + y^{1-q} - 1 > 0; \\ 0; & \text{otherwise.} \end{cases} \quad (22)$$

The q -product recovers the usual product such that $\lim_{q \rightarrow 1} (x \cdot_q y) = xy$. The fundamental properties of the q -product \cdot_q are almost the same as the usual product, but

$$a(x \cdot_q y) \neq ax \cdot_q y \quad (a; x; y \in \mathbb{R}) : \quad (23)$$

The other properties of the q -product are available in [13] and [14].

In order to see one of the validities of the q -product, we recall the well known expression of the exponential function $\exp(x)$ given by

$$\exp(x) = \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n} \right)^n : \quad (24)$$

Replacing the power on the right side of (24) by the n times of the q -product \cdot_q^n :

$$x \cdot_q^n := \underbrace{x \cdot_q \{ \underbrace{x \cdot_q}_{n \text{ times}} \}}_{n \text{ times}}; \quad (25)$$

$\exp_q(x)$ is obtained. In other words, $\lim_{n \rightarrow \infty} \left(1 + \frac{x}{n} \right)^{\cdot_q^n}$ coincides with $\exp_q(x)$:

$$\exp_q(x) = \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n} \right)^{\cdot_q^n} \quad (26)$$

The proof of (26) is given in the appendix of [15]. This coincidence (26) indicates a validity of the q -product. In fact, the present results in the following sections reinforce it.

III. q -MULTINOMIAL COEFFICIENT AND q -STIRLING'S FORMULA

We briefly review the q -multinomial coefficient and the q -Stirling's formula by means of the q -product \cdot_q . As similarly as for the q -product, q -ratio is introduced as follows:

Definition 2 For two positive numbers x and y , the inverse operation to the q -product is defined by

$$x \otimes_q y := \begin{cases} [x^{1-q} y^{1-q} + 1]^{\frac{1}{1-q}}; & \text{if } x > 0; y > 0; x^{1-q} y^{1-q} + 1 > 0; \\ 0; & \text{otherwise} \end{cases} \quad (27)$$

which is called q -ratio in [14].

\otimes_q is also derived from the following satisfactions, similarly as for \otimes_q [13][14].

$$\ln_q(x \otimes_q y) = \ln_q x + \ln_q y; \quad (28)$$

$$\exp_q(x) \otimes_q \exp_q(y) = \exp_q(x + y); \quad (29)$$

The q -product and q -ratio are applied to the definition of the q -multinomial coefficient [15].

Definition 3 For $n = \sum_{i=1}^k n_i$ and $n_i \in \mathbb{N}$ ($i = 1, \dots, k$); the q -multinomial coefficient is defined by

$$\binom{n}{n_1, \dots, n_k}_q := \frac{n!_q}{n_1!_q \dots n_k!_q} = \frac{n!_q}{n_1!_q \otimes_q \dots \otimes_q n_k!_q}; \quad (30)$$

From the definition (30), it is clear that

$$\lim_{q \rightarrow 1} \binom{n}{n_1, \dots, n_k}_q = \binom{n}{n_1, \dots, n_k} = \frac{n!}{n_1! \dots n_k!}; \quad (31)$$

In addition to the q -multinomial coefficient, the q -Stirling's formula is useful for many applications such as our main results. By means of the q -product (22), the q -factorial $n!_q$ is naturally defined as follows.

Definition 4 For a natural number $n \in \mathbb{N}$ and $q \in \mathbb{R}^+$, the q -factorial $n!_q$ is defined by

$$n!_q := 1 \otimes_q \dots \otimes_q n; \quad (32)$$

Using the definition of the q -product (22), $\ln_q(n!_q)$ is explicitly expressed by $\ln_q(n!_q) = \sum_{k=1}^n \frac{k^{1-q}}{1-q}$. If an approximation of $\ln_q(n!_q)$ is not needed, this explicit form should be directly used for its computation. However, in order to clarify the correspondence between the studies $q = 1$ and $q \neq 1$, the approximation of $\ln_q(n!_q)$ is useful. In fact, using the following q -Stirling's formula, we obtain the unique generalized entropy corresponding to the q -exponential function $\exp_q(x)$, shown in the following sections.

Theorem 5 Let $n!_q$ be the q -factorial defined by §2). The rough q -Stirling's formula $\ln_q(n!_q)$ is computed as follows:

$$\ln_q(n!_q) = \begin{cases} < \frac{n}{2} \ln_q n - \frac{n}{2} + O(\ln_q n) & \text{if } q \neq 2; \\ : n \ln n + O(1) & \text{if } q = 2: \end{cases} \quad (33)$$

The proof of the above formulas (33) is given in [15].

IV. TSALLIS ENTROPY UNIQUELY DERIVED FROM THE q -MULTINOMIAL COEFFICIENT AND q -STIRLING'S FORMULA

In this section we show that Tsallis entropy is uniquely and naturally derived from the fundamental formulations presented in the previous section. In order to avoid separate discussions on the positivity of the argument in (30), we consider the q -logarithm of the q -multinomial coefficient to be given by

$$\ln_q \binom{n}{n_1, n_2, \dots, n_k}_q = \ln_q(n!_q) - \sum_{i=1}^k \ln_q(n_i!_q) \quad (34)$$

The unique generalized entropy corresponding to the q -exponential is derived from the q -multinomial coefficient using the q -Stirling's formula as follows [15].

Theorem 6 When n is sufficiently large, the q -logarithm of the q -multinomial coefficient coincides with Tsallis entropy (36) as follows:

$$\ln_q \binom{n}{n_1, n_2, \dots, n_k}_q \approx \begin{cases} \frac{n^2}{2} S_q \left(\frac{n_1}{n} \right); & \frac{n_k}{n}; \text{ if } q > 0; q \neq 2 \\ S_1(n) + \sum_{i=1}^k S_1(n_i) & \text{if } q = 2 \end{cases} \quad (35)$$

where S_q is Tsallis entropy [16]:

$$S_q = \frac{1}{q} \frac{\sum_{i=1}^n p_i^q}{1} \quad (36)$$

and $S_1(n)$ is given by $S_1(n) = \ln n$:

The proof of this theorem is given in [15].

Note that the above relation (35) reveals a surprising symmetry: (35) is equivalent to

$$\ln_{1-(1-q)} \left(\frac{n}{n_1} \right)^{1-q} = \frac{n^{1+(1-q)}}{1+(1-q)} \ln_{(1-q)} \left(\frac{n_1}{n} \right); \quad \frac{n_k}{n}; \quad (37)$$

for $q > 0$ and $q \neq 2$. This expression represents that there exists a symmetry with a factor $1 - q$ around $q = 1$ in the algebra of the q -product. Substitution of some concrete values of q into (35) or (37) helps us understand the meaning of this symmetry.

Remark that the above correspondence (35) and the symmetry (37) reveals that the q -exponential function (6) derived from (1) is consistent with Tsallis entropy only as information measure.

V. THE GENERALIZED SHANNON ADDITIVITY DERIVED FROM THE q -MULTINOMIAL COEFFICIENT

This section shows another way to uniquely determine the generalized entropy. More precisely, the identity derived from the q -multinomial coefficient coincides with the generalized Shannon additivity which is the most important axiom for Tsallis entropy.

Consider a partition of a given natural number n into k groups such as $n = \sum_{i=1}^k n_i$. In addition, each natural number n_i ($i = 1; \dots; k$) is divided into groups such as $n_i = \sum_{j=1}^{m_i} n_{ij}$ where $n_{ij} \geq 1$. Then, the following identity holds for the q -multinomial coefficient.




FIG. 1: partition of a natural number n

$$\frac{2}{4} n = \frac{3}{5} = \frac{2}{4} n = \frac{3}{5} \frac{2}{4} n_1 = \frac{3}{5} \frac{2}{4} n_k = \frac{3}{5} \frac{2}{4} n_k \quad (38)$$

It is very easy to prove the above relation (38) by taking the q -logarithm of the both sides and using (34).

On the other hand, the above identity (38) is reform ed to the generalized Shannon additivity in the following way. Taking the q -logarithm of the both sides of the above relation (38), we have

$$\ln_q 4^n = \ln_q 4^{n_1} + \sum_{i=1}^{X^k} \ln_q 4^{n_{i1}} \quad (39)$$

From the relation (35), we obtain

$$S_{2-q} \left(\frac{n_{11}}{n}; \frac{n_{km_k}}{n} \right) = S_{2-q} \left(\frac{n_1}{n}; \frac{n_k}{n} \right) + \sum_{i=1}^{X^k} \frac{n_i}{n} S_{2-q} \left(\frac{n_{i1}}{n_i}; \frac{n_{im_i}}{n_i} \right) \quad (40)$$

Then, by means of the following probabilities de ned by

$$p_{ij} := \frac{n_{ij}}{n} \quad (i=1; \dots; k; \quad j=1; \dots; m) \quad (41)$$

$$p_i := \sum_{j=1}^{X^i} p_{ij} = \sum_{j=1}^{X^i} \frac{n_{ij}}{n} = \frac{n_i}{n} \quad * \quad n_i = \sum_{j=1}^{X^i} n_{ij} \quad (42)$$

the identity (40) becomes

$$S_q(p_{11}; \dots; p_{km_k}) = S_q(p_1; \dots; p_k) + \sum_{i=1}^{X^k} p_i S_q \left(\frac{p_{i1}}{p_i}; \dots; \frac{p_{im_i}}{p_i} \right) \quad (43)$$

The formula (43) obtained from the q -multinomial coefficient is exactly same as the generalized Shannon additivity (See [GSK3] given below) which is the most important axiom for Tsallis entropy [17].

In fact, the generalized Shannon-Khinchin axioms and the uniqueness theorem for the nonextensive entropy are already given and rigorously proved in [17]. The present result (43) and the already established axiom [GSK3] perfectly coincide with each other.

Theorem 7 Let Δ_n be de ned by the n -dimensional simplex:

$$\Delta_n = \left\{ (p_1; \dots; p_n) \mid p_i \geq 0; \sum_{i=1}^n p_i = 1 \right\} \quad (44)$$

The following axioms [GSK1] [GSK4] determine the function $S_q : \Delta_n \rightarrow \mathbb{R}^+$ such that

$$S_q(p_1; \dots; p_n) = \frac{1 - \sum_{i=1}^n p_i^q}{(q)} \quad (45)$$

where (q) satisfies properties (i) (iv):

(i) q is continuous and has the same sign as $q - 1$; i.e.,

$$(q - 1)(q - 1) > 0; \quad (46)$$

(ii)

$$\lim_{q \rightarrow 1} (q) = (1) = 0; \quad (q) \in (0, 1); \quad (47)$$

(iii) there exists an interval $(a; b) \subset \mathbb{R}^+$ such that $a < 1 < b$ and (q) is differentiable on the interval

$$(a; 1) \cup (1; b); \quad (48)$$

and

(iv) there exists a constant $k > 0$ such that

$$\lim_{q \rightarrow 1} \frac{d}{dq} (q) = \frac{1}{k}; \quad (49)$$

[GSK 1] continuity: S_q is continuous in n and $q \in \mathbb{R}^+$,

[GSK 2] maximality: for any $q \in \mathbb{R}^+$, any $n \in \mathbb{N}$ and any $(p_1, \dots, p_n) \in \mathcal{P}_n$,

$$S_q(p_1, \dots, p_n) \geq S_q\left(\frac{1}{n}, \dots, \frac{1}{n}\right); \quad (50)$$

[GSK 3] generalized Shannon additivity: if

$$p_{ij} \geq 0; \quad \sum_{j=1}^n p_{ij} = 1; \quad \sum_{i=1}^m p_{ij} = 1; \quad i=1, \dots, n; \quad j=1, \dots, m \quad (51)$$

then the following equality holds:

$$S_q(p_{11}, \dots, p_{nm}) = S_q(p_1, \dots, p_n) + \sum_{i=1}^n p_i S_q\left(\frac{p_{i1}}{p_i}, \dots, \frac{p_{im}}{p_i}\right); \quad (52)$$

[GSK 4] expandability:

$$S_1(p_1, \dots, p_n, 0) = S_1(p_1, \dots, p_n); \quad (53)$$

Note that, in order to uniquely determine the Tsallis entropy (36) in the above set of the axioms, $\lim_{q \rightarrow 1}$ should be removed from (49), that is, $\frac{d}{dq} (q) = \frac{1}{k}$ (i.e., $(q) = \frac{1}{k} (q - 1)$) should be used instead of (49). The general form (q) perfectly corresponds to Tsallis' original

introduction of the so-called Tsallis entropy in 1988 [16]. See his original characterization shown in page 9 of [1] for the detail (q) corresponds to a in his notation. His simplest choice of a coincides with the simplest form of q i.e., $\frac{d(q)}{dq} = \frac{1}{k}$).

When one of the authors (H.S.) submitted the paper [17] in 2002, nobody presented the idea of the q -product. However, as shown above, the identity on the q -multinomial coefficient [15] which was formulated based on the q -product [13][14] coincides with one of the axioms (GSK3): generalized Shannon additivity in [17]. This means that the whole theory based on the q -product is self-consistent. Moreover, other fundamental applications of the q -product, such as law of error [18] and the derivation of the unique non self-referential q -canonical distribution [19][20], are also based on the q -product.

VI. CONCLUSION

Starting from a fundamental nonlinear equation $dy/dx = y^q$, we present the scaling property and the algebraic structure of its solution. Moreover, we prove that the algebra determined by its solutions is mathematically consistent with Tsallis entropy only as the corresponding unique information measure based on the following 2 mathematical reasons:

- (1) derivation of Tsallis entropy from the q -multinomial coefficient and q -Stirling's formula,
- (2) coincidence of the identity derived from the q -multinomial coefficient with the generalized Shannon additivity which is the most important axiom for Tsallis entropy.

These mathematical discussions result in the self-consistency between the mathematics derived from the q -exponential and Tsallis entropy.

Recently, we have presented the following fundamental results in Tsallis statistics:

1. Axioms and the uniqueness theorem for the nonextensive entropy [17]
2. Law of error in Tsallis statistics [18]
3. q -Stirling's formula in Tsallis statistics [15]
4. q -multinomial coefficient in Tsallis statistics [15]
5. Central limit theorem in Tsallis statistics (numerical evidence only) [15]
6. q -Pascal's triangle in Tsallis statistics [15]

7. The unique non self-referential q -canonical distribution in Tsallis statistics [19][20]
8. Scaling property characterized by the fundamental nonlinear differential equation [the present paper].

All of the above fundamental results are derived from the algebra of the q -product and mathematically consistent with each other. This means that the q -product is indispensable to the formalism in Tsallis statistics. More important point is that the q -product originates from the fundamental nonlinear differential equation (1) with scale-invariant solutions.

-
- [1] M. Gell-Mann and C. Tsallis, eds., *Nonextensive Entropy: Interdisciplinary Applications* (Oxford Univ. Press, New York, 2004).
 - [2] C. Tsallis, What should a statistical mechanics satisfy to reflect nature?, *Physica D*, 193, 3-34 (2004).
 - [3] In any manuscripts and papers up to now, this constant C is set to be 1 , i.e., $\exp_q(C) = 1$ (See the equation (17) at page 5 of [1] and the equations (22)-(23) at page 8 of [2]). But the arbitrariness of this constant C plays a very important role in scaling (9).
 - [4] S. Abe and Y. Okamoto, eds., *Nonextensive Statistical Mechanics and Its Applications* (Springer-Verlag, Heidelberg, 2001).
 - [5] C. Tsallis, A. R. Plastino and W.-M. Zheng, Power-law sensitivity to initial conditions - New entropic representation, *Chaos, Solitons and Fractals* 8, 885-891 (1997).
 - [6] V. Latora, M. Baranger, A. Rapisarda and C. Tsallis, The rate of entropy increase at the edge of chaos, *Phys. Lett. A* 273, 97-103 (2000).
 - [7] F. Baldovin and A. Robledo, Universal renormalization-group dynamics at the onset of chaos in logistic maps and nonextensive statistical mechanics, *Phys. Rev. E* 66, 045104(R) (2002).
 - [8] F. Baldovin and A. Robledo, Sensitivity to initial conditions at bifurcations in one-dimensional nonlinear maps: Rigorous nonextensive solutions, *Europhys. Lett.* 60, 518-524 (2002).
 - [9] A. Robledo, Criticality in nonlinear one-dimensional maps: RG universal map and nonextensive entropy, *Physica D* 193, 153-160 (2004).
 - [10] F. Baldovin and A. Robledo, Nonextensive Pesin identity. Exact renormalization group analytical results for the dynamics at the edge of chaos of the logistic map, *Phys. Rev. E* 69,

045202(R) (2004).

- [11] U. Timakli and C. Tsallis, Chaos edges of z -logistic maps: Connection between the relaxation and sensitivity entropic indices, *Phys. Rev. E* 73, 037201 (2006).
- [12] M. E. J. Newman, Power laws, Pareto distributions and Zipf's law, *Contemporary Physics* 46, 323-351 (2005).
- [13] L. N. Ivanen, A. Le M ehaute, Q. A. Wang, Generalized algebra within a nonextensive statistics, *Rep. Math. Phys.* 52, 437-444 (2003).
- [14] E. P. Borges, A possible deformed algebra and calculus inspired in nonextensive therm ostatistics, *Physica A* 340, 95-101 (2004).
- [15] H. Suyari, Mathematical structures derived from the q -multinomial coefficient in Tsallis statistics, *Physica A*, 368, pp.63-82 (2006).
- [16] C. Tsallis, Possible generalization of Boltzmann-G ibbs statistics, *J. Stat. Phys.* 52, 479-487 (1988).
- [17] H. Suyari, Generalization of Shannon-Khinchin axioms to nonextensive systems and the uniqueness theorem for the nonextensive entropy, *IEEE Trans. Inform .Theory* 50, 1783-1787 (2004).
- [18] H. Suyari and M. Tsukada, Law of error in Tsallis statistics, *IEEE Trans. Inform .Theory* 51, 753-757 (2005).
- [19] T. Wada and A. M. Scarfone, A non self-referential expression of Tsallis' probability distribution function, *EurPhysJB* 47 557-562 (2005).
- [20] H. Suyari, The unique non self-referential q -canonical distribution and the physical temperature derived from the maximum entropy principle in Tsallis statistics, *Prog. Theor. Phys. Suppl.*, 162, pp.79-86 (2006). (See the author's website (<http://www.ne.jp/asahi/hiroki/suyari/publications.htm>) for the detail derivation.)