

On the dynamics of spin systems in the Landau-Lifshitz theory

U. Krey

Inst. für Physik II, Universität Regensburg, 93040 Regensburg, Germany

November 30, 2006

Abstract

In the framework of the Landau-Lifshitz equations without any dissipation (an approximation which may also be helpful for finite but weak Gilbert damping), with all interactions included (specially always with exchange and magnetic dipole interactions), for general ground states, geometries and domain structures, and for many types of effective fields (including contributions from spin currents etc.) the dynamics of the spin-precession around this ground state is considered.

At first the precession is treated in the linear approximation. For the (small amplitude) eigenmodes of the precession one has a 'rule of geometric mean' for the eigenfrequencies. For the eigenmodes pseudo-orthogonality relations are obtained, which reflect the gyrotropic and elliptic character of the spin precession and differ from those known from the Schrödinger equation. Moreover, in contrast to quantum mechanics, they are defined locally, not through a global average. Thus pseudo-orthogonality statements are valid everywhere (for example simultaneously both in the outer region and also in the core region of a magnetic vortex line).

Then also some aspects of the nonlinear mode coupling with emphasis on 'confluence' and 'splitting' processes of elementary magnetic spin-wave excitations are considered. At the same time these processes contribute to the Gilbert damping.

There are thus essential differences to quantum mechanics, although at a first glance one discovers many similarities. From the results one may also get insights of why these systems are so complex that (although the essential quantities depend only on the local values of the partially long-ranged effective magnetic fields) practically only detailed experiments and computer simulations make sense.

PACS numbers: 75. (Magnetic properties); 75.40 GB (Dynamic properties)

Keywords: Landau-Lifshitz Eqs., Dynamics of Magnetic Systems

1 Introduction

Recently, for possible applications in spintronics, the dynamic behaviour of ultrathin planar magnetic nano-structures is strongly studied ([1,2,...,9], and references therein). For example, one considers circular structures with nontrivial ground states (notably with vortex states).

Theoretically, on the spatial scale of typically ten (or more) nanometers (nm) and on the time scale of ten (or more) picoseconds (ps) the Landau-Lifshitz equations, [10],

e-mail: uwe.krey@physik.uni-regensburg.de

with small phenomenological Gilbert damping', [11], is very reliable and has found many applications in present-day simulation software (for example, in the well-known OOMF, [12], or LLG programs, [13]). It is 'general folklore' that there are considerable similarities between the Landau-Lifshitz-equation without Gilbert damping and the quantum mechanical Schrodinger equation. This would be advantageous, since for weak Gilbert damping one could then exploit a lot of quantum mechanical relations.

Unfortunately, however, difficulties arise, since (as we will see) there are considerable differences from the Schrodinger case.

The present communication intends to make these points clearer; it also serves more general purposes, for example in the intention to go beyond numerical simulation as far as possible.

2 Basic theory

In the framework of the Landau-Lifshitz theory, [10], the local value of the magnetic polarisation $\mathbf{J}(\mathbf{r};t)$ (as a function of position \mathbf{r} and time t) is described by the following ansatz:

$$\mathbf{J}(\mathbf{r};t) = J_s \cdot \hat{\mathbf{n}}(\mathbf{r};t); \quad (1)$$

where J_s is the saturation magnitude and $\hat{\mathbf{n}}(\mathbf{r};t)$ the direction of the magnetic polarisation vector \mathbf{J} . The vector $\hat{\mathbf{n}}$ is thus of unit length.

This fact is important throughout the paper. Essentially due to this constraint on $\hat{\mathbf{n}}(\mathbf{r};t)$ the Landau-Lifshitz theory has the same topological excitations (Bloch lines, Bloch point singularities, vortex excitations, and all kinds of 'topological constrictions') as in field-theory the so-called 'non-linear sigma-model', [14], has; or as in elementary particle physics the corresponding phenomenological Nambu and Jona-Lasinio model has, [15]. However, since the magnetic dipole field (see below) is far-ranged, the Landau-Lifshitz theory is in principle even more complicated than the above-mentioned field theories (which usually do not treat interactions corresponding to the dipole-dipole case).

In any case, the role of the topological constraints in applications (for example, in the wall-motion of magnetic bubble-domains, [16], or in the change of the polarization of the core of a magnetic vortex-line, [17]) should not be underestimated.

However the present paper does not concentrate on topological aspects!

Rather, in the linear approximation (neglecting the damping unless otherwise stated) we derive a 'rule of geometric mean' for the eigenfrequencies and a pseudo-orthogonality relation for the eigenmodes of the precession around the nontrivial ground state. The relations correspond to a gyrotropic and elliptic motion of the magnetization; they differ from those known from quantum mechanics. In particular they are defined locally.

Nonlinear terms are treated in the present paper rather phenomenologically. We only give some schematic arguments for damping processes based on 'confluence' and 'splitting' in connection with the so-called mode-mode coupling of excitations in vortex structures. The results are in favour of the experiments of [2] and [3].

3 Landau-Lifshitz equations

In the absence of any dissipation (see above), on the above-mentioned spatial and temporal scales, the equation of motion for the vector $\mathbf{m}(\mathbf{r};t)$ is the undamped Landau-Lifshitz equation, [10], which can be written as

$$\frac{d\mathbf{m}(\mathbf{r};t)}{dt} = \gamma \mathbf{m}(\mathbf{r};t) \times \mathbf{H}_e(\mathbf{r};t); \quad (2)$$

where the positive quantity $\gamma = g \frac{e\hbar}{2m_e}$ is the so-called Lande factor g (≈ 2) multiplied by the gyromagnetic ratio; e ($= -1.6 \times 10^{-19}$ C) is the electron's charge (a negative quantity), m_e its mass, and \hbar the permability of the vacuum; $\mathbf{H}_e(\mathbf{r};t)$ is an effective field, namely the sum

$$\mathbf{H}_e = \mathbf{H}^{\text{ext}} + \mathbf{H}^{\text{dipole}} + \sum_{i,j,k=1}^3 \frac{2A_{ijk}}{J_s} \frac{\partial \mathbf{m}}{\partial x_i} \times \frac{\partial \mathbf{m}}{\partial x_k} + \text{grad } F_0(\mathbf{m}) + \mathbf{H}^{\text{therm}} + \mathbf{H}^{\text{J}}; \quad (3)$$

In (3), \mathbf{H}^{ext} is the external magnetic field; the A_{ijk} are the so-called exchange constants of the crystal; $F_0(\mathbf{m})$ is the anisotropy energy, representing the easy axes of the magnetization; $\mathbf{H}^{\text{therm}}$ is a fluctuating thermal field $\propto \sqrt{T}$, where T is the Kelvin temperature; this field is neglected together with the friction, to which it is intrinsically related. Finally, \mathbf{H}^{J} is an effective field proportional to the density of the electric current, \mathbf{J} , or to other sources of torques acting on the magnetization. For stationary currents all these contributions to \mathbf{H}_e do not depend explicitly on the time nor on the time derivative, [19]. Often they all can be neglected with respect to the second and/or third term on the r.h.s. of equation (3).

The remaining second term on the r.h.s. of (3), $\mathbf{H}^{\text{dipole}}$, is the cumbersome quantity in micromagnetism, namely the magnetic dipole field. In contrast, the third term represents the exchange interactions, which at short wavelengths are much stronger, such that practically they alone determine the critical temperature T_c of the ferromagnet considered. But for applications at room temperatures and below, at distances which are much larger than the atomic lattice constants (typically 0.2 nm) the dipole fields are most important.

Therefore even in the μm and nm ranges, as mentioned, the exchange interactions can often be neglected (we don't neglect them here!), whereas the field $\mathbf{H}^{\text{dipole}}$, which is, for example, mainly responsible for the domain structure of a ferromagnet, can be derived from the relation $\mathbf{H}^{\text{dipole}} = -\text{grad } \phi_m$.

The magnetostatic potential ϕ_m is determined as follows:

(i) either from the magnetic moment representation

$$\phi_m(\mathbf{r};t) = \frac{1}{4\pi} \int_V d^3x' \frac{\mathbf{J}(\mathbf{r}';t) \cdot (\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^3} \quad (4)$$

(here the integration volume V is that of the magnetic sample),

(ii) or from the equivalent representation of the fictitious magnetic charges :

$$\mathbf{m}(\mathbf{r};t) \stackrel{(ii)}{=} \int_V d^3\mathbf{x} \frac{\text{div} \mathbf{J}(\mathbf{x}^0; t)}{4\pi} + \int_{\partial V} d^2A \frac{\mathbf{J}(\mathbf{x}^0; t) \cdot \mathbf{n}}{4\pi} : \quad (5)$$

Here ∂V is the (oriented) boundary of V ; the vector \mathbf{n} is the its outer normal; the surface measure is d^2A . (Retardation effects do not play a role, unless the frequencies f ($= \frac{1}{2}$) are larger than $\frac{c}{x}$ ($= 3 \cdot 10^{10}$ Hz for $x = 100$ nm).)

In any case, the local value $\mathbf{H}^{\text{dipole}}(\mathbf{r};t)$ of the dipole field depends on the magnetization distribution of the whole sample; i.e., in the context of the Landau-Lifshitz equation the dipole field is a long-ranged, but local quantity, which sounds as a contradiction in itself, but becomes nonetheless essential below.

Unfortunately, the magnitude of this field can also be very large, namely at surfaces typically as large as $\frac{J_s}{0}$, which corresponds to 2.2 Tesla in Fe, 1.7 Tesla in Co and 1.0 Tesla in perm alloy. So for applications the magnetic dipole field is as important as the exchange.

4 Basic relations for the dynamics

4.1 General equations

In the following we write $\sim = \sim_0 + \tilde{\sim}$, where $\sim_0(\mathbf{r})$ is the static ground state, corresponding, e.g., to a non-trivial domain structure of the sample, whereas the vector $\tilde{\sim}(\mathbf{r};t)$ describes the dynamics of the sample. Here we use the linear approximation with respect to the ground state (i.e., we assume that \tilde{j}^j is $\ll 1$, while $\sim_0(\mathbf{r}) \cdot \tilde{\sim}(\mathbf{r};t) = 0$ and $\sim_0^2(\mathbf{r}) = 1$). As a consequence in linear approximation also $\sim^2(\mathbf{r};t) = 1$, as desired.

Furthermore we don't exclude topological singularities, e.g., Bloch points or vortex lines, but assume that these are exclusively contained in the statics of the system (i.e., in $\sim_0(\mathbf{r})$), whereas $\tilde{\sim}(\mathbf{r};t)$ is topologically trivial (but see [20]).

In the linear approximation one gets the following equation of motion :

$$\frac{d\tilde{\sim}}{dt} = \sim_0(\mathbf{r}) \times \mathbf{H}_e^{\sim} + \tilde{\sim}(\mathbf{r};t) \times \mathbf{H}_e^{\sim_0} : \quad (6)$$

Here we have used the reduced time $t \equiv \tau$ (in the following we replace τ by 1, unless otherwise stated).

But $\mathbf{H}_e^{\sim_0}(\mathbf{r})$ has the same direction as $\sim^{(0)}(\mathbf{r})$ (the reader is reminded to the relation $\sim^{(0)} \times \mathbf{H}_e = 0$). Therefore the final term in (6) can be replaced by a term with a Lagrange parameter function $\lambda^{\sim_0}(\mathbf{r})$:

$$\frac{d\tilde{\sim}}{dt} = \sim_0(\mathbf{r}) \times \mathbf{H}_e^{\sim}(\mathbf{r};t) - \lambda^{\sim_0}(\mathbf{r}) \sim(\mathbf{r};t) : \quad (7)$$

This is a linear relation in $\tilde{\sim}$. Generally only the local and instantaneous values of the effective field \mathbf{H}_e^{\sim} (and of $\tilde{\sim}$) are involved. It should also be noted that (7) does not explicitly depend on t , see (4) or (5).

Now we assume the existence of local cartesian coordinates such that the local 3-axis is $\sim_0(\mathbf{r})$, whereas $\mathbf{e}_1(\mathbf{r})$ and $\mathbf{e}_2(\mathbf{r})$ are perpendicular to $\sim_0(\mathbf{r})$ and to each other, but otherwise not yet fixed: Then in linear approximation the following equation for the vector $\sim(\mathbf{r};t)$ (which also must be perpendicular to \sim_0) results:

$$\frac{d}{dt} (\alpha_1(\mathbf{r};t) \mathbf{e}_1(\mathbf{r}) + \alpha_2(\mathbf{r};t) \mathbf{e}_2(\mathbf{r})) = \overset{\S}{K}(\mathbf{r}) (\alpha_1(\mathbf{r};t) \mathbf{e}_1(\mathbf{r}) + \alpha_2(\mathbf{r};t) \mathbf{e}_2(\mathbf{r})) : \quad (8)$$

Here $\overset{\S}{K}(\mathbf{r})$ is an antisymmetric real 2×2 -matrix, i.e., with imaginary eigenvalues (see below), such that (after suitable rotation of the axes perpendicular to $\sim_0(\mathbf{r})$) the equations for $\alpha_1(\mathbf{r};t)$ and $\alpha_2(\mathbf{r};t)$ become of the simple form

$$\frac{d}{dt} \alpha_1 = +h_2^{\text{eff}}(\mathbf{r}) \alpha_2 \quad (9)$$

and

$$\frac{d}{dt} \alpha_2 = -h_1^{\text{eff}}(\mathbf{r}) \alpha_1 ; \quad (10)$$

where $h_1^{\text{eff}}(\mathbf{r})$ and $h_2^{\text{eff}}(\mathbf{r})$ are extremal off-diagonal values of the 2×2 -matrix $\overset{\S}{K}(\mathbf{r})$, see below, which we assume to be non-negative, unless otherwise stated.

The equations (9) and (10) describe an elliptic spin precession around the local equilibrium spin direction $\sim_0(\mathbf{r})$, and the quantities h_1^{eff} and h_2^{eff} are local effective fields corresponding to the lengths of the principal axes of the local precession ellipse.

In the following we skip the superscripts of h_1^{eff} and h_2^{eff} .

4.2 The 'rule of the geometric mean' for the eigenfrequencies

We now make the ansatz $\alpha_1(\mathbf{r};t) = \alpha_1^{(0)}(\mathbf{r}) \cos(!t)$ and $\alpha_2(\mathbf{r};t) = \alpha_2^{(0)}(\mathbf{r}) \sin(!t)$, where all quantities are real, $!$ is positive, unless otherwise stated, and both polarisations enter (the upper (resp. lower) sign in the ansatz for α_2 correspond to mathematically positive (+) (resp. negative, (-)) circularly polarised precession; typically one has a superposition of both polarisations, which results in the above-mentioned elliptic precession. If the external field $H_3^{\text{ext}}(\mathbf{r})$ and/or the exchange field dominate the local effective field, see below, then one has purely positive (for positive H_3^{ext}) (resp. negative, for negative H_3^{ext}) circular polarisation.).

In any case, by the product of (9) and (10) one gets finally the eigenfrequency equation

$$(!)^2 = \frac{\sin !t}{\cos !t} \left(\frac{\alpha_2^{(0)}(\mathbf{r})}{\alpha_1^{(0)}(\mathbf{r})} \frac{h_1^{(0)}(\mathbf{r})}{h_2^{(0)}(\mathbf{r})} - \frac{\alpha_1^{(0)}(\mathbf{r})}{\alpha_2^{(0)}(\mathbf{r})} \frac{h_2^{(0)}(\mathbf{r})}{h_1^{(0)}(\mathbf{r})} \right) ; \quad (11)$$

where the gyromagnetic ratio γ has been restored. For the eigenfrequencies $!^{(i)}$ this yields the (well-known, see [21]) rule of the geometric mean (the index counts the eigenstates):

$$!^{(i)} = \sqrt{h_1^{(i)}(\mathbf{r}) h_2^{(i)}(\mathbf{r})} : \quad (12)$$

Here one should note that the l.h.s. of the result does not depend on \mathbf{r} , in contrast to the r.h.s. In fact in linear approximation, in the Landau-Lifshitz theory, one can

perform any position-averaging of the time-dependent signal (for example, over all spins, but usually with different weights at different positions): the eigenfrequency spectrum will always be the same. (This is useful for applications, [3], and it also allows that even from a distant position there may be contributions to $\tilde{H}_e(\mathbf{r};t)$.)

The relation (12) is actually at the same time more complex and also more simple than in quantum mechanics: on the one hand one has two functions, $h_1(\mathbf{r})$ and $h_2(\mathbf{r})$, instead of only one in quantum mechanics (i.e., the potential energy $V(\mathbf{r})$), whereas in quantum mechanics, on the other hand, not all space-averages, but only the full-space average with the eigenfunction, $\int d^3\mathbf{r} (\mathbf{r}) H(\mathbf{r}) (\mathbf{r})$, corresponds to $\hbar!$ (in the last two equations all quantities have their usual meaning).

If we multiply the canonical representation of the 2 × 2-matrix $\hat{K}(\mathbf{r})$,

$$\hat{K}(\mathbf{r}) = \begin{pmatrix} 0 & ; & h_2(\mathbf{r}) \\ h_1(\mathbf{r}) & ; & 0 \end{pmatrix}; \quad (13)$$

(where the 1- and 2-directions are perpendicular to $\sim^{(0)}(\mathbf{r})$ and extremal (see below)) by the imaginary number $\frac{1}{i}$, then equation (8) becomes similar to a Schrodinger equation with two-component vectors, (e.g., with spin-orbit interaction), although in our case the quantities $\frac{h_j(\mathbf{r})}{i}$ (for $j = 1;2$) are independent imaginary functions, not necessarily fulfilling the identity $(\frac{h_2(\mathbf{r})}{i}) = \frac{h_1(\mathbf{r})}{i}$ (i.e., hermiticity): only for $h_2 = h_1$ (circular precession) one would obtain $\frac{\hat{K}(\mathbf{r})}{i} = \hat{y} \cdot \hbar(\mathbf{r})$ (i.e., a 2 × 2-Schrodinger equation involving the (hermitian) Pauli matrix $\hat{y} = \begin{pmatrix} 0 & ; & i \\ +i & ; & 0 \end{pmatrix}$). Instead, for elliptic precession one obtains $\frac{\hat{K}(\mathbf{r})}{i} = \hat{y} \cdot \frac{h_2(\mathbf{r}) + h_1(\mathbf{r})}{2} + \hat{x} \cdot \frac{h_1(\mathbf{r}) - h_2(\mathbf{r})}{2} i$, with $\hat{x} = \begin{pmatrix} 0 & ; & 1 \\ 1 & ; & 0 \end{pmatrix}$.

Thus in the elliptic case, the matrix $\frac{\hat{K}(\mathbf{r})}{i}$ is no longer hermitian, but still pseudo-hermitian¹, in the sense of the just-mentioned identity; see [22]. The hermiticity would be lost from the beginning, if the Gilbert damping were included

4.3 Pseudo-orthogonality of the eigenmodes

Now we complexify the eigenmodes $\tilde{\sim}$ as follows: Starting from $\tilde{\sim}^{(0)}(\mathbf{r};t) = \frac{1}{2} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \cos(!t) + \frac{1}{2} \begin{pmatrix} 1 \\ -1 \end{pmatrix} \sin(!t)$ we write $\cos(!t) = \text{Re}(e^{-i!t})$ and $\sin(!t) = \text{Re}(e^{-i!t}i)$. In this way (by omitting the real part Re) one obtains the primed quantity $\tilde{\sim}^{(0)}(\mathbf{r};t) := \frac{1}{2} \begin{pmatrix} 1 \\ 1 \end{pmatrix} e^{-i!t} + \frac{1}{2} \begin{pmatrix} 1 \\ -1 \end{pmatrix} e^{-i!t}i$ (i.e., $/ e^{-i!t}$). By expansion with the complete set of eigenmodes this can be extended to general vectors $\tilde{\sim}(\mathbf{r};t)$. One also sees by direct inspection that an eigenvector $\tilde{\sim}^0 / e^{-i!t}$ of the equation $d\tilde{\sim}^0/dt = \hat{K} \tilde{\sim}^0$ with the eigenfrequency $!$ is obtained from the compatibility relation $\tilde{\sim}_2^0 = i\hbar_2^{-1} \tilde{\sim}_1^{1=2} \tilde{\sim}_1^0$.

In the Landau-Lifshitz formalism (denoted by the symbol LL) the pseudo-scalar product of two vectors $\tilde{\sim}$ can now be defined as follows (i.e., locally, and with the

¹ It can be shown that this is equivalent to the statement that the diagonalisation can only be performed with a non-trivial Bogoliubov-Valatin transformation, see [21].

primed quantities on the r.h.s.):

$$h^{(\prime)}(\mathbf{r};t)j^{(\prime)}(\mathbf{r};t)i_{LL} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}^0(\mathbf{r};t) h(\mathbf{r}) \begin{pmatrix} 1 \\ 0 \end{pmatrix}^0(\mathbf{r};t) \begin{pmatrix} 1 \\ 0 \end{pmatrix}^0(\mathbf{r};t) h(\mathbf{r}) \begin{pmatrix} 1 \\ 0 \end{pmatrix}^0(\mathbf{r};t); \quad (14)$$

where the $*$ -symbol denotes the hermitian-conjugate quantity.

In this way one gets $\sim / e^{+i\Omega t}$, and with (13)

$$h^{(\prime)}(\mathbf{r};t)j^{(\prime)}(\mathbf{r};t)i_{LL} = \sim^{(\prime)0}(\mathbf{r};t) \int K(\mathbf{r}) \sim^{(\prime)0}(\mathbf{r};t) : \quad (15)$$

This generalizes a relation of W. Fuller Brown, Jr., [18], which was restricted to homogeneous ground-states.

The main result of the present formalism is of course the following statement:

Two eigenmodes $\sim^{(\prime)}$ resp. $\sim^{(\prime)}$ with different eigenfrequencies, $\Omega^{(\prime)} \neq \Omega^{(\prime)}$, are everywhere pseudo-orthogonal.

E.g., the so-called radial and azimuthal eigenmodes, respectively, of [1] are everywhere pseudo-orthogonal, both in the outer region and also in the core region of a magnetic vortex line; although in the outer region $\sim = e$, $\sim_1 = z$, $\sim_2 = r$, whereas in the core region $\sim = e$, $\sim_1 = r$, $\sim_2 = z$.

This pseudo-orthogonality, which reflects the gyrotropic and elliptic motion of the spins, is also valid for any spatial average, see [23].

Similarly to the formalism of quantum mechanics, but with the additional freedom of locality, one should thus be able to estimate the eigenfrequencies from variational calculations, for example by exploiting equation (12) for points with suitable properties.

The proof of the above-mentioned main result is straightforward (here the superscript t means transposed, and it should be noted that the primed quantities are used):

$$\begin{aligned} & \sim^{(\prime)0}(\mathbf{r};t) \int K(\mathbf{r}) \sim^{(\prime)0}(\mathbf{r};t) d\mathbf{r} \sim^{(\prime)0}(\mathbf{r};t) \sim^{(\prime)0}(\mathbf{r};t) \\ & = \int (K^t(\mathbf{r}) \sim^{(\prime)0}(\mathbf{r})) \sim^{(\prime)0}(\mathbf{r}) = \left(\int K(\mathbf{r}) \sim^{(\prime)0}(\mathbf{r}) \right) \sim^{(\prime)0}(\mathbf{r}) \sim^{(\prime)0}(\mathbf{r}); \quad (16) \end{aligned}$$

ie., one minus sign is produced by the transposition, and a second one by the hermitian-conjugation. As a consequence, the states are pseudo-orthogonal if the eigenfrequencies differ.

Another essential point of the result is not the above-mentioned strange 'precessional' form of the pseudo-scalar product, but rather the fact that in contrast to quantum mechanics, which is nonlocal (see also [24]), here the local values of classical effective quantities count.

We can again apply the pseudo-orthogonality relations to vortex states. Here, sufficiently far from the vortex core, one has $\sim(\mathbf{r};t) / e$, as already mentioned. For $h_1(\mathbf{r})$ and $h_2(\mathbf{r})$ one should see the remarks below.

5 Further remarks on the relation to quantum mechanics

Some statements on the relation to quantum mechanics have already been presented. The following is more subtle and at the same time more basic: the Landau-Lifshitz

equations form a nonlinear set of classical integro-differential equations. The real quantities $\alpha_1(\mathbf{r};t)$ and $\alpha_2(\mathbf{r};t)$ are real (only if one wants to introduce the above-mentioned pseudo-orthogonality one is apparently forced to make the ansatz $\alpha_i \propto e^{-i\epsilon_i t}$) and in linear approximation they can be taken from a real Hilbert space. In contrast, in quantum mechanics, which is totally linear, these observables are represented by hermitian operators acting on the elements of a complex Hilbert space. Although the Landau-Lifshitz equations can be derived from a quantum mechanical model (e.g., the Heisenberg model) this involves some kind of molecular-field approximation (e.g., products of expectation values, $\langle \hat{A} \hat{B} \rangle$, instead of expectation values of products of operators, $\langle \hat{A} \hat{B} \rangle$). One should of course also consider metallic ferromagnets, which in quantum mechanics are better described by an itinerant model instead of the simpler Heisenberg one. But in any case both correspond to the same phenomenological Landau-Lifshitz theory. In the 'itinerant case' the atomistic derivation of the phenomenological theory is more delicate (for the exchange field the problems have been overcome by Korenman and coworkers, [25]). Therefore, although it was shown above that the equation of motion for the classical vector $\tilde{\alpha}(\mathbf{r};t)$ in linear approximation looks very similar to the Schrodinger equation, the correspondence to the Bogoliubov-Valatin approach (see [21]) is actually only true if one performs a quantum mechanical approach from which the Landau-Lifshitz equations can be derived.

A quantum mechanical model is also the basis of the following section.

6 Nonlinearity : mode-coupling, coalescence and splitting

Now the third-order processes, which come into play for higher oscillation amplitudes, are treated rather schematically in an approximation corresponding in quantum mechanics to the Bogoliubov-Valatin formalism : if \hat{b}_1^\dagger and \hat{b}_1 are the well-known creation and destruction operators for (Bogoliubov-Valatin) quasi-particle mode i , then in the next order the Hamilton operator contains a sum (or integral) of additional terms of the form $\hat{H} \approx \hbar \hat{b}_3 (\hat{b}_1^\dagger \hat{b}_2^\dagger) + \hbar_+ \hat{b}_3 (\hat{b}_1) \hat{b}_3^\dagger$, while \hbar and its hermitian-conjugate \hbar_+ are complex amplitudes of physical dimension 'energy'. (In the following it is not necessary to calculate these quantities from $\tilde{\alpha}(\mathbf{r};t)$, although just this is implicitly done in the computer simulations.)

According to Fermi's 'Golden Rule' the relaxation times for the corresponding processes derive from integrals involving the expression

$$\left(\frac{1}{\tau}\right) \approx \frac{2}{\hbar} \hbar^2 \delta(E - E_1 - E_2); \quad (17)$$

where splitting and coalescence processes correspond to the $-$ or $+$ cases, respectively. (In principle this is also not new, see [29].) Of course one has $\hbar_+ \hbar = \hbar \hbar_+$. However, the populations are different (see below).

In the above \hbar factor and the $(E_3 \dots)$ argument, respectively, one has both a spatial and also a temporal constraint corresponding to the mode form and frequency, symbolized by $\mathcal{V} \propto 1 + 2'$ ('splitting') and $\mathcal{V} \propto 1 + 2 \propto 3'$ ('coalescence'), with the energy corresponding to the frequency (through the usual relation $E = \hbar \omega$). These

contribute to the 'Stokes' and 'Antistokes' populations, respectively. At low temperatures the Stokes population ('splitting') is frozen, since excitations to be split do not exist. Therefore through the temperature dependence one can check the essentials of the mode-coupling theory.

In fact, in this way one can simultaneously understand parts of the above-mentioned phenomenological generalisation of the Landau-Lifshitz equation by Gilbert's damping term, [11]. Moreover, (only) if after integration the typical relaxation processes do not distinguish a particular axis, the Gilbert damping is isotropic per ansatz.

7 Application to vortex states (radial and azimuthal modes)

In the following we consider in more detail a flat circular disk with radius R of the order of $1 \mu\text{m}$ and thickness $t_h \ll R$. Outside a central region of radius $O(\epsilon) = O(\frac{J_s}{J_s^2})$ ($\epsilon = \frac{J_s}{J_s^2} \approx 5.7 \text{ \AA}$ for Permalloy) the ground state \sim_0 of the disc is the vortex state (i.e., $\sim_0(\mathbf{r}) = \hat{\mathbf{e}}$, where $\hat{\mathbf{e}}$ is the azimuthal unit vector). The precession around the vortex state is strongly elliptic, almost in-plane. The transverse fields entering the rule of the geometric mean are partially known: The out-of-plane component is $h_1 = \frac{J_s}{J_s^2} \approx 1 + O(\frac{t_h}{R})$ (if no external field is applied). In magnitude, this is a very large field ($\approx 1 \text{ T}$ for perm alloy) driving the spins backwards into in-plane direction, and practically independent of the mode index (but see [26]). This is already an exceptional case: to keep $\sim(\mathbf{r}; t)$ stationary (i.e., ! real), also h_2 must be negative.

Moreover, in contrast to the out-of-plane value h_1 , the in-plane component h_2 depends strongly on \mathbf{k} . If the spatial pattern of the mode corresponds to a plane wave with (in-plane) polarisation parallel to the magnetization, i.e., $\tilde{\mathbf{k}} = \sim_0(\mathbf{r})$ (or for an azimuthal excitation of a vortex configuration, $\tilde{\mathbf{k}} = \hat{\mathbf{e}}$) then h_2 almost vanishes, since there are no (effective) magnetic charges generated in the bulk (again: but see [26] concerning the so-called backward modes); whereas in case of an (in-plane) variation corresponding to a propagation $\tilde{\mathbf{k}} \perp \sim_0(\mathbf{r})$ (or for a radial excitation) the effective magnetic field $h_2(\mathbf{r})$ is maximal.

As already mentioned, the experiments of [3] show the existence of radial and azimuthal modes, respectively. In agreement with the above statements the eigenfrequencies of the azimuthal modes are relatively low. In fact, for the lowest modes they are less than half as high as that of the mode corresponding to the main radial peak. Thus, energetically, there can be coexistence processes (at low temperatures and at room temperature) and splitting processes (only at room temperature, where they are equally probable as the coexistence processes), involving for example a splitting of a radial main mode into two azimuthal modes. (Of course this tentative suggestion of the quenching of these processes at low temperatures has to be carefully checked experimentally).

8 Conclusion

In this communication some results (mostly not new, but nonetheless probably useful) for the dynamical properties contained in the Landau-Lifshitz equation for ferromag-

netic systems with general ground state have been demonstrated. It was shown that the linear approximation of these equations has strong similarities but also strong differences compared with the quantum mechanical Schrodinger equation. The essential difference is

- 1) perhaps not the above rule of the geometric mean,
- 2) also not the complications arising from the magnetic dipole interaction,
- 3) but the simple fact that quantum mechanics is nonlocal (e.g., the state function ψ comprises all values $\psi(\mathbf{r})$ throughout the space), whereas the Landau-Lifshitz theory is a classical local theory (i.e., the local values $\mathbf{H}_e(\mathbf{r})$ of the effective fields count, although the sources of these fields may be at a very large distance).

Generally one has got relations which are more complicated than in the quantum mechanical case. We mention (i) again the 'rule of the geometric mean' for the eigenfrequencies and (ii) a local pseudo-orthogonality of the eigenmodes, reflecting the gyrotropic and elliptic properties of the precession. Moreover, the Gilbert damping was partially traced back to microscopic three- (or more-) particle processes which correspond to confluence or splitting of elementary magnetic excitations as seen in recent experiments, [3].

Essential results depend on the local properties of the magnetization structure (for example, a domain structure). This can be at the same time both an advantage and also a disadvantage of the formalism; i.e., in addition to the known problems of calculating the magnetic dipole fields one sees at this place explicitly why these systems are so complex that one can hardly avoid a comparison with detailed numerical simulations.

Acknowledgements

The author acknowledges stimulating discussions and communications from C.H. Back, M. Buess, I. Neudecker and K. Perzlmaier.

References

- [1] M. Buess, Y. Acrem an, A. Kashuba, C.H. Back, and D. Pescia, J. Phys. CM 15 (2003) R1093
- [2] M. Buess, T.P.J. Knowles, R. Hollinger, T. Haug, U. Krei, D. Weiss, D. Pescia, M.R. Scheinfein, and C.H. Back, Phys. Rev. B 71 (2005) 104415; M. Buess, R. Hollinger, T. Haug, U. Krei, D. Weiss, D. Pescia, M.R. Scheinfein, C.H. Back, Phys. Rev. Lett. 93 (2004) 077207
- [3] M. Buess, T. Haug, M.R. Scheinfein, and C.H. Back, Phys. Rev. Lett. 94 (2005) 127205
- [4] K. Perzlmaier, M. Buess, and C.H. Back, V. Demidov and B. Hillebrands, S.O. Demokritov, Phys. Rev. Lett. 94 (2005) 057202

- [5] A . Puzic, B . Van W aye nberge, K ang-W eiChou, P . Fischer, G . Schutz, T . Ty liszczak, K . Rott, H . B ruckl, G . Reiss, I . Neudecker, T . Haug, M . Buess, C H . Back, J . Appl . Phys. 97 (2005) 10E 704
- [6] G . W oltersdorf, M . Buess, B . Heinrich, C H . Back, F . Nolting, S . Johnson, C . Buehler, Phys . Rev . Lett. 94 (2005) 217204
- [7] I . Neudecker, M . K laui, K . Perzlm aier, D . Backes, L . J . Heyderm an, C A F . Vaz, J A C . Bland, U . Rudiger, C H . Back, Phys . Rev . Lett. 96 (2006) 057207
- [8] I . Neudecker, K . Perzlm aier, F . Ho m ann, G . W oltersdorf, M . Buess, D . W eiss, C H . Back, Phys . Rev . B 73 (2006) 134426
- [9] O . Portm ann, M . Buess, A . V indigni, A . Vaterlaus, D . Pescia, C H . Back, Thin Solid Film s 505 (2006) 2
- [10] L D . Landau, E . Lifshitz, Phys . Z . Sow jetunin 8 (1935) 153
- [11] T L . G ilbert, Phys . Rev . 100 (1955) 1243
- [12] M . J . D onahue, O bject O riented M icrom agnetic Fram ework,
<http://math.nist.gov/oommf>
- [13] M R . Scheinfein, <http://llgmicrohome.mindspring.com>
- [14] J . Zinn-Justin, Q uantum eld theory and criticalphenom ena, O xford, C larendon Press 2003
- [15] Y . Nambu, G . Jona-Lasinio, Phys . Rev . 122 (1961) 345; 124 (1961) 246
- [16] W . Jantz, J C . Slonczewski, B E . A rgye, J . M agn . M agn . M ater. 23 (1981) 8
- [17] B . Van W aeyenberge, A . Puzic, H . Stoll, K W . Chou, T . Ty liszczak, R . Hertel, M . Fah nle, H . B ruckl, K . Rott, G . Reiss, I . Neudecker, D . W eiss, C H . Back, G . Schutz, Nature 444 (2006) 461
- [18] W . F . B rown, Jr., M icrom agnetics . John W iley, New York 1963
- [19] It was the merit of T L . G ilbert, see [11], to add to the effective eld a (nonherm i-tian but simple) contribution $a_G \frac{\partial \vec{m}}{\partial t}$, to include the damping in a phenom enological way instead of an equivalent elder approach of Landau and Lifshitz, which looked much more complicated. But since the 'G ilbert constant' a_G is very sm all, e.g., $a_G = 0.008$ for perm alloy, we neglect it unless otherw ise stated.
- [20] Recently the eigenm odes corresponding to the m ovements of the core of a m agnetic vortex structure have been studied in V . Novosad, F . Y . Fradin, P E . Roy, K . Buchanan, K . Yu . G uslienکو, D S . Bader, Phys . Rev . B 72 (2005) 24455 . H owever these eigenfrequencies are rather low , outside the range of m ost experim ents. See also K . Yu . G uslienکو . W . Scholz, R W . Chantrell, and V . Novosad, Phys . Rev . B 71 (2005) 144407 .

- [21] In principle the rule of the geometric mean should be well-known, but according to the experience of the author this is not always the case. The rule dates back to Kittel, [27], and to Holstein and Primako, [28]; it has also been stressed in the paper U.Krey, J.Magn.Mater. 268 (2004) 277, where it is mentioned that the rule is also obtained from the quasiparticle transformation of Bogoliubov and Valatin (the essential point is that the product $h_1 h_2$ of the present approach corresponds exactly to the Bogoliubov-Valatin result, if one writes $h_{1,2} = (A \ B)$, where A and B are the canonical expressions of the Bogoliubov-Valatin approach. But see the third-last section, 5, before the conclusions. (However, even the Bogoliubov-Valatin approximation should be attributed to an earlier reference, namely to Holstein and Primako, see [28]).
- [22] Here one should remember the fact that the matrices involved are applied to 2-vectors with components of the form $\psi_1(\mathbf{r}) e^{i\mathbf{k}\cdot\mathbf{r}}$ or $\psi_2(\mathbf{r}) e^{i\mathbf{k}\cdot\mathbf{r}}$ where however at the end the real part is taken, such that the i -factor in front of ψ_x in the text simply means that instead of the $\cos(\mathbf{k}\cdot\mathbf{r})$ -function one is working with $\sin(\mathbf{k}\cdot\mathbf{r})$.
- [23] I. Neudecker, G. Woltersdorf, B. Heinrich, T. Oikuno, G. Gubbiotti, C.H. Back, J. Magn. Mater. (2006), in press
- [24] The nonlocality of quantum mechanics is well-known and also seen in other quantities, e.g., the Aharonov-Bohm effect (Y. Aharonov, D. Bohm, Phys. Rev. 115 (1959) 485) and the Bell inequalities (J.S. Bell, Physics 1 (1964) 195).
- [25] V. Korenman, J.L. Murray, R.E. Prange, Phys. Rev. B 16 (1977) 4033, 4048, 4058
- [26] That for vortex configurations in flat circular disks of radius R the effective field h_1 is approximately $\frac{J_s}{2}$ and varies only weakly with \mathbf{r} can be seen from the ansatz $\psi_1 = e^{i\mathbf{k}\cdot\mathbf{r}} e^{i\mathbf{k}\cdot\mathbf{r}}$ (i.e., a wave propagating in the \mathbf{r} direction; here at the end the real part should be taken). Simultaneously the equilibrium magnetization shall be azimuthal, $\psi_0(\mathbf{r}) = e^{i\mathbf{k}\cdot\mathbf{r}}$ (i.e., $\mathbf{k}_m = \frac{m}{R} \mathbf{e}_z$ is parallel to the magnetization, while \mathbf{e}_z is perpendicular to the plane). In this case the potential of the magnetic field is not simply $\psi_m(\mathbf{r}; z) = e^{i\mathbf{k}\cdot\mathbf{r}} z e^{i\mathbf{k}\cdot\mathbf{r}}$, but something like $\psi_m(\mathbf{r}; z) = e^{i\mathbf{k}\cdot\mathbf{r}} \frac{\sinh(\mathbf{k}_m \cdot \mathbf{r})}{\mathbf{k}_m \cdot \mathbf{r}} \exp(i\mathbf{k}\cdot\mathbf{r})$, with $z \in (\frac{t_h}{2}; \frac{t_h}{2})$. As a consequence, also in this case the dipole fields depend on $y = R \cdot \mathbf{r}$ and z in such a way that one obtains so-called 'backward modes' (i.e., with negative dispersion; of course one should assume at this place that $\frac{J_s}{R} \ll 1$, where t_h is the thickness). Further details can be found in B. Hillebrands, K. Onodera (Eds.), Spin dynamics in confined magnetic structures I, Topics in Applied Physics 83, Berlin, Heidelberg, Springer 2001.
- [27] C. Kittel, Phys. Rev. 73 (1948) 155
- [28] T. Holstein, H. Primako, Phys. Rev. 58 (1949) 1098
- [29] C. Warren Haas, H. B. Callen, Ferromagnetic Relaxation and Resonance Line Width, in: Magnetism (G.T. Rado, H. Suhl, eds.), Vol. I, Academic Press, New York and London, 1963, p. 449