

Dilution Effects in Two-dimensional Quantum Orbital System

Takayoshi Tanaka and Sumio Ishihara

Department of Physics, Tohoku University, Sendai 980-8578, Japan

(Dated: March 23, 2022)

We study dilution effects in a Mott insulating state with quantum orbital degree of freedom, termed the two-dimensional orbital compass model. This is a quantum and two-dimensional version of the orbital model where the interactions along different bond directions cause frustration between different orbital configurations. A long-range correlation of a kind of orbital at each row or column, termed the directional order, is studied by means of the quantum Monte-Carlo method. It is shown that decrease of the ordering temperature due to dilution is much stronger than that in spin models. Quantum effect enhances the effective dimensionality in the system and makes the directional order robust against dilution. We discuss an essential mechanism of the dilute orbital systems.

PACS numbers: 71.10.-w, 71.23.-k, 75.30.-m

Recently, there is much effort to reveal impurity effects on correlated electron systems. One of the famous examples is the non-magnetic impurity doping into the low-dimensional quantum-magnets, such as the spin-Peierls systems and even-leg spin ladders, where the antiferromagnetic (AF) long-range order (LRO) is induced by breaking the quantum spin-singlet correlation.

Randomly dilute correlated electron systems with orbital degree of freedom is another class of disordered systems. Since this degree of freedom corresponds to anisotropy of the electronic clouds, the orbital interaction in a solid is essentially directional. That is to say, Hamiltonian explicitly depends on the bond direction unlike spin models [1, 2]. There are competition and frustration between the different-bond interactions which reduce the effective dimensionality in the system. Dilution effects in such a unique system have been studied experimentally in $\text{KCu}_{1-x}\text{Zn}_x\text{F}_3$ [3]. One of the doubly degenerate e_g orbitals in a Cu ion is occupied by a hole and exhibits LRO in KCuF_3 . With substitution of Cu by Zn, the orbital ordering temperature T_{OO} rapidly decreases and disappears around $x = 0.5$. This decrease of T_{OO} is much stronger than that in dilute magnets and in the percolation theory.

This experimental observation was well explained by the calculation based on the orbital Hamiltonian [5]:

$$\mathcal{H} = 2J \sum_{i,l} (\mathbf{T}_{\mathbf{r}_i} \cdot \hat{\mathbf{m}}^l) (\mathbf{T}_{\mathbf{r}_i+\hat{l}} \cdot \hat{\mathbf{m}}^l) \varepsilon_{\mathbf{r}_i} \varepsilon_{\mathbf{r}_i+\hat{l}}, \quad (1)$$

with a factor $\varepsilon_{\mathbf{r}_i}$ taking one for Cu and zero for Zn, and the bond direction l . Here $\mathbf{T}_{\mathbf{r}_i}$ is the pseudo-spin (PS) operator at site i , and $\hat{\mathbf{m}}^l$ is a unit vector in the $T^x - T^z$ plane. When we take $\hat{\mathbf{m}}^l = [\sin(\frac{2n_l}{3}\pi), \cos(\frac{2n_l}{3}\pi)]$ with $(n_x, n_y, n_z) = (2, 1, 0)$, this model represents the interaction between the e_g orbitals. Essence of the orbital dilution is frustration between the different-bond interactions and a macroscopic orbital degeneracy in the classical ground state due to the interactions.

In this paper, we study dilution effects in the two-dimensional and quantum version of Eq. (1) with $\hat{\mathbf{m}}^x =$

$[1, 0]$ and $\hat{\mathbf{m}}^z = [0, 1]$, termed the quantum orbital-compass model (OCM) [6, 7]. This is a minimal orbital model representing the directional and frustrated interaction between orbitals. One of the major different points from the e_g -orbital model is that the Hamiltonian is invariant under a certain local transformations of the PS operator. Due to this symmetry, degeneracy of the orbital state is more serious than that in the e_g -orbital model, and the conventional site-diagonal order is not expected. Instead, the long-range correlation for one of the PS components appears along the vertical or horizontal direction in a square lattice, termed the directional order (DO). Dilution effects in DO are investigated by means of the quantum Monte-Carlo (QMC) method in addition to some rigorous analytical considerations. We, in particular, focus on the quantum effects in the orbital dilution. It is shown that quantum fluctuation from the classical DO state enhances the effective dimensionality in the system and makes DO robust against dilution.

The explicit form of OCM is given in a two-dimensional square lattice as

$$\mathcal{H} = 2J \sum_{i,l=(x,z)} \mathbf{T}_{\mathbf{r}_i}^l \mathbf{T}_{\mathbf{r}_i+\hat{l}}^l \varepsilon_{\mathbf{r}_i} \varepsilon_{\mathbf{r}_i+\hat{l}}. \quad (2)$$

Since the sign of the interaction J is gauged away by rotating $\mathbf{T}_{\mathbf{r}_i}$ with respect to the y axis, we take it negative. This Hamiltonian without dilution ($\varepsilon_{\mathbf{r}_i} = 1$ for $\forall i$) is invariant under the following two-symmetry operations [7, 8]; (i) The global four-fold symmetry: the pseudo-spins at all sites and a crystal lattice are rotated by $\pi/2$, simultaneously, with respect to the y axis. (ii) The local symmetry at each column and row: the z (x)-component of all PSs at each column (row) along the z (x) axis are flipped, i.e. $T_{r_x, r_z}^{z(x)} \rightarrow -T_{r_x, r_z}^{z(x)}$ for each r_x (r_z). In addition, in the classical ferromagnetic-like ground state, there is the following continuous symmetry: (iii) the global PS direction is arbitrary in the $T^x - T^z$ plane.

Let us explain in more detail the symmetry (ii) which is unique in OCM. This symmetry operation at a row

r_z is done by the operator $P_{r_z} (\equiv \prod_{r_x} \sigma_{r_x, r_z}^z)$, with the Pauli matrices $\sigma_{\mathbf{r}}$, as $P_{r_z}^{-1} \sigma_{r_x, r_z}^x P_{r_z} = -\sigma_{r_x, r_z}^x$ and $P_{r_z}^{-1} \sigma_{r_x, r_z}^z P_{r_z} = \sigma_{r_x, r_z}^z$. On an equal footing, the symmetry operation at a column r_x is done by the operator $Q_{r_x} = \prod_{r_z} \sigma_{r_x, r_z}^x$. The operators P_{r_z} and Q_{r_x} commute with \mathcal{H} , and P_{r_z} s commute with each other as well as Q_{r_x} s, but $[Q_{r_x}, P_{r_z}] \neq 0$. The energy eigen-values are characterized by the eigen-value set of P_{r_z} s (p_1, \dots, p_L) or that of Q_{r_x} s (q_1, \dots, q_L) in a $L \times L (= N)$ square lattice. That is, at least, there are the L conserved quantities. Since $P_{r_i}^2 = 1$ and $P_{r_i}^\dagger = P_{r_i}$, $p_i = \pm 1$. It is easily shown that the conventional site-diagonal order parameter $N^{-1} \sum_{\mathbf{r}} \langle \mathbf{T}_{\mathbf{r}} \rangle$ is not invariant under this local-symmetry operation. According to the generalized Elitzur's theorem, it is shown that the site-diagonal orbital LRO does not occur [9]. Instead, a certain kind of LRO, termed the directional order, appears. This is a long-range correlation of T^z (T^x) at each column (row), and corresponds to a breaking of the global four-fold symmetry (i) explained above. This nematic-like order has been studied by means of the correlation function $C_{r_x}^{zz} \equiv \lim_{r_z \rightarrow \infty} \langle T_{r_x 0}^z T_{r_x r_z}^z \rangle$ at zero temperature [8] and $q = N^{-1} \sum_{\mathbf{r}} (T_{\mathbf{r}}^{z2} - T_{\mathbf{r}}^{x2})$ in the classical OCM [6]. We introduce, as an order parameter of DO, a more convenient form D defined by

$$D = N^{-1} (1 - x)^{-1} \times \sum_{\mathbf{r}} (T_{\mathbf{r}}^z T_{\mathbf{r}+\hat{\mathbf{z}}}^z \varepsilon_{\mathbf{r}+\hat{\mathbf{z}}} - T_{\mathbf{r}}^x T_{\mathbf{r}+\hat{\mathbf{x}}}^x \varepsilon_{\mathbf{r}+\hat{\mathbf{x}}}). \quad (3)$$

This parameter D indicates the correlation of T^z along the z axis or that of T^x along x . Unlike q , this is not a constant for the quantum PS operators. Since D commutes with P_{r_z} and Q_{r_x} , being different from $\mathbf{T}_{\mathbf{r}}$, it is possible for $\langle D \rangle$ to be finite.

To examine OCM numerically with and without vacancies, we adopt QMC method in a finite-size cluster. There is no negative-sign problem in OCM. The QMC calculations have been performed in a square lattice of $L \times L$ sites ($L=14 \sim 18$) with the periodic-boundary condition. We chose the Trotter number n to be $12 \sim 22$, and extrapolate the calculated results to $n = \infty$. At low temperatures of $T < 0.04J$, we perform the simulation with $n=60 \sim 72$. We take, at maximum, 10,000MC steps for measurement after 3,000MC steps for thermalization. Physical quantities are averaged over $20 \sim 60$ MC samples at each parameter set. We have checked by MC calculations with 10^6 MC steps, which is much longer than the integrated autocorrelation time determined by the binning analyses, that the order parameter values in the present calculation are reliable within numerical errors. The ordering temperature (T_{DO}) of DO is estimated from the finite-size scaling analyses of the Binder cumulant for D defined by $Q = 1 - \frac{\langle D^4 \rangle}{3 \langle D^2 \rangle^2}$ [6, 10]. At $T = T_{\text{DO}}$, a value of Q should be size independent.

First we show the numerical results in OCM without




FIG. 1: (color online). (a) Temperature dependence of the directional order parameter $\overline{D}(T)$ at $x = 0$ for various system sizes L . The inset shows $\overline{D}(T)$ at low temperatures obtained in the calculation with the large Trotter numbers (filled circles). Open circles are the same data with those in the main panel. (b) $\overline{D}(T)$ for various impurity concentrations.

vacancies. Temperature dependence of $\sqrt{\langle D^2 \rangle} (\equiv \overline{D}(T))$ for several L is presented in Fig. 1(a). Around $T/J = 0.17$, $\overline{D}(T)$ grows with decreasing T , and is saturated to about 0.13 below $T/J = 0.12$. With increasing L , this dependence becomes steep, but the saturated value of $\overline{D}(T)$ at low temperatures does not change much. Below $T/J = 0.07$, $\overline{D}(T)$ seems to decrease. We have performed careful numerical calculations with the larger Trotter numbers $n = 60 \sim 72$. As shown in the inset of Fig. 1(a), $\overline{D}(T)$ takes about 0.125 down to $T/J = 0.015$. Therefore, the reduction of $\overline{D}(T)$ in low temperatures is an artifact due to the small Trotter number. Temperature dependence of the Binder cumulant Q is presented in Fig. 2. The $Q - T$ curves for the several system size L cross around $T/J = 0.15$ corresponding to the temperature where $\overline{D}(T)$ grows. We have also calculated the susceptibility χ_D of D and the specific heat C . Both of them show peak structures around $T/J = 0.15$, which become steep with increasing L . We conclude in the present calculation that DO is realized in the quantum OCM and $T_{\text{DO}} = 0.150 \pm 0.003$. It is noted that the saturated value of \overline{D} at low temperatures is about a half of the classical value of $1/4$. This reduction is much stronger than that in the quantum AF Heisenberg model with $S = 1/2$ in a square lattice where the saturated staggered moment is about 60% of $1/2$ [11]. We also calculate the site-diagonal order parameter $\overline{M} \equiv \sqrt{\langle \mathbf{M}^2 \rangle}$ with $\mathbf{M} = N^{-1} \sum_{\mathbf{r}} \mathbf{T}_{\mathbf{r}}$. The calculated \overline{M} is almost temperature independent and its magnitude decreases with increasing L . This is, the site-diagonal order does not tend to occur because of the local symmetry (ii), mentioned previously. This situation is in contrast to the e_g -orbital model where the site-diagonal order is realized by the order-by-disorder mechanism.

Now we present the dilution effects in OCM. $\overline{D}(T)$ for




FIG. 2: (color online). Temperature dependence of the Binder cumulant Q at $x = 0$ for various system sizes L . Inset shows temperature dependence of Q at $x = 0.06$.

several x are shown in Fig. 1(b). $\overline{D}(T)$ gradually increases with decreasing T and is saturated, for example, to about 0.1 at $x = 0.06$. The temperature where $\overline{D}(T)$ rises up is reduced with doping. The saturated value of $\overline{D}(T)$ becomes small by doping in spite of a factor $(1-x)^{-1}$ in Eq. (3). This phenomenon was also observed in the dilute e_g -orbital model [5] and will be discussed in more detail later. The analyses for Q work well at $x = 0.06$, as shown in the inset of Fig. 2, where T_{DO}/J is identified to be 0.113 ± 0.005 . We have checked that, at $x = 0.25$, the $Q - T$ curves for several L do not cross with each other down to $T = 0.01J$.

The main results in the present work, the x dependence of T_{DO} in the quantum OCM, are shown in Fig. 3. For comparison, we plot the ordering temperature T_{CL} of DO in the classical OCM and the Curie temperature T_{Ising} in the $S = 1/2$ Ising model in a square lattice. T_{DO} decreases monotonically with x and seems to take zero around $x = 0.15 (\equiv x_c)$. It is worth to note that x_c is much smaller than the percolation threshold 0.41 in a square lattice [4]. The decrease of T_{DO} is much stronger than that in T_{Ising} , but is weaker than T_{CL} . In other words, quantum effect makes DO robust against dilution.

To understand the unusual dilution effects in the quantum OCM, we present a rigorous treatment of the dilute OCM in two-coupled chains along the z axis, that is, a two-leg ladder OCM. As mentioned previously, the energy eigen-states are classified by the eigen-value set (p_1, \dots, p_L) of the operator $P_{r_z} = \sigma_{r_L r_z}^z \sigma_{r_R r_z}^z$ where $r_R(r_L)$ indicates the x coordinates of the right (left) chain in the ladder. We consider one of the two-fold ground state with $p_1 = \dots = p_L = 1$ being degenerate with the state characterized by $p_1 = \dots = p_L = -1$. In a row r_z of the ladder, the eigen states are $|T_{r_L r_z}^z T_{r_R r_z}^z\rangle = |\uparrow\uparrow\rangle$ and $|\downarrow\downarrow\rangle$, or their linear combinations $|\pm\rangle \equiv \frac{1}{\sqrt{2}}(|\uparrow\uparrow\rangle \pm |\downarrow\downarrow\rangle)$. There are relations $(T_{r_L r_z}^x T_{r_R r_z}^x)|\pm\rangle = \pm \frac{1}{4}|\pm\rangle$,




FIG. 3: (color online). Impurity concentration x dependence of the ordering temperature. Closed circles are for T_{DO} in the quantum OCM, and closed squares are for T_{CL} in the classical OCM. For comparison, the ordering temperature in the Ising model (triangles) are also plotted. Bold arrow indicates the percolation threshold x_c in a square lattice. The inset shows the ordering temperature T_C in the quantum XY model (circles) and that in the classical one (squares) in a three-dimensional cubic lattice. System size is chosen to be $N = 23^3$ at maximum. The continuous imaginary-time algorithm is utilized.

$T_{r_L r_z}^z |\pm\rangle = \frac{1}{2} |\mp\rangle$ and $T_{r_R r_z}^z |\pm\rangle = \frac{1}{2} |\mp\rangle$. Therefore, within the subspace of $p_1 = \dots = p_L = 1$, we introduce new pseudo-spin operators \mathbf{S}_{r_z} and show correspondences $(T_{r_L r_z}^x T_{r_R r_z}^x) \rightarrow \frac{1}{2} S_{r_z}^z$ and $T_{r_L(r_R)r_z}^z \rightarrow S_{r_z}^x$. The state $|\uparrow\rangle$ ($|\downarrow\rangle$) is the up- (down-) eigen state of the operator $S_{r_z}^z$. As a result, OCM in this subspace is mapped onto the transverse-Ising model in a single chain [13]: $\mathcal{H}_{\text{eff}} = -4J \sum_{r_z} S_{r_z}^x S_{r_z+1}^x - J \sum_{r_z} S_{r_z}^z$. Now one vacancy is introduced at a site (r_L, r_z^0) . Since, in the row r_z^0 , we have $P_{r_z^0} = \sigma_{r_R r_z^0}^z$, the energy eigen states in this row, $|\uparrow\rangle$ and $|\downarrow\rangle$, belong to the different subspace, and $T_{r_R r_z^0}^z$ is replaced by a C -number in each subspace. Thus, OCM in the two-leg ladder with one vacancy is denoted by the following model:

$$\mathcal{H}_{\text{eff}} = -4J \sum_{r_z} S_{r_z}^x S_{r_z+1}^x - J \sum_{r_z \neq r_z^0} S_{r_z}^z \pm J(S_{r_z^0+1}^x + S_{r_z^0-1}^x), \quad (4)$$

where \sum'_{r_z} indicates a sum of r_z except for r_z^0 and $r_z^0 - 1$. Signs $+$ and $-$ in the third term correspond to the states $|\uparrow\rangle$ and $|\downarrow\rangle$ at the site (r_R, r_z^0) , respectively. This model is the two transverse-Ising chains where the external field $\pm J$ along the S^x axis is applied at edges of the chains, i.e. the $r_z^0 + 1$ and $r_z^0 - 1$ sites. In the original language, this field acts on $T_{r_R(r_L), r_z^0+1}^z$ and $T_{r_R(r_L), r_z^0-1}^z$, and enhances the correlation $\langle T_{\mathbf{r}_i}^z T_{\mathbf{r}_j}^z \rangle$ around the vacancy. The above analyses accord with an intuitive picture for roles of vacancy in OCM; due to the directional interaction in the model, orbitals near the vacant site ad-




FIG. 4: (color online). Contour map of local static correlation function $C^{zz}(\mathbf{r}_i)$ in a 20×20 site cluster. Four vacancies $X_{1(2)}$ and $Z_{1(2)}$ are represented by black circles. Temperature is chosen to be $T/T_{\text{DO}}(x=0) = 1.27$.

just their directions to gain the exchange energy. In other words, vacancy tends to fix the orbital direction around it.

Being based on the above picture, we discuss the difference between T_{DO} and T_{Ising} in Fig. 3. Consider a situation in a two-dimensional square lattice where two vacancies are located closely in a same row, e.g. at sites (r_x-1, r_z) and (r_x+1, r_z) . This probability is of the order of x . Since PS between the vacancies has the Ising-type interaction $T_{r_x r_z}^z (T_{r_x r_z+1}^z + T_{r_x r_z-1}^z)$, the correlation of T^z is enhanced around the vacancies, as suggested above. On an equal footing, the correlation of T^x is enhanced around the vacancies located closely in a same column. It is expected that a certain number of domains/patches, where T^z or T^x correlation is enhanced, coexist in a system. Competition between such kinds of domains reduces the order parameter $\overline{D}(T)$ and the ordering temperature T_{DO} . To confirm this picture, we calculate the local-static PS correlation function around the impurity defined by $C^{zz}(\mathbf{r}_i) = \frac{1}{N(1-x)} \sum_j \langle T_{\mathbf{r}_i}^z T_{\mathbf{r}_j}^z \rangle$ (Fig. 4). Temperature is chosen to be $T/T_{\text{DO}}(x=0) = 1.27$. Two pairs of vacancies introduced in a 20×20 site cluster are denoted by $X_{1(2)}$ and $Z_{1(2)}$ in the figure. It is clearly shown that $C^{zz}(\mathbf{r}_i)$ is enhanced between the impurities X_1 and X_2 , and almost vanishes between Z_1 and Z_2 . These results are not seen in dilute spin models.

Next we focus on quantum effects in dilution, i.e. discrepancy between T_{DO} and T_{CL} . In the classical DO state at $T = 0$, the system is considered to be decoupled into the independent Ising chains along the x or z axis, although DO is the two-dimensional order where the four-fold symmetry is broken. The rapid decrease of T_{CL} by dilution reflects the quasi-one dimensional nature of DO, and at finite temperatures, weak two-dimensionality is recovered due to the thermal fluctuation. The quantum fluctuation from the classical DO state brings about a coupling between the independent chains and increases

effective dimensionality of DO even at low temperatures. This is a kind of resonant states between the independent chains and DO becomes robust against dilution. This remarkable quantum effect is unique in the orbital system; we compare the x dependence of the Curie temperatures T_C in the quantum and classical XY models in a cubic lattice (the inset of Fig. 3). A magnitude of spin is chosen to be $1/2$ in the quantum model. It is seen that difference between the two are much smaller than that in OCM, and both T_C 's in the XY model seem to disappear at the percolation threshold ($x_c=0.69$).

Before summarizing the results, we briefly discuss implications of the dilute e_g -orbital model. In our previous studies [5], we have suggested that a remarkable decrease of T_{OO} is related to the tilting of PS around impurities. Roles of this PS tilting become clearer by the present analyses of OCM. The orbital interaction in the e_g -orbital model is essentially of the Ising-type along one direction in a cubic lattice, and there is frustration between them, as well as OCM. Introduced vacancies break the local symmetry and release the frustration. According to vacancy configurations, a specific type of the orbital correlation is stabilized around vacancies. For example, when two vacancies are located closely at a same row along the z axis, the orbital correlation of $d_{3x^2-r^2}/d_{y^2-z^2}$ or $d_{3y^2-r^2}/d_{z^2-x^2}$ are stabilized around them. Competition between these domains/patches with different PS correlations reduce the order parameter and may cause the glass-like phase.

In summary, we study the dilution effects in a quantum and two-dimensional version of the orbital model, i.e. the orbital compass model, where the vertical or horizontal correlation of a PS component, termed the directional order, is expected from the local symmetry. Reduction of the ordering temperature of DO is much stronger than that in the spin model. Quantum fluctuation increases the effective dimensionality of the system and makes DO robust against dilution. Competition between domains with different orbital configurations is essence of the dilute orbital system.

The authors thank to M. Matsumoto for several helpful discussion and providing calculation data for the inset in Fig. 3, and H. Matsueda for valuable comments. This work was supported by Grant-in-Aid for Scientific Research on priority Areas, JSPS KAKENHI (16340094, 16104005) and TOKUTEI "High Field Spin Science in 100T" (No.451, 18044001) from MEXT, NAREGI, and CREST. T. T. appreciates financial support from JSPS.

-
- [1] S. Maekawa, *et al.*, *Physics of Transition Metal Oxides* (Springer Verlag, Berlin, 2004), and references therein.
 - [2] K. I. Kugel and D. I. Khomskii, *Sov. Phys. JETP* **37**, 725 (1973).

- [3] N. Tatami, S. Niioka, and Y. Murakami, (to be published).
- [4] D. Stauffer, Phys. Rep. **54**, 1 (1979).
- [5] T. Tanaka, *et al.*, Phys. Rev. Lett. **95**, 267204 (2005).
- [6] A. Mishra, *et al.*, Phys. Rev. Lett. **93**, 207201 (2004).
- [7] Z. Nussinov, *et al.*, Euro. Phys. Lett. **67**, 990 (2004).
- [8] J. Dorier, *et al.*, Phys. Rev. B **72**, 024448 (2005).
- [9] C.D. Batista and Z. Nussinov, Phys. Rev. B. **72**, 045137 (2005).
- [10] K. Binder, Phys. Rev. Lett. **47**, 693 (1981).
- [11] E. Manousakis, Rev. Mod. Phys. **63**, 1 (1991).
- [12] R. B. Stinchcombe, J. Phys. C **13**, 5565 (1980).
- [13] B. Doucot, *et al.*, Phys. Rev. B **71**, 024505 (2005).