

# Escape driven by $\alpha$ -stable white noises

B. Dybiec\* and E. Gudowska-Nowak†

*M. Smoluchowski Institute of Physics, and Mark Kac Center for Complex Systems Research,  
Jagellonian University, ul. Reymonta 4, 30-059 Kraków, Poland*

P. Hänggi‡

*Institute of Physics, University of Augsburg,  
86135 Augsburg, Germany*

and

*National University of Singapore,  
Faculty of Science, Physics Department, Blk S12,  
2 Science Drive 3, 117542 Singapore, Singapore*

(Dated: March 23, 2022)

We explore the archetype problem of an escape dynamics occurring in a symmetric double well potential when the Brownian particle is driven by *white Lévy noise* in a dynamical regime where inertial effects can safely be neglected. The behavior of escaping trajectories from one well to another is investigated by pointing to the special character that underpins the noise-induced discontinuity which is caused by the generalized Brownian paths that jump beyond the barrier location without actually hitting it. This fact implies that the boundary conditions for the mean first passage time (MFPT) are no longer determined by the well-known local boundary conditions that characterize the case with normal diffusion. By numerically implementing properly the set up boundary conditions, we investigate the survival probability and the average escape time as a function of the corresponding Lévy white noise parameters. Depending on the value of the skewness  $\beta$  of the Lévy noise, the escape can either become enhanced or suppressed: a negative asymmetry  $\beta$  causes typically a decrease for the escape rate while the rate itself depicts a non-monotonic behavior as a function of the stability index  $\alpha$  which characterizes the jump length distribution of Lévy noise, with a marked discontinuity occurring at  $\alpha = 1$ . We find that the typical factor of “two” that characterizes for normal diffusion the ratio between the MFPT for well-bottom-to-well-bottom and well-bottom-to-barrier-top no longer holds true. For sufficiently high barriers the survival probabilities assume an exponential behavior. Distinct non-exponential deviations occur, however, for low barrier heights.

PACS numbers: 05.10.Gg, 02.50.-r, 02.50.Ey, 82.20.Uv, 82.20.Wt

## I. INTRODUCTION

The noise driven escape from a deterministically metastable state is a theme that impacts many phenomena in diverse fields of natural sciences [1, 2, 3, 4]. In particular, as a notable model for a chemical reaction Kramers [1] pioneered the problem of an escape of a Brownian particle of mass  $m$  moving in a potential  $V(x)$  with local minima corresponding to an initial reactant and a final product state. In this scenario, both states are assumed to be separated by a barrier located at a position  $x_b$ . In the spatial-diffusion-limited regime, the Kramers rate theory is based on a stochastic dynamics that does not involve inertial effects and thus is described by an overdamped Langevin dynamics, reading

$$\frac{dx}{dt} = v = -\frac{1}{\eta}V'(x) + \sqrt{\frac{k_B T}{\eta}}\tilde{\zeta}(t). \quad (1)$$

Here,  $\tilde{\zeta}(t)$  constitutes a white Gaussian noise process with correlations  $\langle \tilde{\zeta}(t)\tilde{\zeta}(s) \rangle = 2\delta(t-s)$ , representing thermal fluctuations whose intensity is scaled by the friction  $\eta$ . The escape problem then concerns the surmounting of an energetic barrier for stochastic trajectories that predominantly dwell the neighborhood of separating attractors which in this case are made of two neighboring potential wells. The imposed quasi-stationarity condition is realized by assuming special boundary conditions with respect to the time evolution equation for the probability density. In the overdamped regime, the evolution equation for the probability density  $p(x, t)$  follows the Smoluchowski dynamics

$$\frac{\partial p(x, t)}{\partial t} = -\frac{\partial}{\partial x}J(x, t), \quad (2)$$

where

$$J(x, t) = -\frac{1}{\eta}V'(x)p(x, t) - \frac{k_B T}{\eta} \frac{\partial p(x, t)}{\partial x}, \quad (3)$$

and the stationarity approximation describes escape events which correspond to a constant, non-vanishing flux of probability  $J_s = J(x)$ . Those stochastic realizations of the process that have surpassed the barrier top are immediately absorbed and re-inserted into the original attractor region. In this way a steady probability

\*Electronic address: bartek@th.if.uj.edu.pl

†Electronic address: gudowska@th.if.uj.edu.pl

‡Electronic address: hanggi@physik.uni-augsburg.de

flow across the activated barrier state located between the locally stable states of “reactants” and “products” is established. Escaped trajectories, absorbed at a position larger than the barrier location  $x_a > x_b$  require that  $p_s(x_a) = 0$  on the whole half-line  $x_a > x_b$ . The rate formulation is then based on the “flux over the population” method [2, 5], yielding in this case of a Smoluchowski dynamics the celebrated result [1, 2, 3, 4]:

$$k = \frac{\omega_b \omega_w}{\eta \, 2\pi} \exp \left[ -\frac{\Delta V}{k_B T} \right], \quad (4)$$

with  $\Delta V$  denoting the activation energy  $\Delta V = V(x_b) - V(x_w)$  and  $\omega_w, \omega_b$  representing the frequencies of the particle’s motion near the metastable potential minimum  $x_w$  and at the top of the crossed barrier, respectively. This result represents a feasible estimate for the actual reaction rate, if all trajectories ejected by the source properly thermalize before eventual thermal fluctuations drive them out of the initial well and, most importantly, a distinct time-scale separation between escape dynamics and intra-well relaxation holds true. This latter requirement of a clear-cut time-scale separation is at the basis for the description of the escape dynamics in terms of a (time-independent) rate coefficient [2, 5].

Yet another alternative to the approaches discussed above is rooted in the concept of the mean first passage time (MFPT), i.e. the average time that a random walker starting out from a point  $x_0$  inside the initial domain of attraction, assumes in order to leave the attracting domain for the first time [2, 3, 4]. Put differently, the MFPT is the average time needed to cross the deterministic separatrix-manifold for the first time [2, 6, 7]. At weak noise the MFPT becomes essentially independent of the starting point, i.e.  $t(x_0) \approx t_{\text{MFPT}}$  for all starting configurations away from the immediate neighborhood of the separatrix. Given the fact that the crossing of the separatrix in either direction equals for *normal diffusion* one half, the total escape time  $t_e$  equals  $2t_{\text{MFPT}}$  and thus the rate of escape  $k$  itself becomes in this case:

$$k = \frac{1}{2t_{\text{MFPT}}}. \quad (5)$$

Characterization of the escape rate by use of the MFPT is a rather complex notion for a general class of stochastic processes. In particular, the MFPT analysis requires the choice of a correct boundary condition [6, 7]. These are well known for one-dimensional stochastic diffusion Markov processes  $x(t)$  which are of the Fokker-Planck form, Eqs. (2-3) or for one dimensional master equations with birth and death kinetics. With generally non-Gaussian white noise the knowledge of the boundary location alone typically cannot specify in full the corre-

sponding boundary conditions for, say, absorption or reflection, respectively [6, 8]. In particular, the trajectories driven by non-Gaussian white noise depict discontinuous jumps. As a consequence, the location of the boundary itself is not hit by the majority of discontinuous sample trajectories. This implies that regimes beyond the location of the boundaries must be properly accounted for when setting up the boundary conditions. Most importantly, returns (i.e. so termed re-crossings of the boundary location) from excursions beyond the specified state space back into this very finite interval where the process proceeds must be excluded.

Following our reasoning in discussing Lévy-Brownian motion on finite intervals [8], we present in this paper the analysis of escape events of a non-inertial, generalized diffusion process which is driven by Lévy noise dwelling a symmetric double-well potential.

## II. WHITE LÉVY NOISE

The standard definition of the Gaussian white noise specifies the latter as a time derivative of the Wiener process (i.e. a derivative of a stationary process with independent and Gaussian-distributed increments whose covariance is given by  $\langle W(t)W(s) \rangle = \min(t, s)$ ). A classical Brownian motion (the Wiener process) can be therefore represented as a limit in distribution of independent Gaussian jumps taken at infinitesimally short time intervals of non-random length  $1/n$ . Alternatively (following e.g. definition of Feller [9, 10, 11]), the Wiener process can be thought of as a limiting process of random Gaussian jumps at random Poissonian jump times:

$$\begin{aligned} W(t) &= \lim_{n \rightarrow \infty} \sum_{k=1}^{[tn]} W_k \left( \frac{1}{n} \right) = \lim_{n \rightarrow \infty} W \left( \frac{N(nt)}{n} \right) \\ &= \lim_{n \rightarrow \infty} \sum_{k=1}^{N(tn)} W_k, \end{aligned} \quad (6)$$

The symbol  $[tn]$  stands for an integer number of jumps which for a Poisson counting process  $N(nt), t \geq 0$  with them mean  $\langle N(nt) \rangle = nt$  are assumed to be independent (decoupled) of *i.i.d* random variables  $W_k$  sampled from the Gaussian distribution. The equality sign in Eq. (6) denotes a limit in distribution sense. For  $n \rightarrow \infty$ , the Poisson distribution becomes peaked around  $k = nt$  and the limiting process  $W(t), t \geq 0$  tends to a Brownian motion diffusion for which the cumulative distribution function reads [12]

$$\begin{aligned}
\text{Prob}\{W(t) \leq w\} &= \lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} \text{Prob}\left\{W\left(\frac{k}{n} \leq w\right)\right\} \text{Prob}\{N(nt) = k\} \\
&= \lim_{n \rightarrow \infty} \int_{-\infty}^w \sum_{k=0}^{\infty} (2\pi k/n)^{-1/2} \exp(-nx^2/2k) \frac{(nt)^k}{k!} \exp(-nt) dx = \int_{-\infty}^w \frac{e^{-x^2/2t}}{\sqrt{2\pi t}} dx. \quad (7)
\end{aligned}$$

Here, we introduce a non-Gaussian white noise as a derivative of the generalized Wiener process  $W_{\alpha,\beta}(t)$ , i.e. a non-Gaussian random process with stationary and independent increments. The increments of such generalized Wiener process have the  $\alpha$ -stable distribution with the stability index  $\alpha$  and the time increment  $\Delta t^{1/\alpha}$  as a scale parameter:

$$\begin{aligned}
W_{\alpha,\beta}(t) &= \int_0^t \zeta(s) ds = \int_0^t dL_{\alpha,\beta}(s) \\
&\approx \sum_{i=0}^{N-1} (\Delta s)^{1/\alpha} \zeta_i, \quad (8)
\end{aligned}$$

where  $\zeta_i$  are independent random variables distributed with the stable, Lévy probability density function (PDF)  $L_{\alpha,\beta}(\zeta; \sigma, \mu = 0)$  and  $N\Delta s = t - t_0$ . The parameter  $\alpha$  denotes the stability index, yielding the asymptotic power law for the jump length distribution being proportional to  $|\zeta|^{-1-\alpha}$ . The parameter  $\sigma$  characterizes a scale,  $\beta$  defines an asymmetry (skewness) of the distribution, whereas  $\mu$  denotes the location parameter. Throughout the paper, we deal only with strictly stable distributions not exhibiting a drift regime; this implies a vanishing location parameter  $\mu = 0$  in the remaining part of this work. For  $\alpha \neq 1$ , the characteristic function  $\phi(k) = \int_{-\infty}^{\infty} d\zeta e^{-ik\zeta} L_{\alpha,\beta}(\zeta; \sigma, \mu)$  of an  $\alpha$ -stable random variable  $\zeta$  can be represented by

$$\phi(k) = \exp\left[-\sigma^\alpha |k|^\alpha \left(1 - i\beta \text{sign}(k) \tan \frac{\pi\alpha}{2}\right)\right], \quad (9)$$

while for  $\alpha = 1$  this expression reads

$$\phi(k) = \exp\left[-\sigma |k| \left(1 + i\beta \frac{2}{\pi} \text{sign}(k) \ln |k|\right)\right]. \quad (10)$$

The three remaining parameters vary within the allowed regimes  $\alpha \in (0, 2]$ ,  $\beta \in [-1, 1]$ ,  $\sigma \in (0, \infty)$ . Random variables  $\zeta$  corresponding to the characteristic functions (9) and (10) can be generated using the Janicki–Weron algorithm [13, 14].

For  $\alpha = 2$  (and an arbitrary skewness parameter  $\beta$ , cf. Eq. (9)), the generalized Lévy-Brownian motion  $W_{\alpha,\beta}(t)$  becomes a standard Gaussian Wiener process whose time derivative leads to the well known Gaussian white noise limit.

Following this interpretation, the stochastic source term in Eq. (1) can be represented as a sum of independent pulses (having stable distribution) acting on equally

spaced times. As a consequence, such Lévy noise is white, i.e. its autocorrelation function is formally a Dirac-delta function of time.

For the sake of clarification, we mention here that there exist different representations of other Lévy noises that are non-Markovian in nature. As an example, so called fractional Gaussian (or Lévy) noise is sometimes defined in literature as the time derivative of a fractional Brownian motion process [15, 16, 17, 18, 19]. In contrast to the Gaussian (or our Lévy) white noise, the fractional Gaussian noise (fractional Lévy stable motion) may exhibit slowly decaying time-correlations. This is however not the case with the type of Lévy noise addressed in our paper, where the noise source at the level of the Langevin equation is a *white noise process*.

### III. ESCAPE IN A DOUBLE WELL: SURVIVAL PROBABILITY AND MEAN FIRST PASSAGE TIME

Let us consider a Brownian, overdamped particle in an external potential  $V(x)$ , i.e.,

$$V(x) = -\frac{a}{2}x^2 + \frac{b}{4}x^4. \quad (11)$$

which is driven by Lévy stable white noise. The sample trajectories are then obtained by a direct integration of Eq. (1):

$$x(t) = x_0 - \int_{t_0}^t V'(x(s)) ds + \int_{t_0}^t dL_{\alpha,\beta}(s), \quad (12)$$

using the standard techniques of integration of stochastic differential equation with respect to the Lévy stable PDFs [8, 13, 14, 20, 21, 22, 23]. The first passage time problem has been analyzed as  $\tau = \inf\{t \geq 0 \mid x(t) \geq x_b\}$  with trajectories  $x(t)$  starting at  $x = x_0$  and subject to corresponding boundary conditions. In particular, an absorbing boundary condition is realized by stopping the trajectory whenever it reaches the boundary, or, more typically, whenever it has jumped beyond that very boundary location. The role of reflection, which in the case of a free diffusion [8] has been assured by wrapping the hitting (or crossing) trajectory around the boundary location, while preserving its assigned length, see in Ref. [8, 23], is taken over naturally here by the confining potential walls of the symmetric double well. The details of

our employed numerical scheme for stochastic differential equations driven by Lévy white noise has been detailed elsewhere [8]. In the following we shall omit cases when  $\alpha = 1$  with  $\beta \neq 0$ . In fact, this parameter set is known to induce instabilities in the numerical evaluation of corresponding trajectories [13, 14, 20, 21, 23].

In the following we consider several differing situations. For a particle that starts out at a well bottom  $x_0 = -x_w$ , cf. Fig. 1 and makes excursions toward the neighboring well bottom, taken as an absorbing boundary we use the subscript notation  $w-w$ . Likewise, for a particle starting out at well bottom and being absorbed at barrier top location we use the notation  $w-b$ . Using the statistics of the first passage time events we shall next study the two corresponding survival probabilities

$$S_{w-b}(t) = 1 - \mathcal{F}_{w-b}(t) \quad (13)$$

$$S_{w-w}(t) = 1 - \mathcal{F}_{w-w}(t), \quad (14)$$

where  $\mathcal{F}_{\dots}(t)$  is the corresponding cumulative first passage time distribution, i.e.,

$$\mathcal{F}_{w-b, w-w}(t) = \int_{-\infty}^{x_b, x_w} p_{w-b, w-w}(x, t) dx, \quad (15)$$

with  $p_{w-b, w-w}(t)$  being the corresponding first passage time density.

In order for the escape to become dominated by a clear-cut, single time-scale we use a sufficiently high potential barrier, thereby enforcing rare escape events. A too low barrier would involve too many recrossing events with the escape then being ruled by many time scales. Our used parameters for the potential are the parameter set  $a, b$  with  $a = 128, b = 512$ , yielding a barrier height of  $\Delta V = V(0) - V(-\sqrt{a/b}) = \frac{a^2}{4b} = 8$ . This symmetric potential well is schematically depicted with Fig. 1.




FIG. 1: The generic double well potential  $V(x) = -\frac{a}{2}x^2 + \frac{b}{4}x^4$  for inspection of the Kramers problem discussed in the paper. To assure sufficiently high barrier separating the stationary states  $-x_w, x_w$ , the following set of parameters has been chosen:  $a = 128, b = 512, \Delta V = V(0) - V(-\sqrt{a/b}) = 8$ .

## A. Case of normal diffusion

For  $\alpha = 2$ , the process driven by white Lévy stable noise approaches the Brownian limit. The  $\alpha$ -stable white Lévy noise, as used in this study, then leads to the standard Gaussian white noise with intensity  $2\sigma^2$ , i.e.  $\langle \zeta(t)\zeta(s) \rangle_{\alpha=2} = 2\sigma^2\delta(t-s)$ . In this case, the corresponding values of the MFPT from the potential minimum  $-x_w$  to the top of the absorbing potential barrier, i.e.  $x_b$  or to the neighboring, absorbing minimum,  $x = x_w$ , can be calculated from the following quadrature formulas

$$\begin{aligned} \text{MFPT}(-x_w \rightarrow x) &= \frac{1}{\sigma^2} \int_{-x_w}^x \exp[V(z)/\sigma^2] \\ &\times \int_{-\infty}^z \exp[-V(y)/\sigma^2] dy dz. \end{aligned} \quad (16)$$

Note that for a system driven by white Gaussian noise the energy difference  $\Delta V$  measured in units of  $k_B T$  may be directly related to the intensity of the noise  $\sigma^2$ . In contrast, for Lévy stable noises with  $\alpha < 2$ , the scale parameter  $\sigma$  is no longer “thermodynamically” related to the system temperature and consequently becomes a free parameter of the model. We have set throughout this study this value to  $\sigma = \sqrt{2}$ .

## B. Survival probabilities for $\alpha$ -stable noise driven escape

Typical sample trajectories of the stochastic process defined by Eq. (1) are depicted in Fig. 2. We observe that a non-zero  $\beta$  parameter induces a dynamical asymmetry for the escape dynamics occurring in a *symmetric* double well potential. It is reflected in our numerical simulations by the emergence of stochastic trajectories that spend more time in the vicinity of one of the potential minima. As a consequence, one of the attracting states of the process ( $\pm x_w$ ) becomes favored over the other. This kind of behavior can be also detected in our other figures. Notably, for a decreasing value of the stability index  $\alpha$  we observe larger fluctuations of the particle positions. These occasional long jumps of trajectories may be of the order of, or even larger than the distance  $2x_w$  separating the two minima of the symmetric potential  $V(x)$ .

From the ensemble of single trajectories, which are subject to the boundary conditions discussed above, and presented in Fig. 2, we estimated the survival probability densities  $S_{w-b}(t)$  and  $S_{w-w}(t)$ , cf. see Fig. 3. The behavior of the survival probability is consistent with the results as predicted by inspection of the corresponding stochastic trajectories: We clearly detect from Fig. 3 that at a chosen value of  $\alpha$  a decrease of the skewness  $\beta$ -parameter of a driving white noise causes a distinct decrease the rate of escape of the particle from the left




FIG. 2: Sample trajectories of random walks in the generic double well potential subjected to Lévy stable noise with  $\alpha = 1.9$  (left panel) and  $\alpha = 0.9$  (right panel) with various  $\beta$  ( $\beta = 0$ , top panel,  $\beta = -1.0$ , bottom panel). The scale parameter is  $\sigma = \sqrt{2}$ , the chosen time step of integration is  $dt = 10^{-4}$ .

potential minimum; thus stabilizing the starting position at  $x = -x_w$ . Most importantly, at sufficient high barrier heights a visible exponential decay of the survival probability occurs, – according to  $S(t) = \exp(-t/T_{MFPT})$ . This is the familiar behavior for regular Brownian motion and has been detected already previously for symmetric Lévy noises [24, 25] and for totally skewed (with  $\beta = 1$ ) one sided Lévy motions [26]. Here we observe it as well for *skewed stable noises*. Notably, the characteristic exponent  $T_{MFPT}$  clearly depends on both noise parameters, i.e. on the value of the stability index  $\alpha$  and also on the skewness parameter  $\beta$ .

Our numerical analysis indicates that typically the survival probability  $S_{w-w}(t)$  for the “well-bottom-to-well-bottom” setup exceeds the survival probability  $S_{w-b}(t)$  for the “well-bottom-to-barrier-top” setup. This observation can be explained by features of a noise-driven dynamics: A particle performing the motion under the influence of noise needs more time to dwell the neighboring potential minimum than is needed to reach the top of the barrier. This kind of behavior can be nevertheless weakened by diminishing the stability parameter  $\alpha$ . For  $\alpha < 2$  the trajectory of the random particle becomes discontinuous and at sufficiently small  $\alpha$  a particle, on average, completes its escape from the left potential minimum by a long jump which can overpass the right potential minimum. Consequently, both survival curves start to overlap converging to the same function at a given small stability index  $\alpha$ .

For the Cauchy limit, i.e.,  $\alpha = 1$ , an approximate result for the mean crossing time as a function of the noise strength  $\sigma$  has been given in Ref. [24] by use of the fractional Fokker-Planck equation. In order to compare results obtained in this work with the former studies presented by Chechkin *et al.* [24], we have used a re-scaled form of Eq. (1)

$$\frac{dx}{dt} = (x - x^3) + \zeta(t), \quad (17)$$




FIG. 3: Survival probability densities  $S_{w-b}(t)$  (left panel) and  $S_{w-w}(t)$  (right panel) for a particle dwelling the generic double well potential subjected to Lévy stable noise with  $\alpha = 1.9$  (top panel) and  $\alpha = 0.9$  (bottom panel). The simulation parameters are:  $dt = 10^{-5}$ ,  $N = 2 \times 10^4$ ,  $\sigma = \sqrt{2}$ .




FIG. 4: Survival probability densities  $S_{w-b}(t)$  and  $S_{w-w}(t)$  for a particle wandering in the generic double well potential subjected to Cauchy noise, i.e. Lévy stable noise with  $\alpha = 1.0$ .  $MFPT_{w-b} = 1.038 \pm 0.008$  and  $MFPT_{w-w} = 1.086 \pm 0.008$ . The simulation parameters are:  $dt = 10^{-5}$ ,  $N = 2 \times 10^4$ ,  $\sigma = \sqrt{2}$ .

obtained by a set of transformations  $x \rightarrow x/x_w$ ,  $t \rightarrow t/\tau$  with  $x_w^2 = a/b$ ,  $\tau = \eta/a$  and consequently  $\sigma^\alpha = D \rightarrow$

$\sqrt{\frac{k_B T}{\eta}} \tau / x_w^\alpha$ . In these units  $\sigma \rightarrow \frac{2\sigma}{128^{1/\alpha}}$ , thus,  $\sigma_{\alpha=2} = \frac{1}{4}$  and  $\sigma_{\alpha=1} = \frac{\sqrt{2}}{64}$ . From the approximate formula derived in [24] for a strictly weak noise case at  $\alpha = 1$  one obtains  $\text{MFPT} \approx 141.71$ , whereas our simulations for this noise strength yield  $\text{MFPT} \approx 132.86$ , which differs only by ca. 7%. Exemplary survival probabilities  $S_{w-b}(t)$  and  $S_{w-w}(t)$  for this Cauchy-Lévy noise are presented in Fig. 4. The data indicate that for the Cauchy noise the MFPT for well-bottom-to-barrier-top scenario is somewhat smaller than the MFPT for the case of well-bottom-to-well-bottom, cf. Fig. 4.

Our prior studies of a *free* Lévy-Brownian motion driven by  $\alpha$  stable noise has shown that for some parameterizations of the Lévy noise (totally skewed stable distributions with  $\alpha < 1$ ) a non-exponential survival probability density emerges. Therefore, in this study we tested whether for a particle driven by a Lévy-Smirnoff noise, i.e. with setting  $\alpha = 0.5$ ,  $\beta = 1$ , a decrease of the potential barrier may induce visible deviations from the exponential distribution. The results of the test are displayed in Fig. 5 where the survival probability densities  $S_{w-b}(t)$  and  $S_{w-w}(t)$  for various heights of the potential barrier  $\Delta V$  are presented. For decreasing  $\Delta V$  the motion of a particle indeed approaches the behavior of a free diffusion; consequently also distinct deviations from the exponential character of survival distributions do show up.




FIG. 5: Survival probabilities  $S_{w-b}(t)$  (left panel) and  $S_{w-w}(t)$  (right panel) for various barrier heights separating two wells of the potential. The system is driven by Lévy-Smirnoff noise, i.e. Lévy stable noise with  $\alpha = 0.5$  and  $\beta = 1$ . The simulation parameters are:  $dt = 10^{-5}$ ,  $N = 2 \times 10^4$ ,  $\sigma = \sqrt{2}$ .

### C. Behavior for the MFPT

From the ensembles of collected first passage times we have also evaluated directly the mean values of the distributions. Our findings for MFPTs are presented in Figs. 6–7. The depicted results corroborate with the decrease of the escape rate (i.e. the inverse of the MFPT)

upon increasing the skewness  $\beta$ . Moreover, for the stability index  $\alpha = 2$  (normal Brownian white noise) and for any skewness parameter  $\beta$ , the Gaussian case is properly re-confirmed, cf. Figs. 6–7.




FIG. 6: Mean first passage times for well-bottom-to-barrier-top  $\text{MFPT}_{w-b}$  (left panel) and mean first passage times for well-bottom-to-well-bottom  $\text{MFPT}_{w-w}$  (right panel) as a function of the skewness parameter  $\beta$ . Simulation parameters  $dt = 10^{-5}$ ,  $N = 2 \times 10^4$ ,  $\sigma = \sqrt{2}$ . The error bars estimated using bootstrap method with  $N_b = 2 \times 10^3$ . The white Lévy stable noise with the stability index  $\alpha = 2$  and any allowed value of  $\beta$  is equivalent to the white Gaussian noise, what is manifested by the independence of  $\text{MFPT}(\beta)$  for  $\alpha = 2$ . The various symbols represent the various values of the stability index  $\alpha$ : ‘+’  $\alpha = 2.0$ , ‘x’  $\alpha = 1.9$ , ‘o’  $\alpha = 1.5$ , ‘□’  $\alpha = 1.3$ , and ‘■’  $\alpha = 1.1$ .




FIG. 7: Mean first passage times for well-bottom-to-barrier-top  $\text{MFPT}_{w-b}$  (left panel) and mean first passage times for well-bottom-to-well-bottom  $\text{MFPT}_{w-w}$  (right panel) as a function of the stability index  $\alpha$ . Simulation parameters  $dt = 10^{-5}$ ,  $N = 2 \times 10^4$ ,  $\sigma = \sqrt{2}$  and  $N_b = 2 \times 10^3$ . The black squares indicate equality of MFPTs for  $\alpha = 2$  with any  $\beta$ . The various symbols represent the various values of the skewness parameter  $\beta$ : ‘+’  $\beta = 1.0$ , ‘x’  $\beta = 0.5$ , ‘o’  $\beta = 0.0$ , and ‘□’  $\beta = -0.5$ .

### D. MFPT-ratio no longer obeying normal behavior

Next, we investigate the behavior of the value for the ratio  $R = \text{MFPT}_{w-w}/\text{MFPT}_{w-b}$  of the  $\text{MFPT}_{w-w}$  between the case with well-to well and well-to-bottom. Note that from the exact expression in Eq. (16), for a symmetric potential  $V(x)$  with normal Gaussian white noise fluctuations (i.e. for  $\alpha = 2$  and an arbitrary value of the skewness parameter  $\beta$ ) this ratio yields the commonly known factor of  $R = 2$ . Put differently, in this case the random walker needs twice the time to reach the other potential minimum than to reach to the top of the potential barrier. In contrast, with a decreasing value of  $\alpha$  the ratio  $R$  now *distinctly* deviates from 2, being always smaller than for a Gaussian diffusion case. Our numerical results are depicted in Fig. 8. This deviation can be understood by noticing that for  $\alpha < 2$ , the stochastic escape trajectories of the random walks in the double well potential become discontinuous, meaning that the continuous movement of a particle becomes interrupted with long jumps. These occasional jumps are more probable to occur for small stability index  $\alpha$  and explain the observable statistics of the first passage times, implying in turn that  $R < 2$ . In particular, an increasing probability of long jumps over the barrier in the overall statistics of passages tends to equalize  $\text{MFPT}_{w-w}$  with  $\text{MFPT}_{w-b}$ , yielding  $R$  values approaching 1. The effect of the skewness parameter  $\beta$  which is responsible for the asymmetric stochastic dynamics can be interpreted as an additional contribution to the directionality of random impact pulses which push the Lévy-Brownian particle within the potential well. For negative (positive)  $\beta \approx \pm 1$  with  $\alpha < 1$ , the driving Lévy white noise becomes a one-sided Lévy process with strictly negative (positive) increments which stabilize trajectories around the starting position  $x = \pm x_w$ .




FIG. 8: Ratio  $R = \text{MFPT}_{w-w}/\text{MFPT}_{w-b}$  of MFPTs to the neighboring potential well ( $\text{MFPT}_{w-w}$ ) and to the barrier top ( $\text{MFPT}_{w-b}$ ). The simulation parameters are:  $dt = 10^{-5}$ ,  $N = 2 \times 10^4$ ,  $\sigma = \sqrt{2}$  and  $N_b = 2 \times 10^3$ .

## IV. CONCLUSIONS

In this paper we have studied the survival probabilities and the mean values of the mean first passage times for escape from a symmetric double well potential when the overdamped dynamics is driven by general Lévy white noise. The statistics of escaping trajectories is investigated by a numerical analysis of a Lévy noise driven Langevin equation with properly implemented boundary conditions. In contrast to former studies [24, 25] aimed to understand statistics of barrier crossings with symmetric Lévy flights or exploring the first passage problem for one-sided Lévy motions [26], our work provides the first analysis of the Kramers problem with an arbitrary set of  $\alpha, \beta$ -parameters which characterize the white stable noise-source entering the Langevin dynamics. Following the Markovian character of the stochastic dynamics, – at sufficiently high barriers –, the time dependence of the survival probabilities within the potential well assume an exponential law. As can be expected, distinct deviations from the exponential behavior become detectable, however, for low barrier heights when, similarly to a normal Gaussian diffusion case, many recrossing of the barrier are possible.

An asymmetry of the Lévy noise perturbing the dynamics is shown to either enhance or also suppress the escape events and corroborates the intuitive influence of the  $\beta$  parameter which yields an additional, biasing contribution to the particles' motion due to the directionality of stochastic impact pulses. Moreover, the rate of escape is shown to exhibit a non-monotonic behavior as a function of the stability index  $\alpha$ , with a discontinuity occurring at the value  $\alpha = 1$ .

In clear contrast to a normal diffusion behavior as typified by systems driven by white Gaussian noise, for which the random walker requires twice the time to reach the other potential minimum as compared to the mean time to reach to the (absorbing) top of the barrier, Lévy white noise with  $\alpha < 2$  now causes a decrease in this ratio of the mean first passage times: The ratio  $R = \text{MFPT}_{w-w}/\text{MFPT}_{w-b}$  lies consistently below the “normal” value of  $R = 2$ . This remarkable result holds true for any chosen value of the skewness parameter  $\beta$ .

In summary, our numerical considerations demonstrate the richness of the bi-stable kinetics resulting from driven  $\alpha$ -stable white noises. The observed features are rooted in the non-local jump lengths taken from a distribution which exhibit fat tails which in turn rule the random passages between the attracting states of a double well potential.

## Acknowledgments

The Authors acknowledge the financial support from the Polish State Committee for Scientific Research (KBN) through the grants 1P03B06626 (2004–2005) and 2P03B08225 (2003–2006), Marie Curie TOK COCOS

grant (6th EU Framework Programme under contract: MTKD-CT-2004-517186) and the ESF funds (E.G.N. and P.H.) via the STOCHDYN program. Additionally, B.D. acknowledges the financial support from the Foundation for Polish Science through the domestic

grant for young scientists (2005). Computer simulations have been performed at the Academic Computer Center CYFRONET AGH, Kraków. P.H. also acknowledges the support by the German research foundation (DFG) via grant Ha 1517/26-1, 2 (“Single molecules”).

- 
- [1] H. A. Kramers, *Physica (Utrecht)* **7**, 284 (1940).
  - [2] P. Hänggi, P. Talkner and M. Borkovec, *Rev. Mod. Phys.* **62**, 251 (1990).
  - [3] P. Talkner and P. Hänggi, *New Trends in Kramers Reaction Rate Theory*, (Kluwer Academic Publ., Boston, 1995); G. R. Fleming and P. Hänggi, *Activated Barrier Crossing*, (World Scientific, Singapore, 1993); P. Hänggi and J. Troe, *Ber. Bunsen Phys. Chem.* **95**, 225 (1991).
  - [4] P. Hänggi and P. Jung, *Adv. Chem. Phys.* **89**, 239, (1995).
  - [5] P. Reimann, G. J. Schmid and P. Hänggi, *Phys. Rev. E* **60**, R1 (1999).
  - [6] C. Van den Broeck and P. Hänggi, *Phys. Rev. A* **30**, 2730 (1984); P. Hänggi, F. Marchesoni and P. Grigolini, *Z. Phys. B* **56**, 333 (1984); P. Hänggi and P. Riseborough, *Phys. Rev. A* **27**, 3379 (1983); V. Balakrishnan, C. Van den Broeck and P. Hänggi, *Phys. Rev. A* **38**, 4213 (1988).
  - [7] P. Müller, P. Talkner and P. Reimann, *Physica A* **247**, 338, (1997); P. Hänggi and P. Talkner, *Phys. Rev. A* **32**, 1934 (1985);
  - [8] B. Dybiec, E. Gudowska-Nowak and P. Hänggi, *Phys. Rev. E* **73**, 046104 (2006).
  - [9] W. Feller, *An introduction to Probability Theory and its Applications* (John Wiley, New York, 1968).
  - [10] V. M. Zolotarev, *One Dimensional Stable Distributions* (American Mathematical Society, Providence, 1986).
  - [11] B. V. Gnedenko and A. N. Kolmogorov *Limit Distributions for Sums of Independent Random Variables*, (Adison-Wesley, Reading, MA, 1954).
  - [12] The result follows a general observation for separable continuous time random walks (CTRW) which lead to normal stochastic diffusion for finite mean waiting time for a jump to occur and finite step variance [10].
  - [13] R. Weron, *Statist. Prob. Lett.* **28**, 165 (1996).
  - [14] A. Janicki and A. Weron, *Statistical Science* **9**, 109 (1994); see also in: *Simulation and Chaotic Behavior of  $\alpha$ -Stable Stochastic Processes* (Marcel Dekker, New York, 1994); A. Janicki, *Numerical and Statistical Approximation of Stochastic Differential Equations with Non-Gaussian Measures*, (Hugo Steinhaus Centre for Stochastic Methods, Wrocław, 1996).
  - [15] B. B. Mandelbrot, *SIAM Rev.*, **10**, 422 (1968).
  - [16] S. Chaudhury and B. J. Cheravil, *J. Chem. Phys.* **125**, 024904 (2006).
  - [17] M. Shlesinger, G. M. Zaslavsky and U. Frisch, Eds., *Lévy Flights and Related Topics in Physics* (Springer Verlag, Berlin, 1994).
  - [18] I. M. Sokolov, J. Klafter, *Chaos*, **15**, 026103 (2005).
  - [19] S. C. Kou and X. Sunney Xie, *Phys. Rev. Lett.* **93**, 180603 (2004).
  - [20] P. D. Ditlevsen, *Phys. Rev. E* **60**, 172 (1999).
  - [21] A. Weron and R. Weron, *Lecture Notes in Physics* **457**, 379 (1995).
  - [22] M. E. J. Newman and G.T. Barkema, *Monte Carlo Methods in Statistical Physics* (Clarendon Press, Oxford, 1999).
  - [23] B. Dybiec and E. Gudowska-Nowak, *Phys. Rev. E* **69**, 016105 (2004).
  - [24] A. V. Chechkin, V. Yu. Gonchar, J. Klafter and R. Metzler, *Europhys. Lett.* **72**, 348 (2005).
  - [25] P. Imkeller and I. Pavlyukevich, *J. Phys. A: Math. Gen.* **39**, L237 (2006).
  - [26] I. Eliazar and J. Klafter, *Physica A* **336**, 219 (2004).