

Non-perturbative Approach to Critical Dynamics

Leonie Canet

Service de Physique de l'Etat Condense, CEA Saclay, 91191 Gif-sur-Yvette
Cedex, France

Laboratoire de Physique et Modélisation des Milieux Condenses, CNRS, 25
avenue des Martyrs, BP 166 - 38042 Grenoble Cedex, France

Hugues Chaté

Service de Physique de l'Etat Condense, CEA Saclay, 91191 Gif-sur-Yvette
Cedex, France

Abstract. This paper is devoted to a non-perturbative renormalization group (NPRG) analysis of Model A, which stands as a paradigm for the study of critical dynamics. The NPRG formalism has appeared as a valuable theoretical tool to investigate non-equilibrium critical phenomena, yet the simplest and nontrivial models for critical dynamics have never been studied using NPRG techniques. In this paper we focus on Model A taking this opportunity to provide a pedagogical introduction to NPRG methods for dynamical problems in statistical physics. The dynamical exponent z is computed in $d = 3$ and $d = 2$ and is found in close agreement with results from other methods.

1. Introduction

The understanding of non-equilibrium critical phenomena stands as one of the major challenges of statistical physics. Systems far from thermalequilibrium are omnipresent in nature: slow relaxation or external driving forces tend to prevent real systems from ever reaching their equilibrium distribution. Behavior out-of-equilibrium is far richer than at equilibrium, and many intriguing scaling phenomena, such as self-organized criticality (emergence of scaling without re-tuning of a control parameter) [1], or phase transitions between non-equilibrium stationary states [2, 3], have been observed for long. However, despite the considerable achievements of equilibrium statistical physics, the theoretical comprehension of non-equilibrium critical phenomena remains much poorer. The renormalization group (RG), which has appeared as a cornerstone to explain universality in equilibrium continuous phase transitions, has also allowed some breakthroughs out-of-equilibrium [4]. Nonetheless, many non-equilibrium phenomena remain out of range of perturbative approaches because of large coupling constants or because the interesting dimensions lie far from the critical one. Further theoretical progress out-of-equilibrium is hindered by the lack of analytical tools to handle the corresponding models.

Recently, a novel approach (namely the non-perturbative renormalization group (NPRG)) has been proposed to investigate (non-equilibrium) reaction-diffusion processes [5]. It has allowed to overcome the perturbative limitations and to gain physical insights into models such as branching and annihilating random walks [

which are reviewed in [6]. For instance, these studies have captured non-perturbative effects that essentially determine the phase diagram of some systems [7, 8, 5, 9] and have unveiled a genuinely non-perturbative fixed point governing the phase transition belonging to the so-called parity-conserving universality class [10]. A valuable advantage of this method is that it gives a unified description of a model: the very same equations enable one to probe any dimensions or coupling regimes, including non-universal features. Hence, the NPRG appears as a powerful tool to investigate non-equilibrium systems.

However most readers are still largely unfamiliar with these techniques, though they have been introduced more than a decade ago for systems in equilibrium [11, 12]. The aim of this contribution is to give a brief but pedagogical introduction to the NPRG methods for non-equilibrium systems. For this purpose, Model A stands as one of the simplest (yet far from being trivial) dynamical models and it has never been studied within the NPRG framework so far, which we remedy with the present work. We adopt a practical viewpoint and put a particular emphasis on the discussion of the device of a (non-perturbative) approximation scheme.

The paper is organized as follows. In Section 2 we briefly recall the definition of Model A and its critical properties. In Section 3, the principles and the construction of the NPRG are outlined for generic non-equilibrium systems and specialized in Section 4 to Model A. The corresponding non-perturbative flow equations are derived in Section 5 and their numerical integration is dealt with in Section 6. The results are eventually discussed in Section 7 and followed by a brief summary in Section 8.

2. Model A and critical dynamics

The purely dissipative relaxation of a non-conserved field can be described by the Langevin equation:

$$\partial_t \phi(\mathbf{x};t) = -D \frac{\delta H[\phi]}{\delta \phi(\mathbf{x};t)} + \xi(\mathbf{x};t) \quad (1)$$

where D denotes a constant and uniform relaxation rate and H the usual Landau-Ginzburg-Wilson Hamiltonian. On approaching a critical point, the relaxation time of the order parameter starts diverging, which reflects the critical slowing down of the dynamics. The Langevin equation is a ‘mesoscopic’ description of the system which exploits the associated decoupling of time scales: the order parameter, represented by the (coarse-grained) field ϕ , relaxes much slower than all the other microscopic degrees of freedom, which can hence be modeled by a stochastic Gaussian noise with zero mean and variance

$$\langle \xi(\mathbf{x};t) \xi(\mathbf{x}^0;t^0) \rangle = 2D k_B T^{-d} \delta(\mathbf{x} - \mathbf{x}^0) \delta(t - t^0) \quad (2)$$

The strength of the noise is fixed by the Einstein relation which ensures that the system acquires its equilibrium distribution at long time. We here focus on the case of a scalar order parameter with Ising symmetry, described by the Hamiltonian

$$H[\phi] = \int d^d x \left[\frac{1}{2} [\nabla \phi(\mathbf{x})]^2 + U(\phi) - h(\mathbf{x}) \phi(\mathbf{x}) \right]; \quad U(\phi) = \frac{r}{2} \phi^2 + \frac{u}{4!} \phi^4 \quad (3)$$

The Langevin equation (1) corresponds to a Glauber dynamics for the Ising spin and defines Model A in the classification by Halperin and Hohenberg [13]. Besides

the equilibrium critical exponents and ν , the critical dynamics is characterized by the dynamical exponent z that describes the divergence of the relaxation time $\tau \sim |T - T_c|^{-z}$ near the temperature T_c of the second-order phase transition.

Time Reversal Symmetry

In the long-time limit after the initial perturbation, the system is expected to become time translational invariant (TTI) and the time-reversal symmetry (TRS) to hold. TRS then yields the fluctuation-dissipation theorem (FDT) which linearly relates the two-point correlation function $C(x, x^0; t, t^0) = \langle x(t)x(t^0) \rangle$ with the response function $R(x, x^0; t, t^0) = \langle \delta x(t) / \delta x^0(t^0) \rangle$ following:

$$R(x, x^0; t, t^0) = -\frac{1}{T} \partial_{t^0} C(x, x^0; t, t^0). \quad (4)$$

In the early stages of the relaxation process, the system is generally not TTI such that R and C may depend on both t and t^0 and FDT may not hold. The fluctuation-dissipation ratio $TR = \partial_{t^0} C$ becomes of particular interest to characterize the violation of FDT and the associated ageing phenomena (see [14] for a recent review and references therein). For our non-perturbative study, we rather focus on the stationary dynamics where the system satisfies TRS.

Field Theory

Any RG treatment starts out from a field theory. Upon introducing a Martin-Siggia-Rose response field $\tilde{x}(x; t)$ [15], one can average over the Langevin noise and cast the stochastic equation (1) into a dynamic functional [16, 17]:

$$S[\tilde{x}; x] = \int_0^t dt \int d^d x \left[\tilde{x} \partial_t x + D \frac{H[\tilde{x}, x]}{D} \right] \quad (5)$$

(where $k_B T$ has been set to unity). Correlation and response functions can then be expressed as functional averages with the weight $\exp(-S[\tilde{x}; x])$.

In this equivalent field theoretical formulation, TRS can be conveniently expressed as an invariance of the action (5) under a specific field transformation, as stressed in [18]. This transformation writes

$$\begin{aligned} \tilde{x} &\rightarrow \tilde{x} + \frac{1}{D} \partial_t x \\ x &\rightarrow x \end{aligned} \quad (6)$$

Indeed, one can straightforwardly check that after performing a time inversion $t \rightarrow -t$ in (5) which switches the sign of the kinetic term $\tilde{x} \partial_t x$, the field transformation (6) yields additional contributions from the latter term and from the noise term D that cancel out. Besides, the transformation of the Hamiltonian part under (6) produces an additional term $\int \tilde{x} \partial_t x$, which vanishes upon time integration in the stationary regime. We shall rely in the following on this simple expression of TRS to ensure that this invariance is preserved within the non-perturbative formulation.

3. The NPRG formalism in non-equilibrium statistical physics

The NPRG formalism relies on Wilson's RG idea [19], which consists in building a sequence of scale-dependent effective Hamiltonians, that interpolate smoothly between

the short-distance physics at the (microscopic) scale $k = \infty$ and the long-distance physics at the scale $k = 0$, through progressively averaging over fluctuations. Rather than expressing Γ_k as in the original Wilsonian formulation [1] the flow of effective Hamiltonians for the slow modes, one can work out the flow of effective Gibbs free energies βF_k for the rapid ones, following [20, 21]. βF_k thus only includes fluctuation modes with momenta $|\mathbf{q}| < k$. At the scale $k = \infty$, no fluctuation is yet taken into account and βF_∞ coincides with the microscopic action S [20], while at $k = 0$, all fluctuations are integrated out and βF_0 is the analogue of the Gibbs free energy at thermalequilibrium, in that it encompasses the long-distance and long-time properties of the system. Hence, to construct βF_k , one needs at a given scale k to suppress the slow modes with momentum $|\mathbf{q}| < k$. This is achieved by adding to the original action (5) a scale-dependent term [21, 5, 22] which is quadratic in the fields (so as to affect the propagator of the corresponding modes):

$$S_k[\phi; \tilde{\phi}] = \frac{1}{2} \int_{\mathbf{x}, t} \int_{\mathbf{x}', t'} [\phi(\mathbf{x}; t); \tilde{\phi}(\mathbf{x}; t)] \hat{R}_k(\mathbf{x} - \mathbf{x}'; t - t') [\phi(\mathbf{x}'; t'); \tilde{\phi}(\mathbf{x}'; t')] \quad (7)$$

where \hat{R}_k is a symmetric 2×2 matrix of elements R_k^{ij} ($i, j = 1, 2$). These elements (so-called cutoff functions) will be specified in the following, but their general properties can be already stressed. In order to achieve the renormalization procedure outlined above, these cutoff functions must behave at fixed k as $R_k^{ij} \propto k^2$ (in Fourier space) for small momenta $|\mathbf{q}| < k$ so that the slow fluctuation modes acquire a mass $\propto k^2$ and decouple. On the other hand, R_k^{ij} must vanish for large momenta $|\mathbf{q}| > k$ so that the rapid modes remain unaltered and contribute to the functional averages with weight $\exp(-S_k)$. Besides, the additional constraints

$$\lim_{k \rightarrow 0} R_k^{ij} = 0; \quad \lim_{k \rightarrow \infty} R_k^{ij} = 1 \quad \text{at fixed } \mathbf{q} \quad (8)$$

must be satisfied in order to enforce the correct asymptotic behaviors at the scales $k = \infty$ and $k = 0$, respectively $\beta F_\infty = S$ and $\beta F_0 = \beta F$ [21, 5, 22].

With the additional term (7) the partition functions'

$$Z_k[\mathbf{j}; \tilde{\mathbf{j}}] = \int \mathcal{D}\phi \mathcal{D}\tilde{\phi} \exp(-S - S_k + \int \mathbf{j} \phi + \int \tilde{\mathbf{j}} \tilde{\phi}) \quad (9)$$

become k -dependent. Finally, the effective βF_k which is the central object of the NPRG procedure is defined as the (modified) Legendre transform of $\log Z_k[\mathbf{j}; \tilde{\mathbf{j}}]$:

$$\beta F_k[\phi; \tilde{\phi}] + \log Z_k[\mathbf{j}; \tilde{\mathbf{j}}] = \int \mathbf{j} \phi + \int \tilde{\mathbf{j}} \tilde{\phi} - S_k[\phi; \tilde{\phi}] \quad (10)$$

βF_k is a functional of the conjugate fields $\phi = \log Z_k = \mathbf{j}$ and $\tilde{\phi} = \log Z_k = \tilde{\mathbf{j}}$. The additional term $-S_k$ in Eq. (10) is necessary to set the proper microscopic behavior at $k = \infty$: $\beta F_\infty = S$ [21]. The RG flow of βF_k under an infinitesimal change of the scale $k \rightarrow k + s$ or rather $s = \log(k' / k)$ is governed by an exact functional differential equation [21, 22] (which is derived in the Appendix):

$$\partial_s \beta F_k = \frac{1}{2} \text{Tr}_{\mathbf{q}, t} \left[\partial_s \hat{R}_k - \hat{R}_k^{(2)} + \hat{R}_k^{-1} \right] \quad (11)$$

In this equation, $\hat{R}_k^{(2)}[\phi; \tilde{\phi}]$ is the 2×2 matrix of second derivatives of βF_k with respect to (wrt) ϕ and $\tilde{\phi}$ and $[\hat{R}_k^{(2)} + \hat{R}_k]^{-1}$ hence embodies the full (functional) propagator associated with the effective theory $S_k + S_k$.

Obviously, Eq. (11) cannot be solved exactly and one has to resort to some approximations [21]. However, as the approximations used do not rely on the smallness of a parameter (see next section), the approach remains non-perturbative in essence. In particular, it is not confined to weak-coupling regimes or to the vicinity of critical dimensions and is therefore suitable to overcome the limitations of perturbative RG schemes.

4. NPRG for Model A

To exploit the exact flow equation (11), one has to devise an approximation scheme. This scheme is based on the construction of an Ansatz for χ_k which does not spoil the non-perturbative features of the exact equation (11) and which can be systematically enlarged. The formulation of this Ansatz relies on the physics one wishes to probe, that is some basic physical insights are necessary. The most common truncation consists in expanding χ_k in powers of gradients [20] and time derivatives. The accuracy and convergence of this approximation scheme have been thoroughly studied in the equilibrium context and have shown that quantitatively reliable results can already be obtained at the leading order (r^2) [21]. For instance, for the three-dimensional Ising model, NPRG calculations yield for the critical exponents $\nu = 0.628$ and $\eta = 0.0443$ at order r^2 [21, 23], and $\nu = 0.632$ and $\eta = 0.033$ at order r^4 [23] which are in close agreement with the 6-loop results $\nu = 0.6304(13)$ and $\eta = 0.0335(25)$ [24]. Another useful approximation scheme is the ϵ -field expansion of χ_k . This truncation has the advantage of preserving the momentum structure of higher order vertices but it approximates the functional structure of the effective potential [21]. The derivative expansion is best appropriate for the study of critical physics which is conveyed by the large-distance ($q \rightarrow 0$) and long-time ($\omega \rightarrow 0$) modes. We hence adopt here this approximation scheme and expand χ_k at leading order in derivatives (11) i.e. only terms of order r^2 and ∂_t are retained.

Construction of an Ansatz for χ_k at leading order

The form of the Ansatz for χ_k is dictated by the symmetries. Since we consider the long-time regime where TRS holds, we want the Ansatz to be invariant under the ϵ -field transformation (6) (where D is set to unity), which in turn imposes the following structure:

$$\chi_k(\vec{r}; \omega) = \int d^d x dt \sum_n \tilde{X}_k^a(\vec{r}; \omega) \partial_t^n X_k^b(\vec{r}; \omega) X_k^c(\vec{r}; \omega) \omega^{n-2} : \quad (12)$$

No higher powers of the response ϵ -field are allowed at order ∂_t due to TRS. Indeed, the transformation (6) would connect a generic 'noise' term $\tilde{X}_k^a X_k^b$ ($n > 2$) to higher order kinetic terms $\tilde{X}_k^a (\partial_t)^n X_k^b$, $j = 0 :: n$ which are discarded at order ∂_t . Further constraints on the X_k^i 's, $i = a; b; c$ can be deduced from TRS in the same way as in Section 2. First one must have $X_k^c = X_k^a - X_k^b$ for the additional contributions generated by the transformation (6) of the 'noise' and the 'kinetic' terms to cancel out. As for the linear term in \tilde{X}_k^b , it should write as a (ϵ -field) derivative of a functional $X_k^b(\vec{r}; \omega) = F_k$ for its transform under (6) to vanish upon time integration. We naturally adopt for F_k the usual equilibrium Ansatz at order r^2 for the Ising model

which has been widely studied in the past [21, 23]

$$F_k[\varphi] = \int d^d x \left[\frac{1}{2} Z_k(\varphi) [\nabla \varphi]^2 + U_k(\varphi) \right] \quad (13)$$

In this Ansatz, the functional U_k embodies the effective potential and the renormalization function Z_k encompasses the anomalous dimension of the field (see below).

Furthermore, at the scale $k = 0$, one must identify with the microscopic (bare) action (5) up to the response field rescaling $\tilde{\varphi} = X_k \varphi$ i.e. one has:

$$X_k = \frac{1}{D}; \quad Z_k = 1; \quad U_k = U(\varphi) \quad (U(\varphi) \text{ is defined in Eq. (3)}): \quad (14)$$

The Ansatz for Model A at leading order naturally writes

$$F_k(\varphi; \tilde{\varphi}) = \int_{x,t} \tilde{\varphi} X_k \partial_t \varphi - \int d^d x \left[\frac{1}{2} Z_k(\varphi) (\nabla \varphi)^2 + \tilde{\varphi} U_k(\varphi) X_k \right] \quad (15)$$

This Ansatz constitutes the basis of our work.

Definition of the critical exponents

We now discuss how the critical exponents can be computed within the NPRG approach. In the critical regime, Z_k is expected to endow a scaling form $Z_k \sim k^{-z}$ and accordingly to behave as $k^{(d-2+z)/2}$. The critical exponent hence corresponds to $z = -\partial_s \ln Z_k$ at the critical point [21]. Similarly, X_k is expected to scale as $X_k \sim k^{-x}$ at the critical point such that k^{2-z+x} according to a scaling analysis of Eq. (15). Hence, the dynamical exponent z , which by definition characterizes the divergence of the time scale following $\tau \sim k^{-z}$, is given by $z = 2 - z + x$ where $x = -\partial_s \ln X_k$.

Notice that $Z_k(\varphi)$ is a functional of φ whereas the scaling form Z_k should be a mere (k -dependent) number. In general, one defines Z_k as the value of $Z_k(\varphi)$ at a given point φ_0 , $Z_k = Z_k(\varphi_0)$. Similarly, though X_k here is not a functional, its flow equation depends on φ , $U_k(\varphi)$, $Z_k(\varphi)$ and their derivatives. Henceforth, the notation X_k will mean that the corresponding expressions are evaluated for $\varphi = \varphi_0$. Of course, within the exact renormalization flow, the critical exponents should not in general depend on the choice of φ_0 . However any approximation introduces a residual dependence and the choice of φ_0 may become important. The advocated choice (from equilibrium studies) is the (running) minimum of the effective potential U_k which is implicitly defined by $\partial \varphi U_k(\varphi_0) = 0$, for it possesses the best 'stability' properties [21].

Cuto matrix

Our last discussion to complete the settings of the NPRG formalism for Model A concerns the choice of the cuto matrix \hat{R}_k . The previous symmetry requirements obviously also apply for the quadratic term S_k , which must in particular be invariant under TRS (in the stationary regime).

The minimal non-perturbative renormalization scheme consists in performing a space coarse-graining on the propagator mode $\tilde{\phi}$, which amounts to considering an off-diagonal cutoff matrix \hat{R}_k with elements

$$R_k^{12} = R_k^{21} = Q_k(q^2) \quad \text{and} \quad R_k^{11} = R_k^{22} = 0: \quad (16)$$

This form for \hat{R}_k is the most natural extension of the equilibrium case [1] where one introduces the scale-dependent quadratic term $H_k = \frac{1}{2} \int_{-q}^q Q_k(q^2) \phi_q$ in the (equilibrium) partition function Z_k to achieve the splitting of the fluctuation modes [21]. We recall that the cutoff function $Q_k(q^2)$ must decay fastly for large momentum modes and behave as k^2 for slow modes as emphasized in Section 3. A typical cutoff function which has been widely used since it allows for analytical results is the cutoff introduced by Litim [25]

$$Q_k(q^2) / (k^2 - q^2) (k^2 + q^2): \quad (17)$$

It turns out that, even when considering the dynamics, space coarse-graining is enough to achieve a proper non-perturbative renormalization program since the frequency integrals appear to be convergent and need not be regularized [5]. The cutoff matrix (16) has hence been adopted in all previous non-equilibrium studies and more specially for reaction-diffusion processes [6].

Notice that, on the other hand, one could expect that a time coarse-graining on the $\tilde{\phi}$ propagator may improve the procedure, though it has never been tested. A time coarse-graining could be achieved by adding a frequency cutoff $R_k^{12} = \delta_k(i!)$ on $\tilde{\phi}$ modes. But in this case one would have to coarse-grain the noise part correspondingly in order to sustain TRS (i.e. the invariance under the transformation (6) of S_k). This would amount to introducing an additional cutoff on $\tilde{\phi}\tilde{\phi}$ modes, of the form $R_k^{22} = 2i\delta_k(i!)$. The properties of this mixed regularization scheme have never been investigated as yet and is left for future work since it represents a great deal of numerical efforts.

5. Flow equations

The NPRG flow equations for the renormalization functions U_k , Z_k and X_k are drawn from the exact flow of ϕ_k given by Eq. (11). According to the Ansatz (15), $\phi_k U_k$ can be defined by

$$\phi_k U_k = \frac{(2)^{d+1}(0)}{(2)^{d+1}} \lim_{p; \rightarrow 0} \frac{\phi_k}{\tilde{\phi}(p;)} \quad (18)$$

where the limit of vanishing external momentum and frequency $p; \rightarrow 0$ means that the fields are evaluated in uniform and stationary configurations and the prefactor just corresponds to the volume of the system in Fourier space. Similarly, the renormalization functions Z_k and X_k can be defined by:

$$Z_k = \frac{(2)^{d+1}}{(2)^{d+1}(0)} \lim_{p; \rightarrow 0} \phi_{p^2} \frac{\phi_k^2}{\tilde{\phi}(p;)} \quad (19)$$

$$X_k = \frac{(2)^{d+1}}{(2)^{d+1}(0)} \lim_{p; \rightarrow 0} \phi_i \frac{\phi_k^2}{\tilde{\phi}(p;)} \quad (20)$$

Obviously, X_k can alternatively be defined from the noise part as $X_k / Q_k^2 = \tilde{X}_k^2$ for a uniform and stationary configuration. One can check that both definitions lead to the same flow equation $\partial_s X_k$ which in turn reflects that TRS is preserved by the NPRG flow at any scale s .

The flow equations of the renormalization functions $\partial U_k, Z_k$ and X_k are obtained by taking the scale derivative ∂_s of the expressions (18), (19) and (20) respectively. It is convenient to first rewrite the flow equation $\partial_s X_k$ (given by (11)) as

$$\partial_s X_k = \frac{1}{2} \partial_s \text{Tr} \ln \hat{R}_k^{(2)} + \hat{R}_k; \quad (21)$$

where $\partial_s (\cdot) = \hat{R}_k (\cdot) = \hat{R}_k$ only acts on the s -dependence of the cutoff elements R_k^{ij} . It follows that the field derivatives of $\partial_s X_k$ admit simple diagrammatic representations:

$$\begin{aligned} \frac{\partial_s X_k}{\sim(0;0)} &= \frac{1}{2} \partial_s \text{ (diagram: circle with two external legs labeled 1 and 2)} \\ \frac{\partial_s^2 X_k}{\sim(p; \cdot) \quad (\cdot; p)} &= \frac{1}{2} \partial_s \left(\text{diagram: circle with two external legs labeled 1 and 2, and two internal legs labeled 1 and 2} \right) - \frac{1}{2} \partial_s \left(\text{diagram: circle with two external legs labeled 1 and 2, and two internal legs labeled 1 and 2} \right) \end{aligned}$$

In these graphs, the index 1 or 2 on external legs refers to the corresponding field (\cdot or \sim respectively), the n -point vertices correspond to 2×2 matrices of $\hat{R}_k^{(n)}$ with $n \geq 2$ (external) fixed indices and two summed over. The propagator lines stand for $[\hat{R}_k^{(2)} + \hat{R}_k]^{-1}$, which can be easily computed from (15) and (16). We don't detail the full computation of these graphs nor the subsequent explicit frequency integration which are lengthy but straightforward.

Before giving the resulting flow equations, let us put them in a suitable form for the search of fixed point (since we are interested in one in the scale invariant regime). First it is convenient to explicitly express the Ising symmetry by defining the functions $\tilde{U}_k(\cdot) = U_k(\cdot), \tilde{Z}_k(\cdot) = Z_k(\cdot)$ where $\cdot = \pm 1$ is the Z_2 invariant. The derivatives of the functions of $\tilde{U}_k(\cdot)$ and $U_k(\cdot)$ are simply related: $\partial U_k = \partial \tilde{U}_k$, $\partial^2 U_k = \partial U_k + 2 \partial^2 \tilde{U}_k$ and similarly for Z_k and \tilde{Z}_k . As for the flow equation, $\partial_s(\partial \tilde{U}_k) = 1 = \partial_s(\partial U_k)$ for nonzero and $\partial_s \tilde{Z}_k = \partial_s Z_k$. Then to absorb any explicit dependence on the running scale k , we introduce the dimensionless quantities (according to (15))

$$\begin{aligned} \tilde{g} &= k^{2-d} Z_k \\ \tilde{u}(\cdot) &= k^{-d} \tilde{U}_k(\cdot) \\ \tilde{z}(\cdot) &= Z_k^{-1} \tilde{Z}_k(\cdot) \end{aligned} \quad (22)$$

where the subscript k has been dropped on dimensionless functions. We further introduce a dimensionless cutoff function $r(y) = Q_k(q^2) = (Z_k q^2)$ where $y = q^2/k^2$ and hence $\partial_s Q_k = Z_k k^2 s(y)$ with $s(y) = -\int dy r(y) - 2y^2 \partial_y r(y)$. Finally, the flow

equations for the dimensionless functions u^0 and z are given by:

$$\begin{aligned}
 \partial_s u^0 &= u^0 (2 + z) + (2 + d + z) u^0 + \frac{1}{2} (3u^0 + 2 u^0) L_1^d + \frac{z^0}{2} L_1^{2+d} \\
 \partial_s z &= z z + (2 + d + z) z^0 + \frac{1}{2} (z^0 + 2 z^0) L_1^d - 2 z^0 (3u^0 + 2 u^0) L_2^d \\
 &\quad + \frac{1}{d} (1 + 2d) z^0 L_2^{2+d} + 2 (3u^0 + 2 u^0)^2 M_4^d \\
 &\quad + 4 z^0 (3u^0 + 2 u^0) M_4^{2+d} + 2 z^0 M_4^{4+d} \\
 \partial_s \ln X_k &= \frac{1}{2} (3u^0 + 2 u^0)^2 L_3^d + z^0 (3u^0 + 2 u^0) L_3^{2+d} + \frac{1}{2} z^0 L_3^{4+d}
 \end{aligned} \tag{23}$$

where primes denote derivatives wrt s and the so-called threshold functions L and M are defined by:

$$\begin{aligned}
 L_n^d &= \frac{1}{n!} \frac{d^n}{dy^n} \left[\frac{s(y)}{h(y)^{n+1}} \right] \\
 M_n^d &= \frac{1}{n!} \frac{d^n}{dy^n} \left[\frac{(n + n_0) s(y) (\partial_y h(y))^2}{h(y)^{n+1}} + \frac{2 \partial_y s(y) \partial_y h(y)}{h(y)^n} \right]
 \end{aligned} \tag{24}$$

with $v_d^1 = 2^{d-2}$ ($d=2$) and $h(y) = y(z + r(y)) + u^0 + 2 u^0$. By definition (see Section 4), the anomalous dimension $\partial_s \ln X_k$ is obtained by evaluating Eq. (23) at the running minimum u_0 of the potential U_k , or equivalently at the minimum u_0 of u . Similarly, z is obtained by solving at the minimum the equation $\partial_s z|_0 + z^0 \partial_s u_0 = 0$ where $\partial_s z$ is given by (23). The additional contribution $z^0 \partial_s u_0$ is generated by the running of the minimum implicitly defined as $u^0(u_0) = 0$. Indeed, the running of u implies that its minimum flows according to $\partial_s u^0(u_0) = 0 = \partial_s u^0|_0 + \partial_s u^0|_0 u_0$ which, using (23) evaluated at u_0 yields the expression for $\partial_s u_0$.

We emphasize that, as is to be expected from FDT, the dynamics decouples from the statics, i.e. the non-perturbative flow equations (23) for u^0 and z do not depend on X_k (and they identify with those derived in equilibrium with the Ansatz (13) [21]). Note that the threshold functions L and M intervening in the equation for the anomalous dimensions can be computed analytically upon the choice of the cutoff $r(y) = (1-y)^{-1}$ ($1-y$) (corresponding to (17)), which greatly simplifies the numerical resolution. The numerical procedure to integrate the flow equations (23) is detailed in the next section.

6. Numerical integration of the flow equations

We will consider different levels of approximation. In a first step | which will be referred to as local potential approximation (LPA) | the field dependence of the kinetic renormalization function z can be neglected, i.e. only a running coefficient Z_k is considered. When restoring the ϕ -dependence of z , the corresponding approximation will be denoted UZA.

For both approximations, the numerical procedure to determine the fixed point solution of Eq. (23) and to compute the critical exponents is the following. We sample the field on a mesh of spacing $\Delta\phi$ and discretize the flow equations (23) using finite differences at order $\Delta\phi^4$ to calculate the ϕ -derivatives of u (and z). For the integrals, we either use their analytical expression (whenever available) or

calculate them numerically using Simpson's rule. We implement an explicit forward integration scheme to propagate the solution between scales s and $s + \Delta s$, which turns out to be stable for sufficiently small Δs and s . The convergence of the numerical procedure when varying Δs and s has been carefully checked. We start out at the microscopic scale $s = 0$ ($k = \Lambda$) from a quartic bare potential $u(\phi) = \frac{1}{4} \phi^4$ where Λ represents the temperature. We carry through the numerical integration by lowering s towards $s \rightarrow 1$. For large bare Λ , the system flows to the symmetric (high temperature) phase where the (dimensionfull renormalized) minimum of the potential $\phi_0 = k^{d/2} Z_k^{-1} \phi_0$ vanishes, whereas for small Λ , it flows to the broken (low temperature) phase where ϕ_0 acquires a finite value as $s \rightarrow 1$. For a re-tuned initial ϕ^c , the system is in the critical regime, which corresponds to the effective potential u (and z) flowing to a fixed point (scale invariant) form u^* (and z^*). The critical exponents ν and z can then be computed from the fixed point values of z and ν . The critical index β is obtained by linearizing the flow in the vicinity of $(u^*; z^*)$ and determining the (negative) eigenvalue characterizing the unstable (relevant) direction. This procedure is carried out within the LPA and UZA approximations. The results are gathered in tables 1 and 2 and are commented in the next section.

7. Results

The critical exponents for Model A obtained in this work from the NPRG equations (23) are summarized in Tables 1 and 2 for dimensions $d = 3$ and $d = 2$ respectively, and compared with results ensuing from other field theoretical methods and Monte Carlo simulations.

Let us first comment on the equilibrium exponents ν and β . As emphasized in Section 5, the dynamics decouples from the statics in Eqs. (23) as expected from TRS in the stationary regime. As a consequence, the equilibrium exponents computed here should match (up to numerical accuracy) those obtained in earlier NPRG works on the (equilibrium) Ising model. The three-dimensional Ising model has been thoroughly investigated within the NPRG framework as a testing ground of the method [26]. In particular, critical exponents have been computed using the cutoff function (17) in [23, 27] (though with different numerical procedures). The exponents ν and β we obtain in $d = 3$ precisely reproduce these values both at LPA and UZA.

We know from these previous studies that the accuracy can be improved by optimizing the choice of the cutoff function. At order r^2 in derivatives, optimized exponents are $\nu = 0.6281$ and $\beta = 0.0443$ [28], which are already in close agreement with the 6-loop calculations [24] or Monte Carlo simulations [29]. However, since the determination of ν is related to the momentum structure of the two-point correlation function, its accuracy is poorer than that of β . A better accuracy on ν requires to compute the next order r^4 in derivatives which yields $\nu = 0.033$ [23]. In two dimensions, far fewer NPRG results are available. A calculation with cutoff function (17) has been achieved in [27] and both results are in close agreement. As for $d = 3$, the determination of ν remains poorer than that for β at order r^2 in derivatives. However, the two-dimensional Ising model has not been systematically investigated and neither optimized nor order r^4 exponents have been determined in $d = 2$.

We can now come to the new part of this work which concerns the dynamics. The situation for z is very different from that of the equilibrium critical exponents. For the dynamics, no high-loop expansions or exact results in $d = 2$ are available. Furthermore, results from MC simulations appear to be rather scattered especially in

d = 3			z	Ref.
LPA	0.65	0.11	2.05	[this work]
UZA	0.63	0.05	2.14	[this work]
FT	0.6304 (13)	0.0335 (25)		[24]
MC	0.6297 (5)	0.0362 (8)		[29]
FT			2.0237 (55)	[30]
MC			2.032 (4) 2.055 (10)	[31, 32]

Table 1. Critical exponents of Model A in $d = 3$ from the different NPRG approximations (LPA and UZA) computed in this work, compared with results from field theoretical methods (FT) and Monte Carlo simulations (MC).

d = 2			z	
LPA	0.78	0.43	2.15	[this work]
UZA	1.1	0.37	2.17	[this work]
exact	1	1/4		
FT			2.0842 (39)	[30]
MC			2.1667 (5)	[33]

Table 2. Critical exponents of Model A in $d = 2$ from the different NPRG approximations (LPA and UZA) computed in this work, compared with results from field theoretical methods (FT) and Monte Carlo simulations (MC).

$d = 2$. The values reported in Tables 1 and 2 seem to be accepted as reference values [14]. On the field-theoretical side, the determination of z is very sensitive to the choice of the resummation scheme since only a few orders are known. Various resummation schemes have been studied and we quote here the latest results obtained in [30].

Our results are in reasonable agreement with these various estimates. This is one of the key point of the NPRG approach and a central motivation for this work: the leading order in derivatives appears to already provide a reliable determination of physical quantities, as outlined above. Notice that the variation on z between LPA and UZA is not meaningful and merely provides an estimate of the error. The reason is that going from LPA to UZA does not amount to enriching the Ansatz for the dynamical part since in both cases only a running coefficient X_k is allowed by TRS. Hence even at UZA, the accuracy on z remains poorer than that for the equilibrium exponents and the rapidity of convergence on z can not be tested within these approximations. One would need to implement the next order in time derivatives to improve the accuracy on z , which is rather costly. Alternatively, one could modify the regularization scheme and resort to a frequency and noise coarse-graining as mentioned in Section 4, which is likely to yield better results, but is yet to be investigated.

8. Summary

In this work, we have studied the critical dynamics of Model A within the NPRG formalism. We have in particular detailed the device of an appropriate approximation scheme preserving the symmetries, the derivation of the NPRG flow equations and their resolution. Using a very simple Ansatz, that is at the leading order in derivatives,

we have obtained a reliable estimate for the dynamical exponent z : $z = 2.09(4)$ in $d = 3$ and $z = 2.16(1)$ in $d = 2$. The fact that the leading order already yields quantitative results is a generic feature of the NPRG approach [21] which makes it particularly powerful. This feature is the central motivation for the emphasis put throughout this work on the methodological part: a leading order NPRG calculation can already enable one to investigate nontrivial problems. Restricting to non-equilibrium critical phenomena, NPRG studies have indeed brought out new non-perturbative properties of reaction-diffusion processes [10] and allowed to tackle interface growth problems [34]. We hence believe the approach to be useful to investigate many other non-equilibrium scaling phenomena. Of course, as the application of NPRG techniques to non-equilibrium statistical physics and dynamics is very recent, a great deal of systematic studies remain to be done as well to test the efficiency of the different Ansatz and regularisation schemes out-of-equilibrium, which will be the goal of future works.

The authors are indebted to B. Delamotte for fruitful exchanges all through this work. The authors also wish to thank G. Biroli, A. Lefevre and N. Wschebor for enlightening discussions.

9. Appendix: NPRG flow equation for χ_k

This appendix is dedicated to the derivation of the exact flow equation (11) for χ_k . (A similar derivation can be found for equilibrium systems in [21]). In this appendix, we use the shorthand $\chi = \chi(x; t)$, vectors are denoted by capital letters (eg. $\mathbf{J} = [j; \tilde{j}]$, $\mathbf{J} = [j; \tilde{j}]$) and for functional derivatives we introduce the notation $\hat{\mathcal{J}}_x = \mathcal{J}(x); = \mathcal{J}(x)$ (and similarly for $\hat{\mathcal{J}}_x$). Overhead hat symbols are used for (2×2) matrices.

The variation of χ_k (at fixed \mathbf{J}) under an infinitesimal change of the running scale k follows from Eq. (10):

$$\partial_k \chi_k[\mathbf{J}] = \partial_k W_k[\mathbf{J}] + \int_x \partial_k \mathcal{J}(x) : \hat{\mathcal{J}}_k(x) : \frac{1}{2} \int_{x, x^0} (x) : \partial_k \hat{\mathcal{R}}_k(x; x^0) :^t (x^0) \quad (25)$$

where $W_k = \ln Z_k$. The variation with k of W_k at fixed \mathbf{J} is related to that of W_k at fixed \mathbf{J} by:

$$\partial_k W_k = \partial_k W_{k, \mathbf{J}} + \int_y \partial_k \mathcal{J}(y) : \hat{\mathcal{J}}_y W_k : \quad (26)$$

The expression of $\partial_k W_{k, \mathbf{J}}$ is obtained by taking the derivative of (9) wrt k :

$$\begin{aligned} \partial_k \ln Z_{k, \mathbf{J}} &= \int_D \int_n \frac{1}{2} \int_{x, x^0} (x) : \partial_k \hat{\mathcal{R}}_k(x; x^0) :^t (x^0)^0 e^{S_k} S_k + t_{\mathbf{J}} \\ &= \int_n \frac{1}{2} \int_{x, x^0} \hat{\mathcal{J}}_x : \partial_k \hat{\mathcal{R}}_k(x; x^0) :^t \hat{\mathcal{J}}_{x^0}^0 e^{W_k} \\ &= \partial_k W_k e^{W_k} : \end{aligned}$$

After expressing the derivatives $\hat{\mathcal{J}}_J$ and t_J of $\exp(W_k)$ and dividing out by $\exp(W_k)$

one obtains:

$$\partial_k W_k = \frac{1}{2} \int_{x, x^0}^Z \hat{W}_k(x; x^0) : \partial_k \hat{R}_k(x; x^0) : \hat{W}_k(x; x^0) + \text{Tr} \partial_k \hat{R}_k(x; x^0) : \hat{W}_k(x; x^0) : \quad (27)$$

The last term in the right hand side is the matrix of second (functional) derivatives of W_k which we denote $\hat{W}_k^{(2)}(x; x^0)$. The following equation of k follows from inserting (26) and (27) in (25), which yields

$$\partial_k k[\] = \frac{1}{2} \text{Tr} \int_{x, x^0}^Z \partial_k \hat{R}_k(x; x^0) : \hat{W}_k^{(2)}(x; x^0) : \quad (28)$$

This equation can be conveniently expressed in a closed form upon inverting $\hat{W}_k^{(2)}$. The inverse of $\hat{W}_k^{(2)}$ can be obtained by taking a functional derivative \hat{J}_{x^0} of the definition $\hat{W}_k(x) = \hat{J}_x W_k$:

$$\hat{J}_{x^0}^{(d+1)}(x; x^0) = \int_y^Z \hat{J}_{x^0}(y) : \hat{W}_k^{(2)}(y; x) : \quad (29)$$

The matrix $\hat{J}_{x^0}(y)$ is simply given by two successive derivatives \hat{J}_y and \hat{J}_{x^0} of Eq (10), which yields:

$$\hat{J}_{x^0}(y) = \hat{J}_k^{(2)}(x^0; y) + \hat{R}_k(x^0; y); \quad (30)$$

where $\hat{J}_k^{(2)}(x; x^0)$ denotes the matrix of second (functional) derivatives of k . Finally, inserting (30) in the following equation (28) yields the advocated equation:

$$\partial_k k[\] = \frac{1}{2} \text{Tr} \int_{x, x^0}^Z \partial_k \hat{R}_k(x; x^0) : \hat{J}_k^{(2)} + \hat{R}_k^{-1}(x; x^0) : \quad (31)$$

which can be Fourier transformed and underlies the NPRG calculations of this work.

References

- [1] P. Bak, C. Tang, and K. Wiesenfeld, Phys. Rev. Lett. 59, 381 (1987).
- [2] H. Hinrichsen, Adv. Phys. 49, 815 (2000).
- [3] G. Odor, Rev. of Mod. Phys. 76, 663 (2004).
- [4] U. C. Tauber, M. Howard, and B. P. Vollmayr-Lee, J. Phys. A 38, R79 (2005).
- [5] L. Canet, B. Delamotte, O. Delubriere, and N. Wschebor, Phys. Rev. Lett. 92, 195703 (2004).
- [6] L. Canet, J. Phys. A 39, 7901 (2005).
- [7] J. L. Cardy and U. C. Tauber, Phys. Rev. Lett. 77, 4780 (1996).
- [8] J. L. Cardy and U. C. Tauber, J. Stat. Phys. 90, 1 (1998).
- [9] L. Canet, H. Chate, and B. Delamotte, Phys. Rev. Lett. 92, 255703 (2004).
- [10] L. Canet et al, Phys. Rev. Lett. 95, 100601 (2005).
- [11] C. Wetterich, Phys. Lett. B 301, 90 (1993).
- [12] T. R. Morris, Int. J. Mod. Phys. A 9, 2411 (1994).
- [13] P. C. Hohenberg and B. I. Halperin, Rev. Mod. Phys. 49, 435 (1977).
- [14] P. Calabrese and A. Gambassi, J. Phys. A 38, R133 (2005).
- [15] P. C. Martin, E. D. Siggia, and H. A. Rose, Phys. Rev. A 8, 423 (1973).
- [16] H. K. Janssen, Z. Phys. B 23, 377 (1976).
- [17] C. de Dominicis, J. Phys. (Paris) C 1, 247 (1976).
- [18] A. Andreanov, G. Biroli, and A. Lefevre, J. Stat. Mech. P07008 (2006).
- [19] K. G. Wilson and J. Kogut, Phys. Rep. C 12, 75 (1974).

- [20] N. Tetradis and C. Wetterich, Nucl. Phys. B [FS] 422, 541 (1994).
- [21] J. Berges, N. Tetradis, and C. Wetterich, Phys. Rep. 363, 223 (2002).
- [22] L. Canet, Phys. Ann. Fr. 29, (2004).
- [23] L. Canet, B. Delamotte, D. Mouhanna, and J. Vidal, Phys. Rev. B 68, 064421 (2003).
- [24] R. Guida and J. Zinn-Justin, J. Phys. A 31, 8103 (1998).
- [25] D. F. Litim, Phys. Rev. D 64, 105007 (2001).
- [26] C. Bagnuls and C. Bervillier, Phys. Rep. 348, 91 (2001).
- [27] H. Ballhausen, J. Berges, and C. Wetterich, Phys. Lett. B 582, 144 (2004).
- [28] L. Canet, B. Delamotte, D. Mouhanna, and J. Vidal, Phys. Rev. D 67, 065004 (2003).
- [29] M. Hasenbusch, Int. J. Mod. Phys. C 12, 911 (2001).
- [30] A. S. Krinitsyn, V. V. Prudnikov, and P. V. Prudnikov, Theor. Math. Phys. 147, 561 (2006).
- [31] N. Ito, K. Hukushima, K. Ogawa, and Y. Ozeki, J. Phys. Soc. Jap. 69, 1931 (2000).
- [32] P. Grassberger, Physica A 214, 547, [Erratum 217 227] (1995).
- [33] M. Nightingale and H. Blte, Phys. Rev. B 62, 1089 (2000).
- [34] L. Canet, cond-mat/0509541 (2005).