

Single File Diffusion enhancement in a fluctuating modulated 1D channel

Gwennou Coupier,¹ Michel Saint Jean,^{1,*} and Claudine Guthmann¹¹Laboratoire Matière et Systèmes Complexes, UMR 7057 CNRS & Université Paris 7 - 140 rue de Loumel, F-75015 Paris, France

(Dated: March 23, 2024)

We show that the diffusion of a single file of particles moving in a fluctuating modulated 1D channel is enhanced with respect to the one in a bald pipe. This effect, induced by the fluctuations of the modulation, is favored by the incommensurability between the channel potential modulation and the moving file periodicity. This phenomenon could be of importance in order to optimize the critical current in superconductors, in particular in the case where mobile vortices move in 1D channels designed by adapted patterns of pinning sites.

PACS numbers: 05.40.-a

Recent progress in lithography techniques have given rise to many developments to obtain superconducting mesoscopic devices with large critical currents. These currents being dependent upon the vortices pinning, many studies have been focused in the last past years to understand the properties of a mobile file of vortex moving in 1D channels. Some are devoted to the determination of the mobility of vortices moving in a strip geometry in order to investigate the influence of this motion on the generation of noise [1, 2], others explore the vortex flow in modulated channels designed by pinned vortices [3, 4, 5, 6, 7]. In such systems, the vortex mobility depends upon the adequacy between the inter-vortices distance and the periodicity of the pinned vortices array and/or the channel width. For instance, it is well known that this mobility drops when the vortices file disintegrates into a channel with a commensurate configuration of the pinning sites and that it is very sensitive to the accuracy of this matching. Experimentally, these effects have essentially been observed at a macroscopic scale at which these matching effects result in an important increase of the critical current for some precise values of the magnetic field. The high sensitivity of the critical current towards the matching of the characteristic lengths suggests that fluctuations in the channel modulation, originating in the small movements of trapped vortices for instance, could subtly influence the resulting flow from the mobile vortices. From the theoretical point of view, the Langevin equations describing the movements of particles located in a file and diffusing in a fluctuating modulated channel are non linear and their analytical solution is out of reach. Numerical studies have also been performed [4, 5, 6, 7]; however, a complete description of such a behavior introducing these fluctuations is still lacking.

In this letter we evidence that these fluctuations induce an important enhancement of the vortices diffusion, one order of magnitude higher than in the case of a bald channel. To exhibit this very surprising result, we have used an original experimental approach which allows to follow the movement of each particle located

in the file and determine its diffusion. More precisely, we have compared the diffusion of particles moving in a fluctuating modulated channel to the one observed when the same particles are distributed in a pipe without modulation, hereafter called bald pipe.

In these experiments, millimetric stainless-steel balls are located on the bottom electrode of an horizontal plane condenser, while a metallic frame intercalated between the two electrodes and in contact with the bottom electrode connects them. Depending on the experiments, this frame is a disk whose external diameter is 10 mm or a circular bald pipe, its external diameter of 10 mm and its width of 2 mm preventing any crossing between particles (Fig.1). When a voltage $V = 1$ kV is applied between the two electrodes, the balls become monodispersely charged, repel by each other and spreading throughout the whole available space. We have shown that their electrostatic interaction is described by a modified Bessel function of the second kind K_0 with a screening length $\lambda = 0.48$ mm. Notice that this interaction is exactly similar to the inter-vortex interaction in superconductors [8, 9]. To introduce thermal noise, the whole cell is fixed on loudspeakers supplied by a white noise voltage, this mechanical shaking simulating a tunable effective temperature that is measured in-situ [10]. Throughout the experiments, images of the particles are recorded in real time using a camera. The interval between two successive snapshots is 150 ms and ve series of 10000 images have been recorded for each experiment. With this choice, relevant statistics for the long-time behavior of the displacements are obtained, the length of one experiment remaining reasonable since the effective thermal bath is characterized by a relaxation time of about 100 ms [11]. The diffusion of the particles is measured through the evolution with time of their mean square displacements (m.s.d.) $\langle \Delta r^2(t) \rangle$ given by :

$$\langle \Delta r^2(t) \rangle = \langle r^2(t+t_0) \rangle - \langle r^2(t_0) \rangle = \langle r^2(t+t_0) \rangle - \langle r^2(t_0) \rangle \quad (1)$$

where the orthoradial coordinate θ (in radians) is




FIG. 1: Experimental configurations. Wigner islands : (a) $N = 19$, (b) $N = 18$; circular ball pipe (c) $N = 12$. In inset, the corresponding balls trajectories at the same effective temperature.




FIG. 2: Evolution with time of the orthoradial coordinate of a ball in : a ball circular pipe (i), a Wigner island $N = 19$ outer shell (two different balls) (ii), a Wigner island $N = 19$ inner shell (iii).

the cumulated angle, and not the modulo 2π angle in order to explore an unbounded motion. The brackets $\langle \rangle$ denote ensemble averaging over the initial time t_0 and a set of statistically independent trajectories.

In order to observe a 1D movement in a fluctuating modulated potential, we first considered the outer circular shell of a Wigner island, constituting of 19 interacting particles confined in a circular disk, for which the ground configuration corresponds to self-organized pattern constituted of three concentric shells filled by 1, 6 and 12 balls. This configuration is denoted (1-6-12) hereafter (Fig.1) [9]. To observe well-defined shells, the experiments have been performed at an effective temperature equal to roughly a few tens of the inter-ball interaction ; for this temperature range, the balls seldom jump from




FIG. 3: Comparison between the mean square orthoradial displacements $\Delta\theta^2(t)$ of a ball moving in the Wigner island $N = 19$ outer shell and in a ball circular pipe (log-log scale).

one shell to another [10]. The effective thermal agitation only induces orthoradial motion while the radial displacements are reduced as it can be seen on the balls trajectories presented in the insets of Fig.1 [10]. This offers the opportunity to perform long-time orthoradial diffusion experiments. In the outer shell, the relative mean orthoradial displacements of two neighboring balls are about 0.07 radian (resp. 0.15 in the inner shell), much smaller than the angular distance $2\pi/12$ (resp. $2\pi/6$) between them. Thus, each shell can be considered as a periodic ring. This ring presents global angular movements which are coherently followed by the balls as shown in Fig.2 (ii) where the trajectories of two different balls of the outer shell are reported. Their global orthoradial movements are similar but their trajectories present differences of low relative amplitude at the small time-scale. Therefore, the outer twelve-ball shell is a well-adapted realization of a




FIG. 4: Evolution with time of the mean square orthoradial displacement of a ball located in the outer shell of the commensurate Wigner island $N = 19$ and of the incommensurate Wigner island $N = 18$ (log-log scale).

periodic system of particles moving in a fluctuating modulated potential due to the inner shell. Along the orthoradial direction, the angular displacements of the balls have a gaussian distribution. The evolution with time of the corresponding m.s.d. $\Delta\theta^2(t)$ is presented in Fig. 3 for $150 \text{ ms} \leq t \leq 150 \text{ s}$. This angular m.s.d. increases with time with a t dependence where α is close to 0.6.

We have compared these results with those obtained for the same twelve balls moving in a circular bald pipe, the effective temperature and the inter-ball interaction remaining the same (Fig. 1). We emphasize that, even if the movement of the balls is not strictly 1D, no coupling effects resulting from the circular channel geometry were observed. For instance, the orthoradial movement of a single ball in this circular gutter is a free diffusion [11]. As in the Wigner island case, the distribution of the angular movement of the balls is gaussian. Long-time behavior of the angular m.s.d. differs dramatically from the Wigner island case. Instead of t with $\alpha = 0.6$, the movement is subdiffusive and characterized by a growth much slower than in the modulated case, varying from $1/2$ until $\alpha = 0.4$ at the end of the experiment (the decrease of α being even more important at higher temperature). Equally remarkable is that the amplitude of the diffusion is much smaller in this bald pipe case than in the case of a fluctuating modulated channel; 10 times smaller for instance for the last decade. This important difference in the mobilities can be also directly observed on the various trajectories presented on Fig. 2 (i-ii). The role of the periodicity is confirmed when considering the diffusion in the inner shell, which is seven times higher than in the outer shell, with the same power law. Whereas an outer ball moves in a modulated potential with a $2\pi/6$ period, an inner ball is indeed submitted to a potential with a period twice as small; thus the impulsion transfer that

enhances the diffusion is linked to the modulation of the potential.

These behaviors must be discussed in the frame of the Single File Diffusion theory [12, 13] which describes the diffusion of particles in a single channel where the crossings are forbidden. For infinite systems with contact interaction, this theory predicts that correlations between particles induce a subdiffusion with $\alpha = 0.5$ whereas classical free diffusion is characterized by $\alpha = 1$. We suggested that the smaller value of α for the diffusion in the bald pipe resulted from the cyclicity of the system associated to the non linearity of the interaction [11]. From this point of view, the $t^{0.6}$ behavior observed in the outer shell of the Wigner island is all the more striking. Understanding this phenomenon and explaining why the amplitude in a fluctuating modulated channel is higher than in a smooth one is a complex issue, as it is well-known that the diffusion of particles in a static modulated potential is slower than for a free diffusion. Here the Single File Diffusion and the modulation effects are coupled, and there is no theory describing this coupling yet. Nevertheless, some propositions can be given. In a recent paper, Bandyopadhyay et al. [14] have shown that diffusion of a single particle in a modulated potential can be largely enhanced if the particle is excited by a rapid fluctuating force. In the same way, we can attribute the diffusion increase to the fluctuations of the modulated potential felt by the outer balls, the fluctuating part of the force being associated with the impulsion transfer resulting from the oscillations of the balls located in the neighbor inner shell.

To reveal how the channel modulation and their fluctuations can play a role, we have performed diffusion experiments in which we have modified the inter-shell in-

interaction without drastically changing the moving ϕ . We have compared the diffusion of the outer shell of the $N = 19$ system to the one obtained with a $N = 18$ island whose ground configuration is (1-6-11). The advantage of these two systems is that they roughly have the same number of balls in the outer shell whereas their relative symmetries strongly differ: the commensurate system $N = 19$ exhibits a three fold symmetry whereas the $N = 18$ system is incommensurate. This difference of relative symmetry between the two shells modify the inter-shell coupling and thus might influence the effect of this coupling on the diffusion. The variations of the corresponding $\chi^2(t)$ are presented in Fig.4 for three different temperatures T . For long times, the m.s.d. associated with the incommensurate $N = 18$ system increases roughly with the same power law ($\alpha = 0.6$) as for the commensurate island $N = 19$. However, the evolution with T of the diffusion amplitudes differ according to N and a detailed analysis of these differences proves that it is greatly influenced by the relative inter-shell movement. Indeed, we showed in a previous paper that for $11 \cdot 10^{11}$ K $T \leq 12 \cdot 10^{11}$ K, the relative shells' movements are small in the $N = 19$ system [10]. In the diffusion experiments, one can observe that, in the same range of temperature, the diffusion curves are remarkably roughly superposable (Fig.4). Likewise, at higher temperature, we observe a stronger diffusion as the inter-shell movement is increased. Furthermore, for the same temperatures, the shells for $N = 18$ are always unlocked [10] and the amplitudes of the inter-shell displacements grow with temperature. Similar behaviors can be observed on the diffusion curves: whatever the temperature, the corresponding diffusion curves are always well separated and the diffusion amplitude increases also with the effective temperature (Fig.4). Then we can conclude that there exists a strong correlation between the inter-shell movement and the diffusion in each shell.

However, it would be naive to picture the two shells as a set of gear-wheels, the correlation effect being undoubtedly more subtle. Indeed, a gear-wheels picture could suggest that a more important diffusion increase would be obtained in the case of two commensurate shells but this assumption does not correspond to the observations. Indeed, whatever the effective temperature, diffusion for balls of $N = 18$ is always higher than for $N = 19$, the maximum difference being precisely at the temperature at which the $N = 19$ system is locked whereas the $N = 18$ system is unlocked (Fig.4). When the temperature increases, the differences are less important, the ϕ of particles can be considered as coating above the modulated potential. It then only feels an averaged effect where the influence of the peculiar symmetries is hidden. We stress that a similar effect of the commensurability can also be

observed for balls in the inner shells. Thus, even though the fluctuations of the modulated potential induces impulsion transfers, a better transfer is obtained in incommensurate configurations, a too good matching of the periods inducing the locking of the whole system rather to a positive cooperation.

In conclusion, the diffusion of a ϕ of particles can be accelerated when it moves in a fluctuating modulated potential. This enhancement presents two strong characteristics: the orthoradial m.s.d. increases as a power law with $\alpha > 0.5$ whereas the diffusion in a circular Single File Diffusion is characterized by $\alpha = 0.5$. Moreover, the diffusion amplitude in a fluctuating potential is much larger than in a bald pipe. This effect seems to be favored by the incommensurability between the modulation of the potential and the periodicity of the moving ϕ . This effect could be very important in the case of mobile vortices moving between pinned ones in superconducting devices. In particular, it has to be taken into account to design well adapted patterns of traps in order to optimize the critical current. From the theoretical point of view, getting a better understanding of this phenomenon would require to develop a complete SFD theory including a fluctuating potential.

Electronic address: michel.saintjean@paris7.jussieu.fr

- [1] M. H. Thoenissen, E. van der Drift, and P. H. Kes, Phys. Rev. Lett. 77, 159 (1996).
- [2] G. Stan, S. B. Field, and J. M. Martinis, Phys. Rev. Lett. 92, 097003 (2004).
- [3] S. Anders, A. W. Smith, H. M. Jaeger, R. Besseling, P. H. Kes, and E. van der Drift, Physica C 332, 35 (2000).
- [4] R. Besseling, R. Nijmegen, and P. H. Kes, Phys. Rev. Lett. 82, 3144 (1999).
- [5] R. Besseling, P. H. Kes, T. D. Rose, and V. M. Vinokur, New Journal of Physics 7, 71 (2005).
- [6] N. K. Okubo, R. Besseling, V. M. Vinokur, and P. H. Kes, Phys. Rev. Lett. 88, 247004 (2002).
- [7] N. K. Okubo, R. Besseling, and P. H. Kes, Phys. Rev. B 69, 064504 (2004).
- [8] P. Galatola, G. Coudier, M. Saint Jean, J.-B. Fournier, and C. Guthmann, Eur. Phys. J. B 50, 549 (2006).
- [9] M. Saint Jean, C. Even, and C. Guthmann, Europhys. Lett. 55, 45 (2001).
- [10] G. Coudier, C. Guthmann, Y. Noat, and M. Saint Jean, Phys. Rev. E 71, 046105 (2005).
- [11] G. Coudier, M. Saint Jean, and C. Guthmann, Phys. Rev. E 73, 031112 (2006).
- [12] T. E. Harris, J. Appl. Prob. 2, 323 (1965).
- [13] H. van Beijeren, K. W. Kehr, and R. Kutner, Phys. Rev. B 28, 5711 (1983).
- [14] M. Bandyopadhyay, S. Dattagupta, and M. Sanyal, Phys. Rev. E 73, 051108 (2006).