

LiHo_xY_{1-x}F₄ as a random field Ising ferromagnet

Moshe Schechter

Department of Physics & Astronomy, University of British Columbia, Vancouver, B.C., Canada, V6T 1Z1
(Dated:)

As a result of the interplay between the intrinsic off-diagonal terms of the dipolar interaction and an applied transverse field H_t , the diluted LiHo_xY_{1-x}F₄ system at $x > 0.5$ is equivalent to a ferromagnet in a longitudinal random field (RF). At low H_t the quantum fluctuations between the Ising like doublet states are negligible, while the effective induced RF is appreciable. This results in a practically exact equivalence to the classical RF Ising model. By tuning H_t , the applied longitudinal field, and the dilution, the Ising model can be realized in the presence of an effective RF, transverse field, and constant longitudinal field, all independently controlled. The experimental consequences for $D = 1; 2; 3$ dimensions are discussed.

Since its seminal discussion by Imry and Ma [1], the Ising model in random longitudinal magnetic field was found to have many interesting realizations in nature, and has been a subject of intensive theoretical and experimental investigation for both short-range and dipolar ferromagnetic (FM) interactions (see Refs. [2, 3, 4, 5] and references therein). In particular, experiments are mostly done on diluted antiferromagnets (DAFM) [6], which were shown to be equivalent in their static critical behavior to the random field Ising model (RFIM) [6, 7]. In the DAFM the order parameter is given by the staggered magnetization, and a constant applied longitudinal field plays the role of a tunable random field (RF). However, the straightforward realization of the RFIM in a FM system has not found an experimental realization, due to the difficulty of applying a local random magnetic field.

Interestingly, the diluted LiHo_xY_{1-x}F₄ system provides the possibility of producing effective local magnetic fields in the longitudinal direction, due to the interplay of the off-diagonal terms of the dipolar interactions and an applied constant magnetic field H_t in the transverse direction [8, 9, 10, 11]. Importantly, as is further explained below, due to the strong hyperfine interactions in this system, for $H_t < 0.5T$ the applied transverse field does generate an effective longitudinal RF due to fluctuations between each Ising ground state (GS) doublet and its corresponding excited state [10, 12], but the coupling between the two Ising doublet states is negligible [9]. Furthermore, despite the presence of some correlations between the random fields (RFs), and their origin in the long range dipolar interaction, we show below that on length scales larger than the inter-vacancy distance the behavior of the system is equivalent to that of the uncorrelated RF model. Therefore, despite the presence of a finite H_t , and finite correlations on short scales, LiHo_xY_{1-x}F₄ at $H_t < 0.5T$ is a perfect realization of the classical RFIM with a tunable random magnetic field.

As H_t is enhanced, the effective transverse field becomes appreciable. Thus, the Ising model Hamiltonian

$$H = \sum_{ij} J_{ij} \sum_i \sigma_i^z \sum_j \sigma_j^z + \sum_i \sigma_i^z h_i + \sum_i \sigma_i^x h_k + \sum_i \sigma_i^z h_z \quad (1)$$

can be realized in the LiHo_xY_{1-x}F₄ with an effective local RF term h_i , an effective transverse (quantum) term




FIG. 1: Schematic picture of the phase diagram at some dilution $x < 1$. At $H_t < 0.5T$ the quantum fluctuations between the Ising doublet states are negligible, while the effective random field is already appreciable. In this regime the system is equivalent to the classical RFIM. At $T \approx 0.2K$ only the electro-nuclear ground states are occupied, and the quantum phase transition in the presence of the random effective field can be studied.

, and an effective constant field in the longitudinal direction h_k . In the regime considered in this Rapid Communication, where $x > 0.5$, the system is FM at low T ; H_t [13, 14], and J_{ij} correspond to the long range dipolar interaction. In comparison to the DAFM systems, the LiHo_xY_{1-x}F₄ system is different since in the former the effective interaction is short range, and advantageous since in the DAFM system one can realize neither an effective longitudinal field in the staggered magnetization, nor an appreciable quantum temp. Importantly, the three effective fields, h_k and $\text{Var}(\sigma_i)$ can be independently controlled by tuning H_t , the applied magnetic field in the longitudinal direction H_z , and the dilution x . Specially, for $H_t \approx 1T$ and appreciable dilution x , where $H_t > 1.5T$ and $x \approx 1$.

This gives for the first time an opportunity to study experimentally the RFIM in a FM system, in both the classical and quantum regimes. In particular, the realization of the Hamiltonian (1) in a FM system allows both the study of long-standing questions using direct bulk probes such as magnetization and susceptibility, and for

the first time an experimental study of models in which in addition to the random field a constant longitudinal field or a quantum term plays a role. Thus, the present work could be a basis for numerous new experimental possibilities, a few of which are the following: (i) The thermal and quantum phase transitions (PTs) of the RFIM at $D = 3$ can be studied in the whole T/H_t phase diagram (see Fig.1). Experiments studying the classical phase transition in 3D DAFM systems show that the RF prevents the sample from reaching equilibrium on experimentally accessible time scales [15, 16]. The study of the phase transition in diluted dipolar magnets can shed light on this interesting phenomenon, and the effect of the dipolar interaction on the theoretical predictions for the equivalence of the quantum and classical PTs in the RFIM [17, 18] can be tested as well [19]. Of particular interest is the ability, resulting from the novel quantum term, to control the relaxation rate of the system by tuning the effective transverse field. (ii) As a result of the possibility to apply tunable random field and constant field in the longitudinal direction, one could study experimentally the hysteresis in the RFIM as function of H_z . Theoretical studies of this model in the absence [20] and presence [21] of long range interactions have shown an intriguing disorder-driven out of equilibrium PT, a paradigm for the study of crackling noise in self-organized critical systems. (iii) In $D = 1; 2$ dimensions, the predictions of Imry-Ma [1] and Binder [22] for the destruction of long-range FM order [23] and the rounding of the phase transition in 2D can be verified by studying the magnetization as function of the H_t dependent effective RF.

All our analysis and results below can be easily generalized to any dipolar Ising magnet, as is explained in detail in Ref. [10]. In the following we concentrate on the $\text{LiHo}_x\text{Y}_{1-x}\text{F}_4$ compound since this enables us to present quantitative results for this system which is of prime experimental interest.

Realization of the RFIM in the $\text{LiHo}_x\text{Y}_{1-x}\text{F}_4$. Anisotropic dipolar magnets in general, and the $\text{LiHo}_x\text{Y}_{1-x}\text{F}_4$ compound in particular, are considered to be realizations of the Ising model. As a result of the crystal field anisotropy the GS is an Ising like doublet, and all but the longitudinal dipolar interactions are effectively reduced. The application of H_t induces quantum fluctuations (QF) between the Ising ground states (GSs), resulting in an effective transverse field Ising model (TFIM). For the undiluted system both the thermal and the quantum PT to the PM state were observed [24]. The dominant interaction in the $\text{LiHo}_x\text{Y}_{1-x}\text{F}_4$ system is dipolar [25] $H_{\text{dip}} = \sum_{ij} V_{ij} J_i J_j$. In the pure LiHoF_4 , the off-diagonal terms of the dipolar interaction, e.g. $J_i^z J_j^x$, are not only effectively reduced, but cancel by symmetry. However, at finite dilution, $x < 1$, this cancellation does not hold. Furthermore, in Ref. [10] it was shown that in the presence of a constant H_t , the off-diagonal terms, despite being effectively reduced, are significant since their presence changes the symmetry of the system. The system is no longer symmetric under




FIG. 2: $\sigma^2 = x^2 (E_{\text{IM}} = cV_0 x)^2$ in three dimensions is plotted as function of $N = 4L^3$ for dilutions $x = 0.6$ (blue triangles), 0.96 (black squares), and 0.996 (red circles). L is the length of the cube in unit cells. A averaging is done over 10^4 samples for each data point. Where not drawn, errorbars are smaller than the point size. Inset: $j_j = \sigma_x / N$ as function of x for $N = 4000$. $j_j = \sigma_x / N$ for $x \rightarrow 1$, with $\sigma = 4.12$.

the transformation $S_z \rightarrow S_z$, and an effective local longitudinal magnetic RF \tilde{h}_j is generated. In the regime $H_t \gg H_0 = (B/S)$

$$\tilde{h}_j = \frac{2SH_t}{0} \sum_i V_{ji}^{zx}; \quad (2)$$

see Ref. [10]. The random local energy is given by $\tilde{e}_j = \sum_i \tilde{h}_j S_i$. $0 \rightarrow 10\text{K}$ is the measure of the anisotropy, given by the excitation energy between each of the Ising GSs to its relevant excited state [10], and $S = 5/2$ is the magnitude of the effective Ho spin.

Importantly, \tilde{h}_j (2) is independent of the directions of the spins at sites i , and therefore equivalent for the spin-glass (SG) and FM regimes. However, the energy gain due to the effective RF does depend on the spin configuration. Thus, while the randomness in the energy gain in the SG regime is a result of both the random position and the random orientation of the spins, in the FM regime only the former are random. Furthermore, the RFs \tilde{h}_j are not uncorrelated. For $x \rightarrow 1$ the correlations of \tilde{h}_j at distances smaller than the typical inter-vacancy distance, are large. However, as is shown explicitly below, for all dilutions, for distances larger than the inter-vacancy distance, the typical energy gain in flipping a domain $E_{\text{IM}} = \sum_j \tilde{e}_j$ obeys the Imry-Ma behavior. In particular $\text{Var}(E_{\text{IM}}) / N$, where N is the number of sites in a domain.

Let us now calculate explicitly $\sigma^2 = \left(\sum_{ij} V_{ij}^{zx} = V_0 \right)^2 = (1/cV_0)^2 E_{\text{IM}}^2$, where $c = 2B/H_t S^2 = 0$ and V_0 is the magnitude of the dipolar interaction at distance of one unit cell along the x direction. First, note that for a single vacancy i , as a consequence of lattice symmetry,

$\sum_j V_{ij}^{zx} = 0$. This results in $\sum_{ij} V_{ij}^{zx} = \sum_{ij} V_{ij}^{zx}$, where i, j run over occupied sites and i, j over the unoccupied sites. Consider first the case of $x = 1$. Assuming that the interactions causing the RF, V_{ij}^{zx} , are nearest neighbor only, then in leading order E_{IM}^2 is proportional to the number of nearest neighbor vacancy pairs, and therefore to x^2 . In analogy to the case of dipolar interaction in a random system [26] it is expected that for the random interaction leading to the effective RF the dipolar and short range interactions would be equivalent. Thus, we expect $\text{Var}(E_{IM}) / N$ for all dilutions and $\text{Var}(E_{IM}) / x^2 N$ for $x = 1$, where correlations of three or more vacant spins can be neglected.

In Fig 2 we plot $\text{Var}(E_{IM}) = x^2$ for cubes in 3D as function of $N = 4L^3$ for $x = 0.6; 0.96; 0.996$. Here and below lengths are given in terms of lattice unit cells. The dilutions were chosen to include the cases where $x = 1$, and effective fields are strongly correlated on small distances, $x = 0.6$ where correlations due to the relative positions of three or more vacancies are significant, and $x = 0.96$ in between. For each x we obtain a good linear fit, $\text{Var}(E_{IM}) = x^2 = (\langle x \rangle)^2 N$, which corresponds to $\text{Var}(E_{IM}) = (\langle x \rangle c x V_0)^2 N$. In the inset we plot $\text{Var}(E_{IM}) / N$ as function of x for a 3D cube with $N = 4000$, showing $\lim_{x \rightarrow 0} \text{Var}(E_{IM}) / N = 0$ with $c = 4.12$.

We therefore conclude that for distances larger than the inter-vacancy distance the system is equivalent to a RFIM, with an effective uncorrelated RF

$$j = \sum_j \frac{2 B H_t S^2}{0} V_0 x; \quad (3)$$

j being a random number with $\langle j \rangle = 0$ and $\text{Var}(j) = x^2$. Similar results (not shown) were obtained for $D = 1$. Interestingly, j in Eq.(3) is similar to the RF obtained in the SG regime [10], with x replacing x and the constant prefactor being different for the two regimes due to the different spin configuration.

While our results are general to any dipolar Ising magnet, we now use the specifics of the $\text{LiHo}_x\text{Y}_{1-x}\text{F}_4$ system to show that despite the presence of H_t the analogy to the classical RFIM is possible. The Ising like anisotropy is dictated by a crystal field Hamiltonian having large J_z^2 and J_z^4 terms [25]. In addition, a large $(J_z^4 + J_z^4)$ term results in a mixing of free ion states with $J_z = \pm 4$. The electronic GSs are an Ising like doublet, denoted $|j; \uparrow; \uparrow\rangle, |j; \downarrow; \downarrow\rangle$, belonging to the $J_z = (7; 3; 1; 5)$ multiplets. Although the full splitting of the $2J + 1 = 17$ states is $\sim 600\text{K}$, the first excited state $|j; \uparrow; \downarrow\rangle$, belonging to the $J_z = (6; 2; 2; 6)$ multiplet is only $\sim 10\text{K}$ higher than the GS doublet. The electronic Hamiltonian allows a coupling between the electronic GSs which is second order in H_t . However, the low energy physics of the $\text{LiHo}_x\text{Y}_{1-x}\text{F}_4$ system is strongly affected by the strong hyperfine (hf) interaction between the Ho electronic angular momentum and its nuclear spin [9, 27]. In particular, each electronic GS is split to $2I + 1 = 8$ nearly equidistant states with $I_z = 7/2; \dots; 7/2$ and a

separation of 205mK between them [28]. Thus, the relevant Ising like single spin GS doublet is of electronuclear type, having a definite and opposite nuclear spin states, i.e. a $|j; \uparrow; \uparrow\rangle$ and a $|j; \downarrow; \downarrow\rangle$. The tunneling between the states a and a' requires the flipping of the nuclear spin, resulting in a very weak coupling at low fields [less than 10^{-4}K at $H_t < 1\text{T}$ (Ref. [27])]. Only when j and j' become well hybridized with the electronic GSs $|j; \uparrow; \uparrow\rangle, |j; \downarrow; \downarrow\rangle$, QF between the states a and a' become appreciable. This occurs at $H_t = 0 = (B S) \sim 1.5\text{T}$ (see a detailed plot of (H_t) in Fig 2 of Ref.[9]).

We first consider the low T low H_t (LTLH) regime ($H_t < 1\text{T}$ and $T \sim 0.2\text{K}$). In this regime only the states a, a' are appreciably occupied, and the quantum tunneling between these two states is negligible. Thus, the system is effectively a classical Ising model [9]. Importantly, the effective RF (3), originating from fluctuations between each Ising GS and its corresponding excited state [10, 12], can already be made appreciable, since its behavior is linear in H_t . Thus, in this regime, which is of particular interest for the study of the destruction of long range order (LRO) by the effective RF in $D = 1; 2$ (see below), and the equilibrium and nonequilibrium phenomena in the ordered state in $D = 3$, the analogy between the $\text{LiHo}_x\text{Y}_{1-x}\text{F}_4$ system and the RFIM is practically exact.

Phase transitions in three dimensions. As a result of the very different dependence of $\text{Var}(E_{IM})$ and $\text{Var}(E_{IM})$ on H_t , the thermal and quantum PTs can be well separated, as is sketched in Fig 1. Both phase transitions occur outside the LTLH regime. For the classical PT one has to go to $T > 0.2\text{K}$, since $T_c \sim x \sim 1.54\text{K}$ [8]. Therefore, all nuclear states are appreciably occupied near the transition. Still, at low enough transverse field, $H_t < 0.5\text{T}$, QF are small between any of the time reversed states $|j; \uparrow; \uparrow\rangle, |j; \downarrow; \downarrow\rangle$ [27]. Thus, the nuclear spin states can be traced over, and the classical PT is equivalent to that of the RFIM. Note, that the magnitude of the RF can be tuned by varying the dilution, and for each given sample by changing the magnitude of H_t , as can be inferred from Eq. (3).

Considering the quantum phase transition (QPT) we focus on $T \sim 0.2\text{K}$, where only the GS doublet is appreciably occupied. However, we still have to go out of the LTLH regime since strong QF are required. Specifically, in the dilutions of interest the QPT occurs at $H_t > 3\text{T}$. In Ref.[9] it was shown that QF between the electronuclear Ising states become appreciable at $H_t \sim 1.5\text{T}$, and at $H_t \sim 3\text{T}$ the GS has all hf levels well mixed [27]. However, since at $x > 0.5$ the typical magnitude of the dipolar energy $V_0 x S^2 \sim 0.2\text{K}$, it is only when the QF of the electronic spins are larger than the interaction that the PT to the PM phase takes place. At $x = 1$ it was argued that the system is equivalent to the TFIM, with the hf interactions effectively re-normalizing the transition field [25]. For $x < 1$ and $V_0 x S^2 \sim 0.2\text{K}$ we expect the same picture to be valid, with the addition of the effective longitudinal RF.

At the QPT $H_t = 0 = (J_B S)$, Eq. (2) is not valid, and QF between the Ising doublet states enhance the effective RF. The largest RFs that can be obtained are of the order of $V_0 S = J_B$. By changing the dilution x , one can therefore control the magnitude of the typical random magnetic field at the QPT in the regime $0 < h < V_0 S = J_B$, equivalent to $0 < \epsilon < J$, where J is the average nearest neighbor interaction. Thus, for $x \ll 1$ the QPT can be studied in the presence of weak RF, while for a substantial dilution the RF, as well as the fluctuations, are of the order of the interaction. Note, that LRO is destroyed when either QF or the RF are of the order of the interaction, and the situation where all three energy scales are of the same order is of both theoretical and experimental interest.

Measuring the correlation length in one and two dimensions. Two is the critical dimension for the stability of the FM phase to the RF [1, 2], as the energy gain and energy cost to flip a domain are both linear in the domain size. However, Binder [22] has shown that due to the roughening of the domain wall, a two dimensional FM Ising system in a RF has a finite correlation length $\xi^{2D} \sim \exp(V_0 S^2)^{-1}$. Here we consider first a strictly 2D system, where the plane includes the easy z -axis and is chosen as to have a FM phase at zero transverse field. We first consider an in-plane magnetic field H_x transverse to the easy axis. For $H_x = 0 = (J_B S)$ the effective RF is given by Eq. (3), and therefore, for an appreciable dilution, say $0.5 < x < 0.8$, $\xi^{2D} \sim \exp[(J_0 = J_B H_x)^2]$. Since $S \ll 1$, $\xi^{2D} > \exp(S^2)$ in the perturbative regime. As H_x is increased, QF between the Ising like doublet become significant, h increases and ξ decreases. Assuming one can reach the regime where $\xi^{2D} \sim L$, the finite correlation length can be detected by studying M as function of H_x

$$M = M_0(H_x) \frac{S}{L^2} \frac{L^2}{2} = M_0(H_x) \frac{L}{2}; \quad (4)$$

as the ratio $M = M_0$ depends linearly on L . Here $M_0(H_x)$ corresponds to the dependence of the magnetization on H_x in a FM domain due to single spin fluctuations, as can be measured in a 3D sample where the system stays FM for the relevant magnetic fields [$H_t = 0 = (J_B S)$].

Unlike systems such as the DAFM $\text{RB}_2\text{Co}_x\text{Mg}_{1-x}\text{F}_4$ [29], where the two dimensionality is due to the inherent layered structure, for the $\text{LiHo}_x\text{Y}_{1-x}\text{F}_4$ a realistic system would be quasi two-dimensional, with a finite width d . The energy

cost to flip a domain is then linear in d , while the energy gain is $\sim d$. As a result ξ is enhanced, $\xi^{2D} \sim \exp[(J_0 d = (J_B H_x)^2 d)^{-1}]$, and therefore one has to keep d small in order to observe a finite ξ . In comparison to layered systems, the observation of a finite ξ is more difficult. However, the study of the gradual cross-over between two and three dimensions is made possible. Note, that the degree of two dimensionality can be studied by measuring the dependence of the RF on the angle of H_t in the xy plane. This also allows the continuous change of the RF in a given sample. In particular, for the strict 2D case there is no RF when H_t is perpendicular to the plane.

Using simple Imry-Ma [1] arguments one can show that for a quasi-1D system parallel to the z -axis and of cross-section A , the finite correlation length is given by $\xi^{1D} A (J_0 = J_B H_t)^2$, and the magnetization by

$$M = M_0(H_t) \frac{S}{L} \frac{L}{A} = M_0(H_t) \frac{A}{L} \frac{J_0}{J_B H_t}; \quad (5)$$

Thus, the observation of $M = M_0 / L = H_t$ in the regime $J_0 A = L J_B H_t$, $J_0 = S$ is the experimental manifestation of the instability of the FM phase in one dimensional Ising system to RF, as predicted by Imry and Ma.

As explained in detail in Refs. [10, 12], our analysis above applies directly to a general anisotropic dipolar system, with a crystal field Hamiltonian e.g. of the form $H_{CF} = D S_z^2$, and with no hyperfine interactions. Interestingly, in the $\text{LiHo}_x\text{Y}_{1-x}\text{F}_4$ system, if one neglects the hyperfine interactions, the peculiar form of the crystal field Hamiltonian which allows transitions between the two electronic ground states in second order perturbation, leads to different physical results. Most importantly, the effective transverse term becomes appreciable at small H_t , and the RFIM is obtained at the expense of significant quantum fluctuations. Thus, the proper consideration of the hyperfine interactions in the $\text{LiHo}_x\text{Y}_{1-x}\text{F}_4$ system is not only crucial for obtaining the correct effective fields, but also to obtain the qualitative equivalence of the $\text{LiHo}_x\text{Y}_{1-x}\text{F}_4$ system to general diluted anisotropic dipolar magnets.

It is a pleasure to thank Gabriel Aeppli, Amnon Aharony, David Belanger, Nicolas Laorencie, and Alessandro Silva for useful discussions. This work was supported by NSERC of Canada and PITP.

-
- [1] Y. Imry and S.K. Ma, Phys. Rev. Lett. 35, 1399 (1975).
 [2] T. Nattermann, J. Phys. A 21, L645 (1988).
 [3] D.P. Belanger and A.P. Young, J. Mag. Mag. Mater. 100, 272 (1991).
 [4] D.P. Belanger in Spin Glasses and Random Fields, ed.

- A.P. Young, World Scientific (1998).
 [5] T. Nattermann in Spin Glasses and Random Fields, ed. A.P. Young, World Scientific (1998).
 [6] S. Fishman and A. Aharony, J. Phys. C 12, L729 (1979).
 [7] J.L. Cardy, Phys. Rev. B 29, 505 (1984).

- [8] M . J . P . G ingras (private communication).
- [9] M . Schechter and P . C . E . Stam p, Phys. Rev. Lett. 95, 267208 (2005).
- [10] M . Schechter and N . La orencie, Phys. Rev. Lett. 97, 137204 (2006).
- [11] S M A . Tabei, M . J . P . G ingras, Y . J . Kao, P . Stasiak, and Y . J . Fortin, Phys. Rev. Lett. 97, 237203 (2006).
- [12] M . Schechter, P . C . E . Stam p, and N . La orencie, J. Phys.: Condens. Matter 19, 145218 (2007).
- [13] T F . Rosenbaum , J. Phys. Condens. Matter 8, 9759 (1996).
- [14] A . B iltm o and P . Henelius, Phys. Rev. B 76, 054423 (2007).
- [15] R . J . Birgeneau, R . A . Cow ley, G . Shirane, and H . Yoshizawa, Phys. Rev. Lett. 54, 2147 (1985).
- [16] F . Ye, L . Zhou, S . A . M eyer, L . J . Shelton, D . P . Belanger, L . Lu, S . Larochelle, and M . G reven, Phys. Rev. B 74, 144431 (2006).
- [17] A . A harony, Y . Gefen, and Y . Shapir, J. Phys. C 15, 673 (1982).
- [18] T . Senthil, Phys. Rev. B 57, 8375 (1998).
- [19] The dipolar nature of the interaction has no effect on the validity of the Imry-Ma arguments and on the lower critical dimension, but its presence affects the critical behavior and breaks the universality of the DAFM and RFIM systems, both in their low T phases and the characteristics of the critical behavior[2].
- [20] J . P . Sethna, K . A . Dahmen, and C . R . Myers, Nature 410, 242 (2001).
- [21] J . H . Carpenter and K . A . Dahmen, Phys. Rev. B 67, 020412(R) (2003).
- [22] K . Binder, Z. Phys. B 50, 343 (1983).
- [23] M . Aizenman and J . W ehr, Phys. Rev. Lett. 62 2503, (1989).
- [24] D . Bitko, T . F . Rosenbaum , and G . Aeppli, Phys. Rev. Lett. 77, 940 (1996).
- [25] P . B . Chakraborty, P . Henelius, H . K jonsberg, A . W . Sandvik, and S . M . Girvin, Phys. Rev. B 70, 144411 (2004).
- [26] A . J . Bray, M . A . Moore, and A . P . Young, Phys. Rev. Lett. 56, 2641 (1986).
- [27] M . Schechter and P . C . E . Stam p, in preparation.
- [28] R . G iraud, W . W emsdorfer, A . M . Tkachuk, D . Mailly, and B . Barbara, Phys. Rev. Lett. 87, 057203 (2001).
- [29] I . B . Ferreira, A . R . King, V . Jaccarino, J . L . Cardy, and H . J . Guggenheim , Phys. Rev. B 28, 5192 (1983).