
A simple view on the quantum Hall system

Emil J. Bergholtz and Anders Karlhede

Department of Physics, Stockholm University
AlbaNova University Center
SE-106 91 Stockholm, Sweden

Summary. The physics of the quantum Hall system becomes very simple when studied on a thin torus. Remarkably, however, the very rich structure still exists in this limit and there is a continuous route to the bulk system. Here we review recent progress in understanding various features of the quantum Hall system in terms of a simple one-dimensional model corresponding to the thin torus.

1 Introduction

Even though more than twenty years have passed since the experimental discovery [1] of the fractional quantum Hall effect at filling factor $\nu = 1/3$ and its basic explanation due to Laughlin [2], the physics of the quantum Hall regime still continues to surprise us with new novel phenomena. Already from the beginning it was clear that the quasiparticles in the Laughlin state have fractional charge and later on it was realized that they obey fractional statistics [3, 4].

Soon after the first observations at $\nu = 1/3$ many other gapped quantum Hall states were observed, some of them at fractions that could not be explained by Laughlin's wave functions. To explain these new fractions, hierarchical schemes were developed by Haldane, Halperin and Laughlin [5, 3, 6] and Jain constructed wave functions for these states and proposed an intriguing interpretation in terms of composite fermions [7], where each of the electrons captures an even number of magnetic flux quanta, mapping the original problem of electrons partially filling a Landau level onto composite fermions filling an integer number of Landau levels. This gives a nice picture of how the gap responsible for the quantum Hall effect appears at the fractions $\nu = p/(2m p + 1)$ by mapping the system onto the well understood integer quantum Hall effect. Moreover, the composite fermion theory offers an appealing explanation for the existence of the gapless states observed at even denominator fractions such as $\nu = 1/2$, where the system is mapped onto free fermions in no magnetic field. The mean field theory of such states, due to Halperin, Lee and Read [8], has been spectacularly confirmed by surface acoustic wave experiments at $\nu = 1/2$ [9], and by ballistic experiments near this filling factor [10].

However, in our opinion, a microscopic understanding of composite fermions is still lacking [11]. Gapped quantum Hall states have now been observed that fall

outside Jain's main scheme [12], and the microscopic origin of these states is under debate. Also, in higher Landau levels quantum Hall states exist that might possess even more exotic properties. One such example is the Moore-Read state [13], which is believed to describe the quantum Hall system at $\nu = 5/2$ [14, 15, 16]. This state has attracted great interest recently due to the supposed non-abelian statistics of the quasiholes and its possible application to topologically protected q-bits (decoherence free quantum computational devices) [17].

In a recent line of research it has been shown that studying the quantum Hall system on a thin torus allows for both a simple understanding of already established results and for providing new insights [18, 19, 20, 21, 22, 23, 24]. Here, we give a non-technical review of this work. References [25, 26, 27] contain relevant precursors to the work presented here.

We study the quantum Hall system of spin-polarized electrons on a torus as a function of its circumference, L_1 , by mapping the problem onto a one-dimensional lattice model. When L_1 is small, the range of the electron-electron interaction becomes short (in units of the lattice spacing), and we get a systematic expansion of the quantum Hall system around a simple case—the thin torus. The abelian quantum Hall states are manifested as gapped one-dimensional crystals, 'Tao-Thouless states', and their fractionally charged excitations appear as domain walls between degenerate ground states. At half-filling, $\nu = 1/2$, the electrons condense into a Fermi sea of neutral dipoles which connects smoothly to the gapless state in the bulk. The non-abelian pfaffian (Moore-Read) state believed to describe the $\nu = 5/2$ phase is described by six distinct crystalline states, and the non-trivial quasiparticle and quasihole degeneracies that are crucial for the non-abelian statistics follow simply from the inequivalent ways of creating domain walls between these different vacua. This formulation is manifestly particle-hole symmetric and thus allows for the construction of both quasiparticles and quasiholes.

The outline of this paper is the following. In section 2 we set up a one-dimensional lattice model of the lowest Landau level. In section 3 we discuss how ground states and excitations have a very simple and appealing manifestation on the thin torus, and in section 4 we discuss the crucial issue of how the thin torus picture is connected to the experimentally realizable bulk system.

2 1D lattice model

The energy of a charged particle moving in a magnetic field is quantized in macroscopically degenerate Landau levels. In the strong magnetic field limit, the gap between different Landau levels becomes large and the electrons will populate the lowest available states. Hence the kinetic energy effectively freezes out, leaving a strongly interacting problem in the highest partially populated Landau level (LL). Since a single LL is an effectively one-dimensional system, it is possible to map the two-dimensional quantum Hall system onto a one-dimensional problem. It turns out that this mapping is particularly convenient on the torus.

For simplicity we consider the problem of an electron moving in a perpendicular magnetic field on the surface of a cylinder (the torus case is obtained by straight forward periodising). In Landau gauge, $A = B y \hat{x}$, the lowest Landau level states are

$$\psi_m(x) = \frac{1}{\sqrt{L_1}} e^{i m x / L_1} e^{-(y + 2 m / L_1)^2 / 2}; \quad (1)$$

where we use units such that $\hbar = 1$, $h = 1$, and label the states by integers m . The states are centered along the lines $y_m = 2 m / L_1$, given by the momentum in the x direction. This provides an explicit mapping of the two-dimensional electron gas in the lowest Landau level onto a one-dimensional lattice model, where the lines y_m can be thought of as the sites, see Figure 1.




Fig. 1. A cylinder with a magnetic field B perpendicular to its surface. The single particle states are centered along the lines $y_m = 2 m / L_1$ and can be thought of as sites in a one-dimensional lattice.

A general (two-body) interaction Hamiltonian takes the form

$$H = \sum_n \sum_{k > m} V_{km} c_{n+m}^\dagger c_{n+k}^\dagger c_n c_m; \quad (2)$$

where V_{km} are matrix elements that can be calculated for a given real-space interaction. The physics of the interaction can be understood by dividing H into two parts: V_{k0} , the electrostatic repulsion (including exchange) between two electrons separated k lattice constants, and V_{km} , the amplitude for two particles separated a distance $k - m$ to hop symmetrically to a separation $k + m$ and vice versa. The symmetry of the hopping, which is a consequence of conservation of momentum, implies that the position of the center of mass is conserved.

A general N_e particle state in the lowest Landau level is a linear combination of states characterized by the positions (or, equivalently, the momenta) at which they are centered. We represent these (Slater determinant) states in Fock space as $|n_1 n_2 n_3 \dots\rangle$ where $n_i = 0, 1$ according to whether site i is occupied or not. The problem of finding the ground state and the low lying excitations, at filling fraction $\nu = N_e / N_s$, is thus a matter of arranging N_e electrons on N_s sites.

A very important property of the obtained lattice model is that the lattice constant is $2 / L_1$. This means that, for a given real space interaction, the interaction in the one-dimensional lattice model becomes short range in units of the lattice spacing when the torus becomes thin and we can hope to be able to solve the

problem in this limit. The experimental situation, on the other hand, is obtained as $L_1 \rightarrow 1$, where the lattice model becomes infinitely long range measured in units of the lattice constant. When the system is studied as a function of L_1 , we find that many of the characteristic features of the quantum Hall system is independent of L_1 and there is a continuous route between the two extreme cases—we claim that the two cases are adiabatically connected.

3 The thin torus

Here we consider the quantum Hall system at generic filling fractions, $\nu = p/q < 1$, in the limit $L_1 \rightarrow 0$. For reasonable interactions (including Coulomb), the problem becomes a classical electrostatic one-dimensional problem and the ground states are regular lattices of electrons where the particles are as far apart as possible, as shown in Table 1.

<u>1</u> 00100100100100100 :::i	= 1=3
<u>1</u> 0100101001010010100 :::i	= 2=5
<u>1</u> 010010010010100100100 :::i	= 4=11

Table 1. Examples of ground states in the thin limit, $L_1 \rightarrow 0$. The underlined unit cells containing p electrons on q sites are periodically repeated in the $\nu = p/q$ ground state. The q -fold degeneracy on the torus is reflected by q different translations of the unit cell.

The reason that the physics is completely determined by electrostatics in the thin limit is actually rather simple. The single particle states are essentially gaussians extended roughly one unit length (*i.e.* one magnetic length) and separated by the lattice constant $2\pi/L_1$. Consequently, the overlap between different one-particle wave functions becomes very small and the only non-vanishing matrix elements are those where each electron is created and destroyed at the same site, *i.e.* the electrostatic matrix elements V_{k0} . Thus, L_1 is a parameter that controls the strength of the hopping, which can be continuously turned on by increasing L_1 .

The ground states in the thin limit are regular lattices with unit cells containing p electrons and q sites at filling $\nu = p/q$. This is true for any repulsive interaction that is monotonic, with positive second derivative—Coulomb falls into this category. The same ground states were obtained by Hubbard when he investigated generalized Wigner lattices in the seemingly very different context of quasi-one-dimensional salts [28]. It is interesting to note that, at $\nu = 1/3$, the thin limit ground state, see Table 1, is the state originally proposed by Tao and Thouless in 1983 to explain the fractional quantum Hall effect [29]. We call these states, at general filling factor, Tao-Thouless (TT) states, stressing the fact that they are different from ordinary, classical crystals or Wigner crystals. It is important to note that the TT-states have a gap to all excitations—there are no phonons. The reason for this is that once the fluxes through the holes of the torus are fixed, then the positions of the one-particle

states along the torus are fixed, and hence no vibrations of the lattice are possible. Note also that the q fold degeneracy, present for all energy eigenstates on the torus [30], is trivially manifested by the q different translations of the unit cell.

3.1 Gapped fractions and fractional charge

At odd denominator fractions in the lowest Landau level, the TT-states describe (but are extreme forms of) the gapped abelian quantum Hall states observed in the laboratory. In section 4 we discuss this connection further, but let us first consider the structure of ground states and fractional charge that emerge in the thin limit.

At the Jain fractions, $\nu = p/(2pm + 1)$, the unit cells are $10_{2m} (10_{2m-1})_{p-1}$ in chemical notation. At $\nu = 1/3$ the unit cell is 100, at $\nu = 2/9$ it becomes 100001000 and so on. These states are gapped and q fold degenerate.

The low energy excitations of the TT-states at arbitrary filling fractions are domain walls separating sequences of degenerate ground states. These domain walls carry fractional charge and correspond to the quasiparticle and quasihole excitations in the bulk.

At $\nu = 1/q$ a quasihole (quasiparticle) is constructed by inserting (removing) a zero somewhere in the ground state, see Table 2. This is very similar to Laughlin's original concept of creating a quasihole by inserting a flux quantum. At $\nu = p/(2pm + 1)$ the corresponding quasiparticle (quasihole) excitations are obtained by inserting (removing) 10_{2m-1} somewhere in the TT-state with unit cell $10_{2m} (10_{2m-1})_{p-1}$.

```
j100100100100100100100100100100100 :::i
j100101001001001010010010010100100 :::i
j100100010010010001001001000100100 :::i
```

Table 2. The $\nu = 1/3$ ground state, and the corresponding states with three quasiparticles and three quasiholes respectively. Note that the underlined concentration of electrons (or holes) are domain walls between degenerate $\nu = 1/3$ ground states. The charge ($e=3$) of these excitations is determined by Su and Schrieffer's counting argument.

The charge of these excitations is determined by Su and Schrieffer's counting argument [31]. By removing 10_{2m-1} at $2pm + 1$ separated position and adding $2m$ unit cells $10_{2m} (10_{2m-1})_{p-1}$ to keep the number of sites fixed, $2pm + 1$ quasiholes, each with charge $e = e \frac{(2pm + 1) 2pm}{2pm + 1} = \frac{e}{2pm + 1}$, are created. This readily generalizes to generic fillings $\nu = p/q$ where the lowest lying excitations naturally emerge as domain walls carrying charge

$$e = \frac{e}{q} \quad (3)$$

```
j10101001010100101001010100 :::i
j101010010101010010101001010 :::i
j1010100101001010100101010010 :::i
```

Table 3. The $\nu = 3/7$ ground state, and the corresponding states with a quasiparticle and a quasihole respectively. Note that inserting/removing 10 creates domain walls with the correct charge $\nu = 7$. (Inserting/removing 100 would instead create domain walls with charge $2\nu = 7$.)

3.2 The non-abelian pfaffian state

The single particle states differ from (1) in the higher Landau levels, thus the interaction (*i.e.* V_{km}) is different, and as a consequence, the ground states and their excitations may differ from those in the lowest Landau level. Perhaps most notably, the ground state at half-filling in the second Landau level appears to be gapped and is believed to be accurately described by the Moore-Read pfaffian state [13]. This state, which is motivated by conformal field theory, has quasihole excitations with charge $\nu = 4$ that can only be created in pairs, and obey non-abelian statistics.

Here we describe how this state is manifested on the thin torus and give the degeneracies of the quasihole excitations that are crucial for the existence of non-abelian statistics. Moreover, the particle-hole symmetry allows us to construct also quasiparticles, as well as states with general combinations of quasiholes and quasiparticles [22].

The pfaffian states on the torus are known to be the exact ground states of a hyper-local three-body potential [14, 16]. In the thin torus limit, this implies that the electrostatic energy (of this three-body potential) is minimized by separating all triples of particles as much as possible. At half-filling this means that there are no sequences of four consecutive sites containing three electrons (or holes). The six states displayed in Table 4 are the unique states at half-filling that have no such sequences.

```
j010101010101 :::i 2 translations
j001100110011 :::i 4 translations
```

Table 4. The six degenerate pfaffian ground states on a thin torus.

The extra freedom created by the additional pfaffian ground states allows for the creation of domain walls carrying charge $\nu = \nu = 4$ —*i.e.* half of the fractional charge $\nu = \nu = 2$ that is implied by the center of mass degeneracy. The domain walls that achieve this are those between the two different kinds of ground states $j10101010 :::i$ and $j11001100 :::i$, as shown in Table 5. Again this charge is readily determined by Su and Schrieffer’s counting argument. Note also that, because of the periodic boundary conditions, these excitations can only be created in pairs.

j01010 <u>1001</u> 110011 <u>0010</u> 10101 :::i	two quasiholes
j01010 <u>1011</u> 0011001 <u>1101</u> 0101 :::i	two quasiparticles
j01010 <u>1011</u> 0011001010101 :::i	a quasiparticle-hole pair

Table 5. Examples of domain walls with fractional charge $e=4$.

The degeneracy of these excitations is readily determined by considering the various ways of matching the domains. In Ref. [22] we derived that the degeneracy of a state with $2n - k$ quasiholes and k quasiparticles with fixed positions is 2^{n-1} . Results similar to ours have also been obtained by Haldane [21], and subsequently also by Seidel and Lee [23] for the closely related bosonic pfaffian state at $\nu = 1$.

3.3 The half-filled Landau level

The physics of the half-filled lowest Landau level is known to be very different from the gapped fractions discussed above. There is strong experimental and numerical evidence that the system is gapless. In the composite fermion picture, all magnetic flux is attached to the electrons and the system becomes a free Fermi gas of composite fermions in no magnetic field [7, 8, 32]. Furthermore, it has been proposed that the quasiparticles are dipoles [33, 34, 35].

In the thin limit, the $\nu = 1/2$ ground state is $|010101010\rangle$ and the (gapped) low lying excitations are the fractionally charged excitations described above. In fact, the $\nu = 1/2$ state has a larger energy gap than the $\nu = 1/3$ state on the thin torus. This is clearly different from the observed gapless state in the bulk.

In order to explain this discrepancy we consider the situation when L_1 increases from zero. Short range hopping terms will now become important and start to compete with the electrostatic terms. However, the shortest range hopping V_{21} annihilates the TT state $|01010101\rangle$. Also, from early numerical investigations it was clear that there is a sharp transition from the TT-state $|01010101\rangle$ at $L_1 = 53$ to a gapless homogeneous state [27].

It is interesting to contrast $\nu = 1/2$ with $\nu = 1/3$. At $\nu = 1/2$, the ground state is the TT-state $|j101010\rangle$ when $L_1 \rightarrow 0$. As noted, this state is annihilated by the shortest range hopping term V_{21} which favours hoppable states of the type $|j11001100\rangle$. Thus there is a competition between the electrostatic terms and the hopping term and this leads to a phase transition to a gapless state when L_1 grows. For $\nu = 1/3$ on the other hand, the TT-state $|j1001100\rangle$ is favoured by electrostatics is also a maximally hoppable state favoured by the short range hopping term. In this case there is no competition between electrostatics and hopping and there is no phase transition as L_1 grows.

We now briefly discuss a solvable model that accurately describes the system at L_1 slightly larger than 53. The low-energy sector of the model consists of free one-dimensional neutral fermions (dipoles) [18]. The crucial part in the hamiltonian turns out to be the hopping term V_{21} —the other terms can be treated as perturbations yielding an interacting Luttinger liquid.

We start with the hamiltonian

$$H = \sum_n V_{21} C_n^y C_{n+1} C_{n+2} C_{n+3}^y + H_{\text{sc}} : \quad (4)$$




Fig. 2. The evolution of the one-dimensional density $\langle n_k^y \rangle$ from the small L_1 TT-state (triangles) to the homogenous state (squares) at $\beta = 1/2$. At $L_1 = 5/3$ there is a sharp transition from the TT-state to a homogenous state that is described by our solvable model, and corresponds to a Luttinger liquid of neutral dipoles. At the transition the quantum numbers change.

This provides a good approximation of the interaction on a thin, but not infinitely thin, torus ($L_1 = 6$) as discussed in Ref. [18].

We define a subspace \mathcal{H}^0 of the full Hilbert space by requiring each pair of sites $(2p-1; 2p)$ to have charge one (the equivalent grouping of the sites $(2p; 2p+1)$ gives a trivial copy of our solution). In Ref. [18] it is argued that \mathcal{H}^0 contains the low-energy sector under fairly general conditions. It agrees with what we find in numerical studies, and \mathcal{H}^0 contains the maximally hopable state $|100110011001\rangle$. Furthermore, \mathcal{H} preserves the subspace \mathcal{H}^0 , thus any other ground state candidate may not mix with the states in \mathcal{H}^0 .

There are two possible states for a pair of sites in \mathcal{H}^0 ,

$$|j\#i\rangle = |j1i\rangle, |j\#i\rangle = |j0i\rangle \quad (5)$$

and it is natural to introduce the spin operators

$$S_p^+ = c_{2p-1}^y c_{2p}; \quad S_p^- = c_{2p}^y c_{2p-1}^\dagger \quad (6)$$

On states in $H^0, s^+; s^-$ describe hard core bosons—they commute on different sites but obey anti-commutation relations on the same site. In this subspace, H is simply the nearest neighbor spin $1/2 \times Y$ -chain,

$$H = V_{21} \sum_p (s_{p+1}^+ s_p + s_{p+1} s_p^+); \quad (7)$$

The (hard core) bosons can be expressed in terms of fermions d using the Jordan-Wigner transformation,

$$s_p = K_p d_p; \quad K_p = e^{i \sum_{j=1}^p d_j^\dagger d_j}; \quad (8)$$

and the Hamiltonian (4) is then that of free fermions.

The ground state is obtained by filling all the negative energy states. The excitations are neutral particle-hole excitations out of this Fermi sea. These excitations have a natural interpretation in terms of dipoles as is seen from (6), and in the limit $N_e \rightarrow 1$, the excitations become gapless. It is also straight forward to show that the state is homogeneous. We would like to stress that this explicitly and exactly maps (the low energy sector of) a system of strongly interacting electrons in a strong magnetic field onto a system of non-interacting particles that are neutral and hence are unaffected by the magnetic field.

By considering the relation between the real system—where the electrons interact via Coulomb repulsion—and our model, we conclude that the $\nu = 1/2$ system is a Luttinger liquid of these dipoles on a thin torus (L_1 slightly larger than 5λ). This conclusion is supported by numerical calculations for both Coulomb [19] and short range interactions [27]. Note also that the obtained solution has striking similarities to the bulk state—both are homogenous gapless states with quasiparticles (dipoles) that do not couple to the magnetic field.

4 Bulk physics

In this section we discuss how the two-dimensional bulk physics is related to the physics in the thin limit. We will argue that the abelian and non-abelian gapped states, as well as the gapless state at $\nu = 1/2$, are adiabatically connected to the states found on the thin torus. The strength of the argument varies with the filling factor but we believe the over all picture of bulk states at generic filling factor being adiabatically connected to simple ground states on the thin torus is firmly established.

Before we proceed with a more detailed account for each of the considered cases we make two important remarks: 1) The TT-states and the bulk QH-states do in fact have the same symmetries and qualitative properties. That the TT-state is not homogenous is not a result of spontaneous symmetry breaking—in fact the Laughlin/Jain states have periodic density variations on any finite torus [36]. 2) As indicated in section 3.3, there is actually a simple way of understanding why the TT-state melts at half-filling while it develops smoothly into the bulk QH-state at *e.g.* $\nu = 1/3$.

4.1 Abelian states

We begin by considering the simplest case, $\nu = 1/q$, q odd. At these filling factors the Laughlin wave functions describe the bulk physics; moreover, they are the exact and unique ground states to a short range pseudo-potential interaction and there is a gap to all excitations [37, 38]. This holds also on a torus (or cylinder) for arbitrary circumference L_1 ¹. This is fairly obvious since it depends only on the short distance property of the electron-electron interaction. In our opinion, this establishes that the ground state develops continuously as L_1 increases, without a phase transition, from the TT-state to the bulk Laughlin state for this short range interaction. This result is implicit in the work of Haldane and Rezayi [26]. The same is then very likely to be true for the Coulomb interaction—this is supported by exact diagonalization where no transition is seen as L_1 varies.

We now show that the Laughlin wave function on a cylinder

$$\Psi_{1=q} = \prod_{n < m} (e^{2iz_n - L_1} - e^{2iz_m - L_1})^q e^{\frac{1}{2} \sum_n y_n^2}; \quad (9)$$

where $z = x + iy$, approaches the TT-state as the radius of the cylinder shrinks [26]. Expanding $\Psi_{1=q}$ in powers of $e^{2iz - L_1}$ and using that the single particle states (1) can be written as $\phi_k = \frac{1}{\sqrt{L_1}} e^{2iz - L_1} e^{i y^2/2} e^{i k^2 L_1/2}$, one finds

$$\begin{aligned} \Psi_{1=q} &= \prod_{n < m} c_{f k_n g} (e^{2iz_n - L_1})^{k_n} e^{\frac{1}{2} \sum_n y_n^2} = \\ &= \frac{1}{N_e! L_1^{N_e/2}} \prod_{f k_n g} c_{f k_n g} e^{i k_n^2 L_1/2} e^{\frac{1}{2} \sum_n y_n^2}; \end{aligned} \quad (10)$$

where $c_{f k_n g}$ are coefficients that are independent of L_1 . The weight of a particular electron configuration is multiplied by the factor $e^{i k_m^2 L_1/2}$, thus in the limit $L_1 \rightarrow 0$ the term with the maximal k_m^2 will dominate (all terms have the same $\sum_n k_n$). The dominant term is the one that corresponds to the TT-state discussed above, where the electrons are situated as far apart as possible. In this case at every q th site. This argument can be generalized to the Jain wave functions describing the ground states at filling factors $\nu = p/(2m p + 1)$ showing that they approach the TT-states above as $L_1 \rightarrow 0$. It can also be generalized to show that the fractionally charged quasiparticles in the TT-state, discussed in Section 3.1, are the $L_1 \rightarrow 0$ limits of the bulk quasiparticles at filling factor $\nu = p/(2m p + 1)$.

The TT-state and the bulk Laughlin/Jain state on the torus at $\nu = p/(2m p + 1)$ have the same quantum numbers. The symmetry generators that commute with the hamiltonian are T_1 and T_2 (T translates all particles in the x -direction). The Laughlin/Jain state is an eigenstate of T_1 and $T_2^{2m p + 1}$, with quantum numbers K_1 and K_2 , whereas T_2^k ; $k = 1, 2, \dots, 2m p$ generate the degenerate states—this is true for any L_1 —and the eigenvalues are independent of L_1 . The state is inhomogeneous for any finite L_1 , although the inhomogeneity decreases very rapidly as L_1 grows.

¹ On the torus, the ground state of course has the trivial q -fold center of mass degeneracy.




Fig. 3. The evolution of the one-dimensional density $\langle n \rangle$ from the small L_1 TT-state (triangles) to the nearly homogenous bulk Laughlin state (circles) at $\nu = 1/3$. This process is smooth and the quantum numbers K remain unchanged as L_1 changes. Results are obtained from exact diagonalization of an unscreened Coulomb potential.

Furthermore, the TT-state and the Laughlin/Jain state both have a gap and have quasiparticles and quasiholes with the same fractional charge.

The conclusion is that there is no phase transition separating the TT-states and the bulk Laughlin states. This result has a long history. The very first observation was made already in 1984 by Su who discussed the TT-state as the 'parent state' of the Laughlin state and observed that the fractionally charged quasiparticles could be thought of as domain walls between the degenerate vacua. Rezayi and Haldane noted that the Laughlin state is the exact ground state for the short range interaction on a cylinder of any circumference and showed that the state approaches a crystal as $L_1 \rightarrow 0$ in 1994 [26]. More recently this was reexamined by the present authors in DMRG calculations [27, 18] and in exact diagonalization [19] and a careful numerical study of the rapid crossover from the TT-state to a virtually homogeneous state was performed by Seidel *et. al.* using Monte Carlo methods [20].

In the case of the Jain states, there is no known interaction which they are the exact and unique ground states of. However, as we have noted above they have the same qualitative properties as the corresponding TT-states: same quantum numbers, gap

and quasiparticles with the same charge. These TT-states, including quasiparticle excitations, are obtained as the $L_1 \rightarrow 0$ limits of Jain's wave functions. Furthermore, exact diagonalization of small systems show a smooth development of the ground state from the TT-state to the Jain state as L_1 grows. No transition is observed and there is a gap for all L_1 [19]. Recent progress strongly suggests that this picture is true also for more general odd denominator fractions in the lowest Landau level, such as the non-Jain state at $\nu = 4/11$; in these cases no phase transition is observed for small systems and a new set of trial wave functions connect the solvable limit to the bulk [24, 39]. We conclude that the adiabatic continuity holds also for the hierarchy states.

4.2 Non-abelian states

Recently, it has been understood that also non-abelian gapped quantum Hall states follow the same pattern as we outlined for the abelian states above [21, 22, 23].

The six Moore-Read pfaffian ground states² are the exact ground states of a hyperlocal three-body interaction on the torus—as in the case of the Laughlin states, this holds for general L_1 as it depends on the local properties only. As L_1 decreases the states continuously approach the TT-states in Table 4.

4.3 The gapless state at $\nu = 1/2$

The $\nu = 1/2$ solution on the thin torus, discussed above, has striking similarities to what is expected from theory and experiment for the bulk state. Based on this, we conjectured [18] that this state develops continuously, without a phase transition, to the bulk state as $L_1 \rightarrow 1$. This is however a much more delicate issue than it is for the states above since the state at $\nu = 1/2$ is gapless.

To investigate this conjecture, we performed exact diagonalization studies of small system for various N_e and L_1 using an unscreened Coulomb potential [19]. The obtained ground states were then compared with the Rezayi-Read state [32], that is expected to describe the bulk state, by calculating overlaps. On the torus the Rezayi-Read wave function takes the form

$$\Psi_{RR} = \det_{ij} [e^{ik_i \cdot R_j}]^{\frac{1}{2}} ; \quad (11)$$

where R is the guiding center coordinates and $[\frac{1}{2}]$ is the bosonic Laughlin state at $\nu = 1/2$. This wave function depends on a set of momenta \mathbf{k}_i/g , which determine the conserved quantum numbers K .

For $L_1 = 5/3$ the ground state is the TT-state $|10101010\rangle$. At $L_1 = 5/3$ there is a sharp transition into a new state that we identify as our Luttinger liquid solution, discussed above. As L_1 is increased further, there is a number of different transitions to new states, but these transitions are all much smoother than the one at $L_1 = 5/3$. As shown in Figure 4 for the case of nine electrons, each of these states corresponds to a given set of momenta \mathbf{k}_i/g in the Rezayi-Read state. The Fermi seas of momenta develop in a very natural and systematic way. Starting from

² There are three distinct pfaffian wave functions on the torus. This together with the two-fold center of mass degeneracy gives all the six states on the thin torus.

an elongated sea, which we identified as the exact solution, a single momentum is moved at each level-crossing, terminating in a symmetric Fermi sea when $L_1 = L_2$.




Fig. 4. 'Phase diagram' showing the ground states for $\nu = 1/2$ as a function of L_1 for nine electrons [19]. The results are obtained in exact diagonalization, using unscreened Coulomb interaction. Overlaps with the Rezayi-Read state with the displayed Fermi seas of momenta are shown above each Fermi sea.

Since our Luttinger liquid solution corresponds to one of the Fermi seas in the Rezayi-Read state and this state develops smoothly towards the bulk, we conclude that the Luttinger liquid of neutral dipoles is continuously connected to the bulk ground state.

5 Conclusions

We conclude that the thin torus provides a simple and accurate picture of both abelian and non-abelian quantum Hall states, and even more surprisingly, also of the gapless state at $\nu = 1/2$. The gapless state is particularly important since it provides an explicit microscopic example of how weakly interacting quasiparticles moving in a reduced (zero) magnetic field emerge as the low energy sector of strongly interacting fermions in a strong magnetic field.

There are strong reasons to believe that the picture presented here is valid also for other quantum Hall states. Indeed, the ground state and quasihole degeneracies of other topological states can be obtained on the thin torus [40, 21].

A one-dimensional picture of the quantum Hall system is very natural, and in some sense almost obvious. After all, a single Landau level is a one-dimensional system. The non-trivial result is, of course, that a model with an interaction that is short range in the one-dimensional sense is relevant. We believe that the evidence reviewed here establishes that this is indeed the case.

Acknowledgement. We thank Hans Hansson, Janik Kailasvuori, Emma Wikberg and Maria Hermanns for interesting discussions and fruitful collaborations. This work was supported by the Swedish Research Council and by NordForsk.

References

1. D.C. Tsui, H.L. Störmer and A.C. Gossard, Phys. Rev. Lett. **48**, 1599 (1982).
2. R.B. Laughlin, Phys. Rev. Lett. **50**, 1395 (1983).
3. B.I. Halperin, Phys. Rev. Lett. **52**, 1583 (1984), 2390(E) (1984).
4. D. Arovas, J.R. Schrieffer, and F. Wilczek, Phys. Rev. Lett. **53**, 722 (1984).
5. F.D.M. Haldane, Phys. Rev. Lett. **51**, 605 (1983).
6. R. B. Laughlin, Surf. Sci. **141**, 11 (1984).
7. J.K. Jain, Phys. Rev. Lett. **63**, 199 (1989).
8. B.I. Halperin, P.A. Lee, and N. Read, Phys. Rev. B **47**, 7312 (1993); see also V. Kalmeyer, and S.-C. Zhang, Phys. Rev. B **46**, R9889 (1992).
9. R.L. Willett, M.A. Paalanen, R.R. Ruel, K.W. West, L.N. Pfeiffer, and D.J. Bishop, Phys. Rev. Lett. **65**, 112 (1990).
10. W. Kang, H.L. Störmer, L.N. Pfeiffer, K.W. Baldwin, and K.W. West, Phys. Rev. Lett. **71**, 3850 (1993); V.J. Goldman, B. Su, and J.K. Jain, Phys. Rev. Lett. **72**, 2065 (1994); J.H. Smet, D. Weiss, R.H. Blick, G. Lütjering, K. von Klitzing, R. Fleischmann, R. Ketzmerick, T. Geisel, and G. Weimann, Phys. Rev. Lett. **77**, 2272 (1996).
11. M.I. Dyakonov, in *Recent Trends in Theory of Physical Phenomena in High Magnetic Fields*, eds. I.D. Vagner, P. Wyder, and T. Maniv, (Kluwer, 2003).
12. W. Pan, H.L. Störmer, D.C. Tsui, L.N. Pfeiffer, K.W. Baldwin, and K.W. West, Phys. Rev. Lett. **90**, 016801 (2003).
13. G. Moore and N. Read, Nucl. Phys. B **360**, 362 (1991).
14. M. Greiter, X.-G. Wen and F. Wilczek, Phys. Rev. Lett. **66**, 3205 (1991); Nucl. Phys. **B374**, 567 (1992).
15. R.H. Morf, Phys. Rev. Lett. **80**, 1505 (1998).
16. E.H. Rezayi and F.D.M. Haldane, Phys. Rev. Lett. **84**, 20 (2000).
17. See e.g. M.H. Freedman, A. Kitaev, M. Larsen and Z. Wang, Bull MAS **40**, 31 (2003); A. Kitaev, Ann. Phys. **303**, 2 (2003).
18. E.J. Bergholtz, and A. Karlhede, Phys. Rev. Lett. **94**, 26802 (2005).
19. E.J. Bergholtz and A. Karlhede, J. Stat. Mech. (2006) L04001.
20. A. Seidel, H. Fu, D.-H. Lee, J.M. Leinaas and J. Moore, Phys. Rev. Lett. **95**, 266405 (2005).
21. F.D.M. Haldane, talk at the 2006 APS March meeting, unpublished.
22. E.J. Bergholtz, J. Kailasvuori, E. Wikberg, T.H. Hansson and A. Karlhede, Phys. Rev. B **74**, 081308(R) (2006).
23. A. Seidel and D.-H. Lee, Phys. Rev. Lett. **97**, 056804 (2006).
24. E.J. Bergholtz, T.H. Hansson, M. Hermanns, and A. Karlhede, arXiv:cond-mat/0702516v1 (2007).
25. W.P. Su, Phys. Rev. B **30**, 1069 (1984).
26. E.H. Rezayi and F.D.M. Haldane, Phys. Rev. B **50**, 17199 (1994).
27. E.J. Bergholtz and A. Karlhede, cond-mat/0304517, (2003).
28. J. Hubbard, Phys. Rev. B **17**, 494 (1978).
29. R. Tao and D.J. Thouless, Phys. Rev. B **28**, 1142 (1983).
30. F.D.M. Haldane, Phys. Rev. Lett. **55**, 20 (1985).
31. W.P. Su, and J.R. Schrieffer, Phys. Rev. Lett. **46**, 738 (1981).
32. E.H. Rezayi and N. Read, Phys. Rev. Lett. **72**, 900 (1994).
33. D.-H. Lee, Phys. Rev. Lett. **80**, 4745 (1998), Erratum **82**, 2416 (1999).
34. N. Read, Phys. Rev. B **58**, 16262 (1998).

- 35. V. Pasquier and F.D.M. Haldane, Nuclear Physics B **516**, 719 (1998).
- 36. F.D.M. Haldane and E. H. Rezayi, Phys. Rev. B **31**, 2529 (1985).
- 37. F.D.M. Haldane in the *The Quantum Hall Effect*, eds. R.E. Prange and S.M. Girvin, (Springer-Verlag, New York, 1990).
- 38. S.A. Trugman and S. Kivelson, Phys. Rev. B **31**, 5280 (1985).
- 39. T.H. Hansson, C.C. Chang, J.K. Jain, and S. Viefers, Phys. Rev. Lett. **98**, 076801 (2007).
- 40. N. Read, Phys. Rev. B **73**, 245334 (2006).