

Evidence of Luttinger liquid behavior in one-dimensional dipolar quantum gases

R. Citro,¹ E. Orignac,² S. De Palo,³ and M. L. Chiofalo⁴¹Dipartimento di Fisica "E. R. Caianiello" and CNISM, Università degli Studi di Salerno, Salerno, Italy²Laboratoire de Physique de l'École Normale Supérieure de Lyon, CNRS-UMR 5672, Lyon, France³DEMOCRITOS INFN-CNR and Dipartimento di Fisica Teorica, Università Trieste, Trieste, Italy⁴Classe di Scienze, INFN and CNISM, Scuola Normale Superiore, Pisa, Italy

The ground state and structure of a one-dimensional Bose gas with dipolar repulsions is investigated at zero temperature by a combined Reptation Quantum Monte Carlo (RQMC) and bosonization approach. A non trivial Luttinger-liquid behavior emerges in a wide range of intermediate densities, evolving into a Tonks-Girardeau gas at low density and into a classical quasi-ordered state at high density. The density dependence of the Luttinger exponent is extracted from the numerical data, providing analytical predictions for observable quantities, such as the structure factor and the momentum distribution. We discuss the accessibility of such predictions in current experiments with ultracold atomic and molecular gases.

PACS: 03.75.Hh, 71.10.Pm, 02.70.Ss

The realization of Bose-Einstein condensation (BEC) in trapped ultracold quantum atomic gases [1] is at the frontier of modern atomic and molecular, optical and condensed-matter physics [2]. Especially fascinating experimental results arise from the possibility of tuning the atomic interactions. Use of Fano-Feshbach resonances [4] to change magnitude and sign of the s-wave scattering length characterizing the contact interactions, has allowed e.g. the observation of collapsing Bose condensates [5] and of the crossover from a BEC to a Bardeen-Cooper-Schrieffer-like transition [6].

More recent experiments have demonstrated that the range of the interactions can also be manipulated. Dipole interactions with long-range anisotropic character have been observed in ^{52}Cr atoms [7] after exploiting the large magnetic moments of this atomic species, that is μ_B with μ_B being the Bohr magneton. A BEC containing up to 50000 ^{52}Cr atoms has then been obtained below a transition temperature $T_c \approx 700\text{nK}$ [8] and its dynamical behavior is being investigated [9]. Promising proposals to tune and shape the dipolar interaction strength in quantum gases of heteronuclear polar molecules have more recently been suggested [10]. Significant theoretical predictions have accompanied such realizations [11]. The stability diagram of anisotropic confined dipolar gases has been predicted to be governed by the trapping geometry [12, 13], as corroborated by Path-Integral QMC studies [14]. Different conclusions are reached by more recent Diffusion QMC simulations including the dependence of α on the dipole interaction [15].

Tuning of the interactions can be combined with the enhancement of quantum fluctuations after reducing their dimensionality by e.g. storing them in elongated traps [16, 17], which could be relevant to applications such as precision measurements [18], quantum computing [19], atomtronic quantum devices, and theoretical investigations of novel quantum phase transitions [20].

In the case of quasi-one-dimensional (1D) condensates with short-range interactions, a rich phenomenology is

known to emerge from the collective character of the single-particle degrees of

freedom, despite the absence of broken symmetries [21]. Bosons are known to arrange in a Luttinger-liquid state, with single particles being replaced by collective density excitations [22, 23]. Strong repulsion may also lead to the fermionization of interacting bosons in the so-called Tonks-Girardeau (TG) regime [24, 25, 26]. Experiments in elongated traps have provided evidence for such 1D situations [16].

In the case of quasi-1D condensates with dipolar interactions, an interesting question arises whether the quantum fluctuations are sufficiently enhanced to drive the BEC in a strong-coupling regime. More recent Diffusion QMC simulations [27] for a homogeneous 1D dipolar Bose gas have revealed a crossover behavior with increasing linear density, from a liquid-like superfluid state to an ordered, normal, state with particles localized at lattice sites. For low values of the density, the system is seen to behave as a Tonks-Girardeau gas [24].

A consistent theoretical description of the whole crossover region would be useful to predict the behavior of observable quantities in the experiment. It is well known [22] that the low-energy properties of a one-dimensional system of interacting bosons can be obtained from an effective quadratic Hamiltonian within a Luttinger-liquid theory, provided that the interaction is sufficiently short ranged. Besides, 1D fermionic models with interactions falling off as $1/x$ have been studied in the past by many authors [35, 36], concluding that for $\alpha > 1$ the long-range part of the interaction decays rapidly enough to preserve the linear energy dispersion $\epsilon(k) = u|k|$ typical of short-range models.

In this Letter we provide evidence for a robust Luttinger-liquid behavior of a 1D dipolar Bose gas in the crossover region, evolving into a Tonks-Girardeau gas at low density and into a classical quasi-ordered state at high-density. We reach this conclusion by comparing theoretical results from a bosonization approach against Reptation Quantum Monte Carlo simulations [28]. By

analyzing the scaling behavior of the static structure factor expected for a Luttinger liquid, we find small Luttinger exponents at intermediate-to-high density values, signalling strong repulsions at short distances. The knowledge of Luttinger parameters is then exploited to determine observable properties of the fluid in a parameter range accessible to experiments. To the best of our knowledge this is the first demonstration of such strongly correlated liquid behavior in atomic Bose gases with dipolar interactions [29].

We consider N atoms with mass M and permanent dipoles moments arranged along a line. For the purposes of the present work, we assume $\epsilon \rightarrow 0$ after e.g. exploiting a Fano-Feshbach resonance. In effective Rydberg units $Ry = \hbar^2/(2M r_0^2)$ the Hamiltonian is

$$H = \sum_i \frac{p_i^2}{2m} + \sum_{i < j} \frac{1}{r_{ij}^3} \frac{1}{r_{ij}^3} \mathbf{A}_i \cdot \mathbf{A}_j; \quad (1)$$

where we have defined the effective Bohr radius $r_0 = M C_{dd}/\hbar^2$ in terms of the interaction strength C_{dd} . One has $C_{dd} = \mu_0^2 d^2$ and $C_{dd} = d^2/\epsilon_0$ for magnetic and electric dipoles respectively, where μ_0 and ϵ_0 are the magnetic and electric dipole moments and μ_0 and ϵ_0 are the vacuum permittivities. The dimensionless parameter governing the system is $r_s = 1/(n r_0)$, with n being the linear particle density and $r_s r_0$ the unit length.

We analyse the low-energy structural properties of the fluid. They can be accessed in experiments with atomic gases by means of e.g. Bragg scattering techniques. Here, we infer them from QMC numerical simulations of the static structure factor $S(k)$ on a box with size L and periodic boundary conditions. In terms of the density operator \hat{n} this is

$$S(k) = \frac{1}{L} \int_0^L dx e^{ikx} \langle \hat{n}(x) \hat{n}(0) \rangle; \quad (2)$$

An accurate size-scaling analysis of $S(k)$ is needed to our purposes. To this aim we resort to the Reptation Quantum Monte Carlo of Baroni and Moroni [28], which in essence is a path-integral technique at zero temperature giving direct access to the ground-state wavefunction. This makes the evaluation of the ground-state properties conceptually simple and the extraction of the correlation functions in imaginary time immediate [33], with the additional possibility of determining the excitation spectrum. We use a trial wave-function made up of a two-body Jastrow factor $\psi_{\text{trial}}(\mathbf{R}) = \sum_{i < j} \exp[u(\mathbf{r}_i - \mathbf{r}_j)]$. Removal of the divergences in $H_{\text{trial}} = H \psi_{\text{trial}}$ due to the potential energy in the $x \rightarrow 0$ limit, is obtained after imposing the cusp condition, which yields $u(x \rightarrow 0) = r_s/x$ [34].

The QMC data for $S(k)$ are reported in Fig. 1 for $N = 40$ and different values of $n r_0 = 0.01; 50; 100$ and 1000. Free fermion-like behavior, typical of the




FIG. 1: $S(k)$ in dimensionless units for a dipolar gas with $N = 40$ particles and different values of $n r_0 = 0.01; 50; 100$ and 1000. Decreasing slopes as $k \rightarrow 0$ and the emergence of additional peaks correspond increasing $n r_0$ values.

Tonks-Girardeau regime, is seen at low density values ($n r_0 = 0.01$ in the Fig. 1). Quasi-Bragg peaks emerge at reciprocal lattice vectors $k = n = 2\pi m$ (m integer). Their number increases with increasing $n r_0$, evidencing the occurrence of a quasi-ordered state. Investigation of the QMC data at different values of N show that the peak heights scale as N^{-m} with $m < 1$ being an exponent depending on m , while their intensity diverges with the system size.

Both these characteristics are reminiscent of Luttinger-liquid behavior and therefore we turn on analysing the data within a bosonization description. In terms of the conjugate bosonic operators ϕ and θ , the Hamiltonian for a Luttinger liquid reads [21]

$$H = \hbar \frac{dx}{2} u K (\partial_x \phi)^2 + \frac{u}{K} (\partial_x \theta)^2; \quad (3)$$

where K is the Luttinger exponent and u the sound velocity. Knowledge of K and u allows analytical expressions for physical quantities of interest. The structure factor $S(k)$ can be analytically calculated inserting in Eq. (2) the bosonized expression of the density operator \hat{n} , that is $\hat{n}(x) = \frac{1}{2\pi} \partial_x \phi + \sum_{m \neq 0} A_m e^{2im(\phi(x) - nx)}$ [21, 22], where A_m are non-universal dimensionless constants. In dimensionless units for k , $S(k)$ turns out to be $S_0(k) = K k/(2\pi)$ for $k \rightarrow 0$. For k close to $k = 2\pi m$ on the scale of the cutoff parameter Λ , we find [37]:

$$S(k) = \sum_{m \neq 0} A_m^2 n^2 \frac{1}{m} (k - 2\pi m); \quad (4)$$

where $\frac{1}{m}(k)$ is explicitly given as

$$\frac{1}{m}(k) = L \left(1 - e^{-2\pi i k} \right)^{2m^2 K} (k; m^2 K; L) {}_2F_1 \left(m^2 K; m^2 K + \frac{1}{2} \right) (2\pi i k; 1 + \frac{1}{2} \pi i k; L) e^{-\frac{4\pi i k}{L}}; \quad (5)$$

where $(k; m^2 K; L) = \prod_{j=1}^{\infty} (1 + \frac{k}{j})^{m^2 K} [(\frac{k}{j})^{m^2 K} (1 + \frac{k}{j})^{-1}]^{\frac{1}{2}} {}_2F_1$ the hypergeometric function.

The low- k linear behavior of $S(k)$ can be fitted to extract the Luttinger K at different densities. The resulting $K(n)$ is displayed in Fig. 2. The inset shows $S(k \rightarrow 0)$ for selected densities, where decreasing slopes correspond to increasing values of $nr_0 = 50; 100; 1000$, the symbols representing the RQMC data and the lines the Luttinger prediction $S(k \rightarrow 0) = K k^{2m}$. K remains below the Tonks-Girardeau limit in a wide range of densities, signalling the occurrence of strong correlations. As $nr_0 < 1$, K tends to the Tonks-Girardeau limit $K = 1$. In the opposite high-density limit, K tends to vanish following the classical-theory prediction $K(n \rightarrow 1) = (6/3)n_0^{-1/2}$.




FIG. 2: Luttinger exponent K obtained from fitting the RQMC data for $S(k)$ in the $k \rightarrow 0$ limit. For comparison, the curves $K(n \rightarrow 0) = 1$ and $K(n \rightarrow 1) = (6/3)n_0^{-1/2}$ are also displayed as dashed lines, corresponding to the Tonks-Girardeau $nr_0 \rightarrow 0$ limit and to the classical prediction for $nr_0 \rightarrow 1$. The solid line is a guide to the eye. The inset zooms on the low- k portion of $S(k)$ at selected nr_0 (see text).

$m(k)$ acquires a simplified form depending on whether $m^2 K < 1/2$ or $> 1/2$ as $L \rightarrow \infty$. For $< 1/2$ we get $m(k) \sim L^{1-2m^2 K} \sin(m^2 K) (K; L)$ with $(K; L) \sim (1 - 2m^2 K) (m^2 K + kL = 2) = (1 - m^2 K + kL = 2)$, indeed revealing the presence in $S(k)$ of quasi-Bragg peaks at $|k| = 2m$ and a scaling behavior in the vicinity of the peak. The peak intensities diverge with the system size L and their number increases with decreasing K (namely increasing n). The peak's height scales as $N^{1-2m^2 K}$. For $> 1/2$ instead we get $m(k \rightarrow 0) \sim 2^{-2} [(2 - 1)] [(k/2)^{-1}]$ and the peaks disappear.

Besides the linear k behavior, evidence of a Luttinger liquid comes also from the scaling behavior of $S(k)$. Fig. 3 displays the comparison between the RQMC $S(k)$ close to $k = 2$ towards the quasi-ordered region with $nr_0 = 50$, and the predictions of the Luttinger-liquid theory Eqs. (4)-(5) with $m = 1$. We actually plot $S(k) = S(2 + n)$ for different N values and verify that they indeed collapse

on a single curve. This signature of Luttinger-liquid behavior against the RQMC data is robust in the whole crossover region. This is the central result of the present work. Similar results are indeed found at the remaining density values above $nr_0 = 1$, while the scaling is absent below $nr_0 = 1$. The same scaling analysis holds for the second peak at $k = 4$ when it is present. At $k = 6$ ($m = 3$), we have $2m^2 K > 1$ for $K < 0.055$ and the divergent peak is seen to show up in the simulation only for the data set with $nr_0 = 1000$, as expected.

The peak's asymmetry visible in Fig. 3 results from band curvature effects. Asymmetric structure factors are for example obtained within exact calculations on Calogero-Sutherland models [32] which indeed possess a non-linear spectrum, and could be also viewed as anharmonic phonon effects [38].




FIG. 3: $S(k) = S(2 + n)$ in n^1 units vs. $N(k/2 - 2)/40$ at $nr_0 = 50$ for different N values (see legend). Symbols represent RQMC data, the solid line is the Luttinger-liquid (LL) prediction.

As a consistency check, we have verified that the K values obtained from the RQMC $S(k)$ agree with those determined from the RQMC energy per particle, as follows. Since Hamiltonian (1) is invariant under Galilean boosts [22], we have $uK = \hbar n/m$. Within the bosonization procedure [21, 22], the compressibility is $\chi = K (\hbar u)^{-1}$. Comparing this result with the definition of χ in terms of density derivatives of ϵ , that is $\chi^{-1} = n^2 (n(n))^{(0)}$, we obtain $K = \hbar n / [n(n)^{(0)}]$. This result stems from the trivial fact that the exact RQMC satisfies the compressibility sum-rule. However, we have also demonstrated on a purely formal basis that our Luttinger-liquid scheme satisfies the sum-rule too.

The knowledge of $K(n)$ allows us to determine other observables such as the momentum distribution $n(k)$, which can be accessed in current atomic BEC's experiments via e.g. the analysis of time-of-flight images. Since in the continuum the boson creation operator is represented by $\psi_B^\dagger(x) = \exp[i \int^x dy \phi(y)]$ [21], we find that $n(k) \sim k^{1-2K}$ as $L \rightarrow \infty$. As the density increases, the

divergence of $n(k)$ at $k = 0$ is expected to be gradually reduced until it disappears for $K < 1/2$.

In conclusion, we have shown that dipolar gases in a reduced 1D geometry are promising candidates to observe a strong-coupling Luttinger-liquid behavior with $K < 1$. The Luttinger-liquid behavior is robust over a wide range of nr_0 values, evolving into a Tonks-Girardeau gas at low density and into a classical quasi-ordered state at high-density. Further testing of the Luttinger-liquid behavior can be provided by the excitation spectrum in the crossover [39] where, as from preliminary data, the same combination of bosonization and RQM C techniques can be successfully applied. Observation of the Luttinger liquid in the crossover region is within the reach of current and future experiments in elongated traps. The effective 1D character of trapped dipolar gases is governed by the condition $n^{1/2} l_{\perp} \gg \lambda_{\perp}$, with $\lambda_{\perp} = (4a_{\perp})^{-1/2} a_z$, ensuring the suppression of forward scattering [10]. Values of the transverse oscillator length $a_{\perp} = \sqrt{\hbar/(M \omega_{\perp})}$ related to the harmonic angular frequency ω_{\perp} can be pushed to 50 nm. Then, dipolar SrO molecular gases in elongated traps are promising candidates, as we might have $l_{\perp} \approx 0.2$ nm while $\lambda_{\perp} \approx 240$ nm, yielding access to nr_0 values up to the quasi-ordered regime. For dipolar ^{52}Cr gases instead we can have $l_{\perp} \approx 31$ nm while $r_0 \approx 4.8$ nm, yielding access to nr_0 values remaining down in the Tonks-Girardeau regime. If ultracold SrO dipolar quantum gases were to be created in such quasi-1D conditions, we predict the observation of a power-law behavior of the peak heights in the structure factor and a flattening of the momentum distribution with increasing nr_0 as fingerprints of Luttinger-liquid behavior.

We thank Dr. Paolo Pedri for very useful suggestions on the quasi-1D realizations. RC, SDP and MLC thank Prof. G. C. La Rocca for useful discussions. SDP is indebted to Dr. S. Moroni for helpful discussions and QMC support.

-
- [1] M. J. Anderson et al., *Science* 269, 198 (1995); K. B. Davis et al., *Phys. Rev. Lett.* 75, 3969 (1995); C. C. Bradley et al., *ibid.* 75, 1687 (1995).
 - [2] Bose-Einstein Condensation in Atomic Gases, Proc. Intl School of Physics "Enrico Fermi", Course CXL, M. Inguscio, S. Stringari, and C. Wieman Eds. (IOS Press, Amsterdam, 1999).
 - [3] H. Feshbach, *Ann. Phys.* 5, 357 (1958); U. Fano, *Phys. Rev.* 124, 1866 (1961).
 - [4] S. Inoué et al., *Nature* 392, 151 (1998).
 - [5] J. Roberts et al., *Phys. Rev. Lett.* 86, 4211 (2001); T. Weber et al., *Science* 299, 232 (2003).
 - [6] C. A. Regal et al., *Phys. Rev. Lett.* 92, 040403 (2004); M. Bartstein et al., *Phys. Rev. Lett.* 92, 120401 (2004); M. W. Zwierlein et al., *Phys. Rev. Lett.* 92, 120403 (2004); T. Bourdel et al., *Phys. Rev. Lett.* 93, 050401 (2004); M. W. Zwierlein et al., *Nature* 435, 1047 (2005).
 - [7] J. Stuhler et al., *Rev. Lett.* 95, 150406 (2005).
 - [8] A. Griemser et al., *Phys. Rev. Lett.* 94, 160401 (2005).
 - [9] S. Giovanazzi et al., *Phys. Rev. A* 74, 13621 (2006).
 - [10] H. P. Büchler et al., *cond-mat/0607294* (2006).
 - [11] See e.g. M. Baranov et al., *Phys. Scripta T* 102, 74 (2002).
 - [12] L. Santos et al., *Phys. Rev. Lett.* 85, 1791 (2000); K. Góra et al., *Phys. Rev. A* 61, 051601(R) (2000).
 - [13] The use of time-dependent external fields for stability has been suggested by S. Giovanazzi et al., *Phys. Rev. Lett.* 89, 130401 (2002).
 - [14] K. Nho et al., *Phys. Rev. A* 72, 023615 (2005).
 - [15] D. C. E. Bortolotti et al., *cond-mat/0604432* (2006).
 - [16] D. Hellweg et al., *Acta Phys. Pol. B*, 73, 781 (2001); A. G. Orłitz et al., *Phys. Rev. Lett.* 87, 130402 (2001); S. Richard et al., *Phys. Rev. Lett.*, 91, 010405 (2003).
 - [17] For a collection of papers: Quantum Gases in low dimensions, Pricoupenko et al. Eds., *J. Physique IV* 116 (2004).
 - [18] See e.g. M. A. Kasevich, *Science* 298, 1363 (2002); I. Canalisotto et al., *Phys. Rev. Lett.* 95, 093202 (2005).
 - [19] See e.g. I. Bloch, *Nature Physics* 1, 23 (2005) and references therein.
 - [20] K. Góra et al., *Phys. Rev. Lett.* 88, 170406 (2002); H. Pu et al., *Phys. Rev. Lett.* 87, 140405 (2001); H. Pu et al., *cond-mat/0203066* (2003).
 - [21] See e.g. T. Giamarchi, in *Quantum Physics in one Dimension*, (Oxford University Press, Oxford, UK, 2004).
 - [22] F. D. M. Haldane, *Phys. Rev. Lett.* 47, 1840 (1981).
 - [23] H. Moritz et al., *Phys. Rev. Lett.* 91, 250402 (2003); M. A. Cazalilla, *J. Phys. B: At. Mol. Opt. Phys.* 37, S1 (2003); D. Petrov et al., *J. de Phys. IV* 116, 3 (2004).
 - [24] M. Girardeau, *J. Math. Phys.* 1, 516 (1960).
 - [25] T. D. Schultz, *J. Math. Phys.* 4, 666 (1963).
 - [26] On the possible realization of the TG regime see B. Paredes et al., *Nature* 429, 277 (2004).
 - [27] A. S. Arkhipov, G. E. Astrakharchik, A. V. Belikov, and Yu. E. Izozov, *JETP Lett.* 82, 39 (2005).
 - [28] S. Baroni and S. Moroni, *Phys. Rev. Lett.* 82, 4745 (1999).
 - [29] Similar results are known in lattice models with hard-core [30] or continuum models with $1=r^2$ repulsions [31, 32].
 - [30] G. Gomez-Santos, *Phys. Rev. Lett.* 70, 3780 (1993); *ibid.* 73, 1053(E) (1994).
 - [31] F. Calogero, *J. Math. Phys.* 10, 2191 (1969); *ibid.* 2197 (1969); B. Sutherland, *J. Math. Phys.* 12, 246 (1971); *ibid.* 251 (1971).
 - [32] E. R. Mucciolo, B. S. Shastri, B. D. Simons, and B. L. Altshuler, *Phys. Rev. B* 49, 15197 (1994).
 - [33] S. De Palo, S. Conti and S. Moroni, *Phys. Rev. B* 69, 035109 (2004).
 - [34] We obtain the same results also using either the Luttinger form $u(x)/\sin(x=L)$ or gaussians centered on the lattice sites as trial wavefunctions.
 - [35] Y. T. Sukamoto and N. Kawakami, *J. Phys. Soc. Jpn.* 69, 149 (1999).
 - [36] H. Inoue and K. Nomura, *J. Phys. A* 39, 2161 (2006).
 - [37] For further details we refer to E. Ognac et al., in preparation.
 - [38] H. J. Schulz, *Phys. Rev. Lett.* 71, 1864 (1993).
 - [39] S. Giovanazzi et al., *cond-mat/0411678* (2004) predict a roton spectrum within a quasi-1D mean-field calculation.