

Conductance fluctuations in disordered superconductors with broken time-reversal symmetry near two dimensions

S. Ryu,¹ A. Furusaki,² A. W. W. Ludwig,^{3,1} and C. Mudry⁴

¹Kavli Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106, USA

²Condensed Matter Theory Laboratory, RIKEN, Wako, Saitama 351-0198, Japan

³Department of Physics, University of California, Santa Barbara, CA 93106 USA

⁴Paul Scherrer Institute, CH-5232 Villigen PSI, Switzerland

(Dated: April 15, 2024)

We extend the analysis of the conductance fluctuations in disordered metals by Altshuler, Kravtsov, and Lerner (AKL) to disordered superconductors with broken time-reversal symmetry in $d = (2 + \epsilon)$ dimensions (symmetry classes C and D of Altland and Zirnbauer). Using a perturbative renormalization group analysis of the corresponding non-linear sigma model (NL σ M) we compute the anomalous scaling dimensions of the dominant scalar operators with $2s$ gradients to one-loop order. We show that, in analogy with the result of AKL for ordinary, metallic systems (Wigner-Dyson classes), an infinite number of high-gradient operators would become relevant (in the renormalization group sense) near two dimensions if contributions beyond one-loop order are ignored. We explore the possibility to compare, in symmetry class D, the $\epsilon = (2 - d)$ expansion in $d < 2$ with exact results in one dimension. The method we use to perform the one-loop renormalization analysis is valid for general symmetric spaces of Kahler type, and suggests that this is a generic property of the perturbative treatment of NL σ Ms defined on Riemannian symmetric target spaces.

I. INTRODUCTION

A. Conductance fluctuations and the non-linear sigma model

The non-linear sigma model (NL σ M) provides, as first proposed by F. Wegner,¹ an efficient framework within which to formulate (Anderson-type) localization transitions of noninteracting quantum mechanical particles subject to (static) random potentials. This provides some of the simplest models of electrons in disordered solids. In a seminal piece of work, Altshuler, Kravtsov and Lerner (AKL)^{2,3} showed in the 1980's how to incorporate the important phenomenon of conductance fluctuations^{4,5,6} into this framework. Within a perturbative renormalization group (RG) analysis of the weakly coupled NL σ M in $d = 2$ (or $d = 2 + \epsilon$) spatial dimensions, i.e., in the regime of large (dimensionless) conductance g , these authors stressed the importance of terms in the low-energy effective action of the NL σ M which possess an arbitrary large number ($= 2s$) of spatial derivatives but do not violate any symmetries. These terms are called "high-gradient operators" and are generalizations of the kinetic term in the NL σ M which has two derivatives ($2s = 2$). High-gradient operators are highly irrelevant in the RG sense by "power counting" (i.e., without incorporation of fluctuation corrections). Specifically, AKL and Yudson studied the anomalous (RG) dimensions of these operators to one-loop order in the fluctuation corrections in $d = 2$ and in the $\epsilon = (d - 2)$ -expansion for $d > 2$.^{2,3,7,8} It was found that (i): these operators would become strongly relevant (in the RG sense) for a large enough number $2s$ of derivatives, in spite of their strong irrelevance by power counting, if contributions beyond one-loop order are ignored, and that (ii): they would (consequently) dominate the

cumulants of the conductance of sufficiently large order $2s$, in a sample of (large) linear size L . Based on these observations AKL were led to argue that for large mean (dimensionless) conductance g , the main weight of the probability distribution of the conductance arises from a narrow peak at $g = \bar{g}$, while there exist anomalously long tails (close to being log-normal). The latter arise from the high moments of the conductance, which are dominated by the above-mentioned high-gradient operators appearing in the low-energy effective action. Upon approaching the regime of stronger disorder (smaller mean conductance g), and in particular upon approaching a metal-insulator transition, the probability distribution becomes increasingly broad. AKL find similar behavior for the probability distributions of other observables, including, e.g., the local density of states (LDOS), current densities, and long-time tails of relaxation currents. Very recently,^{9,10} the basic observations of AKL for the LDOS were confirmed in a chiral symmetry class (non-enclature of Refs. 11 and 12), and were analyzed more completely in this case.

Ten years ago, in another seminal piece of work, Zirnbauer,¹¹ and Altland and Zirnbauer¹² (AZ) provided a general classification scheme for the behavior of non-interacting quantum mechanical particles subject to random potentials, on length scales much larger than the mean free path in terms of a total of ten symmetry classes.¹³ In every realization of disorder, the Hamiltonian has the same symmetries as a corresponding ensemble of random matrices. This encompasses the three previously known so-called "Wigner-Dyson symmetry classes" relevant for the physics of Anderson localization of electrons in disordered solids (corresponding to "orthogonal, unitary, and symplectic" random matrix ensembles).¹⁴ Altland and Zirnbauer's exten-

sion includes four novel symmetry classes describing the (Anderson-like) localization physics of non-interacting Bogoliubov-De Gennes (BdG) quasiparticles within a mean-field treatment of pairing in disordered superconductors. (Four symmetry classes arise since $SU(2)$ spin-rotational, or time-reversal invariance may be present or absent.) As parameters are varied, the wave functions of the (BdG) quasiparticles may be extended or localized due to disorder, and (Anderson localization type) transitions between such phases may occur. Since BdG quasiparticles do not carry a conserved electric charge, the difference between phases of localized and extended quasiparticle wave functions manifests itself not in the electrical conductivity, but instead in the thermal conductivity, and if spin is conserved, also in the spin conductivity κ_{spin} .¹⁵ In complete analogy with the fluctuations of the electrical conductance in a finite sample of disordered metals (described by the Wigner-Dyson symmetry classes), the system of BdG quasiparticles also exhibits fluctuations of the appropriate conductance in a finite-size sample. These are the fluctuations of the thermal conductance (divided by temperature T), or of the spin conductance. Henceforth, for notational convenience, we will refer to these quantities simply as "the conductance", and denote the corresponding "dimensionless conductance" again by the symbol g .

Just as for disordered metals (i.e., for the Wigner-Dyson symmetry classes), the NL M also provides an efficient framework to formulate (Anderson-type) localization physics of BdG quasiparticles in superconductors. In this article we focus on systems where time-reversal symmetry is broken in every realization of disorder: the corresponding AZ symmetry classes are conventionally called "class D" if $SU(2)$ spin-rotational invariance is broken and "class C" if it is conserved, in every disorder realization.¹⁶ In parallel, and for comparison, we will also discuss here localization physics of ordinary electrons in the absence of time-reversal symmetry, which is described by the Wigner-Dyson symmetry class conventionally called "unitary class" or "class A". (See also Table I.) When disorder averaging is implemented using fermionic replicas, the target manifolds of the resulting NL M s are the symmetric spaces $O(2N)=U(N)$ for symmetry class D,¹⁷ and $Sp(N)=U(N)$ for symmetry class C.¹⁸ We remind the reader that for the familiar unitary symmetry class A, the corresponding target space¹⁹ is $U(2N)=U(N) \times U(N)$. (In all three cases the number N of replicas is taken to zero at the end of calculations.)

The aim of this paper is to investigate perturbatively, in the spirit of the original work of AKL for ordinary metals, the conductance fluctuations for the novel AZ symmetry classes D and C, i.e., for BdG quasiparticles in superconductors lacking time-reversal symmetry, in the absence (class D) or presence (class C) of spin-rotational symmetry. Our main results are briefly summarized in Sec. IB below. We also point out that it might be possible in the case of symmetry class D to make contact with exact nonperturbative results obtained in $d = 1$ dimen-

RM T	T	SU (2)	NL M	target space	#	N
D	no	no		$O(2N)=U(N)$	+1	N
C	no	yes		$Sp(N)=U(N)$	1	N
A	no			$U(p+q)=U(p) \times U(q)$	0	$p+q$

TABLE I: Fermionic replica target spaces of the NL M describing weakly disordered superconductors and metals when time-reversal symmetry (T) is broken. Symmetry classes D, C, and A refer to the symmetric spaces defined by the RM T limits of the NL M. The number of replicas N is N for the two BdG symmetry classes whereas it is $p = q = N$ for the unitary symmetry class. The index $\# = 0; 1$ distinguishes the behavior of the beta function for the NL M in symmetry classes D, C, and A.

sions from the so-called Dorokhov-Mello-Pereyra-Kumar (Fokker-Planck) analysis²⁰ of the probability distribution of the conductance.

We end this section by recalling, as an aside, that the NL M s in all three symmetry classes, D, C, and A, share certain essential features associated with time-reversal symmetry breaking. First, in $d = 2$ dimensions, all three target manifolds allow for a topological term (a term that does not modify the equations of motion), the so-called "theta- or P nruisken term". Its presence is responsible for the appearance of thermal, spin, or charge Hall insulating phases, and corresponding plateau transitions in $d = 2$ dimensions.^{18,21,22,23,24} Second, the target manifold for symmetry class D stands out in that it is not simply connected.^{11,25,26} Consequently, there are certain discrete Z_2 domain wall excitations that may give rise to a rich phase diagram for certain types of disorder.^{23,27,28} In this paper we shall ignore the physics associated with such Z_2 domain walls, which requires an extension of the NL M description in symmetry class D. These features are not present in certain realizations of this symmetry class, on which we focus in this paper. Finally, all three target spaces have in common that they are Kähler manifolds from a geometrical point of view. This means that each target space defines a complex manifold endowed with a Hermitian metric derived from a Kähler potential.^{29,30} This property will allow us to perform the perturbative RG analysis in a unified way for symmetry classes D, C, and A. (See Sec. ID for a brief summary.)

B. Summary of main results

We begin with the summary of our main results by quoting the standard one-loop result^{32,36} for the (RG) beta function³³

$$\beta(t) = -t + (N - \#)t^2 + O(t^3) \quad (1.1)$$

in $d = 2 + \nu$ spatial dimensions, obeyed by the dimensionless coupling constant t of the relevant NL M [defined in (2.1b) below] whose inverse is proportional to the mean dimensionless conductance g , once the replica




FIG. 1: Infrared (IR) renormalization group (RG) flows for symmetry classes C and D after taking the replica limit depending on whether dimensionality of space d is larger or smaller than 2. The open circle at $t = 0$ is the metallic fixed point. The filled circle at $t = jj$ is the metal-insulator critical point in symmetry class C whereas it describes a stable fixed point for symmetry class D. Up to one loop, $t = jj$ where $jj = d - 2$ is positive for symmetry class C while it is negative for symmetry class D.

limit $N \rightarrow 0$ has been taken. To treat the symmetry classes D, C, and A on equal footing, we have introduced the replica index N that takes the value N for symmetry classes D and C and $p + q$ for symmetry class A. The index $\#$ is also needed to distinguish the one-loop beta function for the NL M corresponding to symmetry classes D ($\# = +1$), symmetry classes C ($\# = -1$), and symmetry classes A ($\# = 0$). The correspondence between the pair $(\#; N)$ and the symmetry classes D, C, and A is summarized in Table I. The beta function in Eq. (1.1) possesses a trivial fixed point,

$$t = 0; \quad (1.2a)$$

which is infrared (IR) stable for $d = 2 + \gamma > 2$ (i.e., $\gamma > 0$), while it is IR unstable when $d = 2 + \gamma < 2$ (i.e., $\gamma < 0$). Upon taking the replica limit $N \rightarrow 0$ this describes a disusive metallic phase. Whenever $0 < \gamma = (N - \#) < 1$, there is a nontrivial zero t of the beta function (1.1)^{34,35,36,37} at

$$t = \frac{\gamma}{N - \#} + O(\gamma^2); \quad (1.2b)$$

For symmetry classes C and A this describes in the replica limit and in $d = 2 + \gamma > 2$ dimensions, a metal-insulator transition separating a metallic from an insulating phase. For symmetry class D, on the other hand, this describes (in the replica limit) a stable fixed point ("critical phase") in $d = 2 - \gamma < 2$ dimensions. The IR flows for symmetry classes C and D are summarized in Fig. 1. A possible evolution of the stable fixed point in symmetry class D for dimensionality $d = (2 - \gamma)$ between one and two dimensions, to be discussed in the following subsection, is sketched in Fig. 2.

In this paper we find the dominant anomalous scaling dimension $x^{(s)}$ among all high-gradient operators con-




FIG. 2: Possible evolution of the stable fixed point for symmetry class D in $d = 2 - \gamma < 2$ dimensions between $d = 1$ and $d = 2$ (after taking the replica limit). In this sketch it is assumed that the zero $t = jj + O(\gamma^2)$ of the beta function, found to one-loop order in the expansion in $d = 2 - \gamma$ dimensions, defines a curve of stable critical points which interpolates between the metallic phase in $d = 2$ dimension (at zero coupling $t = 0$), and the behavior of the corresponding symmetry class D theory in $d = 1$ dimension, which can be solved exactly.

taining $2s$ gradients, to one-loop order. These are

$$x^{(s)} = 2s - 2(s^2 - s)t + O(t^2) \quad \text{for } U(p+q) = [U(p) \times U(q)]; \quad (1.3a)$$

$$x^{(s)} = 2s - 2(s^2 - s)t + O(t^2) \quad \text{for } Sp(N) = U(N); \quad (1.3b)$$

$$x^{(s)} = 2s - (s^2 - s)t + O(t^2) \quad \text{for } SO(2N) = U(N); \quad (1.3c)$$

when evaluated in the vicinity of the zero-coupling fixed point (1.2a) where t is small. At the corresponding fixed points in $d = 2 + \gamma$ dimensions, Eq. (1.2b), these anomalous scaling dimensions become zero, consequently [for generic numbers of replicas, $(p; q)$ or N]

$$x^{(s)} = 2s - 2(s^2 - s) \frac{\gamma}{p+q} + O(\gamma^2) \quad \text{for } U(p+q) = [U(p) \times U(q)]; \quad (1.4a)$$

$$x^{(s)} = 2s - 2(s^2 - s) \frac{\gamma}{N+1} + O(\gamma^2) \quad \text{for } Sp(N) = U(N); \quad (1.4b)$$

$$x^{(s)} = 2s - (s^2 - s) \frac{\gamma}{N-1} + O(\gamma^2) \quad \text{for } SO(2N) = U(N); \quad (1.4c)$$

Equations (1.3a) and (1.4a) were derived a long time ago by Lerner and Wegner.^{38,39} Equations (1.3) and (1.4) are, otherwise, new and are the main results of this paper.

Taking the replica limit of Eq. (1.4) yields the dominant anomalous scaling dimensions at the nontrivial fixed points of the localization problems in $d = 2 + \gamma$ dimensions.³⁴ In particular, in symmetry class C one obtains, in the replica limit, the one-loop anomalous scaling dimensions

$$x^{(s)} = 2s - 2(s^2 - s) + O(\gamma^2) \quad (1.5a)$$

of dominant $2s$ -gradient operators at the metal-insulator transition in $d = 2 + \epsilon > 2$ dimensions ($\epsilon > 0$). On the other hand, in symmetry class D in $d = 2 - \epsilon < 2$ dimensions, the dominant $2s$ -gradient operators have the anomalous scaling dimensions

$$x^{(s)} = 2s - (s^2 - s)j + O(\epsilon^2) \quad (1.5b)$$

at the stable, non-trivial fixed point (in the replica limit).

C. Discussion

When taken at face value, the one-loop expressions in Eqs. (1.3), (1.4), and (1.5) would appear to imply that the anomalous scaling dimension $x^{(s)}$ always turns negative for sufficiently large s .⁴⁰ If this behavior were to persist when all higher orders in ϵ are included, an infinite number of high-gradient operators would be relevant at the nontrivial fixed points of these NLMs, i.e., in principle many coupling constants would have to be re-tuned to reach these nontrivial fixed points. The perturbative treatment of these NLMs would then appear to contradict the belief that the underlying microscopic models undergo continuous phase transitions whose approach obeys single-parameter scaling laws. Of course, higher order contributions in ϵ to these anomalous scaling dimensions, not computed in the present or earlier work, may well change this property and could render the actual anomalous scaling dimensions $x^{(s)}$ larger than the spatial dimension $d = 2 + \epsilon$, thereby making them irrelevant.^{41,42,43,44,45,46,47} [Indeed, similar behavior is known to occur in other random disordered systems (see e.g. Refs. 42 and 47). { See also the discussion in Ref. 46.}]

Nonperturbative properties of the metal-insulator transitions in $d = 3$ dimensions have been obtained numerically for the most part.^{48,49,50,51} They support the one-parameter scaling hypothesis for metal-insulator transitions in $d = 3$ dimensions, whose critical behaviors are presumed to be described by the non-trivial fixed point of the replica limit of the NLM in the corresponding symmetry classes. Whether the apparent breakdown of one-parameter scaling due to the relevance of high-gradient operators at one loop is an artifact of the $2 + \epsilon$ expansion or has a deeper meaning remains an outstanding problem for the description of Anderson localization using the NLM approach.

Symmetry classes D and D III, on the other hand, may perhaps offer a somewhat different opportunity for the following reason. On the one hand, there exists a critical phase described by a stable fixed point in $d = 2 - \epsilon < 2$ dimensions whose properties are accessible perturbatively within the ϵ expansion. These include the properties of the conductance fluctuations and of the high-gradient operators discussed above. On the other hand, a critical phase of these models is also known to exist in $d = 1$ dimensions. Therefore, one might expect that the results obtained from the ϵ expansion in $d = 2 - \epsilon < 2$ dimensions would eventually, when $\epsilon \rightarrow 1$, turn into the

results obtained from the exact solutions of the models in $d = 1$ dimensions, as is schematically depicted in Fig. 2. If this was the case, then one could "test" the properties of the high-gradient operators and the nature of the conductance fluctuations obtained from them following the work of AKL, against exact results in $d = 1$ dimensions. Of course, as with all ϵ expansions, one would not be able to predict exactly the actual numerical values of critical exponents directly at $\epsilon \rightarrow 1$. However, based on experience with the ϵ expansion in, say, ordinary Landau-Ginzburg theory, one might expect that the structure of the entire probability distribution of the conductance would evolve, in some sense, "smoothly" with ϵ , possibly all the way down to $d = 1$ dimensions ($\epsilon = 1$). Let us now pursue this possibility.

Using perturbation theory of the NLM, AKL demonstrate that in general (i.e., for any NLM) there are two contributions to the s th cumulant of the (dimensionless) conductance.² (Below, we write the corresponding expressions appearing directly at the stable fixed point in $d = 2 - \epsilon < 2$ dimensions.) The "standard" contribution reads⁵²

$$\langle \text{hg}^s \rangle_{\text{std}} / (g)^{2s} \quad (1.6)$$

where $\langle \text{hg} \rangle$ is the mean conductance. However, when a high-gradient operator with $2s$ derivatives is added to the action of the NLM with coupling constant z_s , then there is an additional contribution

$$\langle \text{hg}^s \rangle_{\text{add}} / z_s \frac{L^{d - x^{(s)}}}{\nu} \quad (1.7)$$

to this cumulant. Here, L is the linear system size and ν the characteristic microscopic length scale (the "mean free path") for the disorder. We see that it is when $d - x^{(s)} > 0$ that the additional contribution in Eq. (1.7) dominates over the standard contribution in Eq. (1.6). This hence happens precisely when the high-gradient operator with $2s$ derivatives becomes a relevant operator in the action of the NLM.

Now, AKL argue that, if one was to take the one-loop expressions for the anomalous scaling dimensions of the high-gradient operators at face value, which for symmetry class D would amount to the result in Eq. (1.5b) derived by us in this paper, then for cumulants of sufficiently higher order $s > \frac{2}{j} [1 + O(\epsilon)]$ the additional contribution would dominate over the "standard" one, yielding

$$\langle \text{hg}^s \rangle \sim A_s \frac{L^{d - x^{(s)}}}{\nu}; \quad s > \frac{2}{j} [1 + O(\epsilon)] \quad (1.8)$$

as $L \rightarrow \infty$, where A_s are certain amplitudes. Following AKL, the lower cumulants of order $s < \frac{2}{j} [1 + O(\epsilon)]$, on the other hand, would be determined by Eq. (1.6). In summary, the reasoning of AKL implies, when specialized to the stable fixed point in symmetry class D in $d =$

2 $j < 2$ spatial dimensions, the following form of the conductance cumulants

$$\ln g^{(s)} \sim \begin{cases} (g)^{2^{-s}}; & \text{if } s < s_0; \\ A_s (L^{-1})^{d \times (s)}; & \text{if } s > s_0; \end{cases} \quad (1.9a)$$

where

$$s_0 = \frac{2}{j} [1 + O(\epsilon)] \quad (1.9b)$$

and

$$d \times (s) = j \binom{2}{s} (2 + j)s + (2 - j)j + O(\epsilon^2); \quad (1.9c)$$

Let us now discuss the known exact results for the $d = 1$ dimensional case. Specifically, the leading dependence of all cumulants of the conductance of a very long but thick quasi one-dimensional wire of length L , believed to be described by a NL M in $d = 1$ dimensions, has been extracted exactly from the DMPK approach for symmetry classes D and D III.^{53,54,55} For both symmetry classes, these cumulants decay algebraically with system size,

$$\ln g^{(s)} / (L^{-1})^{1-2s}; \quad s = 1, 2, \dots; \quad (1.10)$$

with the same exponent for all cumulants (ϵ is again the mean free path). Moreover, the logarithm of the conductance exhibits the behavior

$$\ln \ln g / (L^{-1})^{1-2s}; \quad \text{var} \ln g \sim L^{-1}; \quad (1.11)$$

Thus, the typical conductance exhibits stretched exponential behavior, in contrast with the mean conductance. This indicates a broad conductance distribution. For symmetry class D III the result (1.10) for the mean conductance has been reproduced by Lamcraft, Simons, and Zimbauer,⁵⁶ using the one-dimensional NL M (which is a principal chiral model for symmetry class D III) and instanton methods.]

If one assumes that the exact results in $d = 1$ describe the end point (at $\epsilon = 1$) of the line of critical points in $2 - j$ dimensions (see Fig 2), there are then two possibilities. First, the one-loop relevance of high-gradient operators is an artifact of the ϵ -expansion and these operators are in fact irrelevant once all orders in the ϵ -expansion are taken into account (see Refs. 42,46). Second, this is not the case and high-gradient operators are truly relevant. In this case, however, deducing the scaling of the cumulants based on the one-loop anomalous scaling dimensions does not appear to be sufficient to determine their scaling behavior. Indeed, in this case the running coupling constants z_s of all the high-gradient operators in the effective action of the NL M grow rapidly, more so the higher the number $2s$ of gradients. Under these conditions, then, there is no rationale to neglect the non-linear coupling between the running coupling constants z_s of the high-gradient operators as was done to reach

the conclusion (1.9), when extracting the scaling of the cumulants of the conductance. In effect, one needs to implement a functional RG analysis which can treat an infinite set of running (and increasing) coupling constants on equal footing. We have performed such an analysis (on which we will report elsewhere) for symmetry class A, but have so far been unable to solve the resulting RG equations. Since in the present case the coupling constants z_s of the high-gradient operators all become large, it is clear (as already mentioned) that they will not flow independently of each other in the IR limit, but will strongly mix with each other under the RG flow. In this situation, their flow does not need to result in the breakdown of single parameter scaling. Indeed, a precisely analogous situation is encountered when computing the local DOS of the two-dimensional random hopping problem,¹⁰ in which case one-parameter scaling for the local DOS is recovered using a functional RG analysis.

D. Summary of the method of calculation

The RG scheme that we adopt in this paper is the covariant background field method.⁵⁷ In Refs. 44,45 this method was used to compute, up to two loop order, the RG flows of the high-gradient operators for the $O(N) = O(N - 1)$ NL M, relevant to statistical models such as the classical Heisenberg magnet. The covariant background field method consists of two key steps. First, the fields of the NL M are separated into short wavelength ("fast") and long wavelength ("slow") degrees of freedom where the slow degrees of freedom are defined as the solutions to the classical equations of motion. Second, the fast fields are integrated out. When using this method, no redundant operator⁵⁸ is generated in the second step of the covariant background field method, a great advantage when one needs to consider mixing of many scaling operators by the RG transformation (as we do here). Another advantage is that explicit invariance under a coordinate reparametrization of the group manifold is maintained, allowing for a geometrical formulation at all stages of the calculation.

In this paper, we shall also extend the covariant background field method to the cases when the target spaces of the NL M are Riemannian or Kähler supermanifolds.⁵⁹ We derive a master formula, Eq. (4.3), in the Kähler case for the RG flow of high-gradient operators, all of which possess a vanishing covariant derivative. [See Eq. (C 19) for the super Riemannian⁵⁹ and Eq. (C 40) for the super Kähler⁵⁹ cases, respectively. To obtain the corresponding formula for the non-supersymmetric cases (for fermionic replicated NL M, i.e., for compact NL M) we set all factors $(-1)^p$ to $+1$, where $p = 0; 1$ is here the grade in the supersymmetric expression].⁶⁰ This master formula is expressed solely in terms of the geometrical data of the target manifold and should apply to all 10 symmetry classes of AZ,^{11,12} but is not limited to them. The generality of the master formula suggests that high-

gradient operators are generically relevant to 1-loop order at the nontrivial fixed point of a NL M defined on a Riemannian target space.

E. Geometric interpretation of high-gradient operators

All high-gradient operators that enter the computation of the cumulants for the conductance can be expressed in terms of tensor fields on the target manifold. We will show that high-gradient operators together with the metric and curvature tensors on the target manifolds of the NL M form a family of tensors with vanishing covariant derivative.

F. Outline of the paper

This paper is organized as follows. Section II introduces the NL M on Kähler manifolds for disordered superconductors with broken time-reversal symmetry as well as for a disordered metal in the standard unitary symmetry class. Section III describes how to use the covariant background field method to deduce perturbative RG flows. The dominant anomalous scaling dimensions for high-gradient operators are computed in Sec. IV to one-loop order. We relegate to appendices some key intermediary steps in the derivation of Eq. (1.3). The master formula, Eq. (4.3), which expresses the RG transformation law obeyed by high-gradient operators solely in terms of the geometrical data of the target manifold is derived in appendix C for any Riemannian or Kählerian supermanifold on which the high-gradient operators have vanishing covariant derivatives.

II. NL M FOR WEAKLY DISORDERED SUPERCONDUCTORS

A. NL M on Kähler manifolds

We start from the NL M description for the symmetry classes D, C and A of Anderson localization in terms of fermionic replicas. Alternatively, one can use either bosonic replica or supersymmetric methods to perform the disorder average. These three options are equivalent as far as perturbative RG computations are concerned. The field entering the NL M is a map from the base space, a d -dimensional Euclidean space R^d , to the target manifold M defined by the pattern of symmetry breaking. The target manifold has a well-defined natural metric g . A classification of all the target manifolds of NL M relevant to the problem of Anderson localization applied to non-interacting quasi-particles were identified in the seminal work of Zimbauer, and Altland and Zimbauer.¹¹ For the cases of interest in this paper, i.e., disordered superconductors with broken time-reversal symmetry, the

target spaces of the relevant NL M turn out to be Kähler manifolds which are the Hermitian symmetric spaces $SO(2N) = U(N)$ for symmetry class D and $Sp(N) = U(N)$ for symmetry class C.⁶¹ The target space of the NL M that describes a disordered metal when time-reversal symmetry is broken is the Hermitian symmetric space $U(2N) = [U(N) \times U(N)]$, which is isomorphic to the complex Grassmannian $G_{2N, N}(C)$.³⁰

Within the mesoscopic community, the usual representation of the NL M in symmetry classes D, C, and A is in terms of matrices belonging to the relevant symmetric spaces, "the Q-matrix representation" after the seminal work by Efetov et al. in Ref. 19. This representation emphasizes the pattern of symmetry breaking but quickly becomes cumbersome when performing an RG analysis of the flow of composite operators. We opt instead to emphasize more directly the geometrical properties of the target spaces, namely the fact that they are examples of Kähler manifolds for symmetry classes D, C, and A. We refer the reader to appendix A for a relation between the representation of the NL M that we shall introduce below and the more common Q-matrix representation thereof.

We now present the definition of a NL M on a Kähler manifold starting from a NL M on a $2M$ -dimensional Riemannian manifold^{30,31} (M, g) ,

$$Z = \int_{\mathcal{Z}} D[\psi] e^{S[\psi]}; \quad (2.1a)$$

$$S[\psi] = \frac{1}{4} \int_{\mathcal{Z}} g_{AB}(\psi) \partial^A \psi^B \partial^A \psi^A; \quad (2.1b)$$

Our conventions are here the following. We reserve the greek alphabet to label the d coordinates r where $R = 1; \mathbb{R}^d; d$ of $r \in R^d$, and use the short-hand notation $\int_{\mathcal{Z}} = \int_{\mathbb{R}^d} d^d r = \int_{\mathbb{R}^d} d^d r = \int_{\mathbb{R}^d} d^d r$ for the integration over R^d with the short-distance cut-off a . The dimensionless coupling constant of the NL M is here denoted by t . The dimensionless conductance shares with $(a^d t)^{-1}$ the same engineering dimension. To each point r from the base space we have assigned a point p on the manifold M with the coordinates $(A) \in R^{2M}$. In the definition of the NL M on a Riemannian manifold, Eq. (2.1), the $2M \times 2M$ components of the metric tensor field g of rank $(0;2)$ are represented by (g_{AB}) . We reserve the capital latin alphabet to label the $2M$ coordinates $A = 1; \dots; 2M$ of $p \in M$.

The components of the metric tensor field g_{AB} on the Riemannian manifold $(M; g)$ are real and symmetric under the exchange of the indices

$$g_{AB} = g_{BA}; \quad A, B = 1; \dots; 2M; \quad (2.1c)$$

There is a distinction between upper/lower capital latin indices. The components of g^{-1} are denoted by g^{AB} and satisfy

$$g_{AC} g^{CB} = \delta_A^B; \quad g^{AC} g_{CB} = \delta^A_B; \quad (2.2)$$

and so on for $A, B, C = 1; \dots; 2M$. On the other hand, there is no distinction between upper and lower greek indices and we will always choose them to be lower indices.

A Hermitian manifold (M, g) is a $2M$ -dimensional Riemannian manifold endowed with a rank $(1;1)$ tensor field J that is defined globally on M and is called a complex structure. On a Hermitian manifold, we can always make a coordinate transformation from (y^a) to $(x^a; y^{\bar{a}})$ that satisfies, for each point $p \in M$,

$$J_p \frac{\partial}{\partial x^a} = \frac{\partial}{\partial y^{\bar{a}}}; \quad J_p \frac{\partial}{\partial y^{\bar{a}}} = \frac{\partial}{\partial x^a}; \quad (2.3)$$

where $a = 1; \dots; M$. It is then natural to introduce complex coordinates $(z^a; \bar{z}^{\bar{a}})$ through

$$z^a = x^a - iy^{\bar{a}}; \quad \bar{z}^{\bar{a}} = x^a + iy^{\bar{a}}; \quad a = 1; \dots; M; \quad (2.4)$$

A NL M on a Hermitian manifold (M, g) is thus defined by the partition function

$$Z = \int D[z; \bar{z}] e^{-S[z; \bar{z}]}; \quad (2.5a)$$

$$S[z; \bar{z}] = \frac{1}{2} \int g_{ab}(z; \bar{z}) \partial z^b \partial \bar{z}^{\bar{a}}; \quad (2.5b)$$

where we have assigned to each point r from the base space a point p on the manifold M with the coordinates $(z^a; \bar{z}^{\bar{a}}) \in \mathbb{C}^M$. Components of tensor fields on M have now two types of indices, holomorphic $(a; b; \dots)$ and antiholomorphic ones $(\bar{a}; \bar{b}; \dots)$. Holomorphic indices can be contracted with holomorphic components of the coordinates z^a whereas antiholomorphic ones with $\bar{z}^{\bar{a}}$, the complex conjugate of z^a . In the definition of the NL M on a Hermitian manifold, Eq. (2.5), the $M \times M$ components of the Hermitian metric tensor field g of rank $(0;2)$ are represented by (g_{ab}) . We reserve the latin alphabet to label the M holomorphic and antiholomorphic coordinates z^a and $\bar{z}^{\bar{a}}$ where $a; \bar{a} = 1; \dots; M$, respectively, of $p \in M$. The use of a capital latin letter on a Hermitian manifold, say A , as an index denotes either an holomorphic ($A = a$) or an antiholomorphic ($A = \bar{a}$) index.

The components of the metric tensor field g_{AB} on the Hermitian manifold (M, g) are symmetric under the exchange of the indices and Hermitian

$$g_{AB} = g_{BA}; \quad (g_{AB}) = g_{A \bar{B}} \quad (2.5c)$$

with $A = a$ or \bar{a} , $B = b$ or \bar{b} , $A = a$ or \bar{a} , and $B = b$ or \bar{b} running from 1 to M . The components of g^{-1} are denoted by g^{AB} and satisfy

$$g_{AC} g^{CB} = \delta_A^B; \quad g^{AC} g_{CB} = \delta^A_B; \quad (2.6)$$

and so on for $A; B; C = 1; \dots; M$.

A Kähler manifold (M, g) is a Hermitian manifold whose Hermitian metric can be derived from a Kähler potential $K(z; \bar{z})$,

$$g_{ab}(z; \bar{z}) = \partial_a \partial_{\bar{b}} K(z; \bar{z}); \quad (2.7)$$

with $a; \bar{b} = 1; \dots; M$. Not all Hermitian manifolds are Kählerian.

The Kähler manifolds corresponding to the Hermitian symmetric spaces $U(p+q) = [U(p) \times U(q)]$ for symmetry class A, $Sp(N) = U(N)$ for symmetry class C, and $SO(2N) = U(N)$ for symmetry class D are specified once the corresponding Kähler potentials have been constructed. To this end, we begin by parametrizing the complex Grassmannian $U(p+q) = [U(p) \times U(q)]$ for symmetry class A by the stereographic coordinates defined by the independent set

$$f'_{Aa}; f'_{\bar{A}\bar{a}}; \quad A = 1; \dots; p; \quad a = 1; \dots; q; \quad (2.8a)$$

of $p+q$ complex-valued matrices.^{62,63,64} With this coordinate system, the Kähler potential on $U(p+q) = [U(p) \times U(q)]$ is given by

$$K(f'; f') = c \ln \det [I_q + f' f'] \quad (2.8b)$$

where c is an arbitrary constant (the scale of the metric) which is chosen to be unity in the following. (Another choice for c just amounts to a rescaling of t .) The Kähler metric induced by the Kähler potential (2.8) is

$$g_{(Aa)(Bb)}(f'; f') = Z_{ab} Y_{BA} \quad (2.9a)$$

where we have introduced $q \times q$ and $p \times p$ Hermitian matrices,

$$Z_{ab} = [I_q + f' f']^{-1}_{ab}; \quad a; b = 1; \dots; q; \quad (2.9b)$$

$$Y_{AB} = [I_p + f' f']^{-1}_{AB}; \quad A; B = 1; \dots; p; \quad (2.9c)$$

The cases of $Sp(N) = U(N)$ for symmetry class C and $SO(2N) = U(N)$ for symmetry class D are obtained by setting $p = q = N$ and by imposing the constraint

$$f' + \# f'^T = 0 \quad (2.10)$$

on the complex Grassmannian $U(p+q) = [U(p) \times U(q)]$ where $\# = 1$ for $Sp(N)$ (symmetry class C) and $\# = +1$ for $SO(2N)$ (symmetry class D). Due to the constraint (2.10), independent entries of $f'_{Aa}; f'_{\bar{A}\bar{a}}; A; a = 1; \dots; N$ can be chosen to be those with the indices $A = a$ for $Sp(N)$ (symmetry class C) and $A < a$ for $SO(2N)$ (symmetry class D), say. In turn, the metric for $Sp(N) = U(N)$ (symmetry class C) and the metric for $SO(2N) = U(N)$ (symmetry class D) are given by

$$g_{(Aa)(Bb)}(f'; f') = \delta_{Aa} \delta_{Bb} S_{(Aa)(Bb)}^{\#} Z_{ab} Z_{AB}; \quad (2.11a)$$

where

$$\delta_{Aa} = 1 \quad \delta_{Aa} = 2; \quad (2.11b)$$

$$S_{(Aa)(Bb)}^{\#} = S_{(Aa)}^{\#} S_{(Bb)}^{\#}; \quad (2.11c)$$

$$S_{(Aa)}^{\#} T_A = \delta_a^A T_A \quad \# T_a; \quad (2.11d)$$

and

$$Z_{ab} = [I_N + f' f']^{-1}_{ab}; \quad a; b = 1; \dots; N; \quad (2.11e)$$

Observe that Z in Eq. (2.9b) and Y in Eq. (2.9c) are not independent anymore because $Y = Z^T$ when $p = q = N$ and the constraint (2.10) holds. By a slight abuse of notation we will sometimes refer to the complex Grassmannian case by $\# = 0$.

B. High-gradient operators

The seminal contribution by Al'tshuler, Kravtsov, and Lerner,² was to observe that the perturbative RG flow obeyed by the cumulants of the conductance necessarily involves an infinite number of composite operators and to identify a criterion for the selection of this infinite family of composite operators. The selection criterion is that the flow must be closed within a family of composite operators characterized by the most dominant scaling dimensions. Applied to the three cases at hand, we thus seek a family of composite operators that transform as scalars under reparametrization of the target manifold and contain as many derivatives as possible subject to the condition that their anomalous scaling dimension be dominant. We are led to the family of high-gradient operators

$$T_{IJK} \dots (z; z) \left\{ \frac{\partial z^I \partial z^J \partial z^K}{z^{2s}} \right\}; \quad (2.12)$$

where $T_{IJK} \dots (z; z)$ is a tensor on the target manifold with s holomorphic and antiholomorphic indices; we do not yet specify how the greek indices must be contracted. Although composite operators containing derivatives of the form $\partial^n z$ ($n \geq 2$) are generated by the RG flow for the cumulants of the conductance they are neither expected to contribute to the dominant RG anomalous scaling dimensions,³⁸ nor needed to close the RG flows of the high-gradient operators (2.12) as we shall see below.

As the simplest example of a high-gradient operator belonging to the family (2.12) is the Lagrangian $g_{IJ} \partial z^J \partial z^I$ of the NLM action. This is in fact the only scalar under both reparametrization of the target manifold and global rotations in the base space with $s = 1$ that belongs to the family (2.12). To see this, observe that the number of independent tensors with two indices on the target manifold is related to that of independent bilinear forms in the group (coset). For simple Lie groups (and their analogues for cosets), there is only one such bilinear form (the Casimir), and hence g_{IJ} is uniquely fixed. The scaling flow obeyed by $g_{IJ} \partial z^J \partial z^I$, controls the scaling flow obeyed by the mean conductance, i.e., the first cumulant of the conductance once the replica limit has been taken.

Life already becomes more complicated if we seek all the high-gradient operators with $s = 2$ in the family (2.12). On a generic complex manifold more than 2 independent tensors of rank (0;4) are available,

$$g_{IJ} g_{KL}; R_{IJKL}; R_{IJMN} R^{MN}_{KL}; \quad (2.13)$$

and so on, where R_{IJKL} is the curvature (Riemann) tensor on the target manifold. Fortunately, the situation is simpler for the Grassmannian since we only need to contend with two independent tensors of rank (0;4),

$$S_{(Aa)(Bb)(Cc)(Dd)}^{(2)} = Z_{ab} Y_{BC} Z_{cd} Y_{DA}; \quad (2.14a)$$

$$S_{(Aa)(Bb)}^{(1)} S_{(Cc)(Dd)}^{(1)} = Z_{ab} Y_{BA} Z_{cd} Y_{DC}; \quad (2.14b)$$

[See Eq. (1.9).]

Observe that $S_{AB}^{(1)}$ is nothing but the metric tensor g_{AB} .

For symmetry class A and for given $s \geq N$, all the independent tensors $T_{IJK} \dots$ of rank (0;2s) of the complex Grassmannian can be expressed by taking the products of the building blocks,

$$S_{(Aa)(Bb)(Cc)(Dd)}^{(1)} \dots = Z_{ab} Y_{BC} Z_{cd} Y_{DE} \dots Z_{ZA} Y \quad (2.15a)$$

made of l pairs of Z and Y , according to the decomposition

$$T_{I_1 J_1 K_1 \dots} = \prod_i Y_{I_1 J_1 K_1 \dots} S_{I_1 J_1 K_1 \dots}^{(l_i)}; \quad \sum_i l_i = s; \quad (2.15b)$$

For the cases of $Sp(N) = U(N)$ (symmetry class C) and $SO(2N) = U(N)$ (symmetry class D) we need to replace Eq. (2.15a) by

$$S_{(Aa)(Bb)(Cc)(Dd)}^{(1)} \dots = C_c S_c^\# Y_{Bc} Z_{Cd} \dots; \quad (2.15c)$$

Another simplification attached to the Kahler manifolds for symmetry classes A, C, and D is that contraction between $S^{(l_i)}$ and the ∂z^I 's reduces to a trace,

$$S_{(Aa)(Bb)(Cc)}^{(1)} \dots \partial_1^I A_a \partial_2^J Y_{bB} \partial_3^K C_c = \text{tr} A_1^+ A_2^+ A_3^+; \quad (2.16a)$$

where we have introduced

$$A^+ = Y \partial^I; \quad A = Z \partial^I Y; \quad (2.16b)$$

for the complex Grassmannian (symmetry class A). For $Sp(N) = U(N)$ (symmetry class C) and for $SO(2N) = U(N)$ (symmetry class D) one must impose the additional constraint

$$(A^+)^T = \# A^+; \quad (A)^T = \# A; \quad (2.16c)$$

Any member T of the family of tensor fields constructed in Eq. (2.15) has the remarkable property that it vanishes under the action of the covariant derivative on any one of its indices

$$\begin{aligned} r_P T_{IJKL} &= \partial_P T_{IJKL} \\ &\quad - \sum_{Q=1}^Q \Gamma_{PI}^Q T_{QJKL} - \sum_{Q=1}^Q \Gamma_{PJ}^Q T_{IQKL} \\ &= 0; \end{aligned} \quad (2.17)$$

Here, A_{BC} are the components of the Levi-Civita connection whose non-vanishing entries are given by

$$A_{bc}^a = g^{aP} \partial_c g_{pb}; \quad A_{bc}^a = g^{aP} \partial_c g_{pb}; \quad (2.18)$$

This is also true for the NLM on the real Grassmannian $O(p+q) = [O(p) \times O(q)]$ which is related to the symplectic symmetry class of Anderson localization, and for the cases of simpler target manifolds such as $O(N) = O(N-1)$

or CP^{n+m-1} for which the tensor fields T are just products of the metric tensor. In all these cases, the fact that all high-gradient operators have vanishing covariant derivatives greatly simplifies the computation of their one-loop RG flows.

For a given number of derivatives, $2s$, there are many degenerate operators of the type (2.12) with $T_{IJK} \dots$ given by Eq. (2.15b) that share the same anomalous scaling dimensions $2s$ at the metallic fixed point. They are specified by the choice of fl_i in Eq. (2.15b) as well as that of the greek indices in the $@z^I$'s. This degeneracy is, however, lifted in perturbation theory as we depart from the metallic fixed point.

We close this section by noting that it is more convenient to introduce the conformal indices $+$ and

$$@ = @_x \quad i@_y : \quad (2.19)$$

instead of the Euclidean indices $= x; y$ in $2d^{2,41}$

III. THE COVARIANT BACKGROUND FIELD METHOD

A. Kahler normal coordinates for NLM

To perform the RG program, we divide the fields $(z; z)$ in the NLM into slow $(;)$, and fast $(;)$, modes. To this end, we write

$$z^a = a + a; \quad z^a = a + a; \quad (3.1)$$

where we require that $(;)$ satisfy the classical equations of motion. The fast modes $(;)$ represent the quantum fluctuations around the classical background $(;)$. They are integrated out, thereby generating an effective action for the slow modes $(;)$. In practice, this effective action is computed order by order within an expansion, the so-called loop expansion, which is nothing but the cumulant expansion. One drawback with the separation (3.1) into fast and slow modes is that the perturbative expansion of the action in powers of the fast modes $(;)$ with respect to the action evaluated at $(;)$ need not transform covariantly under reparametrization of the manifold. This danger can be avoided by choosing Kahler normal coordinates (KNC) in terms of which we denote the fast modes by $(! ; !)$. The KNC coordinates are defined by demanding the manifest covariance of the Kahler potential when expanded in terms of $(! ; !)$.^{65,66} The perturbative expansion of the action in powers of the KNC about the background field is then manifestly covariant, i.e., only covariant objects such as the metric, the Riemann tensor, the covariant derivative on the manifold, and so on appear in the expansion, as we shall shortly see. One advantage of this RG scheme, the covariant background field method, is that no redundant operator is induced upon renormalization. This is a very useful property when dealing with the mixing of the composite operators induced by an RG transformation.⁴⁵ A

related advantage is that there is no need to break the symmetry of the NLM to tame IR divergences.

The relation between the generic parametrization $(;)$ and the KNC parametrization $(! ; !)$ of the fast modes is non-linear. For examples,

$$@ (a + a) = @ a + D w^a \quad (3.2a)$$

$$+ \frac{1}{2} R^a_{c_1 c_2 c_3} (;) @ \quad \mathbb{S} w^{c_2} w^{c_1} + ;$$

$$g_{ij} (+ ; +) = g_{ij} (;) \quad (3.2b)$$

$$R_{ijkl} w^l w^{k_1} + :$$

As promised, the right-hand side is covariant under reparametrization as the pair $(D ; D)$ denotes the holomorphic and antiholomorphic covariant derivatives

$$D w^j = r_a w^j @ a$$

$$= @ w^j + \quad j_{ab} w^b @ a; \quad (3.3a)$$

$$D w^j = D w^j$$

$$= r_a w^j @ a$$

$$= @ w^j + \quad j_{ab} w^b @ a; \quad (3.3b)$$

respectively, with the components

$$R^a_{bcd} = @_d \quad a_{bc}; \quad R^a_{bcd} = @_d \quad a_{bc}; \quad (3.4)$$

$R_{EDAB} = g_{EC} R^C_{DAB}$ and so on of the curvature tensor. The expansion of the action in powers of the KNC is

$$S [\quad + \quad ; +] S [;] =$$

$$+ \frac{1}{2 t} \int_{Z^r} g_{ij} (;) D w^j D w^i$$

$$+ \frac{1}{2 t} \int_{Z^r} R_{ijkl} (;) w^l w^k @ \quad j @ \quad i$$

$$+ \frac{1}{2} w^k @ \quad j w^i \quad \frac{1}{2} @ \quad w^k w^j @ \quad i$$

$$+ : \quad (3.5)$$

B. Canonical form of the metric tensor and vielbeins

The covariant Gaussian expansion (3.5) about an extremum $(;)$ of the action can still be simplified further by performing a local $GL(M; C)$ transformation of the tangent space at $(;)$ that brings the metric tensor at $(;)$ to the canonical form. To this end, introduce the fast degree of freedom $(;)$ by the linear transformation^{67,68}

$$\hat{a} = w^a_a e^{\hat{a}}; \quad w^a = \hat{a}_a e^{\hat{a}}; \quad (3.6a)$$

$$\hat{a} = w^a_a e^{\hat{a}}; \quad w^a = \hat{a}_a e^{\hat{a}}; \quad (3.6b)$$

by demanding that

$$g_{ab} (;) w^b w^a = \hat{a}_b \hat{a}^a \quad (3.6c)$$

where the transformed metric is given by

$$g_{\hat{a}\hat{b}} = \text{diag } I_{M_1}; I_{M_2}; M_1 + M_2 = M : \quad (3.6d)$$

The two background dependent matrices $(g_a e^a)$ and $(g_{\hat{a}} e^{\hat{a}})$ from $GL(M; \mathbb{C})$ that implement this transformation in the antiholomorphic sector are inverse to each other. The same is true of the matrices $(g_a e^a)$ and $(g_{\hat{a}} e^{\hat{a}})$ in the holomorphic sector. These four matrices are called the vielbeins and must obey

$$g_{ab} e_b^a e_a = g_{\hat{a}\hat{b}}; \quad g_{\hat{a}\hat{b}} e_b^{\hat{a}} e_a = g_{ab}; \quad (3.7)$$

by condition (3.6c). From now on, latin letters with a hat refer to the coordinates of the Kahler manifold in the vielbein basis (3.6). Under transformation (3.6) the covariant expansion (3.5) becomes

$$S[\frac{1}{2} g_{\hat{a}\hat{b}} \dot{x}^{\hat{a}} \dot{x}^{\hat{b}}] = S[\frac{1}{2} g_{ab} \dot{x}^a \dot{x}^b] + \frac{1}{2} \int d\tau \int d^d x \hat{D}^{\hat{a}} \hat{D}^{\hat{b}} \hat{A}^{\hat{a}}_{\hat{b}} + \frac{1}{2} \int d\tau \int d^d x R_{ijkl} (\hat{e}^i_1 e^1) (\hat{e}^k_{\hat{k}} e^k) \hat{e}^j_{\hat{j}} \hat{e}^i_{\hat{i}} + \frac{1}{2} \int d\tau \int d^d x (\hat{e}^i_1 e^1) \hat{e}^k_{\hat{k}} \hat{e}^j_{\hat{j}} (\hat{e}^i_{\hat{i}} e^i) + \frac{1}{2} \int d\tau \int d^d x (\hat{e}^k_{\hat{k}} e^k) (\hat{e}^j_{\hat{j}} e^j) \hat{e}^i_{\hat{i}} + \dots \quad (3.8)$$

where yet another pair of covariant derivatives

$$\hat{D}^{\hat{a}} e^a = \partial^{\hat{a}} e^a - \hat{A}^{\hat{b}}_{\hat{a}} e^a; \quad (3.9a)$$

$$\hat{D}^{\hat{a}} e^{\hat{a}} = \partial^{\hat{a}} e^{\hat{a}} - \hat{A}^{\hat{b}}_{\hat{a}} e^{\hat{a}}; \quad (3.9b)$$

have been introduced. The field A is the $U(M)$ gauge field defined through the $U(M)$ spin connection $\hat{A}^{\hat{b}}_{\hat{a}}$, $\hat{A}^{\hat{b}}_{\hat{a}}$ by

$$\hat{A}^{\hat{b}}_{\hat{a}} e^a = \partial_{\hat{a}} e^a - \hat{A}^{\hat{b}}_{\hat{a}} e^a; \quad (3.10a)$$

$$\hat{A}^{\hat{b}}_{\hat{a}} e^{\hat{a}} = \partial_{\hat{a}} e^{\hat{a}} - \hat{A}^{\hat{b}}_{\hat{a}} e^{\hat{a}}; \quad (3.10b)$$

$$\hat{A}^{\hat{b}}_{\hat{a}} = \hat{A}^{\hat{b}}_{\hat{a}} e^c + \hat{A}^{\hat{b}}_{\hat{a}} e^c; \quad (3.10c)$$

[If the metric is pseudo Riemannian, $U(M)$ should be replaced by $U(M_1; M_2)$.] The emergence of an $U(M)$ gauge structure originates from the fact that the vielbeins are not uniquely specified by condition (3.6c). As the physical result cannot depend on the specific choice of gauge, i.e., the specific choice for the vielbeins, the dependence on the spin connection should occur through the $U(M)$ gauge invariant field strength

$$F^{\hat{b}}_{\hat{a}} = \partial_{\hat{a}} \hat{A}^{\hat{b}}_{\hat{c}} - \partial_{\hat{c}} \hat{A}^{\hat{b}}_{\hat{a}} + \hat{A}^{\hat{d}}_{\hat{a}} \hat{A}^{\hat{b}}_{\hat{c}} - \hat{A}^{\hat{d}}_{\hat{c}} \hat{A}^{\hat{b}}_{\hat{a}} = \hat{A}^{\hat{d}}_{\hat{a}} \hat{A}^{\hat{b}}_{\hat{c}} - \hat{A}^{\hat{d}}_{\hat{c}} \hat{A}^{\hat{b}}_{\hat{a}}; \quad (3.10d)$$

C. One-loop beta functions

We are in position to integrate out the fast degrees of freedom for the complex Grassmannian, $SO(2N) = U(N)$, and $Sp(2N) = U(N)$. The partition function for slow and fast modes is

$$Z = \int D[\dots] D[\dots] e^{S[\dots]}; \quad (3.11a)$$

The action $S[\dots] (= S[\dots] + S_0[\dots] + S_I[\dots])$ is separated into three contributions

$$S[\dots] = S[\dots] + S_0[\dots] + S_I[\dots]; \quad (3.11b)$$

where $S_0[\dots]$ is the free Gaussian part for the fast modes, and $S_I[\dots]$ represents the interactions between the slow and fast modes. Integration over the fast mode (\dots) is performed with the help of the cumulant expansion

$$Z = \int D[\dots] e^{S[\dots] + S_I[\dots]}; \quad (3.12a)$$

$$S[\dots] = h S_{I_0} + \frac{1}{2} (S_I)_0^2 + S_{I_0}^2 + \dots; \quad (3.12b)$$

where

$$h(\dots) = \frac{\int D[\dots] e^{S_0[\dots]} (\dots)}{\int D[\dots] e^{S_0[\dots]}}; \quad (3.12c)$$

To evaluate the cumulant expansion, we introduce the Green function

$$G^{\hat{a}}(x) G^{\hat{b}}(y)_0 = \int_{e_{d1}}^Z \frac{d^d k}{(2\pi)^d} \frac{e^{ik(x-y)}}{k^2} =: \int_{e_{d1}}^Z G_0(x-y); \quad (3.13)$$

where $\Lambda = \Lambda^{-1}$ is the UV cutoff. The leading contribution to the increment of the action of the slow mode (\dots) resulting from integrating out the fast mode (\dots) within the second-order cumulant expansion is

$$S[\dots] = G_0(0) R_{ab} \partial^b \partial^a; \quad (3.14)$$

with

$$R_{ab} = R^c_{abc} = R^c_{cab}; \quad (3.15)$$

the Ricci tensor.

For symmetry classes C and D, the Ricci tensor is proportional to the metric (the target space is an Einstein manifold) and given by

$$R_{(C)(E)e} = (N - \#) g_{(C)(E)e}; \quad (3.16)$$

on the one hand and

$$\begin{aligned}
 \text{tr}_{b_1 \dots b_2}^h T_{b_1 \dots b_2}^i(z; z) &= \text{tdl} \sum_{i=1}^{X^s} R^{1 \dots a}_{b_1 \dots b_2} T_{b_1 \dots b_2}^i(z; z) \\
 &= \text{tdl} \sum_{i=1}^{X^r} R^{k \dots a}_{b_1 \dots b_2} T_{b_1 \dots b_2}^i(z; z); \tag{4.3e}
 \end{aligned}$$

on the other hand. Here, we are using the conformal indices defined in Eq. (2.19). Note also that since holomorphic r_i and antiholomorphic r_j covariant derivatives do not commute in general, we have two distinct representations for $[T_{IJK}]_2$.

B. RG transformation

The RG transformation law (4.3) obeyed by any high-gradient operator for a NLM on the Hermitian symmetric spaces $U(p+q) = U(p) \times U(q)$, $SO(2N) = U(N)$, and $Sp(N) = U(N)$ becomes explicit with the help of the

(a) intra-trace formulae

$$\begin{aligned}
 \text{tr}^D A^{-1} M [A]_1 N^E &= \frac{2 \#^2}{2} \text{tdl}; \quad \#^2 \text{tr} A N A M^T + \text{tr} A N \text{tr} (A M) \\
 &\quad + \text{tr} A M \text{tr} (A N) + \#^2 \text{tr} A M^T A N; \tag{4.4a}
 \end{aligned}$$

$$\begin{aligned}
 \text{tr}^D A^+ M A^{-1} N^E &= + \frac{2 \#^2}{2} \text{tdl}; \quad \text{tr} A^+ A N \text{tr} (M) + \# \text{tr} A^+ A N M^T \\
 &\quad + \# \text{tr} A^+ A M^T N + \text{tr} (N) \text{tr} M A A^+; \tag{4.4b}
 \end{aligned}$$

(b) inter-trace formulae

$$\begin{aligned}
 \text{tr}^D A^{-1} M \text{tr} [A]_1 N^E &= \frac{2 \#^2}{2} \text{tdl}; \quad \# \text{tr} A N^T A M + \text{tr} A N A M \\
 &\quad + \text{tr} A N A M + \# \text{tr} A N^T A M; \tag{4.5a}
 \end{aligned}$$

$$\begin{aligned}
 \text{tr}^D A^+ M \text{tr} A^{-1} N^E &= \frac{2 \#^2}{2} \text{tdl}; \quad \text{tr} A^+ A N M + \# \text{tr} A A^+ M N^T \\
 &\quad + \# \text{tr} A^+ A N^T M + \text{tr} A A^+ M N; \tag{4.5b}
 \end{aligned}$$

and (c) diagonal contributions

$$\text{tr}_{a_1 \dots a_2}^h T_{a_1 \dots a_2}^i(z; z) = \text{tdl} (N \#) T_{a_1 \dots a_2}^i(z; z); \tag{4.6}$$

where M and N represent arbitrary rectangular matrices of the proper size and the matrix-valued A were defined in Eq. (2.16). This result, that follows from substituting the components of the curvature tensor (3.4) when expressed in the stereographic coordinates (2.8) into the master formula (4.3), agrees with the one-loop RG result for symmetry class A ($\# = 0$) in Ref. 38.

C. Diagonalization of the RG equation within the subspace of maximal number of switches

As is shown in Appendix B, the RG transformation law (4.3) has an upper (or lower) triangular structure when appropriate quantum numbers ("number of switches") are defined. Since the number of switches never increases under an RG transformation, we can

solve the eigenvalue and eigenvector problem (4.3) for each subspace with a fixed number of switches. We will limit ourselves to operators with the maximal number of switches, as the eigenscaling operator with the most dominant RG anomalous scaling dimension is known to be among them for symmetry class A.³⁹ In this subspace, any operator can be expressed as a polynomial of the ob-

jects

$$\text{tr} (A_+^+ A_-)^1; \text{tr} (A_+^+ A_+)^1 \quad ; \quad (4.7)$$

ie., conformal indices in a single trace are completely alternating. It can also be shown that the spaces spanned by $\text{tr} (A_+^+ A_-)^1$ and $\text{tr} (A_+^+ A_+)^1$ are not mixed by the RG transformation. We can thus consider these two subspaces separately as long as we do not require composite operators to be Hermitian.

We now compute the action of the RG transformation (4.3) on a composite operator of the form

$$O_{\text{frg}}^{(s)} = \prod_{l=1}^{X_L} r_l \quad ; \quad \text{tr} r_l = s \quad (4.8)$$

where all r_l on the right-hand side are a short-hand notation for objects of type $\text{tr} (A_+^+ A_-)^1$ without loss of generality. Since the diagonal contributions (c) to the RG eigenvalues are trivial, we focus on the intra-trace contributions (a) and inter-trace contributions (b) for the moment. If we define the ϕ -diagonal mixing matrix $\mathbb{R}^{(s)} = \mathbb{R}_{\text{frgfr}^0}^{(s)}$ by

$$O_{\text{frg}}^{(s)}(z; z) \stackrel{D}{=} \text{tr} \mathbb{R}_{\text{frgfr}^0}^{(s)} O_{\text{fr}^0}^{(s)}(z; z); \quad (4.9)$$

ie., we have only collected the ϕ -diagonal contributions (4.4) and (4.5) to the RG transformation law obeyed by $O_{\text{frg}}^{(s)}(z; z)$, the action of $\mathbb{R}^{(s)}$ can be cast into the form

$$\mathbb{R}^{(s)} = \sum_k \frac{X}{k} k^2 \frac{\partial}{\partial k} + N \sum_k k \frac{\partial}{\partial k} + \sum_{l,n} 2 \ln \frac{\partial^2}{\partial l \partial n} + (2 - \#^2) (l+n) \frac{\partial}{\partial (l+n)}; \quad (4.10)$$

The eigenvalue problem (4.10) can be solved by brute force for small s in which cases we have verified that the sign of the eigenvalues of the ϕ -diagonal mixing matrix $\mathbb{R}^{(s)}$ for $SO(2N)=U(N)$ are opposite to those for $Sp(N)=U(N)$. In particular, the largest eigenvalue of $\mathbb{R}^{(s)}$ for $SO(2N)=U(N)$ is given by minus the smallest eigenvalue for $Sp(N)=U(N)$ and vice et versa. This fact is consistent with the group isomorphism (3.18b), and the fact that the beta functions for the corresponding NL M match upon the replacement $t \rightarrow -t$.

More generally, one verifies that

$$\lambda^{(s)} = 2(s^2 - s) + s(p+q); \quad (\# = 0); \quad (4.11a)$$

$$\lambda^{(s)} = s^2 - 2s + sN; \quad (\# = +1); \quad (4.11b)$$

$$\lambda^{(s)} = 2s^2 - s + sN; \quad (\# = -1); \quad (4.11c)$$

is an eigenvalue of the ϕ -diagonal mixing operator $\mathbb{R}^{(s)}$ for any given s with the eigenvector

$$v^{(s)} = \prod_{\text{frg } j} \frac{X}{r_j} \frac{Y}{j^{r_j}} (r_j)^{r_j}; \quad (\# = 0); \quad (4.12a)$$

$$v^{(s)} = \prod_{\text{frg } j} \frac{X}{r_j} \frac{Y}{j^{r_j}} (r_j)^{r_j}; \quad (\# = +1); \quad (4.12b)$$

$$v^{(s)} = \prod_{\text{frg } j} \frac{X}{r_j} \frac{Y}{j^{r_j}} 2^{j-1} (r_j)^{r_j}; \quad (\# = -1); \quad (4.12c)$$

As it is known that $\lambda^{(s)}$ is the largest eigenvalue of $\mathbb{R}^{(s)}$ in symmetry class A, we believe this to be also true for symmetry classes C and D. If so, the largest RG eigenvalues for a given number s of gradients is given by, after adding the diagonal contributions (c),

$$\lambda^{(s)} = \lambda^{(s)} - s(p+q) = 2(s^2 - s); \quad (\# = 0); \quad (4.13a)$$

$$\lambda^{(s)} = \lambda^{(s)} - s(N-1) = s^2 - s; \quad (\# = +1); \quad (4.13b)$$

$$\lambda^{(s)} = \lambda^{(s)} - s(N+1) = 2(s^2 - s); \quad (\# = -1); \quad (4.13c)$$

Note that these results are valid up to one loop before taking the replica limit. Equation (1.3) then follows by combining Eq. (4.13) with Eq. (4.2b).

Acknowledgements

C.M. and A.F. acknowledge hospitality of the Kavli Institute for Theoretical Physics at Santa Barbara during the completion of the manuscript. This research was supported in part by the National Science Foundation under Grants No. DMR-00-75064 (A.W.W.L.) and No. PHY 99-07949 (S.R., A.F., C.M.).

APPENDIX A: GROUP-THEORY VERSUS GEOMETRIC REPRESENTATIONS OF THE NL M

The purpose of this appendix is to bridge the geometrical representation of the NL M used in this paper and the one in terms of the so-called "Q-matrix".

First, recall that an element Q in $U(p+q) = [U(p) \times U(q)]$ can be written as¹⁹

$$Q = U^{-1} U; \quad U = \text{diag}(U_p; U_q); \quad U \in U(p+q); \quad (A.1)$$

As noticed by Lerner and Wegner,³⁸ high-gradient operators in the Q-matrix representation as well as the action of the NL M can be conveniently expressed in terms of the objects

$$A = \frac{1}{2} P^+ @ Q P \quad (A.2a)$$

where $\mathbb{P} = \mathbb{P}^\dagger$ and the projectors \mathbb{P}

$$\mathbb{P} = \frac{1}{2} (1 + \mathbb{Q}); \quad (\mathbb{P})^2 = \mathbb{P} \quad (\text{A } 2\text{b})$$

were introduced. A generic high-gradient operator is then built by taking a product of

$$\text{tr } A^\dagger A A^\dagger : \quad (\text{A } 3)$$

For example, the action of the NLM is

$$\begin{aligned} S &= \frac{1}{16} \text{tr}_{\mathbb{Z}} \mathbb{Q} \mathbb{Q} \\ &= \frac{1}{4} \text{tr}_{\mathbb{Z}} A^\dagger A : \end{aligned} \quad (\text{A } 4)$$

In order to establish explicitly the connection between the \mathbb{Q} -matrix representation (A2)-(A4) and the geometrical representation in the stereographic coordinates (2.8), we parametrize \mathbb{P}^\dagger and $\mathbb{P} = \mathbb{P}^\dagger \mathbb{Q}$ in Eq. (A2b) through a $(p+q) \times q$ matrix y_i according to⁶⁹

$$\mathbb{P}^\dagger = \sum_{i=1}^{p+q} y_i X_i^\dagger \quad y_i = \sum_{i=1}^{p+q} X_i^\dagger y_i \quad (\text{A } 5\text{a})$$

where y_i satisfies the constraint

$$y_i = I_q : \quad (\text{A } 5\text{b})$$

The matrix y_i is thus a collection of q orthonormal vectors $f_a \in \mathbb{C}^{p+q}$, $a=1, \dots, q$. We can use this constraint to fix the upper $q \times q$ block of \mathbb{P} . We write

$$\mathbb{P} = \begin{pmatrix} X & y \\ y^\dagger & X \end{pmatrix} \quad (\text{A } 6\text{a})$$

where X_{ij} , y_i , X_{ij} , and $y_i^\dagger X_{jk}$ are $(p+q) \times q$, $p \times q$, $q \times q$, and $p \times q$ matrices, respectively, and the indices i, I , and j run over the sets $i=1, \dots, p+q$, $I=1, \dots, p$, $j=1, \dots, q$, respectively. Next, we express X in terms of y by making use of the constraint (A5b),

$$X_{ni} Z_{nm}^{-1} X_{mk} = \delta_{ik} \quad (X^\dagger Z^{-1} X = I_q): \quad (\text{A } 6\text{b})$$

Here the Hermitian $q \times q$ matrix Z was introduced in Eqs. (2.9b) and (2.9c). Constraint (A6b) is satisfied if X is chosen to be

$$X = Z^{-1} y : \quad (\text{A } 6\text{c})$$

A generic high-gradient operators (A3) in the \mathbb{Q} -matrix representation is thus expressed in terms of the stereographic coordinate $(y; y^\dagger)$ from Eq. (2.8). For example, one verifies that the action of the NLM (A4) reduces to Eq. (2.5) with the metric (2.9a) derived from the Kahler potential (2.8) once the parametrization (A5) with Eq. (A6) is used. Similarly,

$$\begin{aligned} \text{tr } A^\dagger A A^\dagger A &= \\ \text{tr } Z \mathbb{Q}^\dagger Y Y^\dagger \mathbb{Q} &= \text{tr } Z \mathbb{Q}^\dagger Y Y^\dagger \mathbb{Q} : \end{aligned} \quad (\text{A } 7)$$

Observe that the alternating structure in A^\dagger and A in the high-gradient operators has a one-to-one correspondence to that in Z and Y .

APPENDIX B: UPPER TRIANGULAR STRUCTURE OF THE RG TRANSFORMATION

We prove in this appendix that the RG transformation law obeyed by high-gradient operators has an upper (or lower) triangular structure when appropriate "quantum numbers" (number of switches) are defined. A trace made of an even product of A 's,

$$\text{tr } A_1^\dagger A_2 A_3^\dagger A_4 : \quad (\text{B } 1)$$

is said to have n number of switches from A to A^\dagger when, dividing the sequence into pairs according to

$$\text{tr } A_1^\dagger A_2 A_3^\dagger A_4 : \quad (\text{B } 2)$$

there are n pairs $A A^\dagger$ in the sequence. The number n does not depend on the cyclic permutations of the pairs $A A^\dagger$. For a given trace of the form (B2) we can define

$$n_{\text{tr}} = \sum_{i=1}^n X_i : \quad (\text{B } 3)$$

We are going to show that, for the symmetry classes A, C , and D , the total number

$$n = \sum_{\text{tr}} n_{\text{tr}} \quad (\text{B } 4)$$

of switches in a composite operator of type (2.12) never increases under the RG transformation law (4.3). Here, the summation extends over the product over all traces of the form (B2) making up the high-gradient operator. In other words, the RG transformation law (4.3) is triangular with respect to the quantum number (B4).

1. Symmetry class A

We begin with symmetry class A . This case ($\# = 0$) was already treated in Refs. 38,39,43 and is needed for symmetry classes C ($\# = 1$) and D ($\# = +1$). Consider any string $L_1 A_1 A_0 R_1$ and any string $L_2 A_2 A_0 R_2$ of 4 matrices each that enter a high-gradient operator.

First, we assume that each string belongs to two distinct traces. We seek the one-loop RG transformation law obeyed by the number of switches

$$n_1; \quad n_2 : \quad (\text{B } 5\text{a})$$

contained in

$$\text{tr } L_1 A_1 A_0 R_1 \text{tr } L_2 A_2 A_0 R_2 : \quad (\text{B } 5\text{b})$$

By making use of Eq. (4.5) with $\# = 0$ we deduce that the

one-loop RG transformation law obeyed by Eq. (B 5b) is

$$\text{tr} \begin{matrix} h & i \\ A_1 & A_0 \\ 1 & 1 \end{matrix} R_1 L_1 \text{tr} \begin{matrix} h & i \\ A_2 & A_0 \\ 2 & 2 \end{matrix} R_2 L_2 =$$

$$\text{tdl}_{1; 2}^+ + \text{tdl}_{1; 2}^0 + \text{tdl}_{1; 2}^1 + \text{tdl}_{1; 2}^2 \quad (B 6a)$$

$$\text{tdl}_{1; 2}^+ \text{tr} \begin{matrix} A_2^+ & A_0 \\ R_2 L_2 A_1^+ & A_0 \\ R_1 L_1 \end{matrix} \quad (B 6b)$$

$$+ \text{tdl}_{1; 2}^0 \text{tr} \begin{matrix} R_2 L_2 A_2^+ & A_0 \\ A_2^+ & A_0 \\ R_1 L_1 \end{matrix} \quad (B 6c)$$

$$+ \text{tdl}_{1; 2}^1 \text{tr} \begin{matrix} A_1^+ & A_0 \\ A_1^+ & A_0 \\ R_2 L_2 R_1 L_1 \end{matrix} \quad (B 6d)$$

$$\text{tdl}_{1; 2}^2 \text{tr} \begin{matrix} A_1^+ & A_0 \\ R_2 L_2 A_1^+ & A_0 \\ R_1 L_1 \end{matrix} : \quad (B 6e)$$

Line (B 6a) contains $2_{1; 2}^+ + 2_{1; 2}^0$ switches. Lines (B 6b) to (B 6e) contain $2_{1; 2}^1 + 2_{1; 2}^2$ switches. Hence, only line (B 6a) has the potential to increase the initial number (B 5a) of switches. However, an increase over the initial number (B 5a) of switches is not possible in view of the identity $2_{1; 2}^+ + 2_{1; 2}^0 + 2_{1; 2}^1 + 2_{1; 2}^2 = 2_{1; 2}^+ + 2_{1; 2}^0$. Line (B 6a) only contributes if $2_{1; 2}^+ = 2_{1; 2}^0$ and $2_{1; 2}^1 = 2_{1; 2}^2$ both hold, i.e., when the initial number of switches (B 5a) is maximum (2). Consequently, the number of switches $2_{1; 2}^+$ induced by line (B 6a) equals the initial number of switches 2 when $1 = 2$ while it decreases otherwise.

Second, we assume that each string belongs to the same trace. We seek the one-loop RG transformation law obeyed by the number of switches

$$2_{1; 2}^+ + 2_{1; 2}^0 \quad (B 7a)$$

contained in

$$\text{tr} \begin{matrix} L_1 A_1 & A_0 \\ R_1 L_2 A_2 & A_0 \\ R_2 \end{matrix} : \quad (B 7b)$$

By making use of Eq. (4.4) with $\# = 0$ one verifies again that the number of switches n cannot increase after integrating over the fast modes up to one loop.

We have proven that the RG flows of high-gradient operators made of any given number of switches decouple from the RG flows of high-gradient operators made of a lesser number of switches in symmetry class A. As a corollary, the mixing matrix (4.1) is triangular in symmetry class A.

2. Symmetry classes C and D

We turn our attention to symmetry classes $\#j = 1$, i.e., $SO(2N) = U(N)$ and $Sp(N) = U(N)$. We first consider the inter-trace formula of the RG transformation (4.5) applied to

$$\text{tr} \begin{matrix} A_1 & M \\ 1 & 1 \end{matrix} \text{tr} \begin{matrix} A_2 & A_0 \\ 2 & 2 \end{matrix} N \quad (B 8) \text{ respectively.}$$

where $M = R_1 L_1$ and $N = R_2 L_1$. It is sufficient to consider the contributions proportional to $\#$ in Eq. (4.5) as the remaining terms can be treated along the lines of symmetry class A. There are four such contributions

$$(d):$$

$$2_{1; 2}^0 \text{tr} \begin{matrix} A_2^+ & N^T A_0 \\ A_2^+ & A_0 \\ 1 & 1 \end{matrix} M ; \quad (B 9a)$$

$$1_{1; 2}^0 \text{tr} \begin{matrix} A_1^+ & N^T A_0 \\ A_2^+ & A_0 \\ 1 & 1 \end{matrix} M ; \quad (B 9b)$$

$$1_{1; 2}^0 \text{tr} \begin{matrix} A_2^+ & N^T A_0 \\ A_2^+ & A_0 \\ 1 & 1 \end{matrix} M A_1^+ ; \quad (B 9c)$$

$$2_{1; 2}^0 \text{tr} \begin{matrix} A_1^+ & A_2^+ N^T A_0 \\ A_2^+ & M A_1^+ \end{matrix} ; \quad (B 9d)$$

We assign to

$$A = \text{tr} \begin{matrix} A_1^+ & A_0 \\ 1 & 1 \end{matrix} M \text{tr} \begin{matrix} A_2^+ & A_0 \\ 2 & 2 \end{matrix} N \quad (B 10)$$

the quartet of numbers

$$(n_+^+; n_+^+; n_+; n) \quad (B 11)$$

where n counts the total number of switches of type $A_1 A_1$ in A . In symmetry classes D and C, we can combine $\text{tr} O = \text{tr} O^T$ and the symmetry $(A_1)^T = \# A_1$, to infer that

$$(n_+^+; n_+^+; n_+; n) = (n; n_+; n_+^+; n_+^+): \quad (B 12)$$

As opposed to symmetry class A, individual switches n are not separately conserved but the total number of switches

$$n = \begin{matrix} X \\ n \\ ; \end{matrix} \quad (B 13)$$

will be shown to be conserved.

Without loss of generality we choose $\# = +$ in A. The total number of switches n_+^+ and n^+ are

$$n_+^+ = n_+^+ [A_1^+ A_0] + n_+^+ [M] + n_+^+ [A_2^+ A_0] + n_+^+ [N]; \quad (B 14a)$$

$$n^+ = n^+ [A_1^+ A_0] + n^+ [M] + n^+ [A_2^+ A_0] + n^+ [N]; \quad (B 14b)$$

respectively. The total number of switches n_+ and n follow from using $A = \text{tr} \begin{matrix} A_0 M A_1^+ \\ 1 \end{matrix} \text{tr} \begin{matrix} A_0 N A_2^+ \\ 2 \end{matrix}$ and are given by

$$n_+ = n_+ [A_0 M A_1^+] + n_+ [A_0 N A_2^+]; \quad (B 14c)$$

$$n = n [A_0 M A_1^+] + n [A_0 N A_2^+]; \quad (B 14d)$$

The number of switches (B 14) in A changes under the RG transformation (B 9) as follows. The total number of switches n_+^+ and n_+^- from line (B 9a) is given by

$$\begin{aligned} n_+^+ [(a)] &= n_+^+ [A_2^+ N^T A_0] + n_+^+ [A_1^+ A_0] + n_+^+ [M] \\ &= n_+^+ [A_2 N A_2^+] + n_+^+ [A_1^+ A_0] + n_+^+ [M]; \end{aligned} \quad (\text{B 15a})$$

$$\begin{aligned} n_+^- [(a)] &= n_+^- [A_2^+ N^T A_0] + n_+^- [A_1^+ A_0] + n_+^- [M] \\ &= n_+^- [A_2 N A_2^+] + n_+^- [A_1^+ A_0] + n_+^- [M]; \end{aligned} \quad (\text{B 15b})$$

respectively, where we again made use of the symmetry condition $(A^+)^T = \#A^-$, and the fact that the number of A in a trace is always even. Similarly, the total number of switches n_+ and n_- from line (B 9a) is given by

$$\begin{aligned} n_+ [(a)] &= n_+ [N^T] + n_+ [A_2 A_1^+] + n_+ [A_1 M A_2^+] \\ &= n_+ [N] + n_+ [A_2 A_1^+] + n_+ [A_1 M A_2^+]; \end{aligned} \quad (\text{B 15c})$$

$$n_- [(a)] = (\text{as above with } !); \quad (\text{B 15d})$$

respectively. Thus, the total number of switches changes from

$$\begin{aligned} n &= n_+^+ [A_1^+ A_0] + n_+^+ [M] + n_+^+ [A_2^+ A_0] \\ &\quad + n_+^+ [N] + n_+^+ [A_1 M A_1^+] + n_+^+ [A_2 N A_2^+] \end{aligned} \quad (\text{B 16})$$

to

$$\begin{aligned} n [(a)] &= n_+^+ [A_2 N A_2^+] + n_+^+ [A_1^+ A_0] + n_+^+ [M] \\ &\quad + n_+^+ [N] + n_+^+ [A_2 A_1^+] + n_+^+ [A_1 M A_2^+] \end{aligned} \quad (\text{B 17})$$

for the contribution from line (B 9a). The net change

$n_j(a) = n_+^+ [(a)]$ is

$$\begin{aligned} n_j(a) &= n_+^+ [A_2^+ A_0] - n_+^+ [A_2 A_1^+] \\ &\quad + n_+^+ [A_1 M A_1^+] - n_+^+ [A_1 M A_2^+]; \end{aligned} \quad (\text{B 18})$$

When $\epsilon_2; \frac{0}{2} \epsilon_1; \epsilon_2 = 1$, we find

$$\begin{aligned} n_j(a) &= n_+^+ [A_2^+ A_0] - n_+^+ [A_2 A_2^+] \\ &\quad + n_+^+ [A_1 M A_1^+] - n_+^+ [A_1 M A_1^+] \\ &= 0 \end{aligned} \quad (\text{B 19})$$

with the help of the prefactor $\epsilon_2; \frac{0}{2} \epsilon_1; \epsilon_2 = 1$. When $\epsilon_2; \frac{0}{2} \epsilon_1; \epsilon_2 = 1$, we find

$$\begin{aligned} n_j(a) &= n_+^+ [A_2^+ A_0] - n_+^+ [A_2 A_2^+] \\ &\quad + n_+^+ [A_1 M A_1^+] - n_+^+ [A_1 M A_1^+] \\ &= 1 + n_+^+ [A_1 M A_1^+] - n_+^+ [A_1 M A_1^+] \\ &= 0; \end{aligned} \quad (\text{B 20})$$

We conclude that the total number of switches does not increase under the RG transformation (B 9a).

Repeating the same analysis for the contributions (b) (d) in Eq. (B 9) yields that the change in the total number of switches is 0.

We leave it to the reader to verify that the same conclusion holds for the intra-trace formula (4.4).

This completes the proof for the upper triangular structure of the RG equation with respect to the total number of switches in symmetry classes D and C.

APPENDIX C: RG FLOW OF HIGH-GRADIENT OPERATORS FOR NL M ON SUPERMANIFOLDS

We derive in this appendix two "master formulae" needed to perform the one-loop RG analysis of high-gradient operators for NL M defined on an arbitrary Riemannian or Kahlerian supermanifold. Master formulae (C 19) and (C 40) are supersymmetric versions of Eq. (4.3) and can be used for any target supermanifold once the geometrical data of the manifold are known. An expansion in terms of normal coordinate of tensor fields plays again an essential role. For the non-supersymmetric Riemannian case, an expansion in terms of Riemann normal coordinate (RNC) based on the notion of geodesics was used for the computation of the beta function in all generality^{32,57} and the RG analysis of high-gradient operators for the special case of the $O(N) = O(N-1)$ NL M^{44,45}. The RNC expansion can also be applied to Hermitian manifolds. For the (non-supersymmetric) Kahlerian case, however, one can also use the Kahler potential to find a normal coordinate system, the Kahler normal coordinates (KNC) introduced in Refs. 65 and 66. In this Appendix, we will generalize these normal coordinate expansions to the case of supersymmetric target manifolds. All the results that follow reduce to the non-supersymmetric ones if we ignore the commuting-anticommuting nature of the objects and fermionic sign factors.

1. NL M on Riemannian supermanifolds

a. Notations and conventions for Riemannian supermanifolds We begin with a quick review of concepts and

notation. We refer the reader to Ref. 59 for a textbook on the geometry of supermanifolds.

A tensor field of rank $(r;s)$ on a supermanifold has r contravariant (upper) and s covariant (lower) indices, $T^{ab\dots c\dots}$. On a supermanifold, components of tensor fields can be either commuting or anti-commuting supernumbers.

In order to define their mutual statistics, the grade $\epsilon(O) = 0;1$ is assigned to each object O on a supermanifold, where O can be a supernumber, a supertensor field, a component of a supertensor field, etc. When the ordering of two objects O_1 and O_2 is exchanged, a factor $(-1)^{\epsilon(O_1)\epsilon(O_2)}$ arises, $O_1O_2 = (-1)^{\epsilon(O_1)\epsilon(O_2)}O_2O_1$. In order to define the grade for the component of a tensor field of arbitrary rank, we first assign a grade to each of its index. If index a corresponds to a commuting component of the coordinates of a point on the supermanifold $\epsilon(a) = 0$, while $\epsilon(a) = 1$ otherwise. For simplicity, a statistical factor $(-1)^{\epsilon(a)\epsilon(b)}$ is often abbreviated by $(-1)^{ab}$, e.g., $X^aY^b = (-1)^{ab}Y^bX^a$. The grade of the component $T^{ab\dots c\dots}$ of the tensor field T is then $\epsilon(T^{ab\dots c\dots}) = \epsilon(a) + \epsilon(b) + \dots + \epsilon(c) + \dots$.⁷⁰

It is convenient to introduce a shifting rule for the leftmost index of tensors,

$${}^aT^{b\dots c\dots} := T^{ab\dots c\dots}; \quad (C1a)$$

$${}_aT^{b\dots c\dots} := (-1)^aT_a{}^{b\dots c\dots}; \quad (C1b)$$

As with all other objects on a supermanifold, derivatives also carry a grade. Thus, derivatives from the left ∂_a and the right ∂_a need to be distinguished. They are related by

$$\overset{\uparrow}{\partial}_a f = (-1)^{\epsilon(f)+a} f \partial_a; \quad (C2)$$

where f is a function over the manifold. In this appendix we will mainly use derivatives ∂_a and covariant derivatives r_a that act from the right.

Supertransposition of matrices, denoted by $\overset{\uparrow}{\cdot}$, is defined by

$${}^iK_j = (-1)^{j(i+j)} {}_jK^i; \quad (C3a)$$

$${}_jM_i = (-1)^{i+j+ij} {}_iM_j; \quad (C3b)$$

$${}^jN_j = (-1)^{ij} {}_iN^j; \quad (C3c)$$

For illustration, we shall apply these rules to the components g_{ab} of the metric tensor field, and components g^{ab} of the inverse of the metric tensor field. From the shifting rule

$${}^a g^b = g^{ab}; \quad {}_a g_b = (-1)^a g_{ab}; \quad (C4a)$$

$${}_a g_c {}^c g^b = {}_a g^b; \quad {}^a g_c {}^c g_b = {}^a g_b; \quad (C4b)$$

The supersymmetry of the metric tensor field implies

$$\begin{aligned} {}_a g_b &= (-1)^a g_{ab} \\ &= (-1)^{a+ab} g_{ba} \\ &= (-1)^{a+b+ab} {}_b g_a; \end{aligned} \quad (C5a)$$

$$\begin{aligned} {}^a g^b &= g^{ab} \\ &= (-1)^{ab} g^{ba} \\ &= (-1)^{ab} g^a; \end{aligned} \quad (C5b)$$

An index of a component X^a of a supervector field can be raised, lowered, and shifted according to the rules

$${}^a X = X^a; \quad {}_a X = (-1)^a X_a; \quad (C6a)$$

$${}_a X = {}_a g_b {}^b X; \quad X_a = X^b {}_b g_a; \quad (C6b)$$

$${}^a X = {}^a g_b {}^b X; \quad X^a = X_b {}^b g^a; \quad (C6c)$$

b. The NL M on a Riemannian supermanifold The NL M on a Riemannian supermanifold is defined by the partition function

$$Z = \int D[\varphi] e^{S[\varphi]}; \quad (C7a)$$

$$S[\varphi] = \frac{1}{4t} \int d^4x \partial_a \varphi_b \partial^a \varphi^b; \quad (C7b)$$

where $\varphi^a(r) \in \mathbb{R}_c^M \times \mathbb{R}_a^N$ represents the $M+N$ components of the coordinates of point r on the target manifold, with M referring to the number of commuting coordinates whereas N that of the number of anti-commuting ones,⁵⁹ and t is the NL M coupling constant.

c. The covariant background field method and the super Riemann normal coordinate expansion In the background field method, the covariant vector field $\varphi^a(r)$ is separated into two parts,

$$\varphi^a(r) = \bar{\varphi}^a(r) + \delta\varphi^a(r); \quad (C8)$$

whereby $\bar{\varphi}^a(r)$ is assumed to be a slowly varying solution to the classical equations of motion which transforms like a contravariant vector and $\delta\varphi^a(r)$ represents fluctuations around the slow degrees of freedom $\bar{\varphi}^a(r)$.

Two steps are needed to perform the one-loop RG program in a covariant fashion. First, we need to trade the expansion in terms of $\delta\varphi^a$, which is not covariant, for that of fields φ^a that transform like contravariant vectors. Second, we need to rotate the RNC by a vielbein to evaluate the functional integral over the fast modes.

The RNC is based on the notion of geodesics on Riemannian manifolds. Since almost all the notions on a Riemannian manifold such as forms, connection, the Riemann tensor, geodesics, etc., have their counterpart on a Riemannian supermanifold⁵⁹, it is straightforward to develop the super extension of the RNC expansion. The expansion of a covariant tensor field T in terms of the super RNC φ^a is given by

$$T_{b_1 \dots b_{i+1}}(b) = T_{b_1 \dots b_i}(b) + T_{b_1 \dots b_i} \kappa_{bc}(\cdot)^c + \frac{1}{2!} T_{b_1 \dots b_i} \kappa_{c_1 c_2} \dots$$

$$\frac{1}{3} X^1 T_{b_1 \dots b_{i+1}}(b) \dots + (1)^{b_1 c_1} R_{c_1 b_1 c_2} \dots + (1)^{b_1 c_1} R_{c_1 b_1 c_2} \dots + \dots \quad (C9)$$

The RNC expansion for $\mathcal{E}(a + a)$ is given by

$$\mathcal{E}(a + a) = \mathcal{E}^a + a^D + \frac{1}{3} R^a_{c_1 c_2 c_3} \mathcal{E}^{c_3 c_2 c_1} + \dots \quad (C10)$$

where D is the right covariant derivative for a

$$a^D = \mathcal{E}^a + a_{bc} \mathcal{E}^{cb}; \quad (C11)$$

and a_{bc} is a component of the connection. For a Riemannian supermanifold the connection can be derived from the metric. With the covariant expansion Eqs. (C9) and (C10) the action is expanded in terms of a according to

$$S[\dots] = \frac{1}{4} \int d^4x \dots + R_{abcd} \mathcal{E}^{dcb} \mathcal{E}^a \dots \quad (C12)$$

d. Vielbeins on a Riemannian supermanifold To integrate over the fermi modes in the expansion (C12), we rotate a to \hat{a} with the linear transformation

$$\hat{a} = a_a \hat{e}^a(\cdot); \quad a = \hat{a}_{\hat{a}} \hat{e}^{\hat{a}}(\cdot); \quad (C13a)$$

$$\hat{a} = \hat{a}_{\hat{a}} \hat{e}^{\hat{a}}(\cdot)^{\hat{a}}; \quad a = a_a \hat{e}^a(\cdot)^{\hat{a}}; \quad (C13b)$$

where

$$a_a \hat{e}^{\hat{a}} \hat{e}^{\hat{b}} = a^b; \quad \hat{a}_{\hat{a}} \hat{e}^{\hat{a}} \hat{e}^{\hat{b}} = \hat{a}^{\hat{b}}; \quad (C14a)$$

$$a_a \hat{e}^{\hat{a}} \hat{e}^{\hat{b}} = a^b; \quad \hat{a}_{\hat{a}} \hat{e}^{\hat{a}} \hat{e}^{\hat{b}} = \hat{a}^{\hat{b}}; \quad (C14b)$$

$$(1)^{\hat{a}(\hat{a}+a)} \hat{a}^{\hat{a}} = a_a \hat{e}^a; \quad (1)^{a(a+\hat{a})} \hat{a}^{\hat{a}} = \hat{a}_{\hat{a}} \hat{e}^{\hat{a}}; \quad (C14c)$$

Here, the vielbeins (and their inverse) \hat{e} are defined by

$$a_a \hat{e}^{\hat{a}} \hat{e}^{\hat{b}} = a^b; \quad \hat{a}_{\hat{a}} \hat{e}^{\hat{a}} \hat{e}^{\hat{b}} = \hat{a}^{\hat{b}}; \quad (C15a)$$

with

$$a_{\hat{b}} = \begin{pmatrix} D_M & 0 \\ 0 & D_N \end{pmatrix}; \quad (C15b)$$

$$D_M = \text{diag} \left\{ \underbrace{1; \dots; 1; 1; \dots}_M \right\}; \quad (C15c)$$

$$D_N = \begin{pmatrix} 0 & +i \\ i & 0 \end{pmatrix}; \quad \dots; \quad (C15d)$$

Observe that we are allowing the bosonic part of \mathcal{E} to be pseudo Riemannian, i.e., is allowed to have -1 as well as $+1$ as its diagonal elements. From now on, latin letters with a hat refer to the coordinates of the Riemannian manifold in the vielbein basis.

The covariant expansion of the action in terms of \hat{a} is given by

$$S[\dots] = \frac{1}{4} \int d^4x \dots + R_{\hat{a}\hat{b}\hat{c}\hat{d}} \hat{\mathcal{E}}^{\hat{d}\hat{c}\hat{b}} \hat{\mathcal{E}}^{\hat{a}} \dots \quad (C16a)$$

where yet another covariant derivative from the right

$$\hat{a}^{\hat{D}} = \mathcal{E}^{\hat{a}} + \hat{a}^{\hat{A}}(\cdot)_{\hat{b}}^{\hat{D}}; \quad (C16b)$$

$$\hat{a}^{\hat{D}} = \mathcal{E}^{\hat{a}} + \hat{a}^{\hat{A}}(\cdot)_{\hat{b}}^{\hat{D}}; \quad (C16c)$$

has been introduced together with the $O Sp(M, N)$ spin connection [if the bosonic part of \mathcal{E} is pseudo Riemannian, $O Sp(M, N)$ should be replaced by $O Sp(M_1; M_2; N)$ where $M_{1=2}$ is the number of -1 in the diagonal part of \mathcal{E} with $M_1 + M_2 = M$],

$$\hat{a}^{\hat{D}}(\cdot)_{\hat{b}} = \hat{a}^{\hat{D}} e_a(\cdot)^{\hat{c}\hat{b}} \hat{e}_{\hat{b}}^{\hat{c}} + \hat{a}^{\hat{D}} e_a(\cdot)^{\hat{c}\hat{b}} \hat{e}_{\hat{b}}^{\hat{c}}; \quad (C16d)$$

the $O Sp(M, N)$ gauge field

$$\hat{a}^{\hat{A}}(\cdot)_{\hat{b}} = (1)^{\hat{c}\hat{b}} \hat{a}^{\hat{A}}(\cdot)_{\hat{c}}; \quad (C16e)$$

$$\hat{a}^{\hat{A}}(\cdot)_{\hat{b}} = (1)^{\hat{b}(\hat{a}+\hat{b})} \hat{a}^{\hat{A}}(\cdot)_{\hat{b}}; \quad (C16f)$$

and the $O Sp(M, N)$ field strength tensor

$$\hat{a}^{\hat{F}}(\cdot)_{\hat{b}} = \mathcal{E}^{\hat{a}}(\hat{A})_{\hat{b}} - \mathcal{E}^{\hat{a}}(\hat{A})_{\hat{b}} + \hat{a}^{\hat{A}}(\hat{A})_{\hat{c}}^{\hat{D}}(\hat{A})_{\hat{b}}^{\hat{E}} - \hat{a}^{\hat{A}}(\hat{A})_{\hat{c}}^{\hat{D}}(\hat{A})_{\hat{b}}^{\hat{E}} = \hat{a}^{\hat{A}} R^{\hat{a}}_{\hat{b}\hat{c}\hat{d}} \mathcal{E}^{\hat{d}} \mathcal{E}^{\hat{c}} \hat{e}_{\hat{b}}^{\hat{a}}; \quad (C16g)$$

The components of the Riemann tensor are here $R^{\hat{a}}_{\hat{b}\hat{c}\hat{d}}$.

e. One-loop beta function The integration over the fast modes is performed with the help of the cumulant expansion. The effective action for the slow modes is, up to one-loop,

$$Z = \int D[\varphi] e^{S[\varphi] + S[\varphi]}, \quad (C17a)$$

$$S[\varphi] = \frac{1}{2} G_0(0) \varphi^i (1) R_{ij} \varphi^j; \quad (C17b)$$

where

$$R_{ij} = (1)^{k(i+1)+1} g^{kl} R_{likj} \quad (C17c)$$

is the Ricci tensor.

f. RG flows of high-gradient operators The covariant expansion in terms of the super RNC applies to the high-gradient operators

$$T_{ijk...}(\varphi) \varphi^i \varphi^j \varphi^k \quad (C18)$$

as well and can be used to compute their anomalous scaling dimensions. We shall assume that all the covariant derivatives of the tensor field $T_{ijk...}(\varphi)$ vanish. If so we infer the RG transformation law

$$\begin{aligned} D T_{ij...}(\varphi) \varphi^i \varphi^j &= D T_{ij...}(\varphi) \varphi^i \varphi^j \\ &+ T_{ij...}(\varphi) \varphi^i \varphi^j \\ &+ T_{ij...}(\varphi) \varphi^i \varphi^j \end{aligned} \quad (C19a)$$

Thus, all we need are

$$D \varphi^i \varphi^j = \frac{tdl}{2} R^i_{bcd} \varphi^d \varphi^c \varphi^b \varphi^j \frac{tdl}{2}; \quad (1)^{j(k+1)} R^i_{lk_1} \varphi^j \varphi^k \varphi^l; \quad (C19b)$$

$$D \varphi^i \varphi^j = \frac{tdl}{3} R^i_{bcd} \varphi^d \varphi^c \varphi^b; \quad (C19c)$$

on the one hand and

$$T_{b_1 \dots b_s}(\varphi) = \frac{tdl}{2! 3!} X^1 T_{b_1 \dots b_{s+1}}(\varphi) (1)^{(b_1+b) (b_{s+1}+1) \dots} R^b_{b_1 c_1 c_2} + (1)^{b_1} R^b_{c_1 b_1 c_2} \varphi^{c_1} \varphi^{c_2}; \quad (C19d)$$

on the other hand. Here, we are using the conformal indices defined in Eq. (2.19).

2. NL M on Kahler supermanifolds

g. Notations and conventions for Hermitian supermanifolds Some additional notation is needed to deal with Hermitian and Kahlerian manifolds.

With the convention that complex conjugation of a composite objects reverts the ordering, $(O_1 O_2 O_3) \Rightarrow (O_3 O_2 O_1)$, it is useful to introduce sign factors associated to the operation of complex conjugation. For covariant and contravariant holomorphic and antiholomorphic vectors, the action of the complex conjugation is defined by

$$(V^a) = V^a; \quad (V_a) = (1)^a V_a; \quad (C20a)$$

For tensor fields of interest in this paper, the com-

plex conjugation of a tensor field is given by

$$T^{A_1 \dots A_{r+s}} = \quad (C20b)$$

$$(1)^{r+s(A) + (A_{r+1} \dots A_s)} T^{A_1 \dots A_{r+s}};$$

$$\text{with } r+s(A) = \prod_{t,u=1, t < u}^q A_t A_u.$$

The grade for an antiholomorphic index is equal to that of its holomorphic counterpart, $(a) = (a)$.

For illustration, we shall apply these rules to the components $g_a g_b$ of the metric tensor field. Combining the rule (C20b) with the supersymmetry of the metric (C5),

$$g_a g_b = (1)^{a+b} g^{ab} (g_a g_b); \quad g_a g_b = (1)^{a+b} g^{ab} g_a g_b; \quad (C21)$$

we infer that the metric tensor field is superhermitian

$$g_{b a} = (g_a b); \quad (C22)$$

h. The NL M on a Hermitian supermanifold The NL M on a Hermitian supermanifold is defined by the partition function

$$Z = \int D[z; \bar{z}] e^{-S[z; \bar{z}]}; \quad (C23a)$$

$$S[z; \bar{z}] = \frac{1}{2} \int \bar{t}_r \partial_z z^a g_b(z; \bar{z}) \partial_{\bar{z}} \bar{z}^b; \quad (C23b)$$

where $(z^a; \bar{z}^a)$ represents coordinates on the complex supermanifold. For a Kählerian supermanifold, the metric is derived from the Kähler potential $K(z; \bar{z})$,

$$g_{ab} = K(z; \bar{z}) \partial_a \partial_{\bar{b}}; \quad (C24)$$

[Here, observe that the shifting rule for indices (C1b) and the relation between the left- and right-derivatives (C2)

are compatible.]

i. The covariant background field method and the super Kähler normal coordinate expansion In the background field method, the covariant vector field $(z^a; \bar{z}^a)(x)$ is separated into two parts,

$$z^a = \bar{z}^a + \delta z^a; \quad \bar{z}^a = \bar{z}^a + \delta \bar{z}^a; \quad (C25)$$

whereby $(\bar{z}^a; \bar{z}^a)(x)$ is assumed to be a slowly varying solution to the classical equations of motion which transforms like a contravariant vector and $(\delta z^a; \delta \bar{z}^a)(x)$ represents fluctuations around the slow degrees of freedom $(\bar{z}^a; \bar{z}^a)(x)$.

The expansion in terms of $(\delta z; \delta \bar{z})$ is traded off for one in terms of normal coordinates $(w; \bar{w})$, the super version of the KNC. The super KNC expansion for a tensor field of rank $(0; r+s)$ with r holomorphic and s antiholomorphic indices $T_{b_1 \dots r b_p s}(\bar{z}; z)$ is

$$\begin{aligned} T_{b_1 \dots r c p s}(\bar{z}; z) &= T_{b_1 \dots r c p s}(\bar{z}; \bar{z}) + T_{b_1 \dots r c p s} \bar{t}_{k_1} w^{k_1} + T_{b_1 \dots r c p s} \bar{t}_{l_1} \bar{w}^{l_1} \\ &+ \frac{1}{2} T_{b_1 \dots r c p s} \bar{t}_{k_1} \bar{t}_{k_2} w^{k_2} w^{k_1} + \frac{1}{2} T_{b_1 \dots r c p s} \bar{t}_{l_1} \bar{t}_{l_2} \bar{w}^{l_2} \bar{w}^{l_1} + T_{b_1 \dots r c p s} \bar{t}_{k_1} \bar{t}_{l_1} w^{k_1} \bar{w}^{l_1} \\ &+ \sum_{i=1}^{X^r} \frac{1}{i!} \bar{t}_{b_1 \dots i b_{i+1}} \bar{t}_{c_1 \dots c_{i+1}} R_{b_i k_1 l_1}^a w^{k_1} \bar{w}^{l_1} + \dots \\ &= T_{b_1 \dots r c p s}(\bar{z}; \bar{z}) + T_{b_1 \dots r c p s} \bar{t}_{k_1} w^{k_1} + T_{b_1 \dots r c p s} \bar{t}_{l_1} \bar{w}^{l_1} \\ &+ \frac{1}{2} T_{b_1 \dots r c p s} \bar{t}_{k_1} \bar{t}_{k_2} w^{k_2} w^{k_1} + \frac{1}{2} T_{b_1 \dots r c p s} \bar{t}_{l_1} \bar{t}_{l_2} \bar{w}^{l_2} \bar{w}^{l_1} + T_{b_1 \dots r c p s} \bar{t}_{l_1} \bar{t}_{k_1} w^{k_1} \bar{w}^{l_1} \\ &+ \sum_{i=1}^{X^s} \frac{1}{i!} \bar{t}_{b_1 \dots i b_{i+1}} \bar{t}_{c_1 \dots c_{i+1}} R_{c_i l_1 k_1}^a w^{k_1} \bar{w}^{l_1} + \dots \end{aligned} \quad (C26)$$

The super KNC expansion for $\partial(\bar{z}^A + w^A)$ is given by

$$\begin{aligned} \partial(\bar{z}^a + w^a) &= \partial \bar{z}^a + w^a D + \frac{1}{2} R_{c_1 c_2 c_3}^a(\bar{z}; \bar{z}) \partial \bar{z}^{c_3} w^{c_2} w^{c_1} + \dots \\ \partial(\bar{z}^a + w^a) &= \partial \bar{z}^j + w^a D + \frac{1}{2} R_{c_1 c_2 c_3}^a(\bar{z}; \bar{z}) \partial \bar{z}^{c_3} w^{c_2} w^{c_1} + \dots \end{aligned} \quad (C27)$$

where D is the right covariant derivative for $(w^j; \bar{w}^j)$

$$\begin{aligned} w^j D &= w^j r_a \partial^a \\ &= \partial w^j + \bar{t}_{ab}^j w^b \partial^a; \end{aligned} \quad (C28a)$$

$$\begin{aligned} \bar{w}^j D &= \bar{w}^j \bar{r}_a \partial^a \\ &= \partial \bar{w}^j + \bar{t}_{ab}^j \bar{w}^b \partial^a \\ &= (\bar{w}^j D); \end{aligned} \quad (C28b)$$

with \bar{t}_{BC}^A the coefficient of the connection that satisfies $(\bar{t}_{bc}^a) = (1)^{c+b+a(b+c)+bc} \bar{t}_{b c}^a$ [see (C20b)]. With the covariant expansion Eqs. (C26) and (C27) the action

is expanded in terms of $(w^a; \bar{w}^a)$ according to

$$\begin{aligned} S[\bar{z} + \frac{1}{2} \bar{t}_{ab}^c w^a w^b; \bar{z} + \frac{1}{2} \bar{t}_{ab}^c \bar{w}^a \bar{w}^b] &= S[\bar{z}; \bar{z}] \\ &+ \frac{1}{2} \int \bar{t}_r \partial_z w^a g_b(\bar{z}; \bar{z}) \partial_{\bar{z}} \bar{w}^b \\ &+ \frac{1}{2} \int \bar{t}_r R_{i j k l} w^k \bar{w}^l \partial^j \partial^i \\ &+ \frac{1}{2} \int \bar{t}_r \bar{t}_{k_1} \bar{t}_{k_2} w^{k_2} w^{k_1} \partial^j \partial^i \\ &+ \dots \end{aligned} \quad (C29)$$

j. Vielbeins on a Hermitian supermanifold Introduce the fast degrees of freedom $(\delta z; \delta \bar{z})$ by the linear

transformations

$$\hat{a} := w^a{}_a \hat{e}^a; \quad w^a = \hat{a} \hat{e}^a; \quad (\text{C 30a})$$

$$\hat{a} := \hat{a} \hat{e}_a{}^a w; \quad b_w = b \hat{e}_b{}^b; \quad (\text{C 30b})$$

and

$$\hat{a} := w^a{}_a \hat{e}^a; \quad w^a = \hat{a} \hat{e}^a; \quad (\text{C 30c})$$

$$\hat{a} := \hat{a} \hat{e}_a{}^a w; \quad b_w = b \hat{e}_b{}^b; \quad (\text{C 30d})$$

where

$$\hat{a} \hat{e}^a \hat{e}^b = \hat{a} \hat{e}^b; \quad \hat{a} \hat{e}^a \hat{e}^b = \hat{a} \hat{e}^b; \quad (\text{C 30e})$$

$$\hat{a} \hat{e}_a \hat{e}_b = \hat{a} \hat{e}_b; \quad \hat{a} \hat{e}_a \hat{e}_b = \hat{a} \hat{e}_b; \quad (\text{C 30f})$$

$$(1)^{\hat{a}(\hat{a}+a)} \hat{a} \hat{e}^a = \hat{a} \hat{e}_a; \quad (1)^{a(\hat{a}+a)} \hat{a} \hat{e}^a = \hat{a} \hat{e}_a; \quad (\text{C 30g})$$

$$\hat{a} \hat{e}^a = (1)^{(\hat{a}+a)a} \hat{a} \hat{e}^a; \quad (\hat{a} \hat{e}^a) = (1)^{(\hat{a}+a)\hat{a}} \hat{a} \hat{e}^a; \quad (\text{C 30h})$$

These definitions of the vielbeins are equivalent to the non-supersymmetric cases except the last two lines. The vielbeins (and their inverse) \hat{e} are defined such that

$$w^a{}_a g_b{}^b = \hat{a} \hat{e}_a \hat{e}^b \quad (\text{C 30i})$$

where the canonical form of the metric is given by

$$\hat{a} \hat{e}_b = \text{diag } \mathbb{I}_{M_1}; \mathbb{I}_{M_2}; \mathbb{I}_N; \quad (\text{C 30j})$$

Note that we are allowing the bosonic part of \hat{e} to be pseudo Riemannian, i.e., is allowed to have -1 as well as $+1$ as its diagonal elements. From now on, latin letters with a hat refer to the coordinates of the Hermitian manifold in the vielbein basis.

The covariant expansion of the action in terms of $(\hat{a}; \hat{a})$ is given by

$$\begin{aligned} S[\hat{a}; \hat{a}] &= S[\hat{a}; \hat{a}] + \frac{1}{2} \int d^4z \hat{D}^a \hat{e}_a \hat{e}^b \hat{D}^b \\ &+ \frac{1}{2} \int d^4z R_{ijkl} (\hat{e}^i \hat{e}^j) (\hat{e}^k \hat{e}^l) \hat{e}^m \hat{e}^n \\ &+ \frac{1}{2} \int d^4z (\hat{e}^i \hat{e}^j) \hat{e}^k \hat{e}^l (\hat{e}^m \hat{e}^n) \\ &+ \frac{1}{2} \int d^4z (\hat{e}^i \hat{e}^j) (\hat{e}^k \hat{e}^l) \hat{e}^m \hat{e}^n; \end{aligned} \quad (\text{C 31})$$

where yet another covariant derivative from the right

$$\hat{D}^a = \hat{e}^a + \hat{a} (\hat{A})_b{}^b; \quad (\text{C 32a})$$

$$\hat{D}^a = \hat{e}^a + \hat{a} (\hat{A})_b{}^b; \quad (\text{C 32b})$$

$$\hat{D}^a = \hat{e}^a + \hat{a} (\hat{A})_b{}^b; \quad (\text{C 32c})$$

$$\hat{D}^a = \hat{e}^a + \hat{a} (\hat{A})_b{}^b; \quad (\text{C 32d})$$

has been introduced together with the $U(M_1; M_2; N)$ gauge field \hat{A} . The gauge field with different index-structures are related by

$$\hat{D}^a = \hat{D}^a (\hat{A})_b{}^b = (1)^{\hat{b}(\hat{a}+\hat{b})} \hat{D}^a (\hat{A})_b{}^b; \quad (\text{C 33a})$$

$$\hat{D}^a = \hat{D}^a (\hat{A})_b{}^b = \hat{D}^a (\hat{A})_b{}^b; \quad (\text{C 33b})$$

The gauge field is skew superhermitian,

$$(\hat{A})_c{}^c = -(\hat{A})_b{}^b \quad (\text{C 34})$$

where indices of \hat{A} are raised and lowered by \hat{e} . The explicit form of the gauge field in terms of the vielbeins is given by the $U(M_1; M_2; N)$ spin connection

$$\hat{A}^{\hat{b}}{}_c = \hat{e}^a \hat{e}_a \hat{e}^b \hat{\omega}_c; \quad (\text{C 35a})$$

$$\begin{aligned} \hat{A}^{\hat{b}}{}_c &= \hat{e}_p \hat{e}^p \hat{e}^b \hat{\omega}_c \\ &= \hat{e}^a (\hat{e}^b \hat{\omega}_c) (1)^{a+\hat{a}+(a+\hat{a})c} \hat{e}_{ac} \hat{e}^b; \end{aligned} \quad (\text{C 35b})$$

$$\hat{A}^{\hat{b}}{}_c = (1)^{c(\hat{b}+\hat{a})} \hat{A}^{\hat{b}}{}_c; \quad (\text{C 35c})$$

whereby

$$\hat{A}^{\hat{b}}{}_c = \hat{A}^{\hat{b}}{}_c + \hat{A}^{\hat{b}}{}_c; \quad (\text{C 36})$$

The $U(M_1; M_2; N)$ field strength tensor is given by

$$\begin{aligned} \hat{F}^{\hat{b}}{}_c &= \hat{e}_a (\hat{A})_c{}^b - \hat{e}_a (\hat{A})_c{}^b \\ &+ \hat{a} (\hat{A})_c{}^b (\hat{A})_c{}^b - \hat{a} (\hat{A})_c{}^b (\hat{A})_c{}^b \\ &= (1)^{a+\hat{a}+ar} \hat{e}^a R^r{}_{adc} \hat{e}^c \hat{e}^d \hat{e}^b \\ &+ (1)^{a+\hat{a}+ar} \hat{e}^a R^r{}_{adc} \hat{e}^c \hat{e}^d \hat{e}^b; \end{aligned} \quad (\text{C 37})$$

k. One-loop beta function The integration over the fast modes $(\hat{a}; \hat{a})$ is performed with the help of the cumulant expansion. The effective action for the slow modes $(\hat{a}; \hat{a})$ is, up to one-loop,

$$Z = \int D[\hat{a}; \hat{a}] e^{S[\hat{a}; \hat{a}] + S[\hat{a}; \hat{a}]}; \quad (\text{C 38a})$$

$$S[\hat{a}; \hat{a}] = G_0(0) R_{ab} \hat{e}^b \hat{e}^a; \quad (\text{C 38b})$$

$$R_{ij} = (1)^{k+1+i+l} g^k R_{ki+l j} \quad (\text{C 38c})$$

is the Ricci tensor.

l. RG flows of high-gradient operators The covariant expansion in terms of the super KNC applies to the high-gradient operators

$$T_{IJK} \dots (z; z) \hat{e}^I \hat{e}^J \hat{e}^K \quad (\text{C 39})$$

as well and can be used to compute their anomalous scaling dimensions. We shall assume that all the covariant derivatives of the tensor field $T_{IJK} \dots (z; z)$ vanish. If so we infer the RG transformation law

$$D \quad E \quad D \quad E \quad D \quad E$$

$$T_{IJ} :: (z ; z) @ z^I @ z^J \quad = \quad [T_{IJ} :: (z ; z)]_2 @ z^I @ z^J \quad \dagger T_{IJ} :: (z ; z) @ z^I @ z^J \quad + \quad :$$

$$(C 40a)$$

Thus, all we need are

$$D \quad E$$

$$@ z^i @ z^j = \text{tdl} ; \quad (1)^{j(k+1)} R^i{}_{lk} g^j @^k @^l ; \quad (C 40b)$$

$$D \quad E$$

$$@ z^i @ z^j = \text{tdl} ; \quad R^i{}_{bc} @^d @^c g^j ; \quad (C 40c)$$

$$D \quad E$$

$$@ z^i @ z^j = \text{tdl} ; \quad R^i{}_{bcd} @^d @^c g^j ; \quad (C 40d)$$

$$D \quad E$$

$$@ z^i @ z^j = \text{tdl} ; \quad (1)^{j(k+1)} R^i{}_{lk} g^j @^k @^l ; \quad (C 40e)$$

on the one hand and

$$h \quad i$$

$$T_{b_1 \dots b_r c p} (z ; z)_2 = \text{tdl} \quad X^r \quad (1)^{(a+b_1)(b_{i+1}+)} \quad s^{\dagger} \hat{T}_{b_1 \dots b_r c a b_{i+1}} R_c^a{}_{b_1 k_1 l_1} g^{k_1 l_1}$$

$$= \text{tdl} \quad X^s \quad (1)^{(a+c_1)(c_{i+1}+)} \quad s^{\dagger} \hat{T}_{b_1 \dots b_r c p i c a c_{i+1}} R_c^a{}_{c_1 l_1 k_1} g^{k_1 l_1} ; \quad (C 40f)$$

on the other hand. Here, we are using the conformal indices defined in Eq. (2.19).

Since a Kählerian supermanifold is a Riemannian supermanifold, one can make use of either the RNC or KNC expansion. Indeed, all the results in this appendix derived from the KNC expansion can actually be obtained

from the RNC expansion as well. A subtle thing here is that the RNC and KNC expansion for a tensor field $T_{IJK} :: (z ; z)$ and $@ z^A$ are different.^{65,66} However, the RNC and KNC expansions are identical when applied to the high-gradient operators (C 39) and thus give us the same results.

¹ F. Wegner, Z. Physik B 35, 207 (1979).

² B. L. Altshuler, V. E. Kravtsov, and I. V. Lerner, in Mesoscopic Phenomena in Solids, edited by B. L. Altshuler, P. A. Lee, and R. A. Webb, North-Holland, Amsterdam, 449 (1991).

³ B. L. Altshuler, V. E. Kravtsov, and I. V. Lerner, JETP Lett. 43, 440 (1986); Sov. Phys. JETP 64, 1352 (1986); Phys. Lett. A 134, 488 (1989).

⁴ B. L. Altshuler, JETP Lett. 41, 648 (1985).

⁵ P. A. Lee and A. D. Stone, Phys. Rev. Lett. 55, 1622 (1985).

⁶ A. D. Stone, Phys. Rev. Lett. 54, 2692 (1985).

⁷ V. E. Kravtsov, I. V. Lerner, and V. I. Yudson, Sov. Phys. JETP 68, 1441 (1988).

⁸ V. E. Kravtsov, I. V. Lerner, and V. I. Yudson, Phys. Lett. A 134, 245 (1989).

⁹ O. Motrunich, K. Damle, and D. A. Huse, Phys. Rev. B 65, 064206 (2002).

¹⁰ C. Mudry, S. Ryu, and A. Furusaki Phys. Rev. B 67, 064202 (2003).

¹¹ M. R. Zirnbauer, J. Math. Phys. 37, 4986 (1996).

¹² A. A. Kland and M. R. Zirnbauer, Phys. Rev. B 55, 1142 (1997).

¹³ The terminology used for these 10 symmetry classes corresponds to the classification of symmetric spaces by the

mathematician E. Cartan.

¹⁴ K. B. Efetov, Adv. Phys. 32, 53 (1983).

¹⁵ The spin conductivity is the linear response coefficient quantifying the amount of spin-current in the z-direction in spin space, owing in response to a uniform gradient of z-component of magnetic field.

¹⁶ The two remaining BdG symmetry classes are called D III and C I by AZ. They correspond to cases for which time-reversal symmetry is respected in all realizations of the disorder whereas spin-rotation symmetry is broken in symmetry class D III but preserved in symmetry class C I.

¹⁷ T. Senthil and M. P. A. Fisher, Phys. Rev. B 61, 9690 (2000).

¹⁸ T. Senthil, M. P. A. Fisher, L. Balents, and C. Nayak, Phys. Rev. Lett. 81, 4704 (1998).

¹⁹ K. B. Efetov, A. I. Larkin, and D. E. Khmelnitskii, Sov. Phys. JETP 52, 568 (1980).

²⁰ O. N. Dorokhov, JETP Letters 36, 318 (1982); P. A. Mello, P. Pereyra and N. Kumar, Ann. Phys. (NY) 181, 290 (1988). A geometrical interpretation of the DMPK equation was pointed out in A. Humann, J. Phys. A 23, 5733 (1990). A review on the DMPK equation can be found in C. W. J. Beenakker, Rev. Mod. Phys. 69, 731 (1997).

²¹ V. Kagalovsky, B. Horowitz, Y. Avishai, and J. T. Chalker, Phys. Rev. Lett. 82, 3516 (1999).

- ²² T. Senthil, J. B. Marston, and M. P. A. Fisher, *Phys. Rev. B* 60, 4245 (1999).
- ²³ J. T. Chalker, N. Read, V. Kagalovsky, B. Horowitz, Y. Avishai, and A. W. W. Ludwig, *Phys. Rev. B* 65, 012506 (2001).
- ²⁴ A. M. M. Pruisken, *Nucl. Phys. B* 235, 271 (1984).
- ²⁵ N. Read and D. Green, *Phys. Rev. B* 61, 10267 (2000).
- ²⁶ M. Boquet, D. Serban, and M. R. Zimbauer, *Nucl. Phys. B* 578, 628 (2000).
- ²⁷ I. A. Gruzberg, N. Read, and A. W. W. Ludwig, *Phys. Rev. B* 63, 104422 (2001).
- ²⁸ I. A. Gruzberg, N. Read, and S. Vishveshwara, *Phys. Rev. B* 71, 245124 (2005).
- ²⁹ S. Helgason, *Differential geometry, Lie groups and symmetric spaces* (Academic Press, New York, 1978).
- ³⁰ M. Nakahara, *Geometry, Topology, and Physics*, (second edition, Institute of Physics Publishing, Bristol, 2003).
- ³¹ See, e.g., S. Kobayashi and K. Nomizu, *Foundations of differential geometry*, (Wiley, New York 1996).
- ³² D. Friedan, *Ann. Phys. (N.Y.)* 163, 318 (1985).
- ³³ The beta function of t encodes the change to t induced by the infinitesimal rescaling of the short distance cutoff, $a \rightarrow a e^{d_1}$.
- ³⁴ In symmetry class A, in the replica limit $p + q \rightarrow 0$ (see Table I), the coefficient of the t^2 term in the beta function happens to vanish, and one needs to compute the beta function up to order t^3 in order to find a metal-insulator fixed point in symmetry class A.^{35,36,37}
- ³⁵ F. Wegner, *Nucl. Phys. B* 316, 663 (1989).
- ³⁶ S. Hikami, *Phys. Lett. B* 98, 208 (1981).
- ³⁷ S. Hikami, *Physica A* 167, 149 (1990); in *Lecture Notes of NATO ASI on Interference Phenomena in Mesoscopic Systems*, 254, edited by B. Kramer, 429 (Plenum Press, New York 1991); *Prog. Theor. Phys. Suppl.* 107, 213 (1992).
- ³⁸ I. V. Lerner and F. Wegner, *Z. Phys. B* 81, 95 (1990).
- ³⁹ F. Wegner, *Nucl. Phys. B* 354, 441 (1991).
- ⁴⁰ The same stability problem is also present to first order in the $1/N$ -expansion of the $O(N) \rightarrow O(N-1)$ NLM below three dimensions. See A. N. Vasilev and A. S. Stepanenko, *Theor. Math. Phys.* 94, 471 (1993); K. Lang and W. Ruhl, *Z. Phys. C* 61, 495 (1994); S. E. Derkachov, S. K. Kehrein, and A. N. M. Anashov, *Nucl. Phys. B* 493, 660 (1997); and E. Brezin and S. Hikami, *Phys. Rev. B* 55, R10169 (1997).
- ⁴¹ F. Wegner, *Z. Phys. B* 78, 33 (1990).
- ⁴² A. W. W. Ludwig, *Nucl. Phys. B* 330, 639 (1990).
- ⁴³ H. M. Alland F. Wegner, *Nucl. Phys. B* 393, 495 (1993).
- ⁴⁴ G. Castilla and S. Chakravarty, *Phys. Rev. Lett.* 71, 384 (1993).
- ⁴⁵ G. Castilla and S. Chakravarty, *Nucl. Phys. B* 485, 613 (1997).
- ⁴⁶ E. Brezin and S. Hikami, *Phys. Rev. B* 55, R10169 (1997).
- ⁴⁷ Ch. Chatelain and B. Berche, *Nucl. Phys. B* 572, 626 (2000); "Phase transitions in two-dimensional random Potts models", in "Order, disorder, and criticality", edited by Yu. Holovatch, World Scientific, (Singapore 2004), page 146.
- ⁴⁸ K. Slevin and T. Ohtsuki, *Phys. Rev. Lett.* 82, 382 (1999); *ibid.*, 82, 669 (1999).
- ⁴⁹ K. Slevin, P. Markos, and T. Ohtsuki, *Phys. Rev. Lett.* 86, 3594 (2001).
- ⁵⁰ K. Slevin and T. Ohtsuki, *Phys. Rev. B* 63, 045108 (2001).
- ⁵¹ K. Slevin, P. Markos, and T. Ohtsuki, *Phys. Rev. B* 67, 155106 (2003).
- ⁵² In general we denote the stochastic cumulant of a random variable by $\langle \dots \rangle_{st}$.
- ⁵³ P. W. Brouwer, A. Furusaki, I. A. Gruzberg, and C. Mudry, *Phys. Rev. Lett.* 85, 1064 (2000).
- ⁵⁴ T. Imamura and K. Hikami, *J. Phys. Soc. Jpn.* 70, 3312 (2001).
- ⁵⁵ P. W. Brouwer, A. Furusaki, C. Mudry, and S. Ryu, *cond-mat/0511622* and *BUTSURI* 60, 935 (2005).
- ⁵⁶ A. Lamacraft, B. D. Simons, and M. R. Zimbauer, *Phys. Rev. B* 70, 075412 (2004).
- ⁵⁷ L. Alvarez-Gaume, D. Z. Freedman, and S. Mukhi, *Ann. Phys.* 134, 85 (1981).
- ⁵⁸ F. Wegner in *Phase transitions and critical phenomena*, volume 6, edited by C. Domb and M. S. Green, 8 (Academic Press, London 1976).
- ⁵⁹ B. Dewitt, *Supermanifolds*, Second edition, (Cambridge University Press 1992).
- ⁶⁰ The antisymmetry of the Riemann tensor R^A_{BCD} has to be used.
- ⁶¹ As we will not include domain walls in symmetry class D, we replace $O(N)$ by its connected counterpart $SO(N)$ in the following.
- ⁶² B. Zumino, *Phys. Lett. B* 87, 203 (1979).
- ⁶³ K. Higashijima, T. Kimura, and M. Nitta, *Nucl. Phys. B* 623, 133 (2002).
- ⁶⁴ K. Higashijima, T. Kimura, and M. Nitta, *Nucl. Phys. B* 645, 438 (2002).
- ⁶⁵ K. Higashijima and M. Nitta, *Prog. Theor. Phys.* 105, 243 (2001).
- ⁶⁶ K. Higashijima, E. Itou, and M. Nitta, *Prog. Theor. Phys.* 108, 185 (2002).
- ⁶⁷ S. Aoyama and J. W. Holtren, *Nucl. Phys. B* 258, 18 (1985).
- ⁶⁸ G. M. Shore, *Nucl. Phys. B* 334, 172 (1990).
- ⁶⁹ E. Brezin, S. Hikami, and J. Zinn-Justin, *Nucl. Phys. B* 165, 528 (1980).
- ⁷⁰ Here, we are considering c-type (commuting) tensor fields only. If the tensor field itself is a-type (anticommuting), components of the tensor field pick up an extra statistical factor.⁵⁹
- ⁷¹ In this paper, we are always dealing with real c-type vector and tensor fields and their components in the so-called standard basis.⁵⁹